DETECTION OF ANTIBODIES AGAINST BOVINE HERPES VIRUS 1, BOVINE VIRAL DIARRHEA VIRUS AND BOVINE RESPIRATORY SYNCYTIAL VIRUS IN EARLY AND ULTRA-EARLY WEANED BEEF CALVES

¹Diego Daniel Gonzalez, ²Juan Sebastian Vittone, ²Martin Lado, ²Andrea Biolatto, ^{1,3}Marina Valeria Mozgovej, ^{1,3}Alejandra Ferella, ¹Ayelen Sammarruco, ^{1,3}Silvina Maidana, ^{1,3}Sonia Alejandra Romera and ^{1,3}María Jose Dus Santos

¹Instituto de Virología, CICVyA, INTA, Castelar, Buenos Aires, Argentina

²Estación Experimental Agropecuaria Concepción del Uruguay, INTA, Entre Ríos, Argentina

³CONICET, Buenos Aires, Argentina

Received 2013-10-15, Revised 2013-10-28; Accepted 2013-10-29

ABSTRACT

Bovine respiratory disease is the leading cause of morbidity and mortality in weaned calves. In Argentina, two weaning practices have been implemented. In the early weaning, the calf is removed from the cow at 60-70 days of age while in ultra-early weaning the calf is weaned at 30-45 days of age. The purposes of both systems is to improve cow body condition, calf performance, conception rates and forage availability for the cow. In this study we evaluated the antibody response against BVDV and BoHV1 in early and ultra-early weaned calves that had received a conventional vaccination schedule (first dose at weaning and a booster 21 days post-weaning). Passively acquired immunity may provide protection against disease caused by these viruses. The presence of antibodies against BRSV, a virus that was not present in the vaccines used, was also evaluated as an indirect indicator of viral circulation in the herd. At the time of vaccination, calves presented a wide range of maternally-derived antibody titers. Vaccination against BoHV-1 did not evoke seroconversion and antibody titers continued to decay throughout the experience. After vaccination, seroconversion to BVDV could be detected in calves with low antibody titers, while higher antibody titers exerted an inhibitory effect of the active humoral response.

Keywords: Weaning, Bovine Respiratory Disease, Maternal Antibodies, Vaccines

1. INTRODUCTION

In beef herds, calves remain with their cows until they are weaned, which traditionally occurs at six to seven month of age.

Weaning is an inherent husbandry practice in cow-calf beef production systems that imposes physical, psychological and nutritional stressors on calves (Lynch *et al.*, 2010b).

In Argentina, two systems have been implemented for weaning. Early weaning involves the removal of the calf from the cow at 60-70 days of age while in ultra early weaning the calf is weaned at 30-45 days of age.

The purposes of both systems are to improve cow body condition, calf performance, conception rates and forage availability for the cow.

The early weaning of calves would greatly reduce calf-rearing costs by decreasing feed and labor expenditures (Hulbert *et al.*, 2011). The increased nutrient and energy demand in lactating cows associated with lactation can generate to lose body condition. Early and ultra-early weaning managements eliminate these requirements and cows are able to maintain or increase body condition. In addition, they have a greater opportunity to increase the pregnancy rate, since resumption of postpartum ovarian cyclicity is

Corresponding Author: Diego Daniel Gonzalez, Instituto de Virología, CICVyA, INTA, Castelar, Buenos Aires, Argentina

closely related to body condition at partum and the nutritional balance post partum.

Following weaning, dry cows will eat less forage than lactating ones, further reducing the demand placed on the pasture, which can enhance sustainability and forage production.

It has also been determined that ultra-early weaning is an advantageous management due to the lower levels of cortisol in the calves at that age. Moreover, breaking the bond between the cow and calf appears to be easier at 30-45 days post partum than six months later (Valerio, 2007).

By weaning early and providing a highly nutritious diet to ensure a suitable ruminant development, calves can reach their growth potential up to 30% (Galli *et al.*, 2008; Valerio, 2007).

Weaned calves undergo a transition from a monogastric to a ruminant animal, they have almost undeveloped rumens and high quality nutrient requirement; for that reason, they can not feed only forage. An abrupt change from a liquid diet, specifically milk supply from the mother, to a solid food intake could trigger physiological alterations which may result in negative consequences if they are not properly managed. The use of a calf starter is a crucial link to proper ruminal development and successful weaning, since intake of dry feed initiates rumen development. Early and ultra-early weaned calves became ruminant animals at an earlier age and this management results in an increased calf performance.

Following abrupt weaning, beef calves exhibit distress behaviors (Haley *et al.*, 2005; Price *et al.*, 2003) with alterations in hormonal mediators of stress (Blanco *et al.*, 2009) and immune function (Hickey *et al.*, 2003; Arthington *et al.*, 2008; 2005; Lynch *et al.*, 2010a; 2010b) evident up to seven days post-weaning.

Lynch *et al.* (2010b; 2012), showed that weaning together with movement of beef calves from a pasture environment to a housing environment elicited transient neutrophilia, impaired neutrophil phagocytic function, decreased peripheral lymphocyte count and altered immunophenotypes. These results suggested that there was a greater transitory reduction in immune function at housing in abruptly weaned than non-weaned beef calves (Lynch *et al.*, 2010b; 2012).

Moreover, weaning is considered to be a predisposing factor to Bovine Respiratory Disease (BRD) (Callan and Garry, 2002; Snowden, 2009). This syndrome has a complex and multifactorial etiology that can be divided into three major categories: Environmental factors, physiological stress and infection factors (Stilwell *et al.*, 2008; Richeson *et al.*, 2012). BRD is an important disease both during pre-weaning period and shortly after

weaning and the leading cause of morbidity and mortality in weaning calves.

The respiratory viruses commonly responsible for BRD include bovine herpes virus 1 (causing agent of Infectious Bovine Rhinotracheitis or IBR), Bovine Respiratory Syncytial Virus (BRSV), Bovine Viral Diarrhea Virus (BVDV) and Para-Influenza Virus (PI3) (Taylor *et al.*, 2010a). All these viruses are known to be involved in BRD solely or in synergism with other and bacteria.

To ensure the health and well being of early weaned calves, a common practice in beef herds is the vaccination of calves against respiratory viruses at weaning in order to minimize BRD. However, the use of vaccination for the control of BRD in animals on arrival to pens still controversial (Ellis, 2001).

Calves are born agammaglobulinaemic and depend strongly on ingestion of colostrum for survival. In addition, the specific and non-specific immune systems are immature in newborn calves, making them especially vulnerable to infections during the first weeks of life.

Passively acquired immunity may provide protection against disease caused by BoHV1, BVDV and BRSV (Belknap *et al.*, 1991; Bolin and Ridpath, 1995; Howard *et al.*, 1989; Mechor *et al.*, 1987). However, young calves do not produce specific Antibodies (Ab) after vaccination in the presence of maternal antibodies. For this reason it is commonly believed that maternal antibodies can interfere with the efficacy of vaccination.

In this study the efficacy of a conventional vaccination program in weaned calves was evaluated. Calves were vaccinated with a commercial vaccine containing BVDV, BoHV1, PI3, *Haemophilus somnus*, *Moraxella bovis*, *Pasteurella multocida* and *Pasteurella haemolytica* as immunogens. Specific Ab against BVDV, BoHV1 and BRSV (a virus not included in the vaccine) were determined in sera from beef calves that had been early and ultra-early weaned.

2. MATERIALS AND METHODS

2.1. Animals

The experience was conducted in Concepción del Uruguay INTA's Experimental Station (32°29'28"S; 58°20'49"W, 25 m above mean sea level), in the province of Entre Rios Province in Argentina.

Fourty eight Hereford and Polled Hereford cows and their respective calves from the Experimental Station were used.

Cows were vaccinated at 30 day prepartum with a commercial vaccine containing inactivated IBR (BoHV1) and BVDV1a, *Campylobacter fetus*, *Leptospira interrogans pomona* y *Haemophilus somnus*.

These cows were boosted each year. Calves were vaccinated twice, at weaning and 21 days post-weaning (conventional vaccination program) with another commercial vaccine containing inactivated BVDV1a, IBR (HVB-1), PI3, *Haemophilus somnus*, *Moraxella bovis*, *Pasteurella multocida* and *Mannheimia haemolytica* (*Pasteurella haemolytica*).

2.2. Experimental Design

At weaning, calves were housed in four corrals, each covering a 50 m² area and with waterers and feed bunks. Calves were allotted to two experimental groups according to their age and weight. The groups were: Early Weaning calves (EW: 66,2±9,6 days old) and Ultra-Early Weaning (UEW: 40,9±7,3 days old) calves. Each group was placed in two pens (12 animals/corral). One corral from each group was immunized with 2 dosis (at weaning and 21 days post-weaning) of a commercial vaccine against BRD. The animals from the other corral of each group remained unvaccinated.

2.3. Weaning Strategy

Calves were weaned following the management requirements for EW and UEW according to the Manual of early weaning in beef herds (Galli *et al.*, 2005). All calves were offered ad libitum to a calf ruminal starter (Ruter®, Asociación de Cooperativas Argentinas) and water for 16 days.

Breeding rations (16 to 56 days postweaning) were formulated with whole kernel corn and a comercial concentrate (Starter 40% Crude Protein, Asociación de Cooperativas Argentinas).

Daily offer was fixed to 2% of calf weight.

2.4. Sample Collection

Peripheral blood samples from cows were collected via jugular venipuncture at 60 and 30 days pre-calving. Serum was collected after centrifugation at 1500 rpm (Hermle) and stored at -20°C until use.

Sera from all calves were obtained as described for cow sampling at 10 days pre-weaning, at weaning (day 0) and at 7, 21 and 35 days post 2° immunization.

2.5. Detection of Antibodies Against BoHV1 by Indirect ELISA

Total Ab in bovine serum were determined using an indirect ELISA as described previously (Romera *et al.*, 2000). Briefly, polystyrene microtiter ELISA plates (Immulon 1B, Dynatech Laboratories) were coated with 50ul of positive (concentrated and semi-purified BoHV1 reference strain Los Angeles containing 109 DICT50

mL⁻¹) or negative antigen (non-infected MDBK cells) in carbonate buffer of pH 9.6 and incubated for 12 h at 4°C. Plates were blocked with 10% ovalbumin in PBS-Tween 20 0.05%, washed twice and stored at -20°C until use. Samples were tested at 1:4, 1:10 and 1:40 dilutions (Parreno *et al.*, 2010). A peroxidase-labeled affinity purified goat anti-bovine IgG (H + L) (Kirkegaard and Perry Laboratories, KPL) in a 1:4000 dilution was used as the conjugate. After each incubation, plates were washed 4 times. The reaction was developed using hydrogen peroxide/2,2B-azino-bis 3-ethylbenzthiazoline-6 Sulfonic Acid (ABTS) in citrate buffer (pH 5) as the substrate/chromogen system. The Ab titer of each sample was expressed as the log₁₀ of the reciprocal of the highest serum dilution with a corrected optical density OD405C (OD405 in the positive coated wells minus OD405 in the negative coated wells) greater than the cut-off value of the assay (40% positive control value).

2.6. Detection of Neutralizing Antibodies (NA) against BVDV by Virus Neutralization Test

Serum NA from cattle were detected by virus neutralization assay. Briefly, 100 TCID₅₀ of BVDV (Singer strain) were co-incubated for 1 h at 37°C with 75 µL of four-fold dilutions of inactivated serum samples. Then, the mixture was added onto 3×10⁴ MDBK cells/well. Plates were incubated for 72 h at 37°C under 5% CO₂. Control wells without virus were used for each serum sample in order to discard toxicity.

Serum neutralization titer was considered as the highest serum dilution which inhibited 50% of viral cytopathic effect. Titers were expressed as the log₁₀ of the highest dilution able to inhibit 100 TCID₅₀. A BVDV positive serum from an experimentally infected bovine was used as positive control and a BVDV negative serum from a reference animal was considered as negative control.

2.7. Detection of NA against BRSV by Virus Neutralization Test

Serum neutralization assay was carried out as described previously by (Ferella *et al.*, 2012). Inactivated serum samples were fourfold diluted from 1:4 to 1:256. The serum-dilutions were supplied with 100 TCID₅₀ of 51908 BRSV strain and incubated for 1 hour at 37°C in a 5% CO₂ atmosphere. This mixture was inoculated in duplicate onto MDBK cell monolayers (250000 cells mL⁻¹) in 96-well plates. Plates were incubated as above and CPE was observed 5 days post infection.

Samples were considered positive when no CPE was observed. Antibody titers were expressed as the reciprocal of the maximum dilution in which no CPE was observed. Samples with titers equal or higher than 4 were considered positive.

2.8. Statistical Analysis

Differences in Ab titers among groups were evaluated by ANOVA under a model of repeated measures throughout time, followed by a general contrast post-ANOVA test.

Statistical significance was assessed at $p < 0.05$ for all comparisons.

3. RESULTS

Regarding BVDV and BoHV1, there was not evidence of clinical signs of disease. In addition, there

were no BVDV persistently infected calves identified. Passive transfer of maternal antibodies against BVDV and BoHV-1 was detected in all calves.

Table 1 shows the experimental design and the average age and weight of calves according to each treatment.

The distribution of antibodies at weaning for each virus is shown in **Table 2**. BVDV1a specific antibody titers were relatively low in all groups, with 77 % of total calves presenting a NA titer ≤ 32 . Similarly, 77% of total calves presented BoHV-1 specific antibody titers ≤ 1.6 . These titers were presumed not to be protective against both BVDV and BoHV-1 (Bolin and Ridpath, 1995).

The mean antibody titers for each group were determined at 10 days pre-weaning, at weaning and 28, 49 and 56 days post weaning, corresponding to 7, 21 and 35 days post booster respectively (**Fig. 1 and 2**).

Fig. 1. Antibody response against BVDV Experimental groups (EW-UNV, EW-V, UEW-UNV, UEW-V) were immunized with a commercial vaccine against BRD. At different time points animals were bled and specific BVDV antibodies were evaluated by seroneutralization assay. Each bar represents the average titer of the group

Fig. 2. Antibody response against BoHV1 Experimental groups (EW-UNV, EW-V, UEW-UNV, UEW-V) were immunized with a commercial vaccine against BRD. At different time points animals were bled and specific BoHV1 antibodies were evaluated by ELISA. Each bar represents the average titer of the group

Table 1. Experimental design and average age of cattle

Group	Weaning (days of age)	Vaccination	n	Weight at weaning	Weight at 56 days of age (kg)
1	Early (66,6±10,7)	Yes	12	68,1±7,7	90,7±12,3
2	Early (65,8±8,8)	No	12	69,8±9,2	91,7±14,5
3	Ultra-early (40,3±7,8)	Yes	12	56,2±9,7	75,4±16,3
4	Ultra-early (41,4±7,2)	No	12	57,9±11,6	76,3±17,1

Antibody levels detected at 10 days pre-weaning were maternally derived. The BVDV1a mean titers and the antibody range were: 28.51 (4-256) for both EW and UEW group. The BoHV1 mean titers were: 1.54 ± 0.64 for EW calves and 1.38 ± 0.59 for UEW calves. The BRSV mean titers and the antibody range were: 7.8 (2-32) for the EW group and 16 (2-128) for the UEW group. No significant differences could be observed between early and ultra-early weaned calves at that time indicating that passive transfer of antibodies was homogeneous in all groups for the three viruses evaluated.

The kinetic of antibody response showed that the mean antibody titers did not presented significant differences during the time of study. A slight increase of BVDV1a NA titers could be observed at 35 days post

booster in EW-V group, which can be assigned to a few calves that developed high titers of NA (**Table 3, Fig. 1**).

At 35 days post booster (56 days post-weaning) comparisons between EW-V and EW-UNV animals did not shown significant differences ($p = 0.84$) Similar results were obtained between UEW-V and UEW-UNV ($p = 0.97$). This result suggested that immunization of early and ultra-early weaned calves with inactivated vaccines for BVDV1a and BoHV1 did not evoke an active antibody response.

Seroconversion after vaccination with BVDV was evaluated at 35 days post booster in each calf. The distribution of antibody titers at weaning (day 0 for immunization) was considered to determine seroconversion (four-fold or greater increase) (**Table 4**).

Table 2. Distribution of antibody titers to BVDV1a and BoHV1 (number of calves/total calves) at weaning

BVDV Antibody titer at weaning	Experimental group			
	UEW- V	UEW- UNV	EW- V	EW- UNV
4	2/11	4/13	3/12	1/12
8	3/11	2/13	3/12	2/12
16	0/11	0/13	0/12	1/12
32	3/11	4/13	4/12	5/12
64	2/11	0/13	0/12	3/12
128	1/11	3/13	1/12	0/12
256	0/11	0/13	0/12	0/12
512	0/11	0/13	0/12	0/12
1024	0/11	0/13	1/12	0/12
BoHV1 antibody titer at weaning				
<1.6	7/11	7/13	6/12	8/12
1.6	4/11	2/13	2/12	1/12
2.2	0/11	2/13	4/12	3/12
2.8	0/11	2/13	0/12	0/12

Table 3. Distribution of antibody titers to BVDV1a and BoHV1 (number of calves/total calves) at day 35 post booster

BVDV antibody titer 35 days post Booster	Experimental group			
	UEW- V	UEW- UNV	EW- V	EW- UNV
4	1/11	1/13	0/12	0/12
8	3/11	5/13	1/12	1/12
16	0/11	0/13	1/12	1/12
32	2/11	3/13	3/12	5/12
64	0/11	0/13	1/12	0/12
128	1/11	2/13	4/12	5/12
256	0/11	1/13	1/12	0/12
512	1/11	0/13	1/12	0/12
1024	3/11	0/13	0/12	0/12
BoHV1 antibody titer at 35 days post booster				
<1.6	11/11	13/13	12/12	11/12
1.6				1/12
2.2				
2.8				

Table 4. Seroconversion at 35 days post booster (number of calves serconverted/total calves)

BVDV antibody titer at weaning	Experimental group			
	UEW- V	UEW- UNV	EW- V	EW- UNV
4	2/2	0/4	3/3	0/1
8	1/3	1/2	2/3	0/2
16	-	-	-	0/1
32	2/3	0/4	2/4	2/4
64	0/2	-	-	0/3
128	0/1	0/3	0/1	-
256	-	-	-	-
512	-	-	-	-
1024	-	-	0/1	-

Table 5. Distribution of antibody titers to BRSV (number of calves/total calves)

BRSV antibody titer at weaning	Experimental group	
	EW	UEW
2	7/24	2/24
4	2/24	7/24
8	5/24	2/24
16	2/24	8/24
32	5/24	0/24
64	2/24	3/24
128	1/24	2/24
BRSV antibody titer 56 days post		
2	14/24	17/24
4	5/24	3/24
8	3/24	0/24
16	2/24	2/24
32	0/24	0/24
64	0/24	2/24

In the UEW and EW groups 5 out of 11 and 7 out of 12 calves, respectively, showed seroconversion to BVDV after two doses of the vaccine. These vaccinated calves presented at weaning antibody titers that ranged from 4 to 32. Higher antibody titers prevented seroconversion after vaccination in calves from both groups.

Antibody titers to BoHV1, in all groups evaluated, were lower at 35 days post booster than at weaning and no animals showed seroconversion to that virus (data not shown).

Specific antibodies against BRSV were also tested since that virus is involved in BRD but it was not included in the vaccines used in this study (**Table 5**). At weaning the Ab titers in both EW and UEW calves ranged from 2 to 128, with more than 80% of animals presenting an Ab titer less than 32. The geometric mean titers and the antibody range were 9.51 (2-128) and 11.99 (2-128) for EW and UEW groups, respectively. No significant differences were found between both groups at that time ($p = 0.82$). These Abs were supposed to be maternally transferred and declined to very low levels at 56 days post weaning. Calves with Ab titers of 64 or higher at weaning showed Ab titers that ranged from 16 to 64 at 56 days post weaning, while calves with Ab titers lower than 64 became seronegative at 56 after weaning. At that time there were no significant differences between EW and UEW groups ($p = 0.85$) (mean titer for EW group: 3.27; mean titer for UEW: 3.46). Seroconversion due to BRSV infection could not be observed in any calves.

4. DISCUSSION

The impact of early weaning of calves on the reproductive performance of cows has been demonstrated in the last thirty years under research and commercial conditions in Argentina and other countries (Galli *et al.*, 2008).

Early and ultra early weaning implementation has shown to increase the reproductive efficiency, accelerate rebreeding of the dam postpartum and thus increasing pregnancy rates. Therefore, as weaning practices will probably be widely applied, to perform a correct nutritional and sanitary management of weaned calves is a critical issue for the success of the practice.

Respiratory disease in weaning calves is the leading cause of morbidity and mortality. The presence of antibodies against BRD pathogens has been found to be protective, so it would seem intuitive that immunization would be beneficial (Taylor *et al.*, 2010b), especially when the moments of important immunity suppression associated with these viruses can be foreseen, such as during the weaning and grouping of young animals from different herds. However, the use of vaccination for the control of BRD in animals on arrival to feed-lots is still controversial (Ellis, 2001) and the economic benefit of vaccination against BRD is not fully demonstrated. Much of the failure of vaccination may lie in timing of administration, failure of stressed calves to respond properly to vaccination, the multifactorial nature of BRD and the increased susceptibility of stressed calves to all pathogens.

In this study we evaluated the antibody response against BVDV1a and BoHV1 in weaned calves that have received a conventional vaccination schedule (first dose at weaning and booster 21 days post-weaning). The presence of Ab against BRSV, a virus not present in the vaccines used, was also evaluated as an indirect indicator of viral circulation in the herd.

Pregnant cows used in this study were 5 to 6 years old and had been previously vaccinated once or more times according to each cow's age. They had antibodies against the BRD's viruses that could have been due to vaccination or natural infection. Specific colostral IgG1 absorbed by the neonate during the first hours of life would have been mainly transferred intact across the small intestinal absorptive epithelium into the calf circulation. After colostrum intake, maternal antibodies could be detected in all calves. However, colostral antibodies levels measured 10 days before weaning and at weaning were heterogeneous in both groups (EW and UEW); in addition, less than 20% of calves were

expected to have antibody titers capable of protecting against infection. Thus, at the time of vaccination, calves presented a wide range of maternally-derived antibody titers. The duration of this passively derived immunity is dependent on the amount of antibodies ingested, absorbed and the rate of decay. Antibody titers usually decay to a level still high enough to block the active response to vaccination, but not high enough to protect from viral infection. This situation creates a window of susceptibility to viral infections (Chase *et al.*, 2008).

While passive immunity provides protection against BRD, this immunity could also block the development of serum antibodies when immunogens are given to calves which have maternal antibodies. However, other studies have shown that some immunogens induce immunologic memory that is not susceptible to maternal antibody regulation (Endsley *et al.*, 2003; 2004; Ellis *et al.*, 1996; Lemaire *et al.*, 2000; Platt *et al.*, 2009).

Calves used in this study showed low titers of colostral antibodies against BoHV1, no seroconversion could be observed after vaccination and the antibody levels continued to decay throughout the experience. This could be due to a failure in the vaccine used.

When antibody levels against BRSV were analyzed at weaning, a range of antibody titers could be observed but with few animals showing Ab titers of 64 or higher (3 animals in the EW and 5 animals in the UEW group). These animals remained seropositive at 56 days post weaning with antibody titers that ranged from 16 to 64. On the other hand, calves with antibody titer of 32 or lower at weaning became seronegative at the end of the experience. It is important to note that no significant differences were observed between both weaning managements.

BRSV specific colostral antibodies detected in calves indicated that the cow had been infected before; however, calves were not infected throughout the study.

The low titer of colostral antibodies against BRSV detected in EW and UEW calves, pose the need to include that viral agent in the vaccination schedule for both the cow before calving and the calf.

Given that animals were not vaccinated against BRSV, these results indicate that BRSV is probably widespread and may be contributing to respiratory disease in beef herds.

After immunization, seroconversion to BVDV could be detected in calves with low antibody titers (less than 32), while higher antibody titers exerted an inhibitory effect of the active humoral response.

The results of this study are in agreement with those reported by Fulton *et al.* (2004). In their study, the

authors found evidence that the maternal immunity had blocked antibody production, even when the calves received two doses of the vaccine. That effect was more notorious in calves with high maternally-derived antibody levels.

It has been reported that immunization of calves with maternal antibodies with an inactivated vaccine for BoHV1, BVDV and PI3 or with MLV BoHV vaccine might prime for a secondary response after a second dose of the vaccine (Ellis *et al.*, 1996; Fulton *et al.*, 2004). In concordance with that finding, it has been proposed that calves with maternal antibodies could develop protective cell-mediated immunity (T-cell) after vaccination which may protect them against viral challenge (Platt *et al.*, 2009; Vangeel *et al.*, 2007). In a recent study Woolums *et al.* (2013) have shown that the immune response of calves following administration of a 7-way MLV booster vaccination for BRD at weaning was not affected by either the calf's age or the route of priming vaccination (Woolums *et al.*, 2013). In that study, calves failed to develop a detectable immune response after receiving a priming vaccination in the face of maternal antibodies.

In the present work, transfer of passive immunity from the cow to the calf was efficient. However, a wide range of colostral antibody titers could be observed in calves 10 days pre-weaning. In consequence, the calf's age when maternally-derived antibodies do not longer protect against disease or interfere with vaccination was variable. At weaning, when calves are immunized, they may have high, moderate or low antibody titers or they can be seronegative. Under this situation, the design of a unique vaccination program so as to induce protective immunity to all calves becomes a very difficult issue.

A strategy to improve immunity of these calves managed under non-conventional weaning practices would be to enhance the specific immune response in the cow with the purpose of increasing the passive immunity to be transfer to the calf.

Incrementing the number of doses of vaccine and starting vaccination at the earliest possible time may also overcome that difficulty (Fulton *et al.*, 2004), since animals with low colostral antibodies at weaning would respond to the first immunization, while calves with higher colostral antibodies at weaning would seroconvert after the second or third vaccination. However, in beef herds, multiple immunizations of calves are difficult to manage.

These management strategies should be complemented with the screening of calves' immunity to assess the efficacy of the vaccination program. Much of

the failure of vaccination may lie in timing of administration, failure of stressed calves to respond properly to vaccination, the multifactorial nature of BRD and the increased susceptibility of stressed calves to all pathogens. Vaccination protocols should be optimized to protect calves from BRD in these anticipated weaning systems.

5. CONCLUSION

Early and ultra-early calves presented a wide range of maternally-derived antibody titers against BoHV1, BVDV and BRSV. Vaccination against BoHV-1 was not able to evoke seroconversion and antibody titers decayed to very low levels at the end of the experience. BRSV specific colostral antibodies were detected in calves, but only a few numbers of animals showed high antibody titers. Seroconversion to BVDV after vaccination could be detected in calves with low antibody titers, while calves with high antibody titers did not develop an active humoral response.

Much of the failure of vaccination may lie in timing of administration, failure of stressed calves to respond properly to vaccination, the multifactorial nature of BRD and the increased susceptibility of stressed calves to all pathogens. Vaccination protocols should be optimized to protect calves from BRD in these anticipated weaning systems.

6. ACKNOWLEDGEMENT

The researchers thank to Asociación de Cooperativas Argentinas (ACA) for the financial support and cooperation in this research work.

7. REFERENCES

- Arthington, J.D., X. Qiu, R.F. Cooke, J.M. Vendramini and D.B. Araujo *et al.*, 2008. Effects of preshipping management on measures of stress and performance of beef steers during feedlot receiving. *J. Anim. Sci.*, 86: 2016-23. DOI: 10.2527/jas.2008-0968
- Arthington, J.D., J.W. Spears and D.C. Miller, 2005. The effect of early weaning on feedlot performance and measures of stress in beef calves. *J. Anim. Sci.*, 83: 933-939. PMID: 15753350
- Belknap, E.B., J.C. Baker, J.S. Patterson, R.D. Walker and D.M. Haines *et al.*, 1991. The role of passive immunity in bovine respiratory syncytial virus-infected calves. *J. Infect. Dis.*, 163: 470-476. DOI: 10.1093/infdis/163.3.470

- Blanco, M., I. Casasus and J. Palacio, 2009. Effect of age at weaning on the physiological stress response and temperament of two beef cattle breeds. *Animal*, 3: 108-117. DOI: 10.1017/S1751731108002978
- Bolin, S.R. and J.F. Ridpath, 1995. Assessment of protection from systemic infection or disease afforded by low to intermediate titers of passively acquired neutralizing antibody against bovine viral diarrhoea virus in calves. *Am. J. Vet. Res.*, 56: 755-759. PMID: 7653884
- Callan, R.J. and F.B. Garry, 2002. Biosecurity and bovine respiratory disease. *Vet. Clin. North Am. Food Anim. Pract.*, 18: 57-77. PMID: 12064169
- Chase, C.C., D.J. Hurley and A.J. Reber, 2008. Neonatal immune development in the calf and its impact on vaccine response. *Vet. Clin. North Am. Food Anim. Pract.*, 24: 87-104. DOI: 10.1016/j.cvfa.2007.11.001
- Ellis, J.A., 2001. The immunology of the bovine respiratory disease complex. *Vet. Clin. North Am. Food Anim. Pract.*, 17: 535-550. DOI: 10.1016/j.cimid.2012.01.005
- Ellis, J.A., L.E. Hassard, V.S. Cortese and P.S. Morley, 1996. Effects of perinatal vaccination on humoral and cellular immune responses in cows and young calves. *J. Am. Vet. Med. Assoc.*, 208: 393-400. PMID: 8575972
- Endsley, J.J., J.A. Roth, J. Ridpath and J. Neill, 2003. Maternal antibody blocks humoral but not T cell responses to BVDV. *Biologicals*, 31: 123-125. DOI: 10.1016/S1045-1056(03)00027-7
- Endsley, J.J., J.F. Ridpath, J.D. Neill, M.R. Sandbulte and J.A. Roth, 2004. Induction of T lymphocytes specific for bovine viral diarrhoea virus in calves with maternal antibody. *Viral Immunol.*, 17: 13-23. DOI: 10.1089/088282404322875421
- Ferella, A., D. Bellido, P. Chacana, A. Wigdorovitz, M.J. Dus Santos and A. Wigdorovitz, 2012. Chicken egg yolk antibodies against bovine respiratory syncytial virus neutralize the virus in vitro. *Procedia Vaccinol.*, 6: 3-3. DOI: 10.1016/j.provac.2012.04.006
- Fulton, R.W., R.E. Briggs, M.E. Payton, A.W. Confer and J.T. Saliki *et al.*, 2004. Maternally derived humoral immunity to Bovine Viral Diarrhoea Virus (BVDV) 1a, BVDV1b, BVDV2, bovine herpesvirus-1, parainfluenza-3 virus bovine respiratory syncytial virus, *Mannheimia haemolytica* and *Pasteurella multocida* in beef calves, antibody decline by half-life studies and effect on response to vaccination. *Vaccine*, 22: 643-649. DOI: 10.1016/j.vaccine.2003.08.033
- Galli, I., A. Monje, S. Vittone, D. Sampedro and C. Busto, 2005. Destete precoz in cría vacuna. Manual para la toma de decisiones there Ejecucion the técnica.
- Galli, I., G. Teira, F. Perlo, P. Bonato and O. Tisocco *et al.*, 2008. Animal performance and meat quality in cull cows with early weaned calves in Argentina. *Meat. Sci.*, 79: 521-528. DOI: 10.1016/j.meatsci.2007.10.007
- Haley, D.B., D.W. Bailey and J.M. Stookey, 2005. The effects of weaning beef calves in two stages on their behavior and growth rate. *J. Anim. Sci.*, 83: 2205-2214. PMID: 16100076
- Hickey, M.C., M. Drennan and B. Earley, 2003. The effect of abrupt weaning of suckler calves on the plasma concentrations of cortisol, catecholamines, leukocytes, acute-phase proteins and in vitro interferon-gamma production. *J. Anim. Sci.*, 81: 2847-2855. PMID: 14601889
- Howard, C.J., M.C. Clarke and J. Brownlie, 1989. Protection against respiratory infection with bovine virus diarrhoea virus by passively acquired antibody. *Vet. Microbiol.*, 19: 195-203. DOI: 10.1016/0378-1135(89)90066-7
- Hulbert, L.E., C.J. Cobb, J.A. Carroll and M.A. Ballou, 2011. The effects of early weaning on innate immune responses of Holstein calves. *J. Dairy Sci.*, 94: 2545-2556. DOI: 10.3168/jds.2010-3983
- Lemaire, M., V. Weynants, J. Godfroid, F. Schynts and G. Meyer *et al.*, 2000. Effects of bovine herpesvirus type 1 infection in calves with maternal antibodies on immune response and virus latency. *J. Clin. Microbiol.*, 38: 1885-1894. PMID: 10790117
- Lynch, E.M., B. Earley, M. McGee and S. Doyle, 2010a. Characterisation of physiological and immunological responses in beef cows to abrupt weaning and subsequent housing. *BMC Vet. Res.*, 6: 37-37. DOI: 10.1186/1746-6148-6-37
- Lynch, E.M., B. Earley, M. McGee and S. Doyle, 2010b. Effect of abrupt weaning at housing on leukocyte distribution, functional activity of neutrophils and acute phase protein response of beef calves. *BMC Vet. Res.*, 6: 39-39. DOI: 10.1186/1746-6148-6-39
- Lynch, E.M., M. McGee, S. Doyle and B. Earley, 2012. Effect of pre-weaning concentrate supplementation on peripheral distribution of leukocytes, functional activity of neutrophils, acute phase protein and behavioural responses of abruptly weaned and housed beef calves. *BMC Vet. Res.*, 8: 1-1. PMID: 22217360
- Mechor, G.D., C.G. Rousseaux, O.M. Radostits, L.A. Babiuk and L. Petrie, 1987. Protection of newborn calves against fatal multisystemic infectious bovine rhinotracheitis by feeding colostrum from vaccinated cows. *Can. J. Vet. Res.*, 51: 452-459. PMID: 2839276

- Parreno, F., R. Alvarez-Valdes, J.F. Oliveira and J.M. Tamarit, 2010. Neighborhood structures for the container loading problem: A VNS implementation. *J. Heuristics*, 16: 1-22. DOI: 10.1007/s10732-008-9081-3
- Platt, R., P.W. Widel, L.D. Kesi and J.A. Roth, 2009. Comparison of humoral and cellular immune responses to a pentavalent modified live virus vaccine in three age groups of calves with maternal antibodies, before and after BVDV type 2 challenge. *Vaccine*, 27: 4508-4519. DOI: 10.1016/j.vaccine.2009.05.012
- Price, E.O., J.E. Harris, R.E. Borgward, M.L. Sween and J.M. Connor, 2003. Fenceline contact of beef calves with their dams at weaning reduces the negative effects of separation on behavior and growth rate. *J. Anim. Sci.*, 81: 116-121. PMID: 12597380
- Richeson, J.T., E.B. Kegley, J.G. Powell, P.A. Beck and B.L. Ley *et al.*, 2012. Weaning management of newly received beef calves with or without continuous exposure to a persistently infected bovine viral diarrhea virus pen mate: Effects on health, performance, bovine viral diarrhea virus titers and peripheral blood leukocytes. *J. Anim. Sci.*, 90: 1972-1985. PMID: 22648754
- Romera, S.A., L.A. Hilgers, M. Puntel, P.I. Zamorano and V.L. Alcon *et al.*, 2000. Adjuvant effects of sulfolipo-cyclodextrin in a squalene-in-water and water-in-mineral oil emulsions for BHV-1 vaccines in cattle. *Vaccine*, 19: 132-141. DOI: 10.1016/S0264-410X(00)00104-3
- Snowder, G., 2009. Genetics, environment and bovine respiratory disease. *Anim. Health Res. Rev.*, 10: 117-119. DOI: 10.1017/S1466252309990144
- Stilwell, G., M. Matos, N. Carolino and M.S. Lima, 2008. Effect of a quadrivalent vaccine against respiratory virus on the incidence of respiratory disease in weaned beef calves. *Prev. Vet. Med.*, 85: 151-157. DOI: 10.1016/j.prevetmed.2008.02.002
- Taylor, J.D., R.W. Fulton, T.W. Lehenbauer, D.L. Step and A.W. Confer, 2010a. The epidemiology of bovine respiratory disease: What is the evidence for predisposing factors? *Can. Vet. J.*, 51: 1095-1102. PMID: 21197200
- Taylor, J.D., R.W. Fulton, T.W. Lehenbauer, D.L. Step and A.W. Confer, 2010b. The epidemiology of bovine respiratory disease: What is the evidence for preventive measures? *Can. Vet. J.*, 51: 1351-1359. PMID: 21358927
- Valerio, D.A., 2007. Destete precoz e hiper precoz; derrumbando paradigmas. Day E.E.A. INTA Concepción del Uruguay and ACA Animal Nutrition Division.
- Vangeel, I., A.F. Antonis, M. Fluess, L. Riegler and A.R. Peters *et al.*, 2007. Efficacy of a modified live intranasal bovine respiratory syncytial virus vaccine in 3-week-old calves experimentally challenged with BRSV. *Vet. J.*, 174: 627-635. DOI: 10.1016/j.tvjl.2006.10.013
- Woolums, A.R., R.D. Berghaus, L.J. Berghaus, R.W. Ellis and M.E. Pence *et al.*, 2013. Effect of calf age and administration route of initial multivalent modified-live virus vaccine on humoral and cell-mediated immune responses following subsequent administration of a booster vaccination at weaning in beef calves. *Am. J. Vet. Res.*, 74: 343-354. DOI: 10.2460/ajvr.67.7.1179