

# Accepted Manuscript

Effect of iron and ascorbic acid addition on dry infusion process and final color of pumpkin tissue

C. Genevois, S. Flores, M. de Escalada Pla



PII: S0023-6438(14)00170-4

DOI: [10.1016/j.lwt.2014.03.020](https://doi.org/10.1016/j.lwt.2014.03.020)

Reference: YFSTL 3813

To appear in: *LWT - Food Science and Technology*

Received Date: 5 June 2013

Revised Date: 18 March 2014

Accepted Date: 20 March 2014

Please cite this article as: Genevois, C., Flores, S., de Escalada Pla, M., Effect of iron and ascorbic acid addition on dry infusion process and final color of pumpkin tissue, *LWT - Food Science and Technology* (2014), doi: 10.1016/j.lwt.2014.03.020.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1       **EFFECT OF IRON AND ASCORBIC ACID ADDITION ON DRY INFUSION**  
2                   **PROCESS AND FINAL COLOR OF PUMPKIN TISSUE.**

3  
4       Genevois, C.<sup>1,2</sup>; Flores, S.<sup>1,2</sup>; de Escalada Pla, M.<sup>1,2\*</sup>

5  
6       <sup>1</sup> Departamento de Industrias. Facultad de Ciencias Exactas y Naturales. Universidad  
7       de Buenos Aires (UBA). Ciudad Universitaria. (1428), CABA, Argentina.

8       <sup>2</sup> Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

9       \*Corresponding author. e-mail: [marina@di.fcen.uba.ar](mailto:marina@di.fcen.uba.ar). Phone. 54 11 45 76 33 66.

10  
11       **Abstract**

12       In the present study, pumpkin (*Cucurbita moschata* Duchesne ex Poiret) was used as  
13       raw material to produce sweet food fortified with iron (Fe) and ascorbic acid (AA). A  
14       dry infusion process with a subsequent air drying was applied. Response surface  
15       methodology was performed in order to analyze the effect of Fe and AA incorporation  
16       into the formulation on: water loss (WL) and solid gain (SG) during the dry infusion  
17       process, color changes ( $\Delta E$ ) and the dehydration percentage during subsequent air  
18       drying process. The results showed that the presence of Fe and/or AA promoted SG  
19       and WL during the dry infusion and also, weight changes during the air drying process  
20       (PP). An increase of the color changes was also observed. In turn, it was possible to  
21       obtain predictive equations for the parameters studied. The application of edible  
22       coating based on tapioca starch on pumpkin product was also tested showing a  
23       protective effect from the pumpkin color view point.

24  
25       **Key words:** *Cucurbita moschata* Duchesne ex Poiret, functional foods, iron  
26       fortification, edible coating.

28 **Abbreviations**

29

30	Iron	Fe
31	Ascorbic acid	AA
32	Water loss	WL
33	Solid gain	SG
34	Color changes	$\Delta E$
35	Weight changes due to air drying process	PP
36	Micronutrient malnutrition	MM
37	World Health Organization	WHO
38	Central composite design	CCD
39	Revolutions per minute	rpm
40	Recommended Daily Intake	RDI
41	Recommended Dietary Allowance	RDA
42	Non-enzymatic browning	NEB

## 43 1. INTRODUCTION

44 The micronutrient malnutrition (MM) is widespread over the world, but developing  
45 regions are the most affected. From a public health point of view, MM is a concern not  
46 only for the large number of people affected, but also because it remains a risk factor  
47 for many diseases (Ashwell, 2004). Iron (Fe) deficiency is considered the most  
48 prevalent of the MM, showing a continuous increase in its prevalence, representing the  
49 main nutritional deficiency problem in terms of magnitude and spatial distribution (Allen,  
50 Benoist, Dary & Hurrell, 2006; Souto de Olivera, 2009). At present, it is estimated that 2  
51 billion people, or over 30% of the world population, are anemic, mainly due to Fe  
52 deficiency and this situation is further magnified in low-income areas with a high  
53 incidence of infectious diseases that contribute to the high prevalence of anemia  
54 according to World Health Organization (WHO, 2013). Both Fe deficiency and anemia,  
55 even in its moderate form, have serious health consequences for the population,  
56 including stunted growth and cognitive development (WHO, 2013; Zimmermann &  
57 Hurrell, 2007).

58 By the moment, food fortification with Fe is considered the strategy most sustainable  
59 and cost-effective against iron deficiency (Laxmi Narayan, Mills, & Berman, 2006;  
60 Tripathi & Platel, 2013). Nevertheless, there are some technological difficulties to be  
61 solved like changes and unpleasant sensory characteristics of the food matrix due to  
62 this fortification. The Fe compounds that are very soluble in water, for example ferrous  
63 sulfate, provide Fe of high bioavailability and, therefore, would be the primary choice in  
64 food fortification. However, in this type of compounds, Fe is highly reactive, causing  
65 oxidation of fats, vitamins and several amino acids in the food that is fortified (Boccio &  
66 Monteiro, 2004; Gaucheron, 2000) and, consequently, undesirable color and flavor  
67 changes in the food matrix could appear. Rao and Kawamura (2008) reported that the  
68 major technological problems caused by soluble salts of Fe in the production of food  
69 and beverages are the color and flavor alterations.

70 At the same time, there are dietary compounds which positively affect the Fe  
71 absorption, as is the case of ascorbic acid (AA). The presence of this hydrosoluble  
72 vitamin at the intestinal level promotes absorption of non-hemic Fe by means of its  
73 reduction to ferrous ion ( $\text{Fe}^{+2}$ ). In foods, the AA acts as a reducing agent keeping the  
74 Fe in its soluble reduced form (de Escalada Pla, Campos, & Gerschenson, 2009; Souto  
75 de Olivera, 2009), and also acts as an antioxidant through the free radicals  
76 neutralization at the cellular level (Rojas, 1995). Some studies have also shown that  
77 vitamin A and, even more the  $\beta$ -carotene, significantly increase the bioavailability of Fe  
78 (Binaghi, Greco, López, Ronayne, & Valencia, 2005).

79 The policy adopted by some countries was to select as a carrier, those foods widely  
80 consumed by the risk groups. Vegetable and fruit matrices have widely been used to  
81 support vitamins and minerals like  $\text{Ca}^{2+}$  and  $\text{Zn}^{2+}$ , applying impregnation or vacuum  
82 impregnation technology for their enrichment (Gras, Vidal, Betoret, Chiralt, & Fito,  
83 2003). This processing has been proposed by Zhao and Xie (2004) as a pre-treatment  
84 before the final drying step with the purpose of achieving two goals: decreasing  
85 moisture content before final air drying to save energy and incorporating functional  
86 solutes, such as nutrients, antimicrobial, antioxidant, and anti-browning agents to  
87 improve product quality. The impregnation processes of fruits and vegetables with  
88 hypertonic solutions were widely studied and well reported (Gras et al., 2003; Moreno  
89 et al., 2012; Spiazzi & Mascheroni, 1997; Zhao & Xie, 2004). Dry infusion was  
90 recommended as a practical tool for small producers as fruit preservation process that  
91 could be performed in rural areas (Alzamora, Guerrero, Nieto & Vidales, 2003).

92 Edible coatings can have an additive or synergistic effect with other stress factors in  
93 the task of improving the overall quality of foods. The application of coatings on fruits  
94 and vegetables improved color and flavor retention during storage, extending the shelf  
95 life of the product, retarding moisture and/or firmness loss and product senescence  
96 (Campos, Gerschenson & Flores, 2011).

97 Pumpkin *Cucurbita moschata* is one of the most consumed vegetables in Argentina.  
98 Furthermore, an increasing interest in this vegetable has also been reported in other  
99 countries (Gwanama, Botha, & Labuschagne, 2008). Tissue from this kind of pumpkin  
100 was characterized previously (de Escalada Pla, Ponce, Wider, Stortz, Rojas, &  
101 Gerschenson, 2005; de Escalada Pla, Delbon, Rojas, & Gerschenson, 2006; de  
102 Escalada Pla, Ponce, Stortz, Gerschenson, & Rojas, 2007). More recently, the  
103 adequacy of pumpkin mesocarp tissue as a food matrix for Fe supply was reported (de  
104 Escalada Pla et al., 2009). The iron was incorporated after blanching and during the  
105 cooling step. Then, a hypertonic osmotic covering solution was added to storage bags.  
106 The aim of the present work was to study: 1) the possibility of fortifying *Cucurbita*  
107 *moschata* Duchesne ex Poiret tissues with iron through a process of dry infusion, thus  
108 avoiding the use of huge amounts of hypertonic osmotic solutions; 2) the effect of the  
109 joint presence of Fe and AA on process parameters, physical and quality  
110 characteristics in the final product; and 3) the application of an edible coating based on  
111 tapioca starch for protecting pumpkin tissue from possible color detriments due to  
112 Fe/AA contents during the process and food storage.  
113

## 114 2. Material and methods

115

### 116 2.1 Chemicals

117 Food grade sucrose and tapioca starch were employed. The additives:  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$   
118 (Merck, Argentina); potassium sorbate (Sigma, USA); L-(+)-ascorbic acid (Merck,  
119 Argentina); citric acid and glycerol (Sintorgan, Argentina) and other chemicals used  
120 were of analytical grade.

121

### 122 2.2 Preparation of the pumpkin fortified with Fe and AA

123 Pumpkin (*Cucurbita moschata* Duchesne ex Poiret) obtained in a local supermarket  
124 was carefully washed and rinsed with distilled water. Then, cylinders of 15 mm  
125 diameter and 10 mm thickness were cut from the mesocarp using a stainless steel cork  
126 borer. The cylinders were blanched with water vapor for 8 minutes and then rapidly  
127 cooled for 1 minute by immersion in water at 0°C. Finally, they were impregnated with  
128 sucrose (900 g/kg of pumpkin), citric acid (1.5 g/kg of pumpkin) and potassium sorbate  
129 (1.9 g/kg) following a dry infusion process described by Alzamora et al. (2003). Briefly,  
130 pumpkin cylinders were placed in a plastic bowl and sprinkled with powdered sucrose.  
131 Water from vegetal tissue began to flow from the pumpkin cylinder to the surrounding  
132 sucrose concentrate. In that moment, citric acid, potassium sorbate, AA and Fe salt  
133 were added to the liquid solution and the orbital agitation started up. Citric acid was  
134 added in order to decrease pH values below 5; since sorbate and sorbic acid as an  
135 antimicrobial are more effective in this range of pH (Lindsay, 1996). In order to  
136 evaluate the effect of AA and Fe during the preparation process and on the final color  
137 quality, different amounts of AA and  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  were added to the systems according  
138 to a central composite design (CCD) of two factors (independent variables) and five  
139 levels (Table 1). Pumpkin used in all the systems came from a same single lot of raw  
140 product.

141 The dry infusion was carried out at 20°C up to equilibrium on an orbital shaker (Vicking  
142 S.A., Argentina) at 35 revolutions per minute (rpm) to assure good contact of tissue  
143 and the impregnating system. Equilibrium was reached at 72 hours when pumpkin  
144 cylinders and the surrounding solution achieved the same  $a_w$  and pH values. Once the  
145 dry infusion was concluded, the cylinders were drained through a stainless steel  
146 strainer and dried under forced air convection at 40°C for 3 hours, in order to achieve a  
147 water activity ( $a_w$ ) value below 0.85 (Fontana A., 2008).  
148 Finally, the pumpkin cylinders were introduced into low density polyethylene bags of 80  
149  $\mu\text{m}$  thickness, provided with a Ziploc® type closure. Each bag was filled with 5  
150 mesocarp pieces (10 g) and stored in a chamber at 18-20°C.

151

### 152 **2.3 Preparation of the pumpkin fortified and coated**

153 From the results obtained with CCD (see item 3.2), one formulation was chosen and  
154 one additional batch was performed. A dry infusion process, as previously described,  
155 was carried out and after draining, the cylinders were separated into two parts. One  
156 part was dipped into a solution of gelatinized starch in order to generate an edible  
157 coating on pumpkin cylinders, and the other part, pumpkin without coating was also  
158 prepared for comparing purposes in subsequent testing assays. Impregnated pumpkins  
159 with or without coating application were submitted to a drying process with force air  
160 convection at 40°C for 3 hours in order to achieve the following purposes: (1) to  
161 constitute the coating, in the case of coated cylinders and (2) to obtain an additional  
162 reduction of  $a_w$  in both cases (Fontana A., 2008).

163 The edible coating was prepared with native tapioca starch (50 g/kg), glycerol (20 g/kg)  
164 as a plasticizer and potassium sorbate (1 g/kg) as an antimicrobial agent. Samples  
165 were packed and stored as previously explained.

166

### 167 **2.4 Product characterization**



168 In order to analyze the changes during the processing and storage of tissue, the  
 169 samples were taken from blanched pumpkin, equilibrated pumpkin after dry infusion;  
 170 and dried tissue after forced air convection drying. Also, samples of the final product  
 171 after 9 days of storage at 18-20°C were evaluated.

172 The following properties were measured:

173 ♦ pH and  $a_w$ :

174 Pumpkin cylinders were reduced to a puree with the aid of a homogenizer Ultraturrax  
 175 (IKA, USA) at 6500 rpm for 20 seconds. The pH was determined with a pH meter  
 176 (Cole-Parmer, USA).

177 Water activity ( $a_w$ ) was measured with a hygrometer (Aqualab, USA) at 20°C.

178 ♦ Moisture and soluble solids contents:

179 Pumpkin samples were frozen and freeze dried (Christ, Germany) for 48 hours under  
 180 vacuum ( $\approx 1.1$  Pascal) and 25°C, to determine the water content.

181 The percentage of soluble solids (°Brix) was determined with a refractometer with  
 182 automatic temperature compensation (Atago, USA) in the juice extracted from pumpkin  
 183 cylinders by pressing the sample with a spatula.

184 Water loss (WL) and solid gain (SG) in the different systems, during the dry infusion  
 185 step, were calculated according to the following equations (de Escalada Pla et al.,  
 186 2009):

187

$$WL = \frac{M_t \times m_t - M_0 \times m_0}{M_0} \times 100$$

188

189

$$SG = \frac{M_t \times ss_t - M_0 \times ss_0}{M_0} \times 100$$

190

191

192 Where  $M_t$  (g) is the average mass of pumpkin cylinders at time  $t$ ;  $m_t$  is the moisture  
 193 content of tissue at time  $t$  [g water/100 g pumpkin, wet basis];  $M_0$  (g) is the cylinder  
 194 mass average at initial time (before the dry infusion);  $m_0$  is the initial water content of  
 195 tissue [g water / 100 g pumpkin, wet basis];  $ss_0$  and  $ss_t$  are the soluble solid contents in  
 196 tissue at initial time and at time  $t$  [°Brix, or g s s/100 g pumpkin, wet basis], respectively.  
 197 Measurements were performed in duplicate for each system and the average value is  
 198 reported.

199 The water loss during the subsequent air drying process (PP) in wet basis was  
 200 calculated as: 
$$PP = \frac{P_i - P_f}{P_i} \times 100$$

201  $P_i$ : mass of the sample before convective drying.

202  $P_f$ : mass of sample after convective drying.

203

204 ♦ Color

205 Before and after drying, color parameters were evaluated using a photocolormeter  
 206 (Minolta, Japan) in the CIE  $L^*a^*b^*$  space [ $L^*$ : lightness,  $a^*$ : greenness - redness,  $b^*$ :  
 207 blueness - yellowness] under illuminant D65 and with the observer at an angle of two  
 208 degrees. From these parameters, color difference ( $\Delta E$ ) was calculated according to:

209

$$210 \quad \Delta E = \sqrt{(L^* - L_{ref}^*)^2 + (a^* - a_{ref}^*)^2 + (b^* - b_{ref}^*)^2}$$

211

212 Where reference values ( $L_{ref}^*$ ,  $a_{ref}^*$  and  $b_{ref}^*$ ) correspond to the control system,  
 213 impregnated with sucrose in the presence of citric acid and potassium sorbate but  
 214 without addition of Fe and AA in the dry infusion media (system C). In the case of the  
 215 edible coating effect, the color difference was calculated taking as a reference the  
 216 fortified cylinders after infusion and before coating and drying.

217

218 The value of Chroma parameter was also calculated. This parameter describes color  
219 intensity (Olivera et al., 2008) and was calculated as  $\text{Chroma} = (a^{*2} + b^{*2})^{(1/2)}$ . The  
220 averages of three measurements are reported.

221

## 222 **2.5 Experimental design and statistical analysis**

223 In order to evaluate the influence of AA and Fe during dry infusion and drying process  
224 as well as on final color quality, a CCD with two factors (independent variables) and at  
225 five levels (Table 1) was performed. The selection criterion for the lowest levels was, in  
226 the case of Fe, to cover 20% of the Recommended Daily Intake (RDI) with a 100 g  
227 portion, and in the case of AA, to cover 100% of the Recommended Dietary  
228 Allowance (RDA), according to the Argentine Food Code (2012) in its article 1363.  
229 The highest levels used were chosen to cover 100% of the RDA in the case of Fe, and  
230 in the case of AA was considered the level of no observed adverse effects value, with a  
231 maximum of 1000 mg. The central point (0;0) was performed in triplicate. Table 1  
232 shows all experimental runs.

233 Dependent variables WL, SG, PP and  $\Delta E$  were fitted using a second degree  
234 polynomial equation and a multiple regression procedure:

$$235 \quad \psi = B_0 + B_1x_1 + B_2x_2 + B_{11}x_1^2 + B_{22}x_2^2 + B_{12}x_1x_2$$

236 Where,  $\psi$  is the dependent variable analyzed;  $x_1$  and  $x_2$  are independent (Fe and AA  
237 contents) variables that affected  $\psi$  value;  $B_0$  is the value of the fitted response at the  
238 center point of the design, ( $x_1 = 0$  and  $x_2 = 0$ );  $B_1$  and  $B_2$  are the linear coefficients;  $B_{11}$   
239 and  $B_{22}$  are the quadratic coefficients and  $B_{12}$  is the cross coefficient between factors.  
240 This equation permitted to evaluate the effects of linear, quadratic and interaction terms  
241 of independent variables on selected dependent variables. The analysis of variance  
242 (ANOVA) was conducted to assess the adequacy of the model by calculation of the F

243 value for the regression and the determination coefficient ( $R^2$ ), as well as to evaluate  
244 the significance of the equation coefficients. Three dimensional plots were generated  
245 (response surfaces) by fixing investigated variables to the center value of CCD.

246 On the other hand, in order to identify a Fe:AA ratio that minimizes undesirable color  
247 changes, the experimental values of the Chroma parameter were analyzed by the  
248 "Analysis of a central composite experiment (surface response)" module.

249 For color comparative purposes, an additional unfortified system was prepared under  
250 the same conditions as reference.

251 In addition, the significant differences among results were established by analysis of  
252 variance (ANOVA) with a significance level of 0.05 and applying a *post hoc* test, the  
253 Least Significant Difference (LSD) test. The results are reported based on their mean  
254 and standard deviation. Statistica software (version 6, StatSoft, Inc. 2001, USA) was  
255 used for the analysis of the design and generation of the response surfaces and also  
256 for statistical treatment of data.

257

### 258 **3. Results and discussion**

259

#### 260 **3.1 Characteristics of the impregnated and dried product**

261 At the end of dry infusion, the  $a_w$  of pumpkin was in the range of 0.91 and 0.93, while  
262 the initial  $a_w$ , after blanching and before infusion, was  $\sim 1.0$ . Once equilibrated,  
263 samples also showed pH values in the range of 3.4 to 4.6. Neither AA nor Fe exerted  
264 significant effects on  $a_w$  of final product. Nevertheless, the pH decreased, as expected,  
265 when the AA concentration increased ( $p < 0.05$ ), as can be seen in Table 2.

266 Table 2, shows the values of WL and SG measured on tissues submitted to dry  
267 infusion for different contents of Fe and AA. Furthermore, PP values of the  
268 impregnated pumpkin after air drying are also reported. In order to analyze the effect of  
269 Fe and AA contents on the dry infusion and drying process, data were fitted using a

270 second degree polynomial equation. The best fit equation and corresponding plots of  
271 the linear, quadratic and interactive effects of Fe and AA on SG, WL, and PP are  
272 shown in Figure 1, panel a, b and c respectively.

273 It could be seen that SG occurred during the dry infusion process and varied in the  
274 range of 8.8% - 19.7% (Table 2). Figure 1a, for SG, shows the response surface and  
275 the corresponding equation. The linear terms of Fe and AA were significant as well as  
276 the quadratic term of the factor AA (Figure 1a). It would mean that an addition of Fe or  
277 AA promotes the incorporation of solids inside the pumpkin tissue. However, the  
278 presence of both additives simultaneously presents an antagonistic effect because the  
279 interaction term was negative.

280 It could be seen that the WL was varied between 69% and 72.6% (Table 2). In this  
281 case, linear coefficients were both positive, indicating that the presence of Fe or AA  
282 promotes osmotic dehydration in the pumpkin vegetable matrix and it is expected that  
283 the addition of Fe to the formulation exerts the greatest influence on the value of WL,  
284 since the linear coefficient of Fe factor was positive and with a greater magnitude  
285 (Figure 1b). Once again, the presence of both additives simultaneously shows an  
286 antagonistic effect because the interaction term was negative. According to de  
287 Escalada Pla et al. (2009), Fe presence in pumpkin tissue favored the water loss  
288 during an impregnation process with hypertonic solution. Similar results were also  
289 reported by Barrera C., Betoret N. and Fito P. (2004) with vacuum impregnation of  
290 apple tissue fortified with calcium or Fe.

291 Subsequent air drying lowered the water activity about 15%. The final  $a_w$  ranged  
292 between 0.77 and 0.82. The weight changes due to the air drying process (PP) were  
293 approximately 21.9 to 26.9% (Table 2). The predictive equation (Figure 1c) indicated  
294 that the linear terms, the quadratic term for Fe and the interaction term were significant.  
295 A positive effect was observed through linear coefficients, indicating that the presence  
296 of Fe or AA promoted the air dehydration process. Significant negative coefficients for

297 quadratic term of Fe and the interaction term were also observed, indicating a  
298 curvature of the surface.

299

### 300 **3.2 Color evaluation**

301 In Table 3, color attributes can be observed for the final product obtained from the  
302 different treatments. The color difference ( $\Delta E$ ) was determined taking control systems  
303 (unfortified) as reference. Response surface for  $\Delta E$  and the corresponding equation are  
304 shown in Figure 1d. It could be observed that the addition of Fe or AA generated a  
305 darkened color of the pumpkin compared to the control system (Table 3). The second  
306 order equation obtained indicates that the linear and quadratic terms are significant,  
307 being the former positive and the latter ones, negative. The interaction term was not  
308 significant, suggesting that each factor exerts an independent effect on the color  
309 change (Figure 1d).

310 In order to assess the color development in the systems studied, a picture of them is  
311 shown in Figure 2. The control system (C), without fortification was also included for  
312 comparison purposes. It can be observed that system 6 showed the smallest color  
313 alterations due to the fortification and process applied. Based on these observations,  
314 differences in  $L^*$  and in the Chroma parameter due to the final step of process were  
315 also analyzed. Table 3 shows  $L^*$  and Chroma values for impregnated pumpkin, before  
316 and after the air drying process. In general, a reduction of  $L^*$  and Chroma values after  
317 air drying, could be observed. This effect was not evidenced in system 6, neither in the  
318 control system, where no significant changes due to air drying, were observed for  $L^*$   
319 and neither for Chroma.

320 The addition of Fe or AA significant reduced  $L^*$  and Chroma values in comparison with  
321 the control system. For all the systems studied,  $L^*$  ranged between 31 and 38 and the  
322 Chroma presented values from 16.2 to 27 (Table 3). On the other hand, Chroma was

323 the parameter most significantly affected by the drying process. It might be concluded  
324 that the color difference observed was mainly related to chromatic coordinates:  $a^*$  and  
325  $b^*$  changes. The first step of AA destruction is part of the non-enzymatic browning  
326 (NEB) reaction chain (Rojas & Gerschenson, 2001; León & Rojas, 2007). Degradation  
327 of AA through hydrolysis can occur simultaneously to AA oxidation when oxygen is  
328 present, producing 2-keto-L-gulonic acid. It can then be considered that at least two  
329 irreversible parallel or competitive reactions proceed: the AA hydrolysis and the AA  
330 oxidation (De'Nobili, Curto, Delfino, Soria, Fissore, & Rojas, 2013). Some researchers  
331 reported that hydrolytic instability of AA could be responsible for NEB and the decrease  
332 in edible film lightness with storage (De'Nobili et al., 2013; Pérez, De'Nobili, Rizzo,  
333 Gerschenson, Descalzo, & Rojas, 2013). On the other hand, iron in the reduced state  
334 is an active prooxidant, and ascorbate, which could act as a hydrogen donor, in  
335 synergism with iron, serves as an effective chelator (Rosenthal, Rosen, & Bernstein,  
336 1993). However, Hegenauer, Saltman, & Ludwig (1979) indicated that the conversion  
337 of ascorbate to dehydroascorbate and of dehydroascorbate to 2-keto-L-gulonate  
338 occurs rapidly even in unsupplemented milk. Thus, iron supplementation may not affect  
339 materially the vitamin C content of stored milk (Gaucheron, 2000). During the drying  
340 process, the carotenoids can be degraded by exposure to heat and oxygen, with a  
341 consequent increase in cis-isomers (Lago-Vanzela, do Nascimento, Fontes, Mauro, &  
342 Kimura, 2013). Probably, iron contents catalyzed this degradation, altering pumpkin  
343 color. Lightness and Chroma changes observed herein seemed to be related to  
344 independent mechanisms, one associated with AA destruction and the other with  
345 carotenoid oxidation. However, it could be interesting to determine the AA and Fe  
346 contents that minimize these effects.

347 The Chroma parameter was then analyzed in order to detect the Fe and AA  
348 concentration that let us obtain a Chroma value similar to that of the control system.  
349 Response surface obtained for the Chroma value is shown in Figure 3.

350 It must be remarked that all the coefficients of the corresponding second degree  
351 polynomial were significant ( $p < 0.05$ ), except for the coefficient of interaction.

352 In order to define a formulation that allows one to obtain an adequate color, the  
353 Chroma value from the control system (pumpkin without fortification) was taken as the  
354 target value. From equation of prediction, a formulation with 0.3475 g Fe/kg pumpkin  
355 and 0.8745 g AA/kg pumpkin was obtained. It must be remarked that concentration  
356 used on system 6 of CCD, was the most similar to that obtained according to  
357 optimization criteria. Nevertheless, the statistically recommended formulation was  
358 performed and the Chroma of the final product was evaluated recording a value of  $29.6$   
359  $\pm 0.6$  which is not significantly different ( $p < 0.05$ ) from the target value selected (System  
360 C, Table 3).

361

### 362 **3.3 Characteristics of the pumpkin fortified and coated**

363 Based on the formulation proposed, an additional batch was performed and one part of  
364 it was covered with a starch based coating. The other part of the batch was maintained  
365 without coating. All samples were tested evaluating their color (Table 4).

366 Pumpkin cylinders were weighed before and after the edible coating application.  
367 Consequently, it could be estimated that  $\sim 1$ g average of starch gel was deposited on  
368 the surface of each pumpkin cylinder during the dipping process.

369 During the drying process, a water loss of  $\sim 30\%$  was registered, reaching the final  
370 product with an  $a_w$  value of  $\sim 0.8$ . Drying also affected the color of product as can be  
371 observed through the  $\Delta E$  value at the beginning of storage, mainly on system without  
372 coating (Table 4). As can be observed, in Table 4, coating significant reduce product  
373 color changes due to the drying process.

374 Moreover, the processing applied significantly ( $p < 0.05$ ) reduced  $L^*$  values for both  
375 systems, nevertheless, the uncoated system presented a higher reduction. With



376 reference to the Chroma values, no significant differences were observed for system  
377 coated while a significant ( $p < 0.001$ ) reduction was recorded for uncoated one, due to  
378 air drying (Table 4). This suggests a protecting action of the starch coating used during  
379 the air drying process from the point of view of the color. Flores (2006) reported a low  
380 oxygen permeability of tapioca starch coatings, and this property could in part explain  
381 their capacity to protect pumpkin color specially avoiding AA and ferrous iron oxidation.  
382 Lago-Vanzela et al. (2013) assayed edible coatings from native and modified starches  
383 on pumpkin during drying and reported that dehydrated coated products had a better  
384 color and a significantly higher retention of trans- $\alpha$ -carotene and trans- $\beta$ -carotene than  
385 products that did not receive coating. They claimed that the good carotenoid retention  
386 determined in the samples covered with modified cassava starch suggested that the  
387 coating worked as an efficient barrier against oxygen (Lago-Vanzela et al., 2013).  
388 Table 4 also shows values obtained after 9 days of storage. In this case, it could be  
389 observed that for both samples, coated and uncoated, the Chroma value did not  
390 change significantly after nine days of storage at 18-20°C.

391

#### 392 **4. Conclusions**

393 A dry infusion process could be used successfully to incorporate Fe and AA into  
394 pumpkin tissue. It was found that the addition of Fe or AA promoted osmotic  
395 dehydration in pumpkin and water loss during the subsequent air drying process.

396 The presence of Fe or AA intensified color differences of the systems when compared  
397 with the control system (unfortified) and this was mainly detected through the Chroma  
398 evaluation. The dry infusion with Fe and AA with subsequent air drying significantly  
399 decreased the value of the Chroma parameter of the pumpkin matrix with respect to  
400 the value for the unfortified system with the exception of product obtained through the  
401 impregnation in a formulation containing 0.216 g/kg of Fe and 0.80 g/kg of AA (system

402 6), for which the color after the drying process was similar to the one observed for the  
403 control system. From preliminary data herein reported, it might be suggested that  
404 edible tapioca starch coating exerted a protective effect in terms of the color of  
405 pumpkin cylinders during drying.

406 The present study provides important information for the design and processing of a  
407 pumpkin product fortified with Fe and AA which can enlarge the existing background for  
408 the optimization of the production and stability of new functional foods. As a  
409 perspective, a comparison of these results with a test of the consumers' acceptance  
410 could be interesting to perform.

411

412 **Acknowledgements.**

413 The authors acknowledge the financial support from University of Buenos Aires  
414 (UBACyT 20020110200192 y 20020090200693), National Agency of Scientific and  
415 Technological Promotion of Argentina (ANPCyT) and National Scientific and Technical  
416 Research Council of Argentina (CONICET).

417

ACCEPTED MANUSCRIPT

418 **References**

419

420 Allen L., Benoist B., Dary O., & Hurrell R. (2006). Guidelines on food fortification with  
421 micronutrients. World Health Organization and Food and Agriculture Organization of  
422 the United Nations. Retrieved from [http://www.who.int/nutrition/publication/guide\\_food\\_fortification\\_micronutrients.pdf](http://www.who.int/nutrition/publication/guide_food_fortification_micronutrients.pdf). (Last access March 10, 2014).

424

425 Alzamora, S.M., Guerrero, S., Nieto, A.B., & Vidales, S. (2003). Combined  
426 Preservation Technologies for Fruits and Vegetables: Training Manual. In Mejía, D.J  
427 ed. The Agricultural and Food Engineering Technologies Service, The Agricultural  
428 Support Systems Division, FAO. Rome, Italy. Retrieved from [http://www.fao.org/inpho\\_archive/content/documents/vlibrary/ac303e/ac303e00.html](http://www.fao.org/inpho_archive/content/documents/vlibrary/ac303e/ac303e00.html) (Last access March  
429 10, 2014)

430

431  
432 Argentine Food Code (2012). *Dietary or regimen foods*. Chapter XVII. Autonomous City  
433 of Buenos Aires (CABA), Argentina. Retrieved from  
434 [http://www.anmat.gov.ar/alimentos/codigoa/CAPITULO\\_XVII.pdf](http://www.anmat.gov.ar/alimentos/codigoa/CAPITULO_XVII.pdf)  
435 (Last access March 10, 2014).

436

437 Ashwell, M. (2004). *Conceptos sobre los alimentos funcionales*. (Concepts of functional  
438 foods. Washington DC. International Life Science Institute (ILSI) Press.

439

440 Barrera, C., Betoret, N., & Fito, P. (2004).  $Ca^{2+}$  and  $Fe^{2+}$  influence on the osmotic  
441 dehydration kinetics of apple slices (var. Granny Smith). *Journal of Food Engineering*,  
442 65, 9-14.

443

- 444 Binaghi, M.J., Greco, C.B., López, L.B., Ronayne, P.A., & Valencia, M.E. (2005).  
445 Influencia de distintos componentes de la dieta sobre la biodisponibilidad potencial de  
446 hierro (Influence of different dietary components on iron bioavailability potential). In  
447 *Ciencia Actual (Ed.). Actas del X Congreso de Ciencia y Tecnología de Alimentos*  
448 *Proceedings of the X Congress of Food Science and Technology*, Mar del Plata,  
449 Argentina. ISBN: 987-22165-1-7, 2006.
- 450
- 451 Boccio, J. & Monteiro, J.B. (2004). Fortificación de alimentos como estrategia para  
452 prevenir la deficiencia de hierro y zinc. Ventajas y desventajas desde el punto de vista  
453 nutricional y tecnológico. (Food fortification as a strategy to prevent iron deficiency and  
454 zinc. Advantages and disadvantages from the nutritional and technological point of  
455 view). *Revista de Nutrição*, 17 (1), 71-78
- 456
- 457 Campos, C.A., Gerschenson, L.N., & Flores, S.K. (2011). Development of edible films  
458 and coatings with antimicrobial activity. *Food and Bioprocess Technology*, 4, 849–875.
- 459
- 460 Cuppett S.L. (1994). Edible coatings as carriers of food additives, fungicides and  
461 natural antagonists. In Krochta J.M., Baldwin E. & Nisperos-Carriedo M.O, *Edible films*  
462 *and coatings to improve food quality* (pp. 121-137). Technomic Publishing Co. Inc.  
463 Lancaster, PA.
- 464
- 465 Lindsay R.C. (1996). Food Additives. In Fennema O., *Food Chemistry*, pp.767-823.  
466 Third Edition. Marcel Dekker Inc., NY, USA.
- 467
- 468 de Escalada Pla, M., Campos, C., & Gerchenson, L. (2009). Pumpkin (*Cucurbita*  
469 *moschata* Duchesne ex Poiret) mesocarp tissue as a food matrix for supplying iron in a  
470 food product. *Journal of Food Engineering*, 92, 361-369.

471

472 de Escalada Pla, M., Delbon, M., Rojas, A.M., & Gerschenson, L.N. (2006). Effect of  
473 immersion and turgor pressure change on mechanical properties of pumpkin (*Cucumis*  
474 *moschata*, Duch). *Journal of the Science of Food and Agriculture*, 86, (15), 2628-2637.

475

476 de Escalada Pla, M., Ponce, N., Stortz, C., Gerschenson, L., & Rojas, A.M. (2007).  
477 Composition and functional properties of enriched fiber products obtained from  
478 pumpkin (*Cucurbita moschata* Duchesne ex Poiret). *LWT Food Science and*  
479 *Technology*, 40 (7), 1176-1185.

480

481 de Escalada Pla, M., Ponce, N., Wider, M., Stortz, C., Rojas, A.M., & Gerschenson,  
482 L.N. (2005). Chemical and biochemical changes of pumpkin (*Cucumis moschata*,  
483 Duch) tissue in relation to osmotic stress. *Journal of the Science of Food and*  
484 *Agriculture*, 85, (11), 1852-1860.

485

486 De'Nobili M.D., Curto L.M, Delfino J.M., Soria M., Fissore E.N., Rojas A.M. (2013).  
487 Performance of alginate films for retention of l-(+)-ascorbic acid. *International Journal*  
488 *of Pharmaceutics*, 450: 95 -103.

489

490 Flores, S. (2006). *Estudios básicos y aplicados tendientes al desarrollo de películas*  
491 *comestibles que sean soporte del antimicrobiano sorbato de potasio*. Basic and applied  
492 studies for the development of edible films acting as carriers of the antimicrobial  
493 Potassium sorbate. Doctoral thesis, Facultad de Ciencias Exactas y Naturales,  
494 Universidad de Buenos Aires, Argentina.

495

496 Fontana A. (2008). Appendix D: Minimum Water Activity Limits for Growth of  
497 Microorganisms. In "Water Activity in Foods: Fundamentals and Applications". Edited

- 498 by Barbosa-Cánovas G., Fontana A., Schmidt S., & Labuza T. Blackwell Publishing  
499 and the Institute of Food Technologists. Retrieved from:  
500 <http://onlinelibrary.wiley.com/book/10.1002/9780470376454>. (Last access March, 10,  
501 2014).
- 502
- 503 García, M., Bifani, V., Campos, C. A., Martino, M. N., Sobral, P., Flores, S. K.,  
504 C. Ferrero, N. Bertola, N. E. Zaritzky, L. Gerschenson, C. Ramírez, A. Silva, M., &  
505 Menegalli, F. (2008). Edible coating as an oil barrier or active system. In Gutiérrez  
506 Lopez, Barbosa-Cánovas, Welti-Chanes, & Parada Arias (Eds.), *Food Engineering:  
507 Integrated Approaches*, pp. 225-241. NY, Springer.
- 508
- 509 Gaucheron, F. (2000). Iron fortification in dairy industry. *Trends in Food Science &  
510 Technology*, 11, 403-409.
- 511
- 512 González, E., Montenegro, M.A., Nazareno, M.A., & López de Mishima, B.A. (2001).  
513 Carotenoid composition and vitamin A value of an Argentinian squash (*Cucurbita  
514 moschata*). *Archivos Latinoamericanos de Nutrición*, 51(4), 395-399.
- 515
- 516 Gras M.L., Vidal D., Betoret N., Chiralt A., & Fito P. (2003). Calcium fortification of  
517 vegetables by vacuum impregnation. Interaction with cellular matrix. *Journal of Food  
518 Engineering*, 56: 279-284.
- 519
- 520 Gwanama, C., Botha, A. M, & Labuschagne, M.T. (2008). Genetic effects and heterosis  
521 of flowering and fruit characteristics of tropical pumpkin. *Plant Breeding*, 120(3): 271-  
522 272.
- 523
- 524 Han, J.H. (2000). Antimicrobial food packaging. *Food Technology*, 54 (3), 56-65.

525

526 Lago-Vanzela, E.S., do Nascimento, P., Fontes, E.A.F., Mauro, M.A., & Kimura, M.  
527 (2013). Edible coatings from native and modified starches retain carotenoids in  
528 pumpkin during drying. *LWT Food Science and Technology*, 50, 420-425.

529

530 Laxmi Narayan, R., Mills, A., & Berman, J. (2006). Advancement of global health: key  
531 messages from the Disease Control Priorities Project. *The Lancet*, 367, 1193-1208.

532

533 León, P.G., & Rojas, A.M. (2007). Gellan gum films as carriers of L-(+)-ascorbic acid.  
534 *Food Research International*, 40: 565 - 575.

535

536 Moreno J., Simpson R., Baeza A., Morales J., Muñoz C., Sastry S., & Almonacid S.  
537 (2012). Effect of ohmic heating and vacuum impregnation on the osmodehydration  
538 kinetics and microstructure of strawberries (cv. Camarosa). *LWT - Food Science and  
539 Technology*, 45: 148 -154.

540

541 Olivera D., Viña S., Marani C., Ferreyra R., Mugridge A., Chaves A., & Mascheroni R.  
542 (2008). Effect of blanching on the quality of Brussels sprouts (*Brassica oleracea* L.  
543 *Gemmifera* DC) after frozen storage. *Journal of Food Engineering*, 84, 148-155.

544

545 Pérez C., De'Nobili M., Rizzo S., Gerschenson L., Descalzo A, & Rojas A. (2013). High  
546 methoxyl pectin–methyl cellulose films with antioxidant activity at a functional food  
547 interface. *Journal of Food Engineering*, 116: 162 – 169.

548

549 Rao M.V., & Kawamura Y. (2008). Ferrous ammonium phosphate. Chemical and  
550 Technical Assessment (CTA). Retrieved from <http://www.fao.org/fileadmin/templates>



551 /agns/pdf/jecfa/cta/71/ferrous\_ammonium\_phosphate.pdf. (Last access March 10,  
552 2014).

553

554 Rojas, A.M.L. (1995). Destrucción de vitamina C en sistemas modelo de actividad  
555 acuosa reducida. (Destruction of vitamin C in model aqueous systems with reduced  
556 activity). Doctoral thesis, Facultad de Ciencias Exactas y Naturales. Universidad de  
557 Buenos Aires, Argentina.

558

559 Rojas, A.M., & Gerschenson, L.N. (2001). Ascorbic acid destruction in aqueous model  
560 systems: an additional discussion. *Journal of the Science of Food and Agriculture*, 81  
561 (15): 1433 - 1439.

562

563 Rosenthal, I., Rosen, B. & Bernstein, S. (1993). Effects of Milk Fortification with  
564 Ascorbic Acid and Iron. *Milchwissenschaft*, 48: 676 - 679.

565

566 Souto de Olivera, S.M. (2009). Avaliação do impacto do uso do fruto Mangifera indica  
567 L. (manga) no tratamento de crianças anêmicas fazendo uso do sulfato ferroso.  
568 Doctoral thesis, Centro de Ciências da Saúde, Universidade Federal da Paraíba,  
569 Brazil. Retrieved from [http://bdtd.biblioteca.ufpb.br/tde\\_busca/arquivo.php?codArquivo=961](http://bdtd.biblioteca.ufpb.br/tde_busca/arquivo.php?codArquivo=961). (Last access March 10, 2014).

571

572 Spiazia E., & Mascheroni R. (1997). Mass transfer model for osmotic dehydration of  
573 fruits and vegetables. Development of the simulation model. *Journal of Food*  
574 *Engineering*, 34 (4): 387 – 410.

575

576 Tripathi, B., & Platel, K. (2013). Feasibility in fortification of sorghum (*Sorghum bicolor*  
577 *L. Moench*) and pearl millet (*Pennisetum glaucum*) flour with iron. *LWT Food Science*  
578 *and Technology*, 50 (1), 220-225.

579

580 World Health Organization (WHO) (2013). Micronutrient deficiencies: iron deficiency  
581 anaemia. Retrieved from <http://www.who.int/nutrition/topics/ida/en/index.html>. (Last  
582 access March 10, 2014).

583

584 Zhao Y., & Xie J. (2004). Practical applications of vacuum impregnation in fruit and  
585 vegetable processing. *Trends in Food Science & Technology*, 15, 434-451.

586

587 Zimmermann, M., & Hurrell, R. (2007). Nutritional iron deficiency. *The Lancet*, 370,  
588 511-520.

589

590 Captions to figures.

591

592 **Figure 1.** Pumpkin fortification with iron (Fe) and ascorbic acid (AA): Response surface  
593 for variables of dry infusion process a) Solid Gain (SG), b) Water Loss (WL), c) Weight  
594 changes during the air drying process (PP) and d) color changes ( $\Delta E$ ) respect to  
595 control system (without fortification). The best fitted second degree polynomials are:

596  $SG = 80.45 Fe + 0.885 AA + 0.0463 AA^2 - 8.3802 Fe AA$  ( $R^2$ : 0.9843, F: 110)

597  $WL = 307.57 Fe + 6.26 AA - 27.55 Fe AA$  ( $R^2$ : 0.9908, F: 286)

598  $PP = 161.839 Fe + 1.474 AA - 179.523 Fe^2 - 6.746 Fe AA$  ( $R^2$ : 0.9960, F: 433)

599  $\Delta E = 79.438 Fe + 1.267 AA - 181.002 Fe^2 - 0.041 AA^2$  ( $R^2$ : 0.9848, F: 113)

600 Coefficients with significant effect are shown,  $R^2$ : determination coefficient, F: Fisher's  
601 test value.

602

603 **Figure 2.** Pumpkin fortified with iron an ascorbic acid by dry infusion, after air drying.  
604 Numbers corresponds to systems from central composite design. Control system (C),  
605 without fortification, is also included.

606

607 **Figure 3.** Pumpkin fortified with iron (Fe) and ascorbic acid (AA) by dry infusion and air  
608 dried: surface response and the best fitted second degree polynomial for Chroma =

609  $(a^{*2} + b^{*2})^{(1/2)}$ :

610  $Chroma = 40.26 - 1.663 AA - 105.59 Fe + 0.0605 AA^2 + 237.74 Fe^2$

611 ( $R^2=0.755$ , lack of fit p = 0,119).

612

613

614

615

616

617 Captions for Tables.

618

619 **Table 1.** Treatments performed according central composite design for optimization of  
620 pumpkin fortification with iron (Fe) and ascorbic acid (AA). The control system (C) is  
621 also included.

622

623 **Table 2:** Pumpkin fortified with iron (Fe) and ascorbic acid (AA): measured values of  
624 water loss (WL), solid gain (SG), pH after dry infusion process and weight changes  
625 during the air drying (PP).

626

627 **Table 3.** Color difference ( $\Delta E$ ) of pumpkin fortified with iron (Fe) and ascorbic acid (AA)  
628 respect to control system (C) and color parameters: lightness ( $L^*$ ) and chroma before  
629 and after air drying process.

630

631 **Table 4.** Chroma and lightness ( $L^*$ ) parameters of fortified pumpkin with iron and  
632 ascorbic acid, coated and uncoated. Difference of color ( $\Delta E$ ) respect to impregnated  
633 pumpkin before coating and drying.

634

635

636

637

638

1  
2  
3  
4  
5  
6

**Table 1.** Treatments performed according central composite design for optimization of pumpkin fortification with iron (Fe) and ascorbic acid (AA). The control system (C) is also included.

System	Coded		Uncoded	
	Fe <sup>1</sup>	AA <sup>1</sup>	Fe <sup>2</sup>	AA <sup>2</sup>
1	1	1	0.288	15.2
2	1	-1	0.288	5.6
3	-1	1	0.144	15.2
4	-1	-1	0.144	5.6
5	0	0	0.216	10.4
6	0	-2	0.216	0.8
7	0	2	0.216	20
8	-2	0	0.072	10.4
9	2	0	0.360	10.4
10	0	0	0.216	10.4
11	0	0	0.216	10.4
C	NA	NA	NA	NA

<sup>1</sup>Coded levels for Fe and AA

<sup>2</sup>Real values for Fe and AA (g/kg pumpkin)

NA: not added

7  
8  
9  
10  
11  
12

1  
 2 **Table 2:** Pumpkin fortified with iron (Fe) and ascorbic acid (AA): measured values of  
 3 water loss (WL), solid gain (SG), pH after dry infusion process and weight changes  
 4 during the air drying (PP).

5

System	Fe <sup>1</sup>	AA <sup>1</sup>	WL <sup>2</sup>	SG	pH	PP
1	0.288	15.2	72.6±0.1 <sup>a</sup>	8.77±0.05 <sup>a</sup>	3.415±0.007 <sup>a</sup>	25.6±0.1 <sup>a</sup>
2	0.288	5.6	69.0±0.4 <sup>b</sup>	18.2±0.1 <sup>b</sup>	3.760±0.001 <sup>b</sup>	26.6±0.2 <sup>b,c</sup>
3	0.144	15.2	69.3±0.1 <sup>b,c</sup>	14.78±0.06 <sup>c</sup>	3.445±0.007 <sup>a</sup>	26.6±0.2 <sup>b,c</sup>
4	0.144	5.6	69.8±0.1 <sup>b,c</sup>	14.18±0.05 <sup>d,f</sup>	3.885±0.007 <sup>e</sup>	26.3±0.2 <sup>b,c</sup>
5	0.216	10.4	69.9±0.4 <sup>b,c,e</sup>	13.1±0.1 <sup>e</sup>	3.61±0.01 <sup>c</sup>	26.9±0.3 <sup>b</sup>
6	0.216	0.8	69.7±0.3 <sup>b,c</sup>	14.45±0.09 <sup>f</sup>	4.295±0.007 <sup>f</sup>	25.7±0.1 <sup>a</sup>
7	0.216	20	70.2±0.4 <sup>c,d,e</sup>	19.7±0.2 <sup>g</sup>	3.35±0.01 <sup>g</sup>	25.2±0.1 <sup>a</sup>
8	0.072	10.4	71.0±0.1 <sup>d,e,f</sup>	14.06±0.08 <sup>d</sup>	3.73±0.04 <sup>b</sup>	21.9±0.1 <sup>d</sup>
9	0.360	10.4	69.2±0.6 <sup>b</sup>	11.8±0.1 <sup>h</sup>	3.55±0.04 <sup>d</sup>	26.5±0.2 <sup>b,c</sup>
10	0.216	10.4	70.8±0.2 <sup>e,f</sup>	11.86±0.07 <sup>h</sup>	3.59±0.02 <sup>c,d</sup>	26.8±0.2 <sup>b,c</sup>
11	0.216	10.4	71.5±0.3 <sup>f</sup>	12.39±0.08 <sup>i</sup>	3.57±0.01 <sup>c,d</sup>	26.3±0.2 <sup>b,c</sup>
C	NA	NA	64.1±0.2	13.1±0.1 <sup>e</sup>	4.59±0.02	35.6±0.2

6 <sup>1</sup> Contents of Fe and AA (g/kg pumpkin).7 <sup>2</sup> Absolute values were reported.8 Mean and standard deviation ( $n = 3$ ) are reported.9 Different letters in the same column indicate significant differences ( $p < 0.05$ ).

10

11

12

1  
2

3 **Table 3.** Color difference ( $\Delta E$ ) of pumpkin fortified with iron (Fe) and ascorbic acid (AA)  
4 respect to control system (C) and color parameters: lightness ( $L^*$ ) and chroma before  
5 and after air drying process.

System	Fe <sup>1</sup>	AA <sup>1</sup>	$\Delta E$	$L^*_{After}$	$L^*_{Before}$	Chroma <sub>After</sub>	Chroma <sub>Before</sub>
1	0.288	15.2	15.5±2.0 <sup>a</sup>	32.9±0.8 <sup>a</sup>	36.2±0.7	21±2 <sup>a</sup>	27.0±0.4
2	0.288	5.6	9.2±4.1 <sup>b</sup>	38±3 <sup>b,A</sup>	38.91±0.04 <sup>A</sup>	25±3 <sup>a,b</sup>	30.2±0.5
3	0.144	15.2	17.4±0.2 <sup>a</sup>	33.3±0.3 <sup>a,c</sup>	36.3±0.4	17.72±0.06 <sup>a</sup>	27±1
4	0.144	5.6	14.4±0.2 <sup>a,b</sup>	34.6±0.8 <sup>a,c,B</sup>	36.5±0.4 <sup>B</sup>	21.0±0.8 <sup>a</sup>	28±2
5	0.216	10.4	18.5±0.3 <sup>a</sup>	31.7±0.8 <sup>a</sup>	38.8±0.3	17.7±0.5 <sup>a</sup>	30.9±0.8
6	0.216	0.8	10.5±0.6 <sup>b</sup>	36.61±0.02 <sup>b,c,C</sup>	36.53±0.03 <sup>C</sup>	27±3 <sup>b,Z</sup>	26±1 <sup>Z</sup>
7	0.216	20	18.20±0.05 <sup>a</sup>	31±1 <sup>a</sup>	38.3±0.7	19±2 <sup>a</sup>	32±4
8	0.072	10.4	12.0±0.1 <sup>a,b</sup>	35±1 <sup>a,b</sup>	37.3±0.1	24.8±0.8 <sup>a,b,Y</sup>	28.50±0.05 <sup>Y</sup>
9	0.360	10.4	16.5±0.7 <sup>a</sup>	33±1 <sup>a</sup>	36.03±0.06	19.91±0.07 <sup>a</sup>	28±3
10	0.216	10.4	19.6±0.5 <sup>a</sup>	31.26±0.08 <sup>a</sup>	38±1	16.2±0.6 <sup>a</sup>	30±6
11	0.216	10.4	17.9±1.2 <sup>a</sup>	32.6±0.2 <sup>a</sup>	41.9±0.6	18±2 <sup>a</sup>	39.0±0.2
C	0	0	NA	43.6±0.9 <sup>D</sup>	42.4±0.1 <sup>D</sup>	31.1±0.8 <sup>X</sup>	32±2 <sup>X</sup>

6 <sup>1</sup> Contents of Fe and AA (g/kg pumpkin)

7 NA: not applicable.

8  $L^*_{After}$  and Chroma<sub>After</sub> correspond to lightness and chroma after air drying.9  $L^*_{Before}$  and Chroma<sub>Before</sub>, correspond to lightness and chroma before air drying.

10 Mean and standard deviation (n = 3) are reported.

11 Same letters within a column indicate non significant differences among systems (p&lt;0.05).

12 Same capital letters within file indicate non significant differences due to air drying process for a  
13 same system (p<0.05).14  
15 .


1  
 2 **Table 4.** Chroma and lightness ( $L^*$ ) parameters of fortified pumpkin with iron and  
 3 ascorbic acid, coated and uncoated. Difference of color ( $\Delta E$ ) respect to impregnated  
 4 pumpkin before coating and drying.

	Before	After drying			
	drying	Uncoated		Coated	
	(Ref)	$d_0$	$d_9$	$d_0$	$d_9$
<b>Chroma</b>	35.1±0.8 <sup>a</sup>	29.6±0.6 <sup>b***</sup>	28.09±0.07 <sup>b***</sup>	34 ±2 <sup>a</sup>	33.5±0.3 <sup>a</sup>
<b>L*</b>	40.5±0.5 <sup>k</sup>	34.20±0.08 <sup>m***</sup>	37.0±0.4 <sup>l</sup>	36.2±0.9 <sup>l</sup>	40.2±0.9 <sup>k</sup>
<b><math>\Delta E</math></b>	NA	9.05±0.07	7.8±0.2	1.5±0.4 <sup>z</sup>	1.7±0.2 <sup>z</sup>

6 Mean and standard deviation (n = 3) are reported.  
 7 Same letters within a file indicate non significant differences (p<0.05; \*\*\* p<0,001).  
 8 Ref: impregnated system before coating and drying.  
 9  $d_0$ : system at beginning of storage  
 10  $d_9$ : system after 9 days of storage

11  
 12





**Figure 1.** Pumpkin fortification with iron (Fe) and ascorbic acid (AA): Response surface for variables of dry infusion process a) Solid Gain (SG), b) Water Loss (WL), c) Weight changes during the air drying process (PP) and d) color changes ( $\Delta E$ ) respect to control system (without fortification). The best fitted second degree polynomials are:

$$SG = 80.45 Fe + 0.885 AA + 0.0463 AA^2 - 8.3802 Fe AA \quad (R^2: 0.9843, F: 110)$$

$$WL = 307.57 Fe + 6.26 AA - 27.55 Fe AA \quad (R^2: 0.9908, F: 286)$$


$$PP = 161.839 Fe + 1.474 AA - 179.523 Fe^2 - 6.746 Fe AA \quad (R^2: 0.9960, F: 433)$$

$$\Delta E = 79.438 Fe + 1.267 AA - 181.002 Fe^2 - 0.041 AA^2 \quad (R^2: 0.9848, F: 113)$$

Coefficients with significant effect are shown,  $R^2$ : determination coefficient, F: Fisher's test value.



**Figure 2.** Pumpkin fortified with iron and ascorbic acid by dry infusion, after air drying. Numbers correspond to systems from central composite design. Control system (C), without fortification, is also included.



**Figure 3.** Pumpkin fortified with iron (Fe) and ascorbic acid (AA) by dry infusion and air dried: surface response and the best fitted second degree polynomial for Chroma =  $(a^{*2} + b^{*2})^{(1/2)}$ :

$$\text{Chroma} = 40.26 - 1.663 \text{ AA} - 105.59 \text{ Fe} + 0.0605 \text{ AA}^2 + 237.74 \text{ Fe}^2$$

( $R^2=0.755$ , lack of fit  $p = 0,119$ ).

## HIGHLIGHTS

- Pumpkin fortified with iron and ascorbic acid was developed.
- Dry infusion previous to air drying process allowed fortification of pumpkin.
- Iron / ascorbic acid ratio that minimize pumpkin color changes was determined.
- Edible coating based on tapioca starch protects pumpkin from color change.