

Accepted Manuscript

Antimicrobial activity of *de novo* designed cationic peptides against multi-resistant clinical isolates

Facone Diego, Veliz Omar, Corso Alejandra, Noguera Martin, Martínez Melina, Payes Cristian, Semorile Liliana, Maffia Paulo Cesar, PhD



PII: S0223-5234(13)00716-2

DOI: [10.1016/j.ejmech.2013.10.065](https://doi.org/10.1016/j.ejmech.2013.10.065)

Reference: EJMECH 6528

To appear in: *European Journal of Medicinal Chemistry*


Received Date: 23 August 2013

Revised Date: 24 October 2013

Accepted Date: 25 October 2013

Please cite this article as: F. Diego, V. Omar, C. Alejandra, N. Martin, M. Melina, P. Cristian, S. Liliana, M.P. Cesar, Antimicrobial activity of *de novo* designed cationic peptides against multi-resistant clinical isolates, *European Journal of Medicinal Chemistry* (2013), doi: 10.1016/j.ejmech.2013.10.065.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Helical wheel projection of peptide 2

Hemolytic activity

Antibacterial activity

**Structural
characterization by
circular dichroism**

De novo designed alpha helical cationic peptides with antimicrobial activity against multi-resistant clinical isolates

1 Antimicrobial activity of *de novo* designed cationic peptides
2 against multi-resistant clinical isolates.

3 Faccone Diego⁽¹⁾, Veliz Omar⁽¹⁾, Corso Alejandra⁽¹⁾, Noguera Martin⁽³⁾, Martínez Melina⁽²⁾, Payes
4 Cristian^(2,3), Semorile Liliana⁽²⁾, Maffia Paulo Cesar^(2,3).

5 (1) Servicio Antimicrobianos, Instituto Nacional de Enfermedades Infecciosas (INEI)-ANLIS "Dr.
6 Carlos G. Malbran", Buenos Aires, Argentina

7 (2) Laboratorio de Microbiología Molecular, Universidad Nacional de Quilmes, Buenos Aires,
8 Argentina.

9 (3) Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

10

11 Corresponding author: PhD. Maffia, Paulo Cesar

12 E mail: paulo.maffia@unq.edu.ar

13

14 **Abstract.**

15 Antibiotic resistance is one of the main problems concerning public health or clinical practice.
16 Antimicrobial peptides appear as good candidates for the development of new therapeutic
17 drugs. In this study we *de novo* designed a group of cationic antimicrobial peptides, analyzed
18 its physicochemical properties, including its structure by circular dichroism and studied its
19 antimicrobial properties against a panel of clinical isolates expressing different mechanisms of
20 resistance. Three cationic alpha helical peptides exhibited antimicrobial activity comparable to,
21 or even better than the comparator omiganan (MBI-226).

22

23 **Keywords.**

24 Peptides; cationic; antimicrobial; resistance

25 Introduction

26 Antimicrobial peptides (AMPs) are naturally occurring molecules of the innate immune system
27 that play an important role in the host defense of animals and plants [1]. In recent years,
28 natural or designed AMPs have attracted considerable interest as potential candidates for the
29 development of novel antibiotics [2-3]. The main reason for this interest is that its particular
30 mechanism of action is unlikely to induce drug resistance, in part because resistance against
31 AMPs cannot be selected without bacterial cell wall undergoing profound structural changes
32 [4]. However, pathogens can eventually respond to AMPs reducing the negative charge of their
33 cell envelope with specific surface modifications and subvert mechanisms of AMPs [5].
34 Bacteria are capable of adapting and resisting AMPs, through the production of peptidases
35 and proteases that degrade antimicrobial peptides, and the production of compounds that
36 inhibit the action of AMPs [6]

37 The broad activity spectrum and the relative selectivity towards microbial membranes are also
38 two important features that drive the interest of researchers on AMPs as new antibiotic
39 molecules.

40 The cationic AMP omiganan (MBI-226), an analogue of indolicidin, is one of the most studied
41 AMPs and it has recently finished Phase II trials (ClinicalTrials.gov identifier: NCT00608959).
42 Omiganan showed activity against gram-positive and gram-negative bacteria but also *Candida*
43 spp. isolates [7-8]. Therefore, the objectives of this work were to design a group of new
44 peptide sequences, and analyze their physicochemical properties and antimicrobial activities
45 against 82 bacterial strains, including wild type and drug resistant clinical isolates. Omiganan
46 was used in this study as comparator for these peptides.

47

48 Materials and methods

49 *Peptides design and synthesis*

50 The sequences were designed using a combined rational and computer assisted approach.
51 Cationic alpha helical peptides were designed identifying short putative active regions from
52 AMP databases. Then, these regions were combined or modified in order to have cationic
53 sequences with different physicochemical parameters, like alpha helix content and
54 hydrophobicity. For this purpose we used multiple alignment tools and simulators of
55 physicochemical properties like ClustalX, HeliQuest [9] and HydroMCalc [10]. We established
56 specific amino acid positions and identified functionally relevant motifs in natural or designed

57 peptides. Considering all these diverse parameters, a group of peptides was synthesized with
58 or without C terminus amidation. The purity grade of all peptides was >95% by HPLC
59 (GenScript Co., Piscataway, NJ 08854, USA). The peptide sequences: Peptide 1:
60 WPKWWKWKRRWGRKKAKKRRG; peptide2: GLLKKWLKKWKEFKRIVGY; peptide3:
61 FGKEKKAWWRRRKWLK; peptide5: RIVQRIKKWLLKWKKLGY.

62

63 *Bacterial Strains*

64 The panel analyzed included 82 previously well characterized isolates collected at the National
65 Reference Laboratory (INEL) with different mechanisms of resistance: 39 Gram-positive (*vanA*,
66 *vanB*, *vanC*, *mecA*, *ermA*, *ermC*, *msrA*, *InuA* genes) and 43 Gram-negative bacteria (*bla*_{VIM},
67 *bla*_{IMP}, *bla*_{SPM}, *bla*_{KPC-2}, *bla*_{OXA-23}, *bla*_{OXA-58}, *bla*_{CTX-M-2}, *bla*_{PER-2}, *bla*_{GES}, *bla*_{VEB-1}, *bla*_{TEM-1}, *bla*_{CMY}, *bla*_{CIT},
68 *bla*_{SHV-1}, *bla*_{OXA-9}). The panel includes *P. aeruginosa* ATCC27853, *E. coli* ATCC25922, *S. aureus*
69 ATCC29213, *E. faecalis* ATCC51299 and *E. faecalis* ATCC29212 reference strains.

70 *Antimicrobial activity*

71 Minimal inhibitory concentration was determined by standard microdilution assay according to
72 CLSI recommendations [11], using Mueller Hinton Broth (DIFCO) supplemented with Ca²⁺ (20-
73 25mg/L) and Mg²⁺ (10-12.5mg/L). Omiganan[®] was used as comparator.

74 *Hemolytic assay*

75 The cytotoxic activity of the peptides was evaluated according to the method described
76 previously [12]. Briefly, a volume of heparinized human whole blood was diluted 3x in
77 phosphate-buffered saline and then centrifuged 10 min. at 1500 rpm. This procedure was
78 repeated three times. The cellular pellet was resuspended in phosphate-buffered saline to a
79 final dilution of 10% (v/v). The stock cell suspension was further diluted to about 0.5% (v/v).
80 Peptides were then added at different concentrations and incubated at 37 °C for 30 min.
81 Afterwards, tubes were centrifuged and the absorbance of the supernatant was measured at
82 550 nm. The percentage of lysis was then calculated relative to 0% lysis with buffer and 100%
83 lysis with water. The absorbance measurement was repeated three times, and the averaged
84 values were used.

85 *Circular dichroism in the far UV*

86 We studied the secondary structure content by circular dichroism spectroscopy in the far UV,
87 using a JASCO J computer 810 (Jasco Corp., Tokyo, Japan) acid calibrated with (+) 10

88 camphorsulfonic acid. The measurements were performed under nitrogen gas flow of 8 l/h at
89 a temperature of 20 °C, controlled by a Peltier system (JASCO).

90 Spectra were recorded between 185 and 320 nm, using a 0.1 cm cell path length. The peptide
91 concentrations were 40 µM, dissolved in sodium phosphate buffer pH 7.0 or 10 mM in the
92 same buffer with sodium dodecyl sulphate (SDS) 10 mM. The sensitivity was 100 millidegrees.
93 We used a scan speed of 50 nm/min, a response time of 1 s and a bandwidth of 1 nm. We
94 performed an average of five assays for each sample spectra. The average absorption was
95 corrected by buffer and then baseline to zero using the average of readings between 290 and
96 320 nm. Finally, the data were smoothed using a Golay polynomial Savitzky fourth grade, with a
97 window of ten points. The spectra were converted to molar ellipticity residue half by using the
98 relationship: $[\theta] = \theta / (10 \times c \times n \times d)$, where $[\theta]$ is the molar ellipticity (in degrees \times cm² \times
99 dmol⁻¹), θ the ellipticity in millidegrees, n is the number of residues of the peptide and c its
100 molar concentration, d the length of the cell in centimeters.

101 The mean hydrophobicity (H) and the mean hydrophobic moment (μH) were calculated from
102 the amino acid sequences, using the Eisenberg scale for hydrophobicity by the HydroMCalc
103 applet [10]

104

105 **Results.**

106 *Structural analysis of the peptides.* The circular dichroism spectra of peptides in aqueous
107 solution shows that they are all unstructured in aqueous buffer, with a characteristic minimum
108 at approximately 200 nm (Figure 1A). With the addition of SDS micelles (Figure 1B),
109 conformational changes occurred in peptides 2 and 5 that are consistent with the formation of
110 alpha-helix structure with two characteristic minima near 208 and 222 nm. Peptide 1 also
111 underwent such a transition, although the acquired structure level was lower than the one
112 seen for peptides 2 and 5. The circular dichroism spectrum of peptide 3 is almost invariable
113 with the addition of SDS micelles, indicative of the persistence of a disordered conformation.
114 For omiganan, the spectrum was significantly modified in the presence of SDS, the 200 nm
115 band was attenuated and a new band near 230 nm appeared; which could be the result of the
116 interaction between the side chains of tryptophan. Figure 1C shows the helical wheel
117 projection of the peptides, depicting the amphipatic residues and their relative position in the
118 alpha helix.

119 Peptides were designed in order to have different alpha helical content and different
120 amphipathicity, the latter calculated as the hydrophobicity and mean hydrophobic moment
121 with specific software (HydroMCalc and Heliquest). Helical conformation was monitored in SDS
122 micelles, which are generally employed as a simple membrane-mimetic environment. Table 1
123 summarizes the structural analyses and hemolytic activity of the peptides.

124

125 *Hemolytic activity.* The peptides (C-terminus amidated) were incubated with human red blood
126 cells in order to evaluate their hemolytic activity. Table 1 shows the results as a relative value
127 to 100% hemolysis of human red blood cells treated with distilled water. Peptides 1, 3, 4 and 5
128 showed little or negligible hemolytic activity, similarly with omiganan. Peptide 2 displayed an
129 hemolysis of red blood cells almost 4-times higher than omiganan and peptide 5.

130

131 *Antimicrobial activity of the peptides.* Antimicrobial activity of C-terminus amidated and non
132 amidated peptides was evaluated by microdilution test against a first panel with 8 isolates. The
133 panel included 5 clinical (*S. warnerii* M6823, *S. cohnii* M6767, *S. aureus* M6794, *P. aeruginosa*
134 M13513 and *K. pneumoniae* M13540) and 3 ATCC isolates (*S. aureus* ATCC29213, *P. aeruginosa*
135 ATCC27853 and *E. coli* ATCC25922). MICs values of those peptides with amidated C-terminus
136 were equal or lower (up to 3 dilutions) than those peptides with non amidated C-terminus, for
137 the 8 isolates tested (data not shown). Peptide 3, with the lowest hydrophobic moment and
138 helicity, did not show significant antimicrobial activity, except for coagulase negative
139 staphylococci (MIC of 8 and 4 mg/L, respectively). Peptide 4 showed no antimicrobial activity
140 for all the eight isolates tested. On the other hand peptides 1, 2 and 5 showed antimicrobial
141 activity comparable to, or in some cases better than, omiganan.

142 Considering these results, together with the lower hemolytic activity of C-terminus amidated
143 peptides, the antimicrobial activity of the C-terminus amidated peptides 1, 2 and 5 was
144 evaluated against a large panel of 82 well-characterized bacterial isolates, including the 8
145 isolates used in the first panel. Table 2 displays MIC values of peptides 1, 2 and 5 and
146 omiganan against a panel containing 43 gram-negative and 39 gram-positive isolates. This
147 panel included isolates expressing clinically relevant resistance mechanisms to antibiotics, like
148 carbapenemase-producing enterobacteria and *P. aeruginosa*, methicillin-resistant *S. aureus* or
149 vancomycin-resistant enterococci (Table 2). Peptide 1 showed MIC₉₀ values of 128 mg/L for all
150 gram-negative isolates except for *K. pneumoniae* strains (MIC \geq 1024 mg/L). Peptides 2 and 5
151 showed similar performance against gram-negative bacteria with MIC₉₀ values between 32 and

152 128 mg/L, and slightly lower than peptide 1. Peptides 1, 2 and 5 showed a similar activity for
153 each gram-positive species (Table 2). *E. faecalis* isolates displayed higher MIC values than
154 other enterococci species for the three analyzed peptides and omiganan (Table 2). No
155 association between mechanism of resistance and MIC values was observed, similar results
156 were reported for omiganan by Sader *et al.* [7]

157 The omiganan MIC ranges obtained herein were slightly higher (up to three dilutions) than
158 previous reports [7-8]. Omiganan MIC values for ATCC control strains were into the range
159 described by Anderegg *et al.* [13], but on the upper border (Table 2). We suspect that the
160 difference of our results of MIC range for omiganan could be associated to: i) a smaller number
161 of isolates included in our panel, ii) our isolates collection could be strongly biased with
162 antimicrobial resistant strains, and/or iii) intrinsic differences of each population of isolates.

163

164 *Concluding remarks.* We designed a group of peptides with different physicochemical
165 characteristics, and tested their antimicrobial activity against a panel of clinical bacterial
166 isolates. At least seven structural or physical parameters could be considered critical for
167 biological activity: size, sequence, charge, degree of structuring (helicity), hydrophobicity,
168 amphipathicity and angles subtended by hydrophobic and hydrophilic faces of the formed
169 helix [14].

170 Some authors [15] argue that the secondary structure and biological activity are not coupled,
171 and AMPs do not form pores in membranes but rather destabilize them disturbing the
172 organization of the lipids, consistent with the idea that physical chemical and interfacial
173 properties are the critical factors for determining the biological activity; this theory would suit
174 omiganan that is not structured as alpha helix. In any case, helicity seems to be an important
175 parameter for antimicrobial activity in our peptides, since the three peptides that displayed
176 alpha helical content in SDS micelles also showed antibacterial activity. However, other
177 parameters may be involved, for example peptide 1 and 5, although having different helicity,
178 they showed similar antimicrobial activity against Gram-positive strains. But, on the other
179 hand, these two peptides showed different activity when tested on Gram-negative bacteria,
180 especially in *K. pneumoniae* specie (Table 2).

181 Peptide 2 and 5 had similar physicochemical properties, like alpha helix content,
182 amphipathicity and net charge, but also antimicrobial activity, however peptide 2 was highly
183 hemolytic to human red blood cells. Furthermore peptide 1 showed antimicrobial activity

184 against gram-positive and -negative strains, although it did not show high alpha helix content
185 in contact with SDS micelles.

186 Also interesting was the relative low activity of all these peptides against *E. faecalis* isolates
187 (64-256 mg/L), compared to another *Enterococci* species, like *E. faecium* and *E. gallinarum*.
188 This low activity was also observed for omiganan [7], indicating a possible different cell wall
189 composition in *E. faecalis* specie. It is evident that certain differences within the bacterial cell
190 wall are probably associated with these different sensitivities to AMPs.

191 Peptides 1, 2 and 5 showed good antibacterial activity against a broad spectrum of clinical
192 isolates, although peptide 2 displayed high cytotoxicity. These three peptides could become
193 good templates for topical use.

194

195 Acknowledgements

196 This work was supported by grants from ANPCYT FONCYT PICT Start Up 2008-2090 and PIP
197 845/10 from CONICET. Dr. L. Semorile is member of the Research Career of CIC-BA. Dr. P.
198 Maffía is member of the Research Career of CONICET.

199

200 **Figure 1.** Graphical analysis of the peptides structure.

201 **Footnote:** Circular dichroism of peptides in aqueous solution (panel A) and in SDS micelles (panel B). Helical wheel
202 projection diagrams of the peptides, considering the first 18 amino acids (panel C). omi, omiganan; p1, peptide 1;
203 p2, peptide 2; p3, peptide 3; p5, peptide 5.

204

205

206 **Table 1.** Physicochemical properties, structural analyses and hemolytic activity of the peptides

207 **Footnote:** The values for hydrophobicity (H) and mean hydrophobic moment (μH) were obtained from HydroMCalc
208 software. The percent helix values were determined based on circular dichroism spectra calculated as the mean
209 residue molar ellipticity at 222 nm, in SDS micelles. * Isoelectric point and net charge were calculated for the acidic
210 C terminus version of the peptides. Hemolytic activity is shown as a percentage (%) of hemolysis compared to
211 distilled water (100% hemolysis). One representative experiment. Ne: not evaluated

212

213 **Table 2.** Antimicrobial activity of three designed peptides and omiganan against gram-negative
214 and gram-positive bacteria.

215

216 **References**

- 217 [1] Y. Lai, R.L. Gallo, AMPed up immunity: how antimicrobial peptides have multiple roles in
218 immune defense, *Trends Immunol*, 30 (2009) 131-141.
- 219 [2] R.E. Hancock, H.G. Sahl, Antimicrobial and host-defense peptides as new anti-infective
220 therapeutic strategies, *Nat Biotechnol*, 24 (2006) 1551-1557.
- 221 [3] C.D. Fjell, J.A. Hiss, R.E. Hancock, G. Schneider, Designing antimicrobial peptides: form
222 follows function, *Nat Rev Drug Discov*, 11 (2012) 37-51.
- 223 [4] J. Wiesner, A. Vilcinskas, Antimicrobial peptides: the ancient arm of the human immune
224 system, *Virulence*, 1 (2010) 440-464.
- 225 [5] N.Y. Yount, M.R. Yeaman, Immunocontinuum: perspectives in antimicrobial peptide
226 mechanisms of action and resistance, *Protein Pept Lett*, 12 (2005) 49-67.
- 227 [6] A. Peschel, H.G. Sahl, The co-evolution of host cationic antimicrobial peptides and microbial
228 resistance, *Nat Rev Microbiol*, 4 (2006) 529-536.
- 229 [7] H.S. Sader, K.A. Fedler, R.P. Rennie, S. Stevens, R.N. Jones, Omiganan pentahydrochloride
230 (MBI 226), a topical 12-amino-acid cationic peptide: spectrum of antimicrobial activity and
231 measurements of bactericidal activity, *Antimicrob Agents Chemother*, 48 (2004) 3112-3118.
- 232 [8] T.R. Fritsche, P.R. Rhomberg, H.S. Sader, R.N. Jones, Antimicrobial activity of omiganan
233 pentahydrochloride tested against contemporary bacterial pathogens commonly responsible
234 for catheter-associated infections, *J Antimicrob Chemother*, 61 (2008) 1092-1098.
- 235 [9] R. Gautier, D. Douguet, B. Antony, G. Drin, HELIQUEST: a web server to screen sequences
236 with specific alpha-helical properties, *Bioinformatics*, 24 (2008) 2101-2102.
- 237 [10] <http://www.bbcm.univ.trieste.it/~tossi/HydroCalc/HydroMCalc.html>, in.
- 238 [11] Clinical and laboratory standards institute., 2012.
- 239 [12] *Pure Appl. Chem*, 79 (2007) 717-728.
- 240 [13] T.R. Anderegg, T.R. Fritsche, R.N. Jones, Quality control guidelines for MIC susceptibility
241 testing of omiganan pentahydrochloride (MBI 226), a novel antimicrobial peptide, *J Clin*
242 *Microbiol*, 42 (2004) 1386-1387.
- 243 [14] A. Giangaspero, L. Sandri, A. Tossi, Amphipathic alpha helical antimicrobial peptides, *Eur J*
244 *Biochem*, 268 (2001) 5589-5600.
- 245 [15] R. Rathinakumar, W.F. Walkenhorst, W.C. Wimley, Broad-spectrum antimicrobial peptides
246 by rational combinatorial design and high-throughput screening: the importance of interfacial
247 activity, *J Am Chem Soc*, 131 (2009) 7609-7617.
- 248
- 249

Table 1.

	Peptide 1	Peptide 2	Peptide 3	Peptide 4	Peptide 5
Mean Hydrophobicity (<i>H</i>)	-0.67	-0.23	-0.54	-0.12	-0.28
Mean hydrophobic moment (μH)	0.24	0.35	0.14	0.17	0.41
Helicity (%Helix)	27.86	64.2	0.54	8.81	88.43
*Isoelectric Point	13.10	10.89	12.25	9.85	11.75
*Net Charge	+12	+6	+7	+3	+7
Hemolytic activity	3.4	39.8	1.5	1.72	9.5

Omiganan
-0.31
0.28
10.15
12.79
+4
10

ACCEPTED MANUSCRIPT

Table 2 continued,

Specie	Strain	Genes	MIC (mg/L)			
			Peptide 1	Peptide 2	Peptide 5	Omiganan
	M5306	<i>bla</i> _{PER-2} + <i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1}	256	4	64	64
	M9209	<i>bla</i> _{KPC-2}	128	64	128	64
<i>S. aureus</i> (11)	ATCC29213	none	32	64	32	32
	P33	<i>msrA</i>	32	128	64	32
	P28	<i>ermA</i>	32	64	64	32
	P204	<i>ermA</i>	64	64	32	64
	M6276	<i>ermA</i> + <i>lnuA</i>	16	128	64	32
	P239	<i>ermC</i>	32	64	64	32
	M6794	<i>mecA</i>	32	64	64	64
	M2832	<i>mecA</i>	32	128	64	64
	M4046	<i>mecA</i>	32	128	32	32
	M6820	<i>mecA</i>	64	128	64	128
	M6784	<i>mecA</i>	32	32	32	32
<i>S. epidermidis</i> (4)	M2923	none	16	16	16	8
	M2931	none	16	16	8	16
	M2919	<i>mecA</i>	16	16	8	8
	M2921	<i>mecA</i>	8	8	8	8
<i>S. saprophyticus</i> (2)	M4070	<i>mecA</i>	16	32	8	8
	M2981	<i>mecA</i>	16	8	8	8
<i>S. haemolyticus</i> (2)	M2976	<i>mecA</i>	16	8	8	4
	M3014	none	16	8	8	4
<i>S. hominis</i> (2)	M2973	<i>mecA</i>	4	4	4	4
	M2967	<i>mecA</i>	8	8	8	4
<i>S. warnerii</i> (1)	M6823	<i>mecA</i>	8	8	8	8
<i>S. cohnii</i> (1)	M6767	<i>mecA</i>	16	16	8	4
<i>E. faecalis</i> (8)	ATCC 29212	none	64	128	128	128
	ATCC 51299	<i>vanB</i>	256	256	256	256
	M4899	<i>vanB</i>	256	128	256	256
	M6534	<i>vanB</i>	128	256	256	256
	M4992	<i>vanA</i>	128	128	128	128
	M6383	<i>vanA</i>	128	128	128	128
	M4449	<i>vanA</i>	128	128	128	128
	M6983	<i>vanA</i>	64	128	128	128
<i>E. faecium</i> (6)	PZAP95	none	32	16	16	16
	M6261	none	32	16	16	16
	M2619	<i>vanB</i>	32	16	16	16
	M2481	<i>vanB</i>	32	16	16	16
	M2304	<i>vanA</i>	16	16	8	4
	M2664	<i>vanA</i>	16	8	8	8
<i>E. gallinarum</i> (2)	M2723	<i>vanC1</i> + <i>vanA</i>	32	32	16	16
	M2685	<i>vanC1</i> + <i>vanA</i>	16	16	16	16


ACCEPTED MANUSCRIPT

Table 2.

Specie	Strain	Genes	MIC (mg/L)			
			Peptide 1	Peptide 2	Peptide 5	Omiganan
<i>P. aeruginosa</i> (12)						
	ATCC 27853	none	64	64	64	256
	PCOS12	none	64	64	64	512
	M5470	none	128	64	64	512
	M7907	<i>bla</i> _{PER}	128	64	64	128
	M13513	<i>bla</i> _{KPC-2}	32	64	64	256
	M11005	<i>bla</i> _{KPC-2}	64	64	128	512
	M7723	<i>bla</i> _{KPC-2}	64	64	32	256
	M7728	<i>bla</i> _{IMP}	128	64	64	512
	M5109	<i>bla</i> _{VIM} + <i>bla</i> _{GES-1}	128	64	64	256
	M5200	<i>bla</i> _{VIM} + <i>bla</i> _{GES-1}	64	64	64	512
	M7525	<i>bla</i> _{SPM}	64	64	64	512
	M7712	<i>bla</i> _{SPM}	64	64	64	512
<i>Acinetobacter</i> sp. (10)						
	M13523	<i>bla</i> _{OXA-51}	64	4	64	32
	M9665	<i>bla</i> _{OXA-51}	128	32	128	4
	M5282	<i>bla</i> _{OXA-51}	64	8	64	8
	M5179	<i>bla</i> _{OXA-51}	64	32	64	32
	M7489	<i>bla</i> _{OXA-51} + <i>bla</i> _{TEM}	64	16	64	8
	PFAV1	<i>bla</i> _{OXA-51} + <i>bla</i> _{OXA-58} + <i>bla</i> _{PER}	64	16	64	16
	M5277	<i>bla</i> _{PER}	64	8	64	32
	M5949	<i>bla</i> _{OXA-23} + <i>bla</i> _{OXA-GVI}	256	16	256	32
	M7978	<i>bla</i> _{IMP-1}	64	8	64	16
	M9013	<i>bla</i> _{OXA-51} + <i>bla</i> _{IMP}	32	8	32	32
<i>K. pneumoniae</i> (12)						
	PFAV3	none	1024	128	128	128
	M9140	<i>bla</i> _{CIT}	1024	32	32	64
	M9491	<i>bla</i> _{MOX}	1024	64	64	128
	M9170	<i>bla</i> _{OXA-GIII}	>1024	32	64	128
	M5825	<i>bla</i> _{GES-3} + <i>bla</i> _{CTX-M-2}	>1024	8	32	64
	M9310	<i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1} + <i>bla</i> _{SHV-1}	>1024	16	32	64
	M9375	<i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1} + <i>bla</i> _{SHV-1}	1024	64	16	32
	M1803	<i>bla</i> _{PER-2} + <i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1} + <i>bla</i> _{SHV} + <i>bla</i> _{OXA-9}	>1024	32	64	1024
	M7647	<i>bla</i> _{VIM} + <i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1} + <i>bla</i> _{SHV-1}	>1024	32	32	1024
	M13540	<i>bla</i> _{KPC-2}	>1024	16	64	256
	M9885	<i>bla</i> _{KPC-2}	>1024	64	32	256
	M11245	<i>bla</i> _{KPC-2} + <i>bla</i> _{PER-2}	1024	16	8	32
<i>E. coli</i> (9)						
	ATCC 25922	none	128	32	32	64
	M9884	none	128	32	32	64
	M7859	<i>bla</i> _{CIT}	128	32	16	64
	PNEU23	<i>bla</i> _{OXA-GIII} + <i>bla</i> _{TEM-1}	128	32	32	32
	PCOS15	<i>bla</i> _{PER-2} + <i>bla</i> _{TEM-1}	128	64	64	64
	PABC11	<i>bla</i> _{CTX-M-2}	128	32	32	64
	PLCA1	<i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1}	128	64	64	64

	M5306	<i>bla</i> _{PER-2} + <i>bla</i> _{CTX-M-2} + <i>bla</i> _{TEM-1}	256	4	64	64
	M9209	<i>bla</i> _{KPC-2}	128	64	128	64
<i>S. aureus</i> (11)						
	ATCC29213	none	32	64	32	32
	P33	<i>msrA</i>	32	128	64	32
	P28	<i>ermA</i>	32	64	64	32
	P204	<i>ermA</i>	64	64	32	64
	M6276	<i>ermA</i> + <i>lnuA</i>	16	128	64	32
	P239	<i>ermC</i>	32	64	64	32
	M6794	<i>mecA</i>	32	64	64	64
	M2832	<i>mecA</i>	32	128	64	64
	M4046	<i>mecA</i>	32	128	32	32
	M6820	<i>mecA</i>	64	128	64	128
	M6784	<i>mecA</i>	32	32	32	32
<i>S. epidermidis</i> (4)						
	M2923	none	16	16	16	8
	M2931	none	16	16	8	16
	M2919	<i>mecA</i>	16	16	8	8
	M2921	<i>mecA</i>	8	8	8	8
<i>S. saprophyticus</i> (2)						
	M4070	<i>mecA</i>	16	32	8	8
	M2981	<i>mecA</i>	16	8	8	8
<i>S. haemolyticus</i> (2)						
	M2976	<i>mecA</i>	16	8	8	4
	M3014	none	16	8	8	4
<i>S. hominis</i> (2)						
	M2973	<i>mecA</i>	4	4	4	4
	M2967	<i>mecA</i>	8	8	8	4
<i>S. warnerii</i> (1)						
	M6823	<i>mecA</i>	8	8	8	8
<i>S. cohnii</i> (1)						
	M6767	<i>mecA</i>	16	16	8	4
<i>E. faecalis</i> (8)						
	ATCC 29212	none	64	128	128	128
	ATCC 51299	<i>vanB</i>	256	256	256	256
	M4899	<i>vanB</i>	256	128	256	256
	M6534	<i>vanB</i>	128	256	256	256
	M4992	<i>vanA</i>	128	128	128	128
	M6383	<i>vanA</i>	128	128	128	128
	M4449	<i>vanA</i>	128	128	128	128
	M6983	<i>vanA</i>	64	128	128	128
<i>E. faecium</i> (6)						
	PZAP95	none	32	16	16	16
	M6261	none	32	16	16	16
	M2619	<i>vanB</i>	32	16	16	16
	M2481	<i>vanB</i>	32	16	16	16
	M2304	<i>vanA</i>	16	16	8	4
	M2664	<i>vanA</i>	16	8	8	8
<i>E. gallinarum</i> (2)						
	M2723	<i>vanC1</i> + <i>vanA</i>	32	32	16	16
	M2685	<i>vanC1</i> + <i>vanA</i>	16	16	16	16

Figure 1.



Circular dichroism of peptides in aqueous solution (panel A) and in SDS micelles (panel B). Projection of helical wheel of the peptides (panel C). omi, omiganan; p1, peptide 1; p2, peptide 2; p3, peptide 3; p4, peptide 4; p5, peptide 5.

Antimicrobial activity of *de novo* designed cationic peptides against multi-resistant clinical isolates.

Highlights.

- *de novo* designed cationic peptides
- physicochemical properties of peptides
- antimicrobial resistant bacteria