Transmission and Reflection of Strongly Nonlinear Solitary Waves at Granular Interfaces

A. M. Tichler,^{1,2} L. R. Gómez,^{1,3} N. Upadhyaya,¹ X. Campman,² V. F. Nesterenko,⁴ and V. Vitelli^{1,*}

¹*Instituut-Lorentz for Theoretical Physics, Universiteit Leiden, 2300 RA Leiden, The Netherlands*

²*Shell Global Solutions International B.V., Kessler Park 1, 2288 GS Rijswijk, The Netherlands*

³*Department of Physics, Universidad Nacional del Sur—IFISUR—CONICET, 8000 Bahía Blanca, Argentina*

⁴*Jacobs School of Engineering, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA*

(Received 20 March 2013; published 25 July 2013)

The interaction of a solitary wave with an interface formed by two strongly nonlinear noncohesive granular lattices displays rich behavior, characterized by the breakdown of continuum equations of motion in the vicinity of the interface. By treating the solitary wave as a quasiparticle with an effective mass, we construct an intuitive (energy- and linear-momentum-conserving) discrete model to predict the amplitudes of the transmitted solitary waves generated when an incident solitary-wave front, parallel to the interface, moves from a denser to a lighter granular hexagonal lattice. Our findings are corroborated with simulations. We then successfully extend this model to oblique interfaces, where we find that the angle of refraction and reflection of a solitary wave follows, below a critical value, an analogue of Snell's law in which the solitary-wave speed replaces the speed of sound, which is zero in the sonic vacuum.

DOI: 10.1103/PhysRevLett.111.048001

PACS numbers: 45.70.-n, 61.43.Fs, 65.60.+a, 83.80.Fg

The study of solitary waves has over the years led to a paradigmatic shift in our understanding of many-body phenomena characterized by anharmonic effects that manifest themselves in exotic electronic [1] and mechanical states [2,3]. A concrete and technologically relevant [4,5] arena to study strongly nonlinear mechanical waves is the *sonic vacuum* [6,7]—a paradigmatic example of which is an aggregate of grains just in contact. Owing to the vanishing speed of linear sound, even the tiniest strains propagate as supersonic solitary waves, nonlinear periodic waves, and shocklike waves depending on conditions of loading. However, the differential equations describing the propagation of mechanical disturbances around the state of sonic vacuum are both nonlinear and generally not integrable, making it difficult to solve them analytically. Moreover, the continuum approximation itself may fail in the vicinity of a sharp granular interface, where the discrete nature of the granular medium dominates.

So far, most studies of strongly nonlinear granular interfaces have concentrated on what happens when a solitary wave initially propagating in a chain with mass m_1 reaches an interface where the particles mass suddenly changes to m_2 . Depending on the ratio $A = m_2/m_1$, qualitatively different behaviors are observed. When the solitary wave moves from a lighter to a denser medium ($A > 1$), most of its energy gets divided into a reflected and a transmitted pulse, whose respective amplitudes can be estimated by using the conservation of linear momentum and energy [2,6,7]. By contrast, when the incident solitary wave moves from a denser to a lighter medium ($A \ll 1$), a train of (multiple) solitary waves is generated in the lighter medium [2,6]. In this case, a direct application of the two conservation laws (momentum and energy) is not sufficient to predict the amplitude ratios of the *multiple* solitary

waves. Experimental studies of granular chains comprised of steel and polytetrafluoroethylene particles ($A = 0.27$) have shed light on the discrete mechanism responsible for the generation of the train of solitary waves in the polytetrafluoroethylene light chain. Most of the collective motion carried by the incident solitary wave (propagating in the chain of stainless steel particles) is converted into the motion of a single interfacial steel particle whose subsequent motion generates a train of solitary waves [8–10].

In this Letter, we turn to the hitherto unexplored two-dimensional problem of determining the reflection and transmission of a strongly nonlinear solitary-wave front incident upon an interface between two hexagonal lattices both in a sonic vacuum but with different particle masses. We treat the solitary waves as quasiparticles with an effective mass and model the interaction with a two-dimensional granular interface, by assuming an energy- and linear-momentum-conserving scenario validated by simulations. In the $A \ll 1$ case, the last row of “heavy” interfacial beads absorbs on a “fast” time scale the main part of the energy and linear momentum of the incident solitary-wave front (assumed parallel to the interface) and subsequently decelerates on a “slow” time scale, generating a train of (asymptotically) separated solitary waves in the “lighter” sonic vacuum. Crucial to understanding this phenomenon is the role of contact breaking at the interface and the resulting breakdown of the continuum approximation. When a strongly nonlinear wave is incident at an oblique angle to the interface, we find that the angles of refraction and reflection are surprisingly well captured by a granular analog of Snell's law that holds irrespective of the solitary-wave amplitude.

Parallel interfaces.—In order to investigate solitary waves scattering at two-dimensional granular interfaces,

we performed molecular dynamics simulations of an impact experiment performed on hexagonal packings of 10^4 frictionless spherical grains using periodic boundary conditions. As shown in Fig. 1, an interface is introduced by assigning a mass m_1 to the rows of grains on its left (shown in red) and a mass m_2 to rows on its right (shown in yellow). Both portions of the hexagonal lattice are comprised of grains with zero initial overlap and equal diameters. Two grains of radius R and masses $\{m_i, m_j\}$ at positions $\{\vec{x}_i, \vec{x}_j\}$ interact via a one-sided nonlinear repulsive potential following the Hertz law [11]

$$V_{ij} = \frac{K_{ij}}{\alpha} \delta_{ij}^{5/2} \quad (1)$$

only for positive compressional strains $\delta_{ij} \equiv 2R - |\vec{x}_i - \vec{x}_j| > 0$; otherwise, $V_{ij} = 0$, when $\delta_{ij} \leq 0$. Here, the

FIG. 1 (color online). Time sequence leading to the generation of a solitary-wave train in simulations. The (red) beads on the left of the interface constitute the heavier medium with mass m_1 , and the (yellow) beads on the right of the interface constitute the lighter medium with mass m_2 . The mass ratio $A \equiv m_2/m_1 = 0.125$. The velocity field, overlaid in green, denotes the instantaneous speeds of the beads. An incident solitary wave (a) collides with the interface of two sonic vacuums (b)–(d) leading to the formation of a train of transmitted solitary waves (e).

interaction parameter $K_{ij} = (2/3)RE_{ij}^*$ is expressed in terms of the effective Young's modulus of the two particles, E_{ij}^* ; see Ref. [12] for more details. At $t = 0$ we impart to the leftmost row a speed u_p and subsequently integrate Newton's equations of motion numerically subject to periodic boundary conditions perpendicularly to the direction of propagation. The elastic moduli and radii for heavy and light particles are considered to be the same.

As shown in Fig. 1(a), this initial condition leads to the generation of a nonlinear wave front parallel to the interface traveling towards the right with a speed $V_s \sim u_p^{1/5}$ analogously to solitary waves in granular chains composed of 5 particles width [2]. At later times shown in Fig. 1(b), when the solitary wave has interacted with the interface, we see a ruptured interface with one of the interfacial rows of heavy (red) beads “dancing” (in a somewhat oscillatory motion) in contact with the lattice of lighter (yellow) beads, throttling the generation of an oscillatory wave profile in the lighter lattice close to the interface. This oscillatory wave is subsequently disintegrated into a sequence of separate solitary waves, as shown in Fig. 1(c). The separate solitary waves propagate with different speeds (dependent on their amplitude), while a second collision of the dancing interfacial row of particles, shown in Fig. 1(d), generates a second delayed solitary-wave train with smaller amplitudes; see Fig. 1(e) and Movie 1 in Supplemental Material [13].

Quasiparticle collision model.—We take advantage of the isotropic elasticity of the hexagonal lattice to assume that the dynamics of a solitary-wave front parallel to the interface, as in Fig. 1, is effectively one-dimensional and governed, in the continuum limit, by the nonlinear wave equation [2]

$$\xi_{tt} = c^2 \left[\xi^{3/2} + \frac{2R^2}{5} \xi^{1/4} (\xi^{5/4})_{xx} \right], \quad (2)$$

where c is a constant determined by grain material property and geometry as well as their arrangements and $\xi(x, t)$ is the strain field $\xi(x, t) = -\partial_x u(x, t)$ expressed in terms of the particle displacement field $u(x, t)$ along the x direction.

A strongly nonlinear solitary-wave solution of Eq. (2) can be derived analytically [2], and it has been validated by extensive simulations and experiments mostly on granular chains [2,3,7,9,14–18]. Crucially, the total energy $E = P^2/2m_{\text{eff}}$ carried by the solitary wave depends quadratically on its total momentum P , which allows one to define an effective mass $m_{\text{eff}} \approx 1.4m$ for the solitary wave [2,6,7,9,19,20].

The notion of the solitary wave as a quasiparticle allows one to construct a simple quasi-one-dimensional model for the generation of the solitary-wave train, illustrated schematically in Fig. 2. At t_0 , we assume that a chain of light yellow beads is uncompressed and all the energy and linear momentum P_0 , carried by the incident solitary wave, is concentrated in the heavy red interfacial particles. At a

FIG. 2 (color online). (a) Schematic illustration of our model for the formation of a solitary-wave train, side view. (b) Momentum ratios $P_{2,n}/P_{2,1}$ between the n th solitary wave and the leading one in the train for $A = m_2/m_1 = 0.125$. Red circles are the theoretical predictions, while the black squares are the numerical values from the simulations of Fig. 1.

subsequent time t_1 a single solitary wave is generated in the light chain by reducing the energy and linear momentum of the interfacial heavy particles which for simplicity we will consider as a single particle with mass m_1 . We apply conservation of energy and linear momentum to the “collision” process between the dancing bead with mass m_1 (mass of particles in the heavy chain) and the solitary wave, treated as a quasiparticle with mass $m_{2,\text{eff}}$. We calculate the momentum of the dancing interfacial particle $P_{1,1}$ after the first collision as

$$P_{1,1} = \frac{P_0(B-1)}{(B+1)}, \quad (3)$$

where $B \equiv m_1/m_{2,\text{eff}}$. The momentum $P_{2,1}$ carried by the first leading solitary wave at $t = t_1$ is

$$P_{2,1} = \frac{2P_0}{B+1}. \quad (4)$$

At time t_2 , another independent single solitary wave is generated in the light chain, further reducing the energy and linear momentum of the dancing interfacial particle. Upon applying conservation of energy and linear momentum, as before, and assuming that the first solitary wave does not participate in this process, we find the momentum of the dancing particle at $t = t_2$, $P_{1,2}$, and of the second solitary wave, $P_{2,2}$, as

$$P_{1,2} = \frac{P_0(B-1)^2}{(B+1)^2}, \quad P_{2,2} = \frac{2P_0(B-1)}{(B+1)^2}. \quad (5)$$

Upon iterating this process n times, we find that the heavy interfacial row of beads at $t = t_n$ is left with a linear momentum $P_{1,n}$, while the n th solitary wave carries momentum $P_{2,n}$ given by

FIG. 3 (color online). Snapshot of the simulations showing a solitary wave incident upon an interface separating two hexagonal lattices in a sonic vacuum. (a) For $A \equiv m_2/m_1 = 0.125$, the transmitted disturbance propagates in the form of a nonlinear oscillatory wave analogous to the train of solitary waves shown for granular chains in Fig. 1 for $A < 1$. (b) For $A > 1$, we find both a reflected and a transmitted solitary wave as shown in Fig. 1 for $A = 8$.

$$P_{1,n} = \frac{P_0(B-1)^n}{(B+1)^n}, \quad P_{2,n} = \frac{2P_0(B-1)^{n-1}}{(B+1)^n}. \quad (6)$$

Figure 2(b) illustrates the favorable comparison of $P_{2,n}/P_{2,1} = (B-1/B+1)^{n-1}$ against numerical data (red circles) for $A = 0.125$. The amplitudes of the delayed secondary sequence of solitary waves generated is neglected in our approximate model.

Oblique interfaces.—The simulation snapshots in Fig. 3 illustrate the propagation and interaction of a solitary-wave front with an oblique interface separating two hexagonal granular lattices for the case $A \equiv (m_2/m_1) < 1$ in panel (a) and for the case $A > 1$ in panel (b)—see also Movies 2(a) and 2(b) in Supplemental Material [13]. From these simulations, we have determined numerically the angle of refraction θ_{refr} for different values of the angle of incidence θ_i , as shown in the inset in Fig. 4(a).

Inspection of the main panel of Fig. 4(a) suggests that a linear relationship exists between $\sin\theta_{\text{refr}}$ and $\sin\theta_i$ for mass ratios $A \ll 1$ (squares) and $A \leq 1$ (circles). We now show that the measured proportionality coefficient is consistent with a nonlinear analogue of Snell’s law $V_0 \sin\theta_{\text{refr}} = V_{2,1} \sin\theta_i$, where $\{V_{2,1}, V_0\}$ denote, respectively, the speeds of the (leading) refracted solitary wave and of the incident one. To work out explicitly the dependence of $V_{2,1}/V_0$ on the mass ratio $A < 1$, we employ a

FIG. 4 (color online). (a) Angle of refraction θ_{refr} vs angle of incidence θ_i for the hexagonal lattice when $A < 1$. The square (circle) symbols correspond to the case when $A = 0.125$ ($A = 0.9$) and compares the numerically obtained ratio of the speed of the leading transmitted solitary wave to the incident solitary wave against the analytical estimates given by solid curves. (b) Comparison of numerical data (symbols) with the analytical estimate (solid curves) for the angle of refraction (black data) and angle of reflection (blue data) vs the angle of incidence for the hexagonal lattice when $A > 1$. The top panels of (a) and (b) describe the relevant angles, where the interface is shown as the dashed (red) line and arrows represent the direction of propagation of the solitary-wave front (thick dark region).

reasoning similar to the one leading to Eq. (4) that accounts for the discrete mechanism at play near the interface. Upon making use of the scaling relation $P \sim m^3 V_s^5$ between the total momentum P carried by a solitary wave and its speed V_s [9], we find in the limit of small θ_i that

$$\frac{\sin(\theta_{\text{refr}})}{\sin(\theta_i)} \approx \left(\frac{2}{(B+1)A^3} \right)^{1/5}. \quad (7)$$

The right-hand side of Eq. (7) is the slope of the continuous line plotted for $A = 0.125$ and $A = 0.9$ in Fig. 4(a). It matches the data (open symbols) obtained by numerically evaluating the left-hand side of Eq. (7). This agreement corroborates the nonlinear analogue of Snell's law. Note that, for the case $A < 1$, the reflected wave amplitude is negligibly small; see Fig. 3(a).

By contrast, in the case $A > 1$, Fig. 3(b) shows a train of reflected waves with most of the energy concentrated in the leading solitary wave, which allows us to neglect rupturing at the interface. As shown in Fig. 4(b), the angle of reflection θ_{refl} is not equal to the angle of incidence θ_i , since the reflected solitary-wave speed V_{refl} is not the

FIG. 5 (color online). Delayed reflection of a solitary wave for the case $\theta_i > \theta_c$, when $A = 0.125$ and $\theta_i = 1.8\theta_c$.

same as the incident speed V_0 . Instead, we evaluate the ratio $V_{1,1}/V_0$ in analogy with the derivation of Eq. (7). Since, in this case, the appropriate collision model is that between two solitary-wave quasiparticles, the effective mass contribution cancels out, and we obtain an equation analogous to Eq. (3) with the replacement $B \rightarrow 1/A$ and a sign reversal. Thus, Snell's law for reflection assumes the form

$$\frac{\sin(\theta_{\text{refl}})}{\sin(\theta_i)} \approx \left(\frac{A-1}{A+1} \right)^{1/5}. \quad (8)$$

The right-hand side of Eq. (8) is the slope of the continuous (blue) line plotted for $A = 3$ in Fig. 4(b) and matches the numerical data (blue circles) obtained from evaluating the left-hand side of Eq. (8). An agreement is found also for the angle of refraction in this case, as illustrated by the data (black squares) and (black) continuous line in Fig. 4(b). Despite the fact that our granular lattices are in a state of sonic vacuum, the ratio between the sine of the angles in Snell's law is nearly independent of the amplitude of the incident solitary-wave front (see Fig. S1 in Supplemental Material [13]). This seemingly puzzling observation can be rationalized by viewing the incident and (leading) reflected or refracted solitary waves as quasiparticles scattering at the interface whose speeds are proportional to the incoming ones [21].

This analog of Snell's law also implies the existence of a critical angle of incidence θ_c , for which the transmitted solitary wave will propagate in a direction parallel to the line of the interface or $\theta_{\text{refr}} = 90^\circ$ in Eq. (7), with the replacement $B \rightarrow 1/A$. Figure 5 shows the reflection of a solitary wave for the case $\theta_i > \theta_c$. Note that, in this case, the reflected solitary wave is “delayed” with respect to the transmitted train, such that the reflected and transmitted fronts cross the interface at different points.

Conclusion.—We have constructed a discrete model that predicts the amplitudes of transmitted and reflected solitary-wave fronts across a 2D granular interface. We find that the angle of refraction and reflection provides a novel granular analogue of Snell's law where the solitary-wave speed replaces the vanishing speed of sound and could potentially stimulate further experiments.

We acknowledge discussion with S. Ulrich. L. R. G. and N. U. acknowledge financial support from FOM, Shell, ANPCyT, and CONICET. A. M. T., L. R. G., and N. U. contributed equally to this work.

*vitelli@lorentz.leidenuniv.nl

- [1] A. J. Heeger, S. Kivelson, J. R. Schrieffer, and W.-P. Su, *Rev. Mod. Phys.* **60**, 781 (1988).
- [2] V. F. Nesterenko, *Dynamics of Heterogeneous Materials* (Springer-Verlag, New York, 2001), Chap. 1.
- [3] S. Sen, J. Hong, J. Bang, E. Avalos, and R. Doney, *Phys. Rep.* **462**, 21 (2008).
- [4] A. Spadoni and C. Daraio, *Proc. Natl. Acad. Sci. U.S.A.* **107**, 7230 (2010).
- [5] N. Boechler, G. Theocharis, and C. Daraio, *Nat. Mater.* **10**, 665 (2011).
- [6] V. F. Nesterenko, A. N. Lazaridi, and E. B. Sibiriyakov, *Prikl. Mekh. Tekh. Fiz.* **36**, 19 (1995) [*J. Appl. Mech. Tech. Phys.* **36**, 166 (1995)].
- [7] V. F. Nesterenko, *J. Phys. IV (France)* **04**, C8-729 (1994).
- [8] V. F. Nesterenko, C. Daraio, E. Herbold, and S. Jin, *Phys. Rev. Lett.* **95**, 158702 (2005).
- [9] S. Job, F. Melo, A. Sokolow, and S. Sen, *Granular Matter* **10**, 13 (2007).
- [10] A. Sokolow, E. G. Bittle, and S. Sen, *Europhys. Lett.* **77**, 24002 (2007).
- [11] L. D. Landau and E. M. Lifshitz, *Theory of Elasticity* (Pergamon, New York, 1970).
- [12] E. Somfai, J. N. Roux, J. H. Snoeijer, M. van Hecke, and W. van Saarloos, *Phys. Rev. E* **72**, 021301 (2005).
- [13] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.111.048001> for additional figures and movies.
- [14] A. N. Lazaridi and V. F. Nesterenko, *Prikl. Mekh. Tekh. Fiz.* **26**, 115 (1985) [*J. Appl. Mech. Tech. Phys.* **26**, 405 (1985)].
- [15] C. Coste, E. Falcon, and S. Fauve, *Phys. Rev. E* **56**, 6104 (1997).
- [16] C. Daraio, V. F. Nesterenko, E. B. Herbold, and S. Jin, *Phys. Rev. E* **72**, 016603 (2005).
- [17] L. R. Gómez, A. M. Turner, M. van Hecke, and V. Vitelli, *Phys. Rev. Lett.* **108**, 058001 (2012).
- [18] L. R. Gómez, A. M. Turner, and V. Vitelli, *Phys. Rev. E* **86**, 041302 (2012).
- [19] C. Daraio, V. F. Nesterenko, and S. Jin, in *Shock Compression of Condensed Matter 2003, Proceedings of the Conference of the American Physical Society Topical Group on Shock Compression of Condensed Matter*, edited by M. D. Furnish, Y. M. Gupta, and J. W. Forbes (AIP, New York, 2004), Vol. 706, p. 197.
- [20] E. J. Hinch and S. Saint-Jean, *Proc. R. Soc. A* **455**, 3201 (1999).
- [21] It is interesting that shear zones are refracted when they pass the interface between different granular materials where effective coefficients of friction replace wave speeds; see T. Unger, *Phys. Rev. Lett.* **98**, 018301 (2007); H. A. Knudsen and J. Bergli, *Phys. Rev. Lett.* **103**, 108301 (2009).