

Accepted Manuscript

Isolation and selection of yeasts from wine grape ecosystem secreting cold-active pectinolytic activity

María Gabriela Merín, Lucía M. Mendoza, Marta E. Farías, Vilma Inés Morata


PII: S0168-1605(11)00204-2
DOI: doi: [10.1016/j.ijfoodmicro.2011.04.004](https://doi.org/10.1016/j.ijfoodmicro.2011.04.004)
Reference: FOOD 5511

To appear in: *International Journal of Food Microbiology*

Received date: 19 February 2011
Revised date: 17 March 2011
Accepted date: 6 April 2011

Please cite this article as: Merín, María Gabriela, Mendoza, Lucía M., Farías, Marta E., Morata, Vilma Inés, Isolation and selection of yeasts from wine grape ecosystem secreting cold-active pectinolytic activity, *International Journal of Food Microbiology* (2011), doi: [10.1016/j.ijfoodmicro.2011.04.004](https://doi.org/10.1016/j.ijfoodmicro.2011.04.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Isolation and selection of yeasts from wine grape ecosystem secreting cold-active pectinolytic activity

María Gabriela Merín^{a,b}, Lucía M. Mendoza^{b,c}, Marta E. Farías^{b,c}, Vilma Inés Morata^{a,b,*}.

^a*Facultad de Ciencias Aplicadas a la Industria, Universidad Nacional de Cuyo, Bernardo de Irigoyen 375 (5600) San Rafael, Mendoza, Argentina;*

^b*Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET);*

^c*Centro de Referencia para Lactobacilos (CERELA-CONICET), Chacabuco 145 (4000) Tucumán, Argentina.*

*Corresponding author: Facultad de Ciencias Aplicadas a la Industria, Universidad Nacional de Cuyo, Bernardo de Irigoyen 375 (5600) San Rafael, Mendoza, Argentina. Tel./Fax: +54-2627-421947/430673.
E-mail address: vmorata@fcai.uncu.edu.ar (V.I. Morata)

Abstract

The present study was undertaken with the purpose of selecting yeasts from wine grapes that are able to produce extracellular cold-active pectinases. After two consecutive selections yeast isolates were identified by pheno- and genotyping, and pectinolytic activity was preliminarily characterised at proximate winemaking conditions. Out of 1023 indigenous microorganisms isolated from grape skins of D.O. San Rafael (Mendoza, Argentina) viticulture region, 565 (55%) showed pectinolytic activity on plates and, among them, 96 (17%) were chosen in a primary selection. Ten isolates were finally selected for exhibiting the greatest activity at low temperature (12 °C) and identified as *Aureobasidium pullulans*. GM-R-22 strain demonstrated the highest pectinolytic activity (0.751 U/mL) at pH 3.5 and 12 °C. Yeast pectinases were constitutively produced. This study is the first report about strains of *A. pullulans* producing pectinases which are able to show good activity at low temperature. These pectinolytic strains could be of interest in wine production.

Keywords: Cold-active pectinases; *Aureobasidium pullulans*; Wine yeasts.

1. Introduction

Pectinolytic enzymes are widely used in the beverage industry to clarify fruit juices and wine. They can degrade pectic substances, which are mainly composed of partially methyl-esterified galacturonic acid subunits joined by $\alpha(1-4)$ -glycosidic bonds. These enzymes play an important role in winemaking (Kashyap et al., 2001). They can help to improve liquefaction, clarification and filterability, releasing more colour and flavor compounds entrapped in the grape skins, thereby making a positive contribution to the wine bouquet, and facilitating the liberation of phenolic compounds (Van Rensburg and Pretorius, 2000). In a previous work carried out in our laboratory, an autochthonous-microbial pectinolytic preparation was applied to red-skin maceration with very good results in pigment and polyphenols extraction (Cabeza et al., 2009).

Most strains of *Saccharomyces cerevisiae*, usually used in winemaking, do not show the capacity to degrade pectic substrates. A few wild strains have been reported to have the ability to degrade pectin in wine fermentation (Blanco et al., 1994; Fernández-González et al., 2005). Recently, there has been interest in the application of non-*Saccharomyces* wine yeasts, but the ability of these yeasts to secrete pectinases has been scarcely studied (Charoenchai et al., 1997; Fernández et al., 2000; Strauss et al., 2001).

Fermentation conducted at low temperatures (15-20 °C) can increase the production and retention of volatile compounds, improving the aromatic profile of wines (Molina et al., 2007). However, it is well known that colour extraction in red wines is reduced within this temperature range. Consequently, pectinolytic enzymes able to be active at low temperature, both for extraction and clarification, are needed. Few attempts have been made to isolate and characterise cold-active pectinases from yeasts (Birgisson et al., 2003; Nakagawa et al., 2004).

“San Rafael” Designation of Origin (D.O.), a Western Argentinean wine-making area, stands at the foot of Los Andes principal mountain range. It presents a special microclimate that contributes to distinguish its vineyards and wines. Grapes from this region may have a unique biota of wine yeasts with potential to produce pectinases that act at low temperatures.

This work was aimed at the selection and identification of yeasts isolated from wine grapes, which are capable of secreting cold-active pectinases. It reports the preliminary characterisation of pectinolytic activity at some vinification conditions such as low pH (3.5), low temperature (12 °C) as well as at 28 °C.

2. Materials and Methods

2.1. Study area, sampling and yeast isolation

The samples were collected from different districts of “San Rafael” D.O., in Southern Mendoza (34.5°-36° South latitude and 70°-66.5° Western longitude). All samples were taken in 2007 vintage, at optimum ripeness and in good sanitary conditions. Clusters of 11 grape varieties were randomly harvested in vineyards, transported to the laboratory aseptically, and kept cold until their study.

Ten berries of each grape variety were placed in a flask containing 10 mL of sterile peptone water (0.1%). The system was shaken at 165 rpm during 1 h at room temperature. An aliquot was spread in surface on three different media: WL nutrient agar (Merck); YGC (Merck); and grape must-agar medium containing per litre: concentrated grape must (868 g reducing sugar/L), 2.3 mL; yeast extract, 1.0 g; agar, 2.0 g; pH 4.5. The plates were incubated at 28 °C for 24-48 h. Approximately 30 colonies were isolated from each plate according to their macroscopic features and frequencies.

2.2. Screening on plates for pectinolytic yeast selection

2.2.1. Primary selection

The isolates were point-inoculated onto a mineral medium containing citric pectin as a sole carbon source: (g/L) citric pectin, 2.0; yeast extract, 1.0; agar, 15.0; KH_2PO_4 , 0.2; CaCl_2 , 0.05; $(\text{NH}_4)_2\text{SO}_4$, 1.0; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.8; MnSO_4 , 0.05; pH 4.5, reported by Moyo et al. (2003) with slight modifications. The production of pectinolytic enzymes was detected on plates incubated at 28 °C for 48 h. Enzyme activity was indicated by the formation of a clear halo around the colonies against a purple–brown background on pectin plate after lugol's solution addition (Fernandes-Salomão et al., 1996).

2.2.2. Secondary selection

A second selection was carried out according to the capability of pre-selected yeasts to grow and produce pectinases that are active at low temperature (12 °C). Two criteria were taken into account, the first one was to select the yeasts producing the widest halo diameter/colony diameter relation (D_h/D_c) and the second criterion was to choose those that produced the largest colony size (D_c) and, at the same time, secreting important amounts of enzyme, although this originated a medium value of D_h/D_c relation. Moreover, the pre-selected isolates were tested to detect pectinolytic activity at 28 °C with the same criteria, in order to choose the best pectinase-secreting yeasts at this temperature.

2.3. Enzyme assays under proximate oenological conditions

2.3.1. Production of extracellular enzymatic extracts

For enzyme production, the 21 selected yeast strains were inoculated in a basal medium (g/100 mL citric-citrate buffer 50 mM, pH 3.5: D-(+)-glucose, 2.0; soy peptone, 1.0; meat peptone, 1.0; yeast extract, 1.0) at 5% of pre-cultures grown in the same medium.

Pectinolytic strains were incubated with agitation (130 rpm) at 12 °C during 6 days or 28 °C during 3 days (according to previous assays, data not shown). After removing cells by

centrifugation (5000×g, 15 min at 4 °C), the supernatants were filtered through a 0.22 µm pore size membrane. These cell-free supernatants were used to assay the enzymatic activity.

2.3.2. Evaluation of pectinolytic activity by releasing of reducing sugars

Pectinolytic activity was assayed by quantification of reducing sugars released from a pectin dispersion (0.25% pectin in 50 mM citric-citrate buffer, pH 3.5) using 3,5-dinitrosalicylic acid reagent (DNS) (Miller, 1959). Galacturonic acid was used as standard (Sigma Chem. Co.). The reaction mixtures containing 0.45 mL of substrate and 0.05 mL of enzymatic extract were incubated at 12 °C or 28 °C for 30 min. The reaction was stopped by adding 0.5 mL DNS reagent. After cooling, 1.5 mL of distilled water was added and the absorbance measured at 530 nm. One unit of pectinase activity (U) was defined as the amount of enzyme required to release 1 µmol of reducing sugar per min under assay conditions. All tests were carried out in triplicate.

Total protein determination was performed using bovine serum albumin as standard (Bradford, 1976).

2.4. Pheno- and genotypic identification of cold-active-pectinase-secreting yeasts

Based on the purpose of this work only the strains selected at 12 °C as the best pectinolytic yeasts were identified at species level following the taxonomic criteria described by Kurtzman and Fell (2000) based on their morphological and physiological characteristics as well as by PCR-RFLP analysis of the ITS1-5.8S-ITS2 region from the nuclear rDNA gene complex.

PCR reaction was carried out according to protocols described by Esteve-Zarzoso et al. (1999) with some modifications using universal primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') already described by White et al. (1990). PCR products were digested with *Cfo*I, *Hin*fI and

*Hae*III restriction enzymes following the supplier's instructions. Amplified products and their restriction fragments were electrophoresed on 1.4 and 2.2% agarose gels, respectively, in 1× TAE (Tris-acetic acid-EDTA) buffer. Gels were stained with ethidium bromide, visualized, and photographed under UV light. Fragment sizes were estimated by comparison against a DNA standard (100-bp ladder). All PCR-RFLP reagents were from Promega Co. (USA).

2.5. Statistical analysis

ANOVA and Fisher LSD test with $\alpha=0.05$ were performed by mean comparison, using STATGRAPHICS Plus 5.1 (Manugistics, Rockville, MD). The data normality and variance homogeneity in the residuals were verified by modified Shapiro-Wilks and Levene's test, respectively.

3. Results and Discussion

3.1. Screening of pectinolytic activity for yeast selection

One thousand and twenty three microorganisms isolated from wine grape skins were firstly evaluated for their potential to secrete extracellular pectinases on plates. Among them, 565 (55%) showed pectinolytic activity. Finally, 96 pectinolytic yeast isolates producing the greatest clarification halos at 28 °C were pre-selected. These isolates were subjected to a secondary screening according to their capacity to grow and produce pectinases active at low temperature (12 °C) and also at 28 °C.

In the secondary screening the Dh/Dc relations at 12 °C varied between 1.5 and 6.0 units while at 28 °C this parameter differed between 1.8 and 11.2 units. In Table 1 are summarised the selected strains at 12 and 28 °C. Among the best isolates at both temperatures half of them were chosen for their largest Dh/Dc relations, and the rest of the strains were picked for their growth capacity apart from the large halo size.

Out of 96 isolates chosen in the primary selection only GM-Se-320 strain was picked either at 12 or 28 °C, the remaining isolates selected at 12 °C were different from those chosen at 28 °C, indicating that a direct relation was not maintained at both temperatures respect to the best pectinolytic yeasts.

3.2. Evaluation of pectinolytic activities by selected yeasts under proximate oenological conditions

In view of the potential application of these pectinolytic strains to winemaking process, the pectinase production and enzyme activity were assayed at pH and temperatures related to vinification. All selected yeasts showed good pectinolytic activity at pH 3.5 and 12 °C or 28 °C. As shown Fig. 1, the pectinase activity varied slightly among the isolates, although statistically significant differences were observed. According to activity level, five groups were constituted among the strains selected at 12 °C (Fig. 1A), while the yeasts chosen at 28 °C showed four clusters (Fig. 1B). Particularly, among yeasts tested at 12 °C, strains R-22 and Ch-24 exhibited the highest enzymatic activities under the test conditions (0.751 U/mL and 0.696 U/mL, respectively). The other isolates produced values between 44 and 83% of the maximum activity. On the other hand, the yeasts assayed at 28 °C showed pectinolytic activities between 0.673 and 1.698 U/mL, but most of them were similar respect to their net pectinase activity.

Yeast pectinases belonging to different environments described by other authors (Blanco et al., 2002; Moyo et al., 2003; Oliveira et al., 2006) have shown comparable (respect to values at 12 °C) or lower (1.5-2.0 times at 28 °C) activities than those secreted by yeasts studied in this work. However, conditions of enzymatic tests for strains reported in the literature were less extreme (37-45 °C and pH 5.0-5.5) than those applied in this study.

The selected strains showed pectinolytic activity in absence of pectin, suggesting the constitutive nature of these extracellular pectinases, in agreement with other authors. It has been demonstrated that yeasts such as *Kluyveromyces wickerhamii* (Moyo et al., 2003) and moulds such as *Aspergillus niger* sp. CH-Y-143 (Aguillar and Huitron, 1990), are capable of producing polygalacturonases constitutively.

Additionally, the specific enzymatic activity of selected yeasts was evaluated. At 12 °C this activity was considerably high for the strains GM-Ch-24 and GM-R-22 (31.5 and 22.7 U/mg total protein, respectively); hence, those strains exhibiting the greatest net pectinase activity (Fig. 1A) also showed good level of specific activity at low temperature. Regarding strains at 28 °C, all of them demonstrated specific activities similar to those revealed by strains at 12 °C, except for strain GM-R-13 which showed a value significantly higher (data not shown).

3.3. Pheno- and genotypic identification of cold-active-pectinase-producing strains

Pectinolytic isolates selected at 12 °C were pheno- and genotypically characterised and identified as *Aureobasidium pullulans*. All the strains showed the same profile by ITS-PCR-RFLP analysis, with a PCR product size of 600 bp (Fig. 2A) and restriction fragments of (190+180+100) bp, (150+450) bp and (290+180+130) bp for *CfoI*, *HaeIII* and *HinfI* enzymes, respectively (Fig. 2B). Similar restriction profiles were observed by Sangorrín et al. (2007). None of them fermented any of the tested sugars as a sole carbon source, whereas they assimilated most of them, with a rapid consumption of di- and tri-saccharides. Based on colony morphology and sugar assimilation, the 10 strains could be grouped in 4 phenotypic profiles, indicating the possibility of intraspecific differentiation in *A. pullulans* pectinolytic strains (data not shown).

A. pullulans is a ubiquitous species found mainly in soil, water and wood, as well as in osmotically stressed environments (Gunde-Cimerman et al., 2000). From the biotechnological point of view, this species is of commercial interest because its production of extracellular enzymes (Chi et al., 2009).

Previous studies have shown that *A. pullulans* occurs in majority on the surface of grape berries (Fleet, 2003). This species was isolated from immature, mature, damaged and undamaged grapes in different viticultural regions worldwide (Prakitchaiwattana et al., 2004; Renouf et al., 2007; Urso et al., 2008). On the other hand, this species is established as a very good producer of extracellular pectinases (Manachini et al., 1988). According to Biely et al. (1996), its pectinolytic systems are inducible in carbon starvation conditions which might be the case on the grape berry surface.

Although it is known that *A. pullulans* is a species able to produce pectinolytic enzymes, few works have studied pectinase production or activity in *A. pullulans* at pH values close to 4 (Galiotou-Panayotou et al., 1998) or lower than 3.5 (Biely et al., 1996). Our study confirms enzymatic activity at pH 3.5. Nevertheless, with regards to temperature conditions, the literature reports data on the pectinase production by *A. pullulans* or its enzymatic activity at temperature values higher than 25 °C. Therefore, to our knowledge, it is the first report about *A. pullulans* displaying activities at low temperature (12 °C). Taking into account that mentioned conditions are relevant for winemaking, pectinolytic strains selected in the present study could be of interest in wine production.

In addition, it is important to emphasize that, apart from *A. pullulans*, we did not find any other yeast species among the best cold-active-pectinase-producing microorganisms isolated from grape skins in San Rafael D.O. This fact is in agreement with prior studies performed by Charoenchai et al. (1997), who reported that pectinolytic activity was not

found in any of the wine yeasts studied. Nevertheless, various species of *Kluyveromyces*, *Candida*, *Cryptococcus* and strains of *S. cerevisiae*, have been reported to possess pectinolytic activity (Blanco et al., 1994; da Silva et al., 2005; Moyo et al., 2003).

In conclusion, this preliminary study showed that some strains of *A. pullulans* produced pectinolytic enzymes that are able to act effectively at some wine conditions such as low temperature and low pH. Further studies on pectinase characterisation as well as the ability of these strains to degrade pectin *in vivo* are needed. Indeed, microfermentation assays are being carried out with the aim of demonstrating that selected *A. pullulans* strains are appropriate for being incorporated in wine making. These investigations are evaluating their co-culture with commercial strains of *S. cerevisiae* and their impact on the sensory quality of the wine, such as flavour and colour.

Acknowledgments

This research was supported by PIP N° 6472 (CONICET), UNCuyo N° 06/L096 project and I+D (UNCuyo) programme grants.

References

- Aguillar, G., Huitron, C., 1990. Constitutive exo-pectinase produced by *Aspergillus* sp. CH-Y-1043 on different carbon sources. *Biotechnology Letters* 12, 655-660.
- Biely, P., Heinrichová, K., Kružiková, M., 1996. Induction and inducers of the pectinolytic system in *Aureobasidium pullulans*. *Current Microbiology* 33, 6-10.
- Birgisson, H., Delgado, O., García Arroyo, L., Hatti-Kaul, R., Mattiasson, Bo, 2003. Cold-adapted yeasts as producers of cold-active polygalacturonases. *Extremophiles* 7, 185-193.
- Blanco, P., Sieiro, C., Diaz, A., Villa, T.G., 1994. Production of an endopolygalacturonase from *Saccharomyces cerevisiae*. *Canadian Journal of Microbiology* 40, 974-977.

- Blanco, P., Thow, G., Simpson, C.G., Villa, T.G., Williamson, B., 2002. Mutagenesis of key amino acids alters activity of a *Saccharomyces cerevisiae* endo-polygalacturonase expressed in *Pichia pastoris*. FEMS Microbiology Letters 210, 187-191.
- Bradford, M. M., 1976. A rapid and sensitive method for quantification of microgram quantities of protein utilizing the principle of protein-dye binding. Analytical Biochemistry 72, 248-254.
- Cabeza, M. S., Merín, M. G., Martín, M. C., Sabaté, D. C., Audisio, M. C., Morata, V. I., 2009. Effect of pectinase-surfactin system on pigments and total polyphenols extraction from Malbec grape skins. American Journal of Enology and Viticulture 60 (4), 477-483.
- Charoenchai, C., Fleet, G.H., Henschke, P.A., Todd, B.E.N., 1997. Screening of non-*Saccharomyces* wine yeasts for the presence of extracellular hydrolytic enzymes. Australian Journal of Grape and Wine Research 3, 2-8.
- Chi, Z.M., Wang, F., Chi, Z., Yue, L., Liu, G., Zhang, T., 2009. Bioproducts from *Aureobasidium pullulans*, a biotechnologically important yeast. Applied Microbiology and Biotechnology 2, 793-804.
- da Silva, E.G., Borges, M.F., Medina, C., Piccoli, R.H., Schwan, R.F., 2005. Pectinolytic enzymes secreted by yeasts from tropical fruits. FEMS Yeast Research 5, 859-865.
- Esteve-Zarzoso, B., Belloch, C., Uruburu, F., Querol, A., 1999. Identification of yeasts by RFLP analysis of the 5.8S rRNA gene and the two ribosomal internal transcribed spacers. International Journal Systematic Bacteriology 49, 329-337.
- Fernandes-Salomão, T.M., Rodrigues Amorim, A.C., Chaves-Alves, V.M., Cavalcante Coelho, J.L., Olzany Silva, D., Fernandes de Araújo, E., 1996. Isolation of pectinase hyperproducing mutants of *Penicillium expansum*. Revista Microbiologia, São Paulo 27 (1), 15-18.

- Fernández, M., Úbeda, J.F., Briones, A.I., 2000. Typing of non-*Saccharomyces* yeasts with enzymatic activities of interest in wine-making. *International Journal of Food Microbiology* 59, 29-36.
- Fernández-González, M., Úbeda, J.F., Cordero Otero, R.R., Thanvanthri Gururajan, V., Briones, A.I., 2005. Engineering of an oenological *Saccharomyces cerevisiae* strain with pectinolytic activity and its effect on wine. *International Journal of Food Microbiology* 102, 173-183.
- Fleet, G.H., 2003. Yeast interactions and wine flavour. *International Journal of Food Microbiology* 86, 11-22.
- Galiotou-Panayotou, M., Kalantzi, O., Aggelis, G., 1998. Modelling of simultaneous production of polygalacturonase and exopolysaccharide by *Aureobasidium pullulans* ATHUM 2915. *Antonie van Leeuwenhoek* 73, 155-162.
- Gunde-Cimerman, N., Zalar, P., de Hoog, S., Plemenitas, A., 2000. Hypersaline waters in salterns-natural ecological niches for halophilic black yeasts. *FEMS Microbiology Ecology* 32, 235-240.
- Kashyap, D.R., Vohra, P.K., Chopra, S., Tewari, R., 2001. Applications of pectinases in the commercial sector: a review. *Bioresource Technology* 77, 215-227.
- Kurtzman C.P., Fell, J.W., 2000. *The Yeast: A Taxonomic Study*. 4th edition-Fourth revised and enlarged Edition. Elsevier Science Publisher, Amsterdam.
- Manachini, P.L., Parini, C., Fortina, M.G., 1988. Pectic enzymes from *Aureobasidium pullulans* LV 10. *Enzyme and Microbial Technology* 10, 682-685.
- Miller, G.L. 1959. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry* 31, 426-428.

- Molina, A., Swiegers, J.H., Varela, C., Pretorius, I.S., Agosin, E., 2007. Influence of wine fermentation temperature on the synthesis of yeast-derived volatile aroma compounds. *Applied Microbiology and Biotechnology* 77, 675-687.
- Moyo S., Gashe, B.A., Collison, E.K., Mpuchane, S., 2003. Optimising growth conditions for the pectinolytic activity of *Kluyveromyces wickerhamii* by using response surface methodology. *International Journal of Food Microbiology* 85, 87-100.
- Nakagawa, T., Nagaoka, T., Taniguchi, S., Miyaji, T., Tomizuka, N., 2004. Isolation and characterization of psychrophilic yeasts producing cold-adapted pectinolytic enzymes. *Letters in Applied Microbiology* 38, 383-387.
- Oliveira, K. F., Malavolta, L., Souza, C. S., Vicente, E. J., Laluce, C., 2006. Pectinolytic activity secreted by yeasts isolated from fermented citrus molasses. *Journal of Applied Microbiology* 100, 633-640.
- Prakitchaiwattana, C.J., Fleet, G.H., Heard, G.M., 2004. Application and evaluation of denaturing gradient gel electrophoresis to analyse the yeast ecology of wine grapes. *FEMS Yeast Research* 4, 865-877.
- Renouf, V., Claisse, O., Lonvaud-Funel, A., 2007. Inventory and monitoring of wine microbial consortia. *Applied Microbiology and Biotechnology* 75, 149-164.
- Sangorrín, M.P., Lopes, C.A., Giraudo, M.R., Caballero, A.C., 2007. Diversity and killer behaviour of indigenous yeasts isolated from the fermentation vat surfaces in four Patagonian wineries. *International Journal of Food Microbiology* 119, 351-357.
- Strauss, M.L.A., Jolly, N.P., Lambrechts, M.G., van Rensburg, P., 2001. Screening for the production of extracellular hydrolytic enzymes by non-*Saccharomyces* wine yeasts. *Journal of Applied Microbiology* 91, 182-190.

- Urso, R., Rantsiou, K., Dolci, P., Rolle, L., Comi, G., Cocolin, L., 2008. Yeast biodiversity and dynamics during sweet wine production as determined by molecular methods. *FEMS Yeast Res* 8, 1053-1062.
- Van Rensburg, P., Pretorius, I.S., 2000. Enzymes in winemaking: harnessing natural catalysts for efficient biotransformations – A review. *South African Journal of Enology and Viticulture* 21, 52-73.
- White, T.J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungi ribosomal RNA gene for phylogenetics. In: Innis, M.A., Gelfand, D.H. Sninsky, J.J., White, T.T. (Eds.), *PCR protocols. A guide of methods and applications*. Academic Press, San Diego, pp. 315-320.

Figure captions

Figure 1. Pectinolytic activity at 12 °C (A) and 28 °C (B) (means \pm SD, n = 3).

Different lowercase letters mean significant differences, according to the least significant difference (LSD) test ($p < 0.05$). One unit of pectinase activity (U) was defined as the amount of enzyme required to liberate 1 μ mol of reducing sugar per min at pH 3.5.

Figure 2. PCR amplified products of 5.8S rDNA-ITS region (A) and restriction fragments (B) of selected cold-active pectinolytic microorganisms. Lanes 1-10: from left to right PN-321, Ma-16, PN-34, PN-22, CS-123, R-22, Cy-25, Cy-126, Ch-24 and Se-320 strains. MW: 100 bp DNA ladder. The restriction enzymes used in each case are indicated at the top of the picture. It is shown one strain as representative profile of the 10 cold-active pectinolytic microorganisms.

Table 1. Pectinolytic strains selected at 12 °C and 28 °C based on the Diameter halo/Diameter colony relation (Dh/Dc) and colony size (Dc), resulting from screening in solid medium containing pectin.

Strain	Dh (mm)	Dc ^a (mm)	Dh/Dc ^a
12 °C			
PN-321	21	3.5	6.00
Ma-16	11	2.5	4.40
PN-34	19	4.5	4.22
PN-22	21	5.0	4.20
CS-123	12	3.0	4.00
R-22	19	6.0	3.17
Cy-25	17	6.0	2.83
Cy-126	19	7.0	2.71
Ch-24	15	7.0	2.14
Se-320	18	9.0	2.00
28 °C			
Cy-322	28	2.5	11.20
R-28	33	4.0	8.25
Cy-318	20	2.5	8.00
R-13	40	5.5	7.27
N-323	23	3.5	6.57
Se-320	36	6.0	6.00
R-35	30	6.0	5.00
M-113	46	12.0	3.83
Ma-217	30	9.0	3.33
Se-130	29	9.5	3.05
Se-222	30	10.0	3.00

^a Numbers in bold indicate the criterion followed in the selection at both temperatures.




Fig. 1




Fig. 2

Research Highlights

We screened wild yeasts from wine grape ecosystem secreting cold-active pectinases. > We studied pectinase activity under pH and temperature oenological conditions. > *A. pullulans* strains were selected as the best cold-active-pectinase producers. > Good level of pectinolytic activity and constitutive enzyme production were shown. > Enzyme activity at tested vinification conditions is relevant for wine production.

ACCEPTED MANUSCRIPT