



BRIEF REPORT

## Isolation of saprophytic *Leptospira* spp. from a selected environmental water source of Argentina



Exequiel Scialfa<sup>a,c,\*</sup>, Sylvia Grune<sup>b,d</sup>, Bibiana Brihuega<sup>b</sup>, Pablo Aguirre<sup>a</sup>, Marina Rivero<sup>c</sup>

<sup>a</sup> División Zoonosis Rurales, España 770, Azul 7300, Provincia de Buenos Aires, Argentina

<sup>b</sup> Instituto de Patobiología, CCVyA-CNIA, INTA, CC 77, 1708 Morón, Provincia de Buenos Aires, Argentina

<sup>c</sup> Universidad Nacional del Centro de la Provincia de Buenos Aires, Provincia de Buenos Aires, Argentina

<sup>d</sup> National Research Council of Argentina (CONICET), Buenos Aires, Argentina

Received 2 November 2016; accepted 8 August 2017

Available online 29 November 2017

### KEYWORDS

*Leptospira* spp.;  
Isolation;  
Water;  
Argentina

**Abstract** Ten *Leptospira* spp. strains were isolated from water samples from Nievas stream, Olavarría, Buenos Aires province (Argentina). The isolates showed the typical motility and morphology of the genus *Leptospira* under dark field microscopy, developing in liquid EMJH medium after eight days of incubation at 13 °C and 30 °C. All isolates were negative by the Multiple Locus Variable Number Tandem Repeat Analysis (MLVA). Molecular identification by 16S rRNA gene sequencing identified all isolates as nonpathogenic leptospires. Four isolates showed a genetic profile identical to that of the reference strain *Leptospira biflexa* serovar Patoc, and six isolates revealed sequence similarities within the 97–98% range, closely related to *Leptospira yanagawae* and *Leptospira meyeri*, respectively. Strains ScialfaASA42, ScialfaASA45, ScialfaASA44, ScialfaASA47, ScialfaASA49, ScialfaASA50 and ScialfaASA51 possibly represent a novel species of the genus *Leptospira*.

© 2017 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### PALABRAS CLAVE

*Leptospira* spp.;  
Aislamiento;  
Agua;  
Argentina

**Aislamiento de *Leptospira* spp. saprófitas desde fuentes de agua en un ambiente seleccionado de Argentina**

**Resumen** Se aislaron 10 cepas de *Leptospira* spp. a partir de muestras de agua del arroyo Nievas, partido de Olavarría (provincia de Buenos Aires, Argentina). Los aislamientos mostraron motilidad y morfología típica del género *Leptospira* bajo microscopía de campo oscuro y se desarrollaron en medio líquido EMJH después de 8 días de incubación a 13 y 30 °C. Todos los

\* Corresponding author.

E-mail address: [escialfa@yahoo.com.ar](mailto:escialfa@yahoo.com.ar) (E. Scialfa).

aislamientos fueron negativos por MLVA, y mediante la secuenciación del gen 16S del ARNr se identificaron como leptospiras no patógenas. Cuatro de estos aislamientos mostraron un perfil genético idéntico a la cepa *Leptospira biflexa* serovar Patoc de referencia, en tanto que 6 de ellos presentaron similitudes de secuencias estrechamente relacionadas con las especies *Leptospira yanagawae* y *Leptospira meyeri* dentro del intervalo del 97 y 98%, respectivamente. Las cepas ScialfaASA42, ScialfaASA45, ScialfaASA44, ScialfaASA47, ScialfaASA49, ScialfaASA50 y ScialfaASA51 posiblemente representen una nueva especie del género *Leptospira*.

© 2017 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Leptospirosis is a bacterial zoonotic disease caused by pathogenic species of the genus *Leptospira*, which is mainly transmitted to humans by direct contact with rodents and domestic animals, mainly dogs and cattle. However, human outbreaks are frequently produced by contact with water that is contaminated with urine of animal reservoirs<sup>9,13</sup>; and leptospirae penetrate abraded skin or mucous membranes and disseminate through the organism. The genus *Leptospira* includes at least 22 species arranged into three large subgroups based on 16S rRNA phylogeny, ten (10) pathogenic species, seven (7) saprophytic species and five (5) intermediate species<sup>2,6</sup>. In Argentina, there are few studies of leptospirae in environmental sources, mainly from water.

In this paper, we report the circulation of two non-pathogenic species of the genus *Leptospira* isolated from water samples from Nievas stream, Olavarría, Argentina.

Initially, leptospiral strains were isolated from water samples from Nievas stream, Olavarría, Buenos Aires province, Argentina (Fig. 1) by following isolation procedures. Water samples were collected in a recreational area during March 2015, and transported in sterile 500 ml glass bottles. Temperature and pH were monitored in the field. Water samples were filtered through a sterile membrane. In this study, a prefiltering technique was applied using Whatman filter paper before filtration through a 0.22 µm pore size membrane filter. A sample of the filtered water (1 ml) was inoculated into Ellinghausen–McCullough–Johnson–Harris (EMJH medium: Difco Laboratories, Detroit, Michigan, USA) liquid medium with the addition of 5-fluorouracil (200 µg/ml) as selective antimicrobial agent. Cultures were incubated by duplicate at 13 °C and 30 °C for 90 days, and leptospiral growth was monitored weekly using dark field microscopy. On day eight, growth was observed at both temperatures, and ten strains were isolated.

DNA templates were obtained using Chelex Resin-100 (Bio Rad). Multiple Locus Variable Number Tandem Repeat Analysis (MLVA) was performed using two sets of oligonucleotides specific for pathogenic leptospirae (*Leptospira interrogans*, *Leptospira kirschneri* and *Leptospira borgpetersenii*). Oligonucleotides that hybridized to the flanking regions of the VNTR4, VNTR7, VNTR9, VNTR10, VNTR19, VNTR23 and VNTR31 loci were used to discriminate strains of *L. interrogans* and oligonucleotides that hybridized to the flanking regions of the VNTR4, VNTR7, VNTR10; Lb4 and Lb5 loci were used for *L. kirschneri*, *L. borgpetersenii* and *L. interrogans* strains<sup>7,8</sup>. The final volume (50 µl) of each reaction mixture contained polymerase chain reaction (PCR) buffer (20 mM Tris–HCl, pH 8.4, 50 mM KCl),

200 µM deoxynucleoside triphosphates, 2 µM each of the corresponding forward and reverse primers, 2 mM MgCl<sub>2</sub>, 1.25 U of *Taq* DNA polymerase (Invitrogen) and 5 µl of DNA template. PCR amplifications were carried out in a Thermo Scientific PxE 0.2 Thermal Cycler, using the following cycling parameters: 94 °C for 5 min, followed by 35 cycles of denaturation at 94 °C for 30 s, annealing at 55 °C for 30 s and extension at 72 °C for 90 s, with a final cycle at 72 °C for 10 min. The amplified samples were examined by electrophoresis in ethidium bromide-containing 2% agarose gels in TAE buffer (40 mM Tris-acetate, 1 mM EDTA, pH 8.0) at 100 V for 50 min. Amplified DNA bands were visualized upon ultraviolet light exposure (Uvi Tec transilluminator BTS-20.M, Manufacturer UviTec, St. John's Innovation Centre, Cowley Road, Cambridge, England). Amplicon sizes were estimated using CienMarker (Biodynamics) and the GelAnalyzer 2010 program. To calculate repeat copy numbers, the following formula was used: number of repeats (bp) = [(fragment size (bp) – flanking regions (bp)]/repeat size (bp). Repeat copy numbers were rounded down to the closest whole number. If the copy number was less than one, it was rounded to zero.

PCR targeting the 16S rRNA gene was carried out for bacterial identification. The following primers were used: 5'-GGCGGCGCTCTTAAACATG-3' and 5'-GTCCGCCTACGCACCCTTACG-3'; these primers have the ability to amplify all pathogenic and nonpathogenic leptospirae<sup>4</sup>. After verification of the amplicon by electrophoresis (in an ethidium bromide-containing 2% agarose gel) and visualization upon UV light exposure, PCR products were purified using a commercial kit (Embiotech). The sample was sequenced at the Institute of Biotechnology, National Institute of Agricultural Technology (Argentina) using a 3130xl Genetic Analyzer (Applied Biosystems). For alignment and construction of the phylogeny, the program MEGA version 5 was used<sup>12</sup>. The dendrogram based on partial sequences of the 16S rRNA gene was constructed using Neighbor-joining with a bootstrap of 100 (Fig. 2).

At the moment of water sample collection in Nievas stream, we observed a temperature of 24 °C and pH 7.4. Growth was observed after 8 days in culture medium at 13 °C and 30 °C, and the isolates showed the typical motility and morphology of the genus *Leptospira* sp. (cells of 10–13 µm in length and less than 0.22 µm in diameter, characterized by flexuous motility) under dark field microscopy. Strains were maintained in Ellinghausen–McCullough–Johnson–Harris (EMJH) liquid medium.



Figure 1 (A) Colony San Miguel, Olavarría, Buenos Aires province, Argentina. (B) Recreational area of Nievas stream.




Figure 2 Neighbor-joining analysis of the sequence 16S rRNA of all strains isolated. Phylogenetic analysis based of 16S rRNA including 19 representative species of *Leptospira*. The corresponding sequence of *Leptonema illini* strain DSM 21528 was used as an outgroup. The dendrogram was constructed using the Neighbor-joining method. Bootstrap values are displayed as percentages.

All isolates were negative by the MLVA; however, molecular identification by 16S rRNA gene sequencing identified all isolates as non-pathogenic leptospires. Four isolates showed a genetic profile identical to that of the reference

*L. biflexa* serovar Patoc (strains ScialfaASA43, ScialfaASA46, ScialfaASA48 and ScialfaASA52), and six isolates (strains ScialfaASA42, ScialfaASA45, ScialfaASA44, ScialfaASA47, ScialfaASA49, ScialfaASA50 and ScialfaASA51) with sequence

similarities within the 97–98% range using Blast (results not shown), and closely related to species *Leptospira yanagawae* and *Leptospira meyeri* respectively. Strains ScialfaASA42, ScialfaASA45, ScialfaASA44, ScialfaASA47, ScialfaASA49, ScialfaASA50 and ScialfaASA51 possibly represent a novel species of the genus *Leptospira*. The sequence obtained from the strain ScialfaASA42 is available in GenBank under Accession number KT160428.

In Argentina isolates from water sources (rivers, lakes, and streams) have been obtained in Buenos Aires, Santa Fe, Corrientes and Jujuy province. In the period 1961–1996 a total of 47 saprophytes strains and 5 pathogenic strains were isolated from water (two unclassified *L. interrogans*, one *L. interrogans* serovar Icterohaemorrhagiae and one *L. interrogans* serovar Pomona)<sup>1</sup>. Brihuega et al.<sup>3</sup> isolated a pathogenic strain from water samples from Reconquista river, which was classified as *L. interrogans* serovar Icterohaemorrhagiae strain Reconquista. In the year 2011, five strains were isolated from water samples from rural channels in Casilda, Santa Fe province, Argentina, and one was classified as *L. borgpetersenii*<sup>5</sup>. To the best of our knowledge, eight pathogenic *Leptospira* sp. strains have been isolated so far from environmental water sources in Argentina. In the center of Buenos Aires province, there have been no reports of pathogenic leptospires from water sources. However, in previous studies on wild animals captured in the region (*Rattus norvegicus*, *Akodon azarae*, *Lycalopex griseus*, *Conepatus chinga* and *Didelphis albiventris*), we showed the circulation of several strains belonging to *L. interrogans* serogroup Icterohaemorrhagiae: serovar Copenhageni strains Fiocruz L1-130 and M20, belonging to serovar Icterohaemorrhagiae RGA (in peri-urban and rural area). Furthermore, strains belonging to *L. interrogans* serogroup Canicola: serovar CanicolaHound Utrecht IV (in rural area) were found<sup>9–11</sup>. In the present study, we showed that in the selected environment, especially in water, conditions (pH and temperature) are optimal for the survival of non-pathogenic and pathogenic leptospires. In water samples from Nievas stream in Olavarria, Buenos Aires province, we observed the circulation of two saprophytic species of the genus *Leptospira* (*L. biflexa* serovar Patoc and *Leptospira* sp. with 97–98% sequence similarities to *L. yanagawae* and *L. meyeri* respectively). The MLVA assay and the molecular characterization of the 16S rRNA gene sequencing were important tools for the identification of these isolates as non-pathogenic leptospires; however, they were insufficient to define strains ScialfaASA42, ScialfaASA45, ScialfaASA44, ScialfaASA47, ScialfaASA49, ScialfaASA50 and ScialfaASA51 as known saprophytic species of the genus *Leptospira*. Further studies such as pulsed-field gel electrophoresis: PFGE; DNA–DNA hybridization, average nucleotide identity: ANI and genome-to-genome distance calculator, GGD, should be conducted to further study these isolated strains.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this study.

**Confidentiality of data.** The authors declare that no patient data appear in this article.

**Right to privacy and informed consent.** The authors declare that no patient data appear in this article.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## References

- Asociación Argentina de Veterinarios de Laboratorios de Diagnóstico (AAVLD). Comisión Científica sobre Leptospirosis de la República Argentina. Informe sobre Leptospirosis en la República Argentina. Serie Enfermedades Transmisibles. Buenos Aires, Argentina: Editado por Fundación Mundo Sano; 2006, 18 pp.
- Bourhy P, Collet L, Brisse S, Picardeau M. *Leptospira mayottensis* sp. nov., a pathogenic species of the genus *Leptospira* sp. isolated from humans. *Int J Syst Evol Microbiol*. 2014;64:4061–7.
- Brihuega B, Auteri C, Romero G, Samartino L. Aislamiento de una cepa patógena de *Leptospira* de un río urbano: Respuesta frente a quinolonas fluoradas. *Rev Med Vet*. 2006;87:144–6.
- Djadid ND, Ganji ZF, Gouya MM, Rezvani M, Zakeri S. A simple and rapid nested polymerase chain reaction-restriction fragment length polymorphism technique for differentiation of pathogenic and nonpathogenic *Leptospira* spp. *Diagn Microbiol Infect Dis*. 2009;63:251–6.
- Francois BS, Brihuega B, Grune S, Gattarello MV, Coreea PD, Petrakovsky MJ, Gualtieri SC. Aislamiento de *Leptospira borgpetersenii* de fuentes de agua en Argentina. *Rev Cubana Med Trop*. 2013;65:177–84.
- Gravekamp C, Van De Kemp H, Franzen M, Carrington D, Schoone GJ, Van Eys GJJM. Detection of seven species of pathogenic leptospires by PCR using two sets of primers. *J Gen Microbiol*. 1993;139:1691–700.
- Majed Z, Bellenger E, Postic D, Pourcel C, Baranton G, Picardeau M. Identification of variable-number tandem-repeat loci in *Leptospira interrogans* sensu stricto. *J Clin Microbiol*. 2005;43:539–45.
- Pavan ME, Cairo F, Pettinari MJ, Samartino L, Brihuega B. Genotyping of *Leptospira interrogans* strains from Argentina by multiple-locus variable-number tandem repeat analysis (MLVA). *Comp Immunol Microbiol Infect Dis*. 2011;34:135–41.
- Scialfa E, Bolpe J, Bardón JC, Ridao G, Gentile J, Gallicchio O. Isolation of *Leptospira interrogans* from suburban rats in Tandil, Buenos Aires, Argentina. *Rev Argent Microbiol*. 2010;42:126–8.
- Scialfa E, Brihuega B, Morris W, Recavarren M, Grune S, Quintana S, Romero G, Bolpe J, Schettino M. First isolation of *Leptospira interrogans* from *Conepatus chinga* in Argentina. *Afr J App Microbiol Res*. 2012;1:1–5.
- Scialfa E, Brihuega B, Venzano A, Morris W, Bolpe J, Schettino M. First isolation of *Leptospira interrogans* from *Lycalopex griseus* (South American gray fox) in Argentina shows new MLVA genotype. *J Wildl Dis*. 2013;49:168–72.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA 5 Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731–9.
- Vanasco NB, Schmeling MF, Lottersberger J, Costa F, Ko AI, Tarabla HD. Clinical characteristics and risk factors of human leptospirosis in Argentina (1999–2005). *Acta Trop*. 2008;107:255–8.