
MYCOTAXON

A new species of *Scopinella* from pampas grass in Argentina

ROMINA M. SÁNCHEZ, LUCRECIA GIORDANO, FRED A. ANDERSON & M. VIRGINIA BIANCHINOTTI*

*Centro de Recursos Naturales Renovables de la Zona Semiárida-Universidad Nacional del Sur
Camino La Carrindanga Km 7, B8000FWB, Bahía Blanca, Argentina*

* CORRESPONDENCE TO: vbianchi@uns.edu.ar

ABSTRACT — A previously undescribed species of the genus *Scopinella* was found during a survey of the fungi associated with *Cortaderia selloana* (*Poaceae*) in Argentina. The ascospores are strongly asymmetrical and have the shape of a truncated pyramid with a convex quadrangular base. This peculiar ascospore shape sets this species apart from the others of the genus and is therefore described here as new. This is the second record of the genus from South America.

KEY WORDS — ascomycete, grasses, saprophytic fungi

Introduction

Cortaderia selloana (pampas grass) is a perennial tussock-forming grass species that is native to Argentina, Brazil, Chile, Paraguay and Uruguay (Astegiano et al. 1995). Its huge tussocks are very conspicuous and become attractive when they flower to produce showy feathery inflorescences. It has been introduced in many regions of the world as a garden plant, for dryland forage, soil amendment, and as windbreaks. In some of these, it has become invasive (Pausas et al. 2006). In particular, it has become a serious weed in New Zealand, where the feasibility of conducting biological control is being explored (Bellgard et al. 2010).

Most of the records of fungal species found associated with *C. selloana* are the result of studies performed in the countries where it has been introduced (Bellgard et al. 2010, Farr & Rossman 2012, McKenzie et al. 2007, Medd et al. 2003). Studies are now being conducted in Argentina to learn about the associated mycobiota present in its native distribution area.

During a survey of the fungi associated with *C. selloana* leaves in Buenos Aires province, a new species of the genus *Scopinella* Lév. was discovered.

The genus *Scopinella* is characterized by ascomata with long necks and globose bodies, evanescent asci, and cuboid-ellipsoidal ascospores with two prominent longitudinal germ slits (Cannon & Hawksworth 1982, Stchigel et al. 2006). It is a small genus that comprises seven species that grow either on fungi or plant substrates. It belongs to *Hypocreales*, where at present it cannot be assigned to any family (Lumbsch & Huhndorf 2010).

The shape of the ascospores of the specimen found on *C. selloana* distinguishes it from the other species of the genus, so we describe it here as new.

Materials & methods

Partially senescent leaves of *Cortaderia selloana* were carefully inspected under a Wild M5A stereomicroscope. Sections bearing fruiting bodies were hand-made with a razor blade and mounted in either distilled water alone or with the addition of phloxine. Measurements were made in water. A Leica DM2000 dissecting microscope with a Leica EC3 camera was used to capture micromorphological images. For scanning electron microscopy intact ascomata were dehydrated in a graded acetone series from 10% to 90% , and finally 95% for fifteen minutes in each step, followed by three changes in absolute acetone. After critical point-drying, the specimens were mounted on aluminum stubs, sputter coated with gold-palladium, and viewed with a LEO EVO 40 Scanning Electron Microscope (SEM). Recently collected leaves with the fungus were pressed and preserved in “Bahía Blanca Biología” Herbarium (BBB). Herbarium is abbreviated according to Thiers (2012).

Scopinella pyramidospora R.M. Sánchez, L. Giord., F. Anderson & Bianchin., **sp. nov.** PLATE 1

MYCOBANK MB800633

DIAGNOSIS: Differs from all other *Scopinella* species by its strongly asymmetrical, pyramidal ascospores.

TYPE: Argentina, Buenos Aires Province, Provincial Route No. 205, Km 308, (36°06'43"S 60°58'22"W): on senescent leaves of *Cortaderia selloana* (Schult. & Schult. f.) Asch. & Graebn. (*Poaceae*), 4.VII.2011, coll. FE Anderson (**HOLOTYPE**, BBB [MVB 1501]).

ETYMOLOGY: from the Greek πυραμίδος = pyramidal and σπόρος: spore.

ASCOMATA solitary to gregarious, superficial, settled on a dense mass of brown hyphae, globose to subglobose, with long necks, brown to black when dry, brown to paler brown at base when mounted in water, 106–150 µm high, 106–225 µm wide, peridial cells forming *textura intricata*. NECKS cylindrical, 206–313 µm long, 31–75 µm wide at base, ostiolum 19–38 µm wide, cells of neck disposed in *textura porrecta*. PARAPHYSES moniliform, ramified, septate, hyaline, 4–7 µm wide. ASCI 8-spored, unitunicate, clavate to globose, short-stipitate, evanescent, 13–28 × 10–16 µm. ASCOSPORES 1-celled, strongly

asymmetrical, resembling a truncated pyramid with quadrangular, somewhat convex, base, rounded vertices, smooth-walled at optical level, verruculose under SEM ($>7,000\times$) with a dark brown median band, paler at ends, germ slits occasionally present, $7.5\text{--}9.5 \times 5.5\text{--}8.5 \mu\text{m}$ ($x = 8.4 \times 6.7$, $n = 60$). ANAMORPH not known.

DISTRIBUTION— Argentina, known only from the type collection.

COMMENTS— *Scopinella pyramidospora* differs from all other species of the genus in having strongly asymmetrical, pyramidal ascospores. Ascospore asymmetry is best visualized by moving the microslide in three dimensions when mounted in water. Germ slits were seen only once under optical microscopy. This suggests that they are not as conspicuous and/or common as in the other species of the genus. Although the spore size falls in the range of *S. caulicola* (Fuckel) Malloch, ascospores of the latter are strongly flattened bilaterally and almost rectangular in surface view (as shown in von Arx & Müller 1954). The ascomata of *S. caulicola* have much longer necks, up to 2 mm long, with subhyaline short bristles at the apex (fide von Arx & Müller 1954). Ascomatal necks of *S. pyramidospora* are glabrous.

Discussion

The genus *Scopinella* was erected by L veill  (1847) to accommodate the single species *S. barbata* (Pers.) L v. Saccardo (1891) added a second species, *S. pleiospora* (J. Schr t.) Sacc., which had originally been described as *Melanospora pleiospora*. The genus was then ignored for almost a century until Hawksworth (1975) reestablished it by recognizing the synonymy between *S. barbata* and *Chaetoceratostoma hispidum* Turconi & Maffei. Hawksworth (1975) considered *Scopinella* to be monospecific and did not include Saccardo's *Scopinella pleiospora*. Malloch (1976a,b) added three species to the genus (*S. caulicola*, *S. solani*, and *S. sphaerophila*) and provided a key to differentiate these additions and the original *S. barbata*. Two more species were added, *S. gallicola* by Tsuneda & Hiratsuka (1981) and *S. musciformis* by Stchigel et al. (2006). No reference was made to *Scopinella pleiospora* in any of these studies. We failed to access type or authenticated material, and presume the type material of *S. pleiospora* has been lost.




PLATE 1: *Scopinella pyramidospora*: **1.** Ascomata on the substrate (arrows). **2.** Close-up of an ascoma. **3.** Apex of the ascomata neck (SEM, 4500 \times). **4.** Eight-spored ascus. **5.** Asymmetrical ascospores in various views. **6, 7.** Ascospore germ slits (arrows). **8.** Ascospores (SEM, 7000 \times). **9.** Lateral view of a pyramidal ascospore (SEM, 10000 \times). **10.** Detail of ascospore wall (SEM, 20000 \times). Bars: 1 = 200 μ m; 2 = 50 μ m; 3 = 10 μ m; 4, 6, 7 = 5 μ m; 5 = 7 μ m; 8 = 4 μ m; 9 = 3 μ m; 10 = 2 μ m.

Species of *Scopinella* fall into two ecological groups. One comprises species growing on other fungi (*S. gallicola* and *S. sphaerophila*), and the other saprobic species, usually growing on plant substrates (*S. barbata*, *S. caulicola*, *S. musciformis*, *S. pyramidospora*, and *S. solani*). *Scopinella pleiospora* was described growing on rabbit dung. *Scopinella pyramidospora* and *S. solani* are the only two species of the genus known to occur on grasses. *Scopinella solani* was found growing on dead leaves of *Poa nevadensis* in North America (Malloch 1976a) and on overwintered

inflorescences of *Brachypodium pinnatum*, *Dactylis glomerata*, and *Elytrigia repens* in Slovakia (Pastirčák & Pastirčáková 2007). *Scopinella solani* differs from *S. pyramidospora* in having longer ascomatal necks (300–500 µm), narrower asci (7–11.5 µm), and symmetrical and smaller ascospores (4.5–6.5 × 3.8–5.8 µm).

Species of *Scopinella* are difficult to cultivate, and the species from *Cortaderia selloana* was no exception. All attempts to isolate it on artificial media failed. Because of this, and the fact that the available material is at present very scarce, it was decided against performing any molecular work. To our knowledge the only species that have been cultivated are *S. gallicola* (Tsuneda & Hiratsuka 1981) and *S. solani* (Zhang & Blackwell 2002). Tsuneda & Hiratsuka (1982) found that for *S. gallicola* both growth and fructification were stimulated in dual culture with other fungi.

Phylogenetic studies have been performed on *S. solani* which is the only species that has been sequenced. Zhang & Blackwell (2002) set it in a distinct clade in *Hypocreales* apart from *Melanospora* in contrast with what was previously accepted (Hawksworth 1975).

The genus *Scopinella* was reported in Argentina and South America for the first time with the finding of *S. solani* on bark of *Geoffroea decorticans* (Bianchinotti 1998). This is the second record of the genus from this region.

Acknowledgments

This work was funded by “National Biocontrol Collective” and “Future Forests Research”, New Zealand. Lucrecia Giordano is a fellow at CIC. MVB is a researcher at CONICET. Amy Rossman and Alberto Stchigel are thanked for acting as pre-submission reviewers and for their helpful comments.

Literature cited

- Arx JA von, Müller E. 1954. Die Gattungen der amerosporen Pyrenomyceten. Beitr. Kryptfl. Schweiz 11 (1): 1–434.
- Astegiano ME, Anton AM, Connor HE. 1995. Sinopsis del género *Cortaderia* (Poaceae) en Argentina. Darwiniana 33: 43–51.
- Bellgard SE, Winks CJ, Than DJ, Aliaga CC. 2010. Natural enemies of the South American pampas grasses *Cortaderia* spp. in New Zealand. 215–218, in SM Zydembos (ed.), Proceedings of the 17th Australasian Weeds Conference. New Zealand Plant Protection Society.
- Bianchinotti MV. 1998. Contribución al conocimiento de la micobiota argentina. Micromicetes sobre *Geoffroea decorticans* (Leguminosae). III. Bol. Soc. Argent. Bot. 33: 149–155.
- Cannon PF, Hawksworth DL. 1982. A re-evaluation of *Melanospora* Corda and similar Pyrenomycetes, with a revision of the British species. Bot. J. Linn. Soc. 84: 115–160. <http://dx.doi.org/10.1111/j.1095-8339.1982.tb00363.x>
- Farr DF, Rossman AY. 2012. Fungal databases. Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.ars-grin.gov/fungaldatabases/> [accessed: 16 April 2012].

- Hawksworth DL. 1975. *Chaetocerotostoma* Turc. & Maffei, a genus to be rejected. Trans. Brit. Mycol. Soc. 64: 447–453. [http://dx.doi.org/10.1016/S0007-1536\(75\)80143-4](http://dx.doi.org/10.1016/S0007-1536(75)80143-4)
- Léveillé DM. 1847. Mycologie, Mycétologie. 454–496, in: C Orbigny. Dictionnaire Universel d'Histoire Naturelle 8.
- Lumbsch HT, Huhndorf SM. 2010. Part one. Outline of *Ascomycota*—2009. Myconet 14: 1–42.
- Malloch D. 1976a. *Scopinella solani*. Fungi Canadenses 82.
- Malloch D. 1976b. *Scopinella sphaerophila*. Fungi Canadenses 83.
- McKenzie EHC, Thongkantha S, Lumyong S. 2007. *Zygosporium bioblitzii* sp. nov. on dead leaves of *Cortaderia* and *Dracaena*. N.Z. J. Bot. 45: 433–435. <http://dx.doi.org/10.1080/00288250709509724>
- Medd RW, Murray GM, Pickering DI. 2003. Review of the epidemiology and economic importance of *Pyrenophora semeniperda*. Australas. Plant Pathol. 32: 539–550. <http://dx.doi.org/10.1071/AP03059>
- Pastirčák M, Pastirčáková K. 2007. *Scopinella solani* on graminicolous hosts in Slovakia and the Czech Republic. Mycotaxon 102: 383–387.
- Pausas JG, Lloret F, Vilá M. 2006. Simulating the effects of different disturbance regimes on *Cortaderia selloana* invasión. Biol. Conservation 128: 128–135. <http://dx.doi.org/10.1016/j.biocon.2005.09.022>
- Saccardo PA. 1891. Supplementum universale, pars I. Sylloge Fungorum 9. 1141 p.
- Stchigel AM, Umaña L, Guarro J, Mata M. 2006. Two new ascomycetes from rainforest litter in Costa Rica. Mycologia 98: 815–820. <http://dx.doi.org/10.3852/mycologia.98.5.815>
- Thiers B. 2012 [continuously updated]. Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [accessed: February 2012].
- Tsuneda A, Hiratsuka Y. 1981. *Scopinella gallicola*, a new species from rust galls of *Endocronartium harknessii* on *Pinus contorta*. Canad. J. Bot. 59: 1192–1195. <http://dx.doi.org/10.1139/b81-163>
- Tsuneda A, Hiratsuka Y. 1982. Commensal relationship between *Scopinella gallicola* and *Cladosporium* sp. Rept. Tottori Mycol. Inst. (Japan) 20: 63–69.
- Zhang N, Blackwell M. 2002. Molecular phylogeny of *Melanospora* and similar pyrenomycetous fungi. Mycol. Res. 106: 148–155.