


This article was downloaded by:[Colinas, Pedro A.]
On: 2 June 2008
Access Details: [subscription number 793619332]
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

Sulfonamidoglycosylation of Methyl Glycosides Employing Perchloric Acid Supported on Silica Gel

Pedro A. Colinas^a; Nicolás A. Núñez^a; Rodolfo D. Bravo^a

^a LADECOR, Departamento de Química, Facultad de Ciencias Exactas,
Universidad Nacional de La Plata, La Plata, Argentina

Online Publication Date: 01 March 2008

To cite this Article: Colinas, Pedro A., Núñez, Nicolás A. and Bravo, Rodolfo D.
(2008) 'Sulfonamidoglycosylation of Methyl Glycosides Employing Perchloric Acid
Supported on Silica Gel', Journal of Carbohydrate Chemistry, 27:3, 141 — 147

To link to this article: DOI: 10.1080/07328300802030829

URL: <http://dx.doi.org/10.1080/07328300802030829>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Sulfonamidoglycosylation of Methyl Glycosides Employing Perchloric Acid Supported on Silica Gel

Pedro A. Colinas, Nicolás A. Núñez, and Rodolfo D. Bravo

LADECOR, Departamento de Química, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, La Plata, Argentina

The sulfonamidoglycosylation of methyl glycosides in the presence of perchloric acid immobilized on silica gel proceeded effectively to afford the corresponding sulfonamido glycosides with good to high yields with minimal workup and short reaction times. This methodology was applied to the synthesis of some *N*-glycosyl sulfamides, which showed low nanomolar activity against human carbonic anhydrase.

Keywords Sulfonamidoglycosylation, Methyl glycosides, Perchloric acid-silica

Sulfonamides, which have been clinically used for many years, have been found to possess a large number of different biological activities, including antibacterial, antiviral, antidiabetic, diuretic, and antithyroid activities.^[1] They exert their diverse pharmacological effects by interacting with a wide range of different cellular targets.

In the development of new chemotherapeutic agents, several sulfonamides have emerged as useful therapeutics for the treatment of cancer. E7010,^[2] E7070,^[2] ABT751,^[3] and T138067^[4] have been found to be inhibitors of tumor cell proliferation and some of them are under clinical evaluation and will soon be launched as antitumor drugs (Fig. 1). The mechanism of antitumor action of these compounds has been studied in detail and they inhibit

Received August 28, 2007; accepted November 5, 2007.

Address correspondence to Pedro A. Colinas, LADECOR, Departamento de Química, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, 47 y 115 (1900) La Plata, La Plata, Argentina. E-mail: pcolinas@quimica.unlp.edu.ar





Figure 1: Antimitotic sulfonamides.

microtubule assembly by binding to tubulin at the colchicine binding site.^[5] Other sulfonamides possessing a free sulfonamido moiety probably act as strong carbonic anhydrase inhibitors.^[6] It was found that isozyme CA IX is overexpressed in a variety of tumor types and plays an important role in the growth and survival of tumor cells.^[7] Thus, their inhibition is the target for the development of novel antitumor therapies.

Only few reports described the synthesis of sulfonamidoglycosides. Danishefsky's group reported on the reaction of glycols with iodonium di-sym-collidine perchlorate and benzenesulfonamide to afford stereoselectively 2- β -iodo-1- α -sulfonamido-hexoses.^[8] This class of glycosylsulfonamides were used for the preparation of oligosaccharides with the 2-aminohexose subunit.^[9] Recently we reported on the stereoselective synthesis of glycosylsulfonamides via sulfonamidoglycosylation of benzylated glycols using a catalytic amount of triphenylphosphine hydrobromide.^[10] The novel method afforded the β -sulfonamidoglycosides with good to high yields. Later on we developed a novel approach to ribofuranosylsulfonamides using the reaction of methyl glycosides with sulfonamides in the presence of boron trifluoride etherate.^[11] Some of the glycosylsulfonamides prepared by us showed antiproliferative activity against the human hepatocellular carcinoma in the micromolar range.^[10] Unfortunately, our last developed methodology has an important drawback: boron trifluoride etherate is a highly toxic reagent. This prompted us to initiate studies designed to provide an environmentally friendlier route for the synthesis of sulfonamidoribofuranosides. It seems very interesting to study the use of acids supported on silica gel to promote the sulfonamidoglycosylations. The application of heterogeneous catalysts such as inorganic acids in organic synthesis is an attractive area of research.^[12] Noting recent reports of the use of perchloric acid supported on silica gel^[13] as an inexpensive, nontoxic, and recyclable catalyst for various organic transformations, affording the corresponding products in excellent yields with high selectivity,^[14] we decided to investigate $\text{HClO}_4\text{-SiO}_2$ in the sulfonamidoglycosylations.



Scheme 2: Reagents and conditions: (i) $\text{HClO}_4\text{-SiO}_2$, rt, CH_2Cl_2 .

The use of higher quantities of the $\text{HClO}_4\text{-SiO}_2$ only led to lower yields (entry 6) and they were not improved by using higher quantities of the sulfonamide (entry 1). The results showed that tetrafluoroboric acid was not effective for catalyzing the sulfonamidoglycosylation reactions (entry 7). The addition of molecular sieves, as is mentioned in our previous report,^[11] was not necessary. The generated methanol was probably removed by the silica gel from the reaction system.

With this knowledge in hand, the selected conditions were applied to a variety of sulfonamides and methyl glycosides to analyze the scope of our methodology (Sch. 2). The results are shown in Table 2. The anomeric ratios were determined by ^1H NMR of the reaction mixture.

The products were easily purified by flash chromatography. The ^1H and ^{13}C NMR and 2D COSY experiments and mass spectral data of the sulfonamides were in full accordance with their structure.

The results shown in Table 2 indicated additional applications of the new procedure. Thus, hindered sulfonamides were also effectively coupled to the methyl ribofuranosides (entries 3 and 4). However, a sulfonamide with higher

Table 2: Reaction of sulfonamides with methyl glycosides^a.


Entry	Methyl glycoside	Sulfonamide	Time (min)	Ratio $\alpha:\beta^{b,c}$	Yield (%) ^c
1	1	Benzyl	5	45:55 (46:54)	92 (84)
2	1	Ethane	5	50:50 (48:52)	97 (85)
3	1	<i>N</i> -Methyl- <i>p</i> -toluene	10	40:60 (40:60)	89 (74)
4	1	<i>N</i> -Butylbenzyl	10	50:50 (50:50)	78 (64)
5	1	<i>N</i> -Isopropyl- <i>p</i> -toluene	10	47:53 (45:55)	15 (16)
6	2	<i>p</i> -Toluene	5	45:55 (40:60)	95 (75)
7	3	<i>p</i> -Toluene	5	1:99	97 (80, 87 ^d)
8	4	<i>p</i> -Toluene	5	1:99	98 (82, 80 ^d)

^aAll the reactions were performed in CH_2Cl_2 using 1.3 g of $\text{HClO}_4\text{-SiO}_2$ and 1.5 equiv. of sulfonamide at rt.

^bAnomeric ratios were determined by ^1H NMR spectroscopy.

^cRatios and yields previously reported are shown between parenthesis (Ref. 11).

^dRef. (10).



Scheme 3: Reagents and conditions: i) $\text{HClO}_4\text{-SiO}_2$, rt, CH_3CN .

steric hindrance afforded a poor yield of the glycosylsulfonamide (entry 5). The methodology was also applied to methyl glycopyranosides to furnish the corresponding sulfonamidoglycosides in excellent yields (entries 7 and 8).

The stereochemical outcome was in accordance with the results previously described by us^[10,11] and no influence of the promoter in the anomeric selectivity was found.


Very recently we have prepared a series of glycosylated sulfamides, which showed low nanomolar activity against human carbonic anhydrase.^[15] The $\text{HClO}_4\text{-SiO}_2$ was applied to the synthesis of some of these *N*-glycosylsulfamides (Sch. 3). The results are shown in Table 3.

Table 3: Reaction of sulfamide with methyl glycosides^a.

Entry	Methyl glycoside	Time (min)	Product	Ratio $\alpha:\beta^b$	Yield (%)
1	1	5	11	50:50	82
2	3	5	12	1:99	95
3	4	5	13	6:94	94
4	5	10	14	5:95	90
5	6	10	15	8:92	92

^aAll the reactions were performed in CH_3CN using 1.3 g of $\text{HClO}_4\text{-SiO}_2$ and 3 equiv. of sulfamide at rt.

^bAnomeric ratios were determined by ^1H NMR spectroscopy.



Scheme 4: Reagents and conditions: (i) $\text{HClO}_4\text{-SiO}_2$, r t, CH_2Cl_2 .

Our attention next turned to the reaction of D-glycals with *p*-toluenesulfonamide using $\text{HClO}_4\text{-SiO}_2$. The addition reaction afforded the *O*-glycosides (**17**) in poor yields (Sch. 4). (The reactions were performed in methylenechloride using 0.05 equiv of $\text{HClO}_4\text{-SiO}_2$ and 1.2 equiv. of sulfonamide. Higher quantities of the promoter or the sulfonamide showed no effect on the yields (10%–14%). Similar results using boron trifluoride etherate as catalyst were described by us.^[10] The formation of this product could be explained by the reaction of sulfonamide with the carbenium ion at C(3) followed by the addition of the benzyl alcohol to the double bond.

In summary, our present methodology has the following advantages: (a) the promoter is an inexpensive noncorrosive powder, (b) the workup required only filtration of the promoter followed by flash chromatography, and (c) there were short reaction times (in order of minutes) and good to high yields. In conclusion, we have developed a mild and eco-friendly approach for the sulfonamidoglycosylation of methyl glycosides to afford the corresponding *N*-glycosyl sulfonamides. To the best of our knowledge, it is the first report of a sulfonamidoglycosylation promoted by a heterogenous reagent.

ACKNOWLEDGMENTS

The authors wish to thank CIC (Pcia de Buenos Aires) for financial support and Dr. Agustín Ponzinibbio for NMR measurements. P.A.C. is member of CIC of CONICET (Argentina).

REFERENCES

- [1] Scozzafava, A.; Owa, T.; Mastrolorenzo, A.; Supuran, C.T. Anticancer and antiviral sulfonamides. *Curr. Med. Chem.* **2003**, *10*, 925–953.
- [2] Supuran, C.T. Indisulam: an anticancer sulfonamide in clinical development. *Exp. Opin. Invest. Drugs* **2003**, *12*, 283–287.
- [3] Jordan, M.A.; Wilson, L. Microtubules as a target for anticancer drugs. *Nat. Rev. Cancer* **2004**, *4*, 253–265.
- [4] Medina, J.C.; Roche, D.; Shan, B.; Learned, M.; Frankmoelle, W.P.; Clark, D.L.; Rosen, T.; Jaen, J.C. Novel halogenated sulfonamides inhibit the growth of

- multidrug resistant MCF-7/ADR cancer cells. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 1843–1846.
- [5] Shan, B.; Medina, J.C.; Santha, E.; Frankmoelle, W.P.; Chou, T.-C.; Narbut, M.R.; Stott, D.; Wu, P.; Jaen, J.C.; Rosen, T.; Timmermans, P.B.M.W.M.; Beckmann, H. Selective, covalent modification of beta-tubulin residue Cys-239 by T138067, an antitumor agent with in vivo efficacy against multidrug-resistant tumors. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 5686–5691.
- [6] Abbate, F.; Casini, A.; Owa, t.; Scozzafava, A.; Supuran, C.T. Carbonic anhydrase inhibitors: E7070, a sulfonamide anticancer agent, potently inhibits cytosolic isozymes I and II, and transmembrane, tumor-associated isozyme IX. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 217–223.
- [7] Robertson, N.; Potter, C.; Harris, A.L. Role of carbonic anhydrase IX in human tumor cell growth, survival, and invasion. *Cancer Res.* **2004**, *64*, 6160–6165.
- [8] Griffith, D.A.; Danishefsky, S.J. Sulfonamidoglycosylation of glycals. A route to oligosaccharides with 2-aminohexose subunits. *J. Am. Chem. Soc.* **1990**, *112*, 5811–5819.
- [9] For a review see; Danishefsky, S.J.; Bilodeau, M.T. Glycals in organic synthesis: the evolution of comprehensive strategies for the assembly of oligosaccharides and glycoconjugates of biological consequence. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 1380–1419.
- [10] Colinas, P.A.; Bravo, R.D. A novel sulfonamidoglycosylation of glycals. *Org. Lett.* **2003**, *5*, 4509–4511.
- [11] Colinas, P.A.; Bravo, R.D. Sulfonamidoglycosylation of methyl ribofuranosides: a novel approach to furanosylsulfonamides. *Tetrahedron Lett.* **2005**, *46*, 1687–1689.
- [12] Corma, A.; García, H. Lewis Acids: From conventional homogeneous to green homogeneous and heterogeneous catalysis. *Chem. Rev.* **2003**, *103*, 4307–4365.
- [13] Chakraborti, A.K.; Gulhane, R. Perchloric acid adsorbed on silica gel as a new, highly efficient, and versatile catalyst for acetylation of phenols, thiols, alcohols, and amines. *Chem. Commun.* **2003**, 1896–1897.
- [14] For applications of this reagent system see: Narasimhulu, M.; Reddy, T.S.; Mahesh, K.C.; Prabhakar, P.; Rao, Ch.B.; Venkateswarlu, Y. Silica supported perchloric acid: A mild and highly efficient heterogeneous catalyst for the synthesis of poly-substituted quinolines via Friedländer hetero-annulation. *J. Mol. Catal. A Chem.* **2007**, *266*, 114–117 and references therein.
- [15] Colinas, P.A.; Bravo, R.D.; Vullo, D.; Scozzafava, A.; Supuran, C.T. Carbonic anhydrase inhibitors. Inhibition of cytosolic isoforms I and II, and extracellular isoforms IV, IX, and XII with sulfamides incorporating sugar moieties. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5086–5090.