

Cocoa, Chocolate, and Cardiovascular Disease

Monica Galleano, PhD,* Patricia I. Oteiza, PhD,†‡ and Cesar G. Fraga, PhD*†

Abstract: A significant body of evidence demonstrates that diets rich in fruits and vegetables promote health and attenuate, or delay, the onset of various diseases, including cardiovascular disease, diabetes, certain cancers, and several other age-related degenerative disorders. The concept that moderate chocolate consumption could be part of a healthy diet has gained acceptance in past years based on the health benefits ascribed to selected cocoa components. Specifically, cocoa as a plant and chocolate as food contain a series of chemicals that can interact with cell and tissue components, providing protection against the development and amelioration of pathological conditions. The most relevant effects of cocoa and chocolate have been related to cardiovascular disease. The mechanisms behind these effects are still under investigation. However, the maintenance or restoration of vascular NO production and bioavailability and the antioxidant effects are the mechanisms most consistently supported by experimental data. This review will summarize the most recent research on the cardiovascular effects of cocoa flavanols and related compounds.

Key Words: flavonoids, flavanols, hypertension, oxygen radicals, antioxidant, cardiovascular health

(*J Cardiovasc Pharmacol*TM 2009;54:483–490)

CARDIOVASCULAR DISEASE AND DIET

Cardiovascular disease (CVD), including stroke, is the leading cause of death and disability in developed countries. Atherosclerosis, vascular dysfunction, platelet aggregation, and other inflammation associated conditions are central to CVD. Diet is a major factor contributing to the onset and development of CVD, by primarily affecting all of the above-mentioned conditions. Nevertheless, high intake of calories and certain fats increases the risk for CVD,¹ and diet also provides micronutrients that seem fundamental in controlling CVD.² Defining specific modifications of dietary habits in a population can have a major impact on CVD, especially during the long period in which the disease is silent.³ To set

new goals to improve human diets, it is important to understand how macro- and micronutrients can interact with biological systems to enhance health.

CVD AND COCOA

Robust epidemiological evidence demonstrates that diets rich in fruits and vegetables promote health and attenuate, or delay, the onset of CVD (and references therein).^{3–6} The questions that remain open are (1) are all fruits and vegetables equivalent? If not, can we identify those with best health benefits? and (2) how can we recognize the compounds responsible for such effects? A pioneer population study showed an inverse association between flavonoid consumption and the risk of coronary heart disease.⁷ Under this paradigm, cocoa and chocolate have been intensively studied in past years, mostly driven by the large content of flavanols and related compounds in cocoa beans that are generally conserved in commercially available chocolate.

High cocoa and chocolate consumption have been associated with a decreased risk for CVD in a few population studies. A substudy of a population from the Zutphen Elderly Study showed that cocoa consumption was associated with a decrease in blood pressure and overall cardiovascular mortality.⁸ A case–control study done in Italy showed that the risk for myocardial infarction was inversely associated to chocolate consumption, reaching a 77% decrease in risk when comparing the population that ate more than 3 portions of chocolate per day with the population that consumed less than 1.⁹ Several dietary intervention studies in humans and animals demonstrated that cocoa^{10–18} and other flavanol-rich foods/beverages^{19–21} may exert protective vascular effects.

Although data for the above mentioned population and clinical studies show a similar trend suggesting flavanols as cardioprotective agents, the biochemical mechanisms behind that cardioprotection are not conclusively identified. Based on a number of in vitro studies, it has been speculated that biochemical mechanisms contributing to the health effects of flavanols include antioxidant effects,²² modulation of cell signaling and gene expression,^{23,24} and alterations of certain cell membrane properties and receptor functions.^{25,26} In addition to the above mentioned mechanisms, flavanols can inhibit several enzyme activities.^{27–30} These mechanisms are not necessarily independent and could then occur concurrently or synergistically, as will be discussed in the following sections.

CHEMICAL STRUCTURE OF COCOA FLAVANOLS AND PROCYANIDINS

Polyphenols occur as plant secondary metabolites. Their ubiquitous occurrence in plants and plant foods favors animal

Received for publication May 30, 2009; accepted June 23, 2009.

From the *Physical Chemistry-Program of Free Radicals in Biochemistry (PRALIB), School of Pharmacy and Biochemistry, University of Buenos Aires-National Council of Scientific and Technological Research (CONICET), Argentina; †Department of Nutrition, University of California, Davis, CA; and ‡Department of Environmental Toxicology, University of California, Davis, CA.

Supported by National Institute of Health AT2966, Center for Health and Nutrition Research-State of California Vitamin Price Fixing Consumer Settlement Fund, and Universidad de Buenos Aires-Ciencia y Técnica B801-B802, Argentina.

The authors report no conflicts of interest.

Reprints: Cesar G. Fraga, PhD, Department of Nutrition, University of California, Davis, One Shields Avenue, Davis, CA 95616 (e-mail: cfraga@fhyb.uba.ar). Copyright © 2009 by Lippincott Williams & Wilkins

consumption and indeed human and animal tissue presence. Flavonoids are a chemically defined family of polyphenols that includes several thousand compounds. The flavonoids have a basic structure (Fig. 1), and several subclasses of flavonoids are characterized by a substitution pattern in the B and C rings. The main subclasses include flavanols, flavanones, flavones, isoflavones, flavonols, and anthocyanidins.³¹ Flavanols are compounds present in high concentrations not only in cocoa, but also in grapes, apples, pomegranates, and tea, among other widely consumed fruits and vegetables. In cocoa and cocoa products, flavanols are present as (1) monomers, that is, (–)-epicatechin (EC) and (+)-catechin (CT) and (2) oligomers of EC (procyanidins).³² EC (and CT) oligomers are denominated procyanidins, condensed tannins, or proanthocyanidins. Different plants present a particular pattern of monomers and oligomerization derivatives, for example, cocoa procyanidins are mostly of the denominated B type, for example, B2 dimer (Fig. 1), whereas the gallolyated catechins predominate in tea (*Camellia sinensis*).

BIOAVAILABILITY AND METABOLISM OF COCOA FLAVANOLS AND PROCYANIDINS

When discussing the biological activity of flavonoids in general, and flavanols in particular, there are 4 major factors to be considered: (1) bioavailability from foods, (2) absorption and metabolism at the gastrointestinal tract, (3) tissue and cellular distribution after absorption, and (4) which are the chemical form(s) biologically available to the cell/tissue and their potential metabolism at cellular level. As indicated above, flavanols are present as monomers or procyanidins. Although it was initially thought that the procyanidins could not pass the acidic conditions of the stomach,³³ data from human subjects show that flavanols and procyanidins are stable during gastric transit.^{34,35} Once in the mesenteric circulation, flavanols predominately exist in a conjugated form, both as methylated

and glucuronidated flavanols.^{36–38} In the liver, further glucuronidation and methylation can take place, as well as sulfation.^{36–38} Metabolic studies have confirmed the presence of these conjugates in the plasma and urine of rodents and humans^{38–41} and in the bile³⁸ and brain of rats.⁴² It has been reported that colonic microflora can break flavonoid flavan structure to form simple phenolics and ring-fission metabolites that may be physiologically relevant.^{43,44} In summary, nonmetabolized flavanols or metabolites of flavanols can exert biological effects depending essentially on flavanol metabolism and presence in the target tissue.^{28–30}

In humans, plasma concentrations of EC plus EC metabolites can be found in the micromolar range as soon as 1 hour after cocoa consumption,^{12,39,45} with the major metabolite being 4'-O-methyl-epicatechin-7-β-D-glucuronide.¹² There is some evidence that certain flavanols are better absorbed than others. After human subjects consumed a cocoa beverage containing EC and CT in a 1:1 ratio, peak plasma CT concentrations were typically less than 10% of EC (0.16 vs. 5.92 μM).⁴⁵ Part of these differences in plasma flavanol concentrations could be due to procyanidin degradation; for example, dimers has been shown to form EC and methylated EC under certain conditions,^{41,46,47} although the physiological relevance of such degradation remains to be confirmed.

Although several research groups have examined the bioavailability of the monomeric flavanols, there is limited information on the bioavailability and metabolism of procyanidins. B2 dimer has been detected in the plasma of humans and rats^{41,45–49}; B5 dimer (EC-(4β-6)-EC) has been detected in very minute quantities in simulated gastric and intestinal juice⁴⁷; and no other type of dimer has been detected in rats.⁵⁰ It is important to note that the dimers that have been detected in the plasma are those made up of EC, not of CT subunits. A chemical explanation for these discrepancies is the different hydroxyl group orientation in the C3 position of the flavonoid B ring, which affects the interaction between the




FIGURE 1. Chemical structure of flavanols and procyanidins. B2 dimer is a characteristic cocoa procyanidin formed by two (–)-EC units linked by 4→8 bonds.

3-OH on the C ring and the B ring, resulting in dissimilar biological actions.^{51,52}

COCOA FLAVANOLS AS ANTIOXIDANTS: FREE RADICAL SCAVENGERS AND METAL CHELATORS

Plant polyphenols have long been considered physiologically relevant antioxidants based on the facts that (1) polyphenols have chemical structures favoring antioxidant actions, that is, free radical scavenging and chelation of redox-active metals, (2) many polyphenols retain key features of their structure after ingestion and metabolism by mammals, and (3) certain polyphenols can provide physiological benefits in pathological situations associated with high free radical production, for example, hypertension. Flavanols share these properties, as confirmed by the extensive literature, demonstrating that flavanols have free radical scavenging activity in a myriad of biochemical and *ex vivo* systems^{23,25,26,53–61} and also in animal models and in humans.^{62–64} In theory, these antioxidant actions can result in a reduction of the steady state concentration of free radicals and other oxidants, diminishing the subsequent oxidation of target molecules such as lipids, proteins, and nucleic acids. However, one important limitation for the “antioxidant action” hypothesis resides in the relatively low flavanol and procyanidin plasma concentrations observed even after the consumption of foods rich in these compounds.^{45,65} The actual concentrations that can be reached in plasma of humans subjected to realistic polyphenol consumption are in the nanomolar range and are transient in nature (peaking at 2–4 hours).^{12,45} This low bioavailability leads to a kinetically unfavorable condition with respect to other compounds with similar free radical scavenger capabilities, which are present in blood in significantly higher micromolar concentrations, that is, tocopherols and ascorbate. Thus, a function of flavanols as direct free radical scavengers is unlikely to be relevant and could be limited to the blood and other tissues directly exposed after consumption, that is, gastrointestinal tract. It has been suggested that other mechanisms, compatible with the physiological levels reached by flavanols, may explain the observed changes in cell or tissue oxidation levels after flavanol consumption. These mechanisms are beyond the ability of flavanols and other flavonoids to directly prevent free radical mediated tissue damage.^{65,66}

Another relevant detrimental effect of oxidants is the reaction of nitric oxide (NO) with superoxide to form peroxynitrite.⁶⁷ This reaction that occur in a near diffusion-controlled rate, and is important at the vascular level, leads to two undesirable conditions: (1) reduction of the NO availability necessary for proper smooth muscle cell function in vessel relaxation and (2) increased formation of peroxynitrite that will promote oxidative and nitrosative damage (Fig. 2). NO production in mammalian cells is catalyzed by the enzyme nitric oxide synthase, whose activity is by 4 isoforms, that is, endothelial (eNOS), expressed mostly in endothelial cells; neuronal (nNOS), present mainly in neurons; inducible (iNOS), expressed in response a variety of proinflammatory stimuli; and mitochondrial (mtNOS), present in the inner membrane of the mitochondrion.^{68,69} In the vascular environment,




FIGURE 2. Scheme relating flavanols (circles) with NO and superoxide (O₂⁻) metabolism in endothelial cells. Flavanols could act by (1) inhibiting NADPH oxidase–dependent superoxide production, (2) activating eNOS, (3) scavenging superoxide, H₂O₂, and other oxidants that mediate damage to cell components, and (4) modifying membrane-related events, leading to changes in NO and superoxide production. NO generated in endothelial cells will diffuse outside the cell and reach smooth muscle cells where it will induce vascular relaxation.

superoxide is not only originated by the mitochondria respiratory chain,^{70,71} xanthine oxidase, and cytochrome P450,^{72,73} but also by 2 other enzymes: a nonphagocytic NADPH oxidase⁷⁴ and an uncoupled endothelial NOS.⁷⁵ Then, flavanols and procyanidins present in circulation or in the vasculature can improve NO vascular concentration by interfering with the reaction between NO and superoxide by (1) inhibiting NADPH oxidase–dependent superoxide production,³⁰ (2) optimizing NO generation by NOS⁷⁶ and/or (3) scavenging superoxide, H₂O₂, and other oxidants that mediate damage to cell components,⁷⁷ and (4) modifying membrane-related events, leading to changes in NO and superoxide production.

COCOA FLAVANOLS AS ANTIOXIDANTS: PROTEIN AND LIPID INTERACTIONS

Flavanols and procyanidins have multiple phenolic hydroxyl groups (Fig. 1) that favor their interaction with biological membranes, which can occur via the formation of hydrogen bonds. Furthermore, the presence of both hydrophobic and hydrophilic residues within the flavanol molecule allows these compounds to interact with phospholipid head groups and be adsorbed onto the surface of membranes. These interactions can result in changes in a number of membrane properties, leading to alterations in the regulation of membrane associated molecules and events, including, enzymes, receptors, and other functional proteins receptors.^{78–80}

Some enzymes affected by cocoa consumptions are directly associated with CVD and oxidant metabolism, such as 5-lipoxygenase,^{81,82} cyclooxygenase-2,^{83,84} and metalloproteinases.⁸⁵ The interactions of flavanols and proteins can also lead to changes in the modulation of gene expression. A direct interaction between nucleic acids and flavanols is thermodynamically feasible,⁸⁶ but the possibility that these compounds reach the DNA and achieve mechanistically relevant concentrations is rather low. The modulation of signaling pathways by flavanols has been extensively studied.^{87–89} More specifically, the effects of flavanols and procyanidins on the oxidant-regulated nuclear factor κ B (NF- κ B) activation pathway have received special attention. In Jurkat T cells, we demonstrated that EC, CT, and B2 dimer inhibited phorbol myristate acetate-induced IL-2 production and interfere with several steps of the NF- κ B activation cascade.²³ Essentially, because monomers and dimers (and their metabolites) can be transported into the cells, they can act by (1) attenuating intracellular oxidants associated with select stimuli and the subsequent activation of NF- κ B (antioxidant effect) and/or (2) interacting with specific proteins, resulting in the inhibition of the phosphorylation and/or degradation of the inhibitory protein I κ B α , the transport of active NF- κ B from the cytosol into the nucleus, and/or the binding of NF- κ B to κ B DNA.^{23,90} Large procyanidins (with 3 or more units), that are mostly affecting cells from outside, modulate NF- κ B activation by modulating the binding of the ligand (stimuli) to its receptor, as we observed in Caco-2 cells exposed to tumor necrosis factor alpha.⁹¹ Another example of flavanol interaction with membranes is the finding that EC, B2 dimer, and C1 trimer modulate intracellular calcium in Jurkat T cells.⁹²

COCOA, LIPID METABOLISM, AND ATHEROSCLEROSIS

Alterations in plasma cholesterol concentration, especially increased levels of low-density lipoprotein (LDL) cholesterol and decreased levels of high-density lipoprotein (HDL) cholesterol, are associated to the development of atherosclerosis^{93,94} and CVD.¹ Reductions in LDL cholesterol plasma levels have been reported after treatments with polyphenols from different sources.^{95–98} Regarding the action of cocoa, mild hypercholesterolemic subjects lowered significantly (~5%) their LDL cholesterol level after 4 weeks of a dietary intervention with cocoa powder (81–163 mg/d of EC + CT).⁹⁹ Even in normocholesterolemic young subjects, a 15% reduction was observed in LDL cholesterol level after 14 days of a daily consumption of 105 g of flavanol-containing milk chocolate (168 mg of flavanols).¹⁰⁰ Patients with essential hypertension showed 11% of decrease in LDL cholesterol after 15 days receiving 100 g/d dark chocolate (88 mg of flavonols).¹⁰ Under an equivalent protocol, a similar result was observed in patients with glucose-intolerant hypertension (–7.5%).¹⁰¹ Although the above studies did not investigate potential mechanisms, other works have proposed that LDL cholesterol decrease associated to flavanols consumption from different sources include (1) inhibition of cholesterol absorption in the digestive tract,⁹⁶ (2) inhibition of LDL biosynthesis in the liver,⁹⁸ (3) suppression of the hepatic

secretion of apolipoprotein B100,⁹⁵ and/or (4) increased expression of LDL receptors in the liver.⁹⁷ All of these mechanisms should be the result of interactions between flavanols and membranes, as whole structures, or with particular lipids or proteins. An increase in HDL cholesterol has been demonstrated in normocholesterolemic and mildly hypercholesterolemic subjects after dark chocolate or cocoa powder supplementation.^{99,102,103} The mechanisms responsible for these effects on HDL concentrations remain unclear.

It is accepted that oxidized LDL has a role in the development of atherosclerosis.¹⁰⁴ Numerous studies in animals and humans showed that isolated LDLs are more resistant to *in vitro* oxidation after the consumption of cocoa products.^{57,102,105–109} One study showed a decrease in plasma levels of oxidized LDL plasma levels after dietary cocoa powder supplementation.⁹⁹ All these studies suggest a role for cocoa components in the *in vivo* protection of LDL. These effects have been mostly ascribed to the scavenging of oxidants formed in the surface of the LDL and to the chelation of metals that are catalyst of free radical formation, but they could also be the result of changes in the LDL surface rendering LDL less susceptible to oxidation.

COCOA, ENDOTHELIAL FUNCTION, AND HYPERTENSION

The regulation of the vascular tone is the result of a complex network of molecules that includes catecholamines, vasoactive peptides (angiotensin-II or vasopressin), prostaglandins, and importantly NO. Consumption of a high-flavanol cocoa drink (providing 176–185 mg) by patients with cardiovascular risk factors increased the bioavailability of NO and the augmented flow-mediated vasodilation effects that were reversed by the infusion of a NO synthesis inhibitor.^{13–15} Studies with isolated flavonoids showed that EC was able to reproduce the vascular effects observed with cocoa products, suggesting that this flavanol should be responsible for the vascular effects.¹²

Observational and epidemiological studies indicate that diets rich in polyphenols decrease blood pressure and prevent the increase in blood pressure associated to several pathologies. For years, red wine was considered to have beneficial effects on cardiovascular health, a relationship supported by the French paradox.¹¹⁰ Cocoa and cocoa-derived products have gained attention because of their potential antihypertensive effects. A substudy of the Zutphen population showed that cocoa consumption was associated with a decrease in blood pressure.⁸ In addition, an association between the intake of cocoa and a low incidence of hypertension was observed in an indigenous population living on the Kuna Islands.¹¹¹ The Kuna Indians of Panama have a very low incidence of hypertension and CVD, but when members of this tribe moved to urban places, their blood pressure was increased. The relocation led to cultural changes and to a decrease in the consumption of cocoa, suggesting that this dietary change was responsible of the observed changes in blood pressure. These results were complemented by other studies that showed an enhancement of endothelial function including acute positive effects on flow-mediated dilation by cocoa consumption in humans.^{12,13,15–17}

Consistent with the association between cocoa consumption and low incidence of hypertension are the results from several short-term clinical studies showing that the intake of certain chocolates can decrease blood pressure in humans.^{10,11,100,112,113} Grassi et al studied 15 healthy young adults with typical Italian diets isocalorically supplemented daily with 100 g of dark chocolate or 90 g of white chocolate (assuming 500 and 0 mg of polyphenols, respectively). They observed that the dark chocolate supplement was associated with decreased systolic blood pressure, whereas the white chocolate had no effect.¹¹² Results were extended to patients with essential hypertension¹⁰ and more recently to patients with glucose-intolerant hypertension.¹⁰¹ We studied the effects of the regular consumption of a flavanol-containing milk chocolate on blood pressure and on oxidative stress parameters in healthy young soccer players.¹⁰⁰ The consumption of the flavanol-containing milk chocolate was significantly associated with a decrease in blood pressure and an amelioration of oxidative stress. Taubert et al¹¹ studied the effects of low doses of polyphenol-rich dark chocolate in humans during 18 weeks. Dark chocolate intake reduced mean, systolic, and diastolic blood pressure and oxidative stress. Blood pressure decrease was accompanied by a sustained increase of *S*-nitrosoglutathione, suggesting an improved formation of NO. The above studies provide support for the involvement of oxidative stress in the vascular tone regulation by NO availability. A meta-analysis of 5 studies relating cocoa consumption with decreases in blood pressure confirmed the individual results.¹¹⁴ Significantly, the reductions in systolic (4.7 mm Hg) and diastolic (2.8 mm Hg) blood pressure associated to cocoa and chocolate consumption were similar to those obtained with antihypertensive drugs.¹¹⁴ In another study, a dark chocolate and a sugar-free cocoa drink were effective in improving endothelial function and decreasing blood pressure in overweight adults.¹¹⁵ In accordance with these results, several studies have shown significant decreases in blood pressure after the consumption of other flavanol-containing beverages such as wine^{19,110,116} and tea.^{117–120} On the other hand, one study showed that the administration of 900 mg of flavanols per day during 2 weeks enhanced insulin-mediated vasodilation but did not modify blood pressure in patients with essential hypertension. These authors concluded that the treatment was not long enough.¹²¹

COCOA, PLATELET ACTIVATION, AND THROMBOSIS

Platelet activation is a central event in coagulation, but it is also related to the acute development of thrombosis and to the long-term CVD pathogenesis. A thrombi release from an unstable atherosclerotic plaque is very often the first clinical manifestation of a myocardial ischemia or infarction or stroke.^{122,123} Regarding the effects of cocoa on platelet function, Rein et al administered a cocoa beverage (~897 mg of total EC and oligomeric procyanidins) or placebo to healthy subjects. Blood obtained after 2 hours was stimulated with epinephrine and adenosine diphosphate. Platelets present in those samples were studied by the detection of activated conformation of the fibrinogen-binding receptor GPIIb-IIIa

and the expression of CD62P (associated with platelet activation). Both parameters were significantly reduced in cocoa-treated individuals. Cocoa also inhibited coagulation by reducing the formation of hemostatically active platelet micro-particles and increasing platelet-related hemostasis time.¹²⁴ In another study, platelet function was evaluated in smokers 2 hours after receiving 40 g of dark chocolate (~47 mg of EC + CT).¹²⁵ Platelet adherence as a result of a shear stress that mimics severely stenotic or disrupted plaques was significantly reduced (–5%) in association with dark chocolate supplementation. Using a similar assay in a protocol including heart transplant recipients, a reduction in platelet adherence was also found.¹²⁶ Both studies propose that the antioxidant properties of cocoa flavanols can lead with an increase in NO bioavailability, which would be associated with the decrease in platelet reactivity, given the potent action of NO as platelet inhibitor.¹²⁷ In isolated platelets, it was shown that a mixture of CT and quercetin can cause a similar effect, increasing NO levels through the inhibition of protein kinase C–dependent NADPH oxidase.¹²⁸ Other mechanisms for the effects of flavonoids on platelet reactivity that may be associated with a NO availability because of the inhibition of oxidant production include (1) the reduction in phospholipase C activity associated to hydrogen peroxide production,¹²⁹ (2) the modulation of eicosanoid metabolism,¹³⁰ (3) the blockade of platelets TxA2 receptors,¹³¹ and (4) the inhibition of platelet lipooxygenase.^{132,133}

COCOA AND INFLAMMATION

There is increasing evidence that inflammation is pivotal in the induction and perpetuation of CVD. A participation of dietary flavonoids in the modulation of inflammation would contribute to reduce cardiovascular risk. An inverse association was observed between dietary flavonoid intake and serum C-reactive protein, that is, both a biomarker for chronic inflammation and a sensitive risk factor for CVD.¹³⁴ Moreover, data from the National Health and Nutrition Examination Survey 1999–2002 has shown an inverse association between particular flavonols, that is, quercetin and kaempferol and C-reactive protein. A significant association between inflammation and moderate consumption of cocoa products was found in a study comparing subjects who ate chocolate regularly in the form of dark chocolate (n = 824) with subjects who did not eat chocolate for at least 1 year (n = 1317). Serum C-reactive protein concentration in the subgroup having up to 1 serving (20 g of cocoa) every 3 days was significantly lower than in both nonconsumers and subjects having higher consumption.¹³⁵ The possible mechanisms involved in the anti-inflammatory effects of cocoa products have recently been revised.^{136,137} Of interest are cell experiments showing the inhibition of mitogen-activated protein kinase activities by cocoa procyanidins^{138,139} and cocoa extracts.¹⁴⁰

CONCLUDING REMARKS

A full range of health benefits can be associated with the actions of flavanols and procyanidins on vascular function. These benefits are mainly ascribed to diets rich in flavanols and procyanidins, and chocolate and cocoa derivatives are among

the most valuable components of such a diet. Considering the fact that CVD is associated with a series of conditions that can trigger oxidant production and oxidant-regulated cell signaling, it would be logical to relate the free radical scavenging and metal chelating properties of cocoa flavanols to CVD protective effects. However, other biochemical mechanisms related to specific flavanol–lipid and flavanol–protein interactions can partially explain the observed in vitro and in vivo antioxidant effects. These mechanisms are more consistent with the in vivo flavanol and procyanidins levels observed in most human and animal tissues.

REFERENCES

- Lichtenstein AH, Appel LJ, Brands M, et al. Summary of American Heart Association Diet and Lifestyle Recommendations revision 2006. *Arterioscler Thromb Vasc Biol.* 2006;26:2186–2191.
- Knoops KT, de Groot LC, Kromhout D, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA.* 2004;292:1433–1439.
- Appel LJ, Champagne CM, Harsha DW, et al. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *J Am Med Assoc.* 2003;289:2083–2093.
- Appel LJ, Moore TJ, Obarzanek E, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med.* 1997;336:1117–1124.
- Bazzano LA, He J, Ogden LG, et al. Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *Am J Clin Nutr.* 2002;76:93–99.
- Hung HC, Joshipura KJ, Jiang R, et al. Fruit and vegetable intake and risk of major chronic disease. *J Natl Cancer Inst.* 2004;96:1577–1584.
- Hertog MG, Feskens EJ, Hollman PC, et al. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *Lancet.* 1993;342:1007–1011.
- Buijsse B, Feskens EJ, Kok FJ, et al. Cocoa intake, blood pressure, and cardiovascular mortality: the Zutphen Elderly Study. *Arch Intern Med.* 2006;166:411–417.
- Gallus S, Tavani A, La Vecchia C. Response to chocolate, well-being, and health among elderly men: chocolate and acute myocardial infarction in a case-control study from Italy. *Eur J Clin Nutr.* 2009;63:588–589.
- Grassi D, Necozione S, Lippi C, et al. Cocoa reduces blood pressure and insulin resistance and improves endothelium-dependent vasodilation in hypertensives. *Hypertension.* 2005;46:398–405.
- Taubert D, Roesen R, Lehmann C, et al. Effects of low habitual cocoa intake on blood pressure and bioactive nitric oxide: a randomized controlled trial. *J Am Med Assoc.* 2007;298:49–60.
- Schroeter H, Heiss C, Balzer J, et al. (-)-Epicatechin mediates beneficial effects of flavanol-rich cocoa on vascular function in humans. *Proc Natl Acad Sci U S A.* 2006;103:1024–1029.
- Heiss C, Dejam A, Kleinbongard P, et al. Vascular effects of cocoa rich in flavan-3-ols. *J Am Med Assoc.* 2003;290:1030–1031.
- Heiss C, Finis D, Kleinbongard P, et al. Sustained increase in flow-mediated dilation after daily intake of high-flavanol cocoa drink over 1 week. *J Cardiovasc Pharmacol.* 2007;49:74–80.
- Heiss C, Kleinbongard P, Dejam A, et al. Acute consumption of flavanol-rich cocoa and the reversal of endothelial dysfunction in smokers. *J Am Coll Cardiol.* 2005;46:1276–1283.
- Heiss C, Schroeter H, Balzer J, et al. Endothelial function, nitric oxide, and cocoa flavanols. *J Cardiovasc Pharmacol.* 2006;47(Suppl 2):S128–S135; discussion S172–S176.
- Fisher ND, Hughes M, Gerhard-Herman M, et al. Flavanol-rich cocoa induces nitric-oxide-dependent vasodilation in healthy humans. *J Hypertens.* 2003;21:2281–2286.
- Murphy KJ, Chronopoulos AK, Singh I, et al. Dietary flavanols and procyanidin oligomers from cocoa (Theobroma cacao) inhibit platelet function. *Am J Clin Nutr.* 2003;77:1466–1473.
- Diebolt M, Bucher B, Andriantsitohaina R. Wine polyphenols decrease blood pressure, improve NO vasodilatation, and induce gene expression. *Hypertension.* 2001;38:159–165.
- Duffy SJ, Keaney JF Jr, Holbrook M, et al. Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation.* 2001;104:151–156.
- Freedman JE, Parker C III, Li L, et al. Select flavonoids and whole juice from purple grapes inhibit platelet function and enhance nitric oxide release. *Circulation.* 2001;103:2792–2798.
- Rice-Evans C. Flavonoid antioxidants. *Curr Med Chem.* 2001;8:797–807.
- Mackenzie GG, Carrasquedo F, Delfino JM, et al. Epicatechin, catechin, and dimeric procyanidins inhibit PMA-induced NF-kappaB activation at multiple steps in Jurkat T cells. *FASEB J.* 2004;18:167–169.
- Saliou C, Rihn B, Cillard J, et al. Selective inhibition of NF-kappaB activation by the flavonoid hepatoprotector silymarin in HepG2. Evidence for different activating pathways. *FEBS Lett.* 1998;440:8–12.
- Erlejman AG, Verstraeten SV, Fraga CG, et al. The interaction of flavonoids with membranes: potential determinant of flavonoid antioxidant effects. *Free Radic Res.* 2004;38:1311–1320.
- Verstraeten SV, Keen CL, Schmitz HH, et al. Flavan-3-ols and procyanidins protect liposomes against lipid oxidation and disruption of the bilayer structure. *Free Radic Biol Med.* 2003;34:84–92.
- Middleton E Jr, Kandaswami C, Theoharides TC. The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. *Pharmacol Rev.* 2000;52:673–751.
- Steffen Y, Gruber C, Schewe T, et al. Mono-O-methylated flavanols and other flavonoids as inhibitors of endothelial NADPH oxidase. *Arch Biochem Biophys.* 2008;469:209–219.
- Steffen Y, Jung T, Klotz LO, et al. Protein modification elicited by oxidized low-density lipoprotein (LDL) in endothelial cells: protection by (-)-epicatechin. *Free Radic Biol Med.* 2007;42:955–970.
- Steffen Y, Schewe T, Sies H. (-)-Epicatechin elevates nitric oxide in endothelial cells via inhibition of NADPH oxidase. *Biochem Biophys Res Commun.* 2007;359:828–833.
- Crozier A, Jaganath IB, Clifford MN. Phenols, polyphenols and tannins: an overview. Crozier A, Clifford MN, Ashihara H, eds. *Plant Secondary Metabolites-Occurrence, Structure, and Role in the Human Diet.* Oxford, United Kingdom: Blackwell Publishing Ltd; 2006:1–24.
- Adamson GE, Lazarus SA, Mitchell AE, et al. HPLC method for the quantification of procyanidins in cocoa and chocolate samples and correlation to total antioxidant capacity. *J Agric Food Chem.* 1999;47:4184–4188.
- Spencer JP, Chaudry F, Pannala AS, et al. Decomposition of cocoa procyanidins in the gastric milieu. *Biochem Biophys Res Commun.* 2000;272:236–241.
- Mullen W, Borges G, Donovan JL, et al. Milk decreases urinary excretion but not plasma pharmacokinetics of cocoa flavan-3-ol metabolites in humans. *Am J Clin Nutr.* 2009;89:1784–1791.
- Rios LY, Bennett RN, Lazarus SA, et al. Cocoa procyanidins are stable during gastric transit in humans. *Am J Clin Nutr.* 2002;76:1106–1110.
- Spencer JP. Metabolism of tea flavonoids in the gastrointestinal tract. *J Nutr.* 2003;133:S3255–S3261.
- Piskula MK, Terao J. Accumulation of (-)-epicatechin metabolites in rat plasma after oral administration and distribution of conjugation enzymes in rat tissues. *J Nutr.* 1998;128:1172–1178.
- Donovan JL, Crespy V, Manach C, et al. Catechin is metabolized by both the small intestine and liver of rats. *J Nutr.* 2001;131:1753–1757.
- Baba S, Osakabe N, Yasuda A, et al. Bioavailability of (-)-epicatechin upon intake of chocolate and cocoa in human volunteers. *Free Radic Res.* 2000;33:635–641.
- Donovan JL, Bell JR, Kasim-Karakas S, et al. Catechin is present as metabolites in human plasma after consumption of red wine. *J Nutr.* 1999;129:1662–1668.
- Baba S, Osakabe N, Natsume M, et al. Absorption and urinary excretion of procyanidin B2 [Epicatechin-(4b-8)-epicatechin] in rats. *Free Radic Biol Med.* 2002;33:142–148.
- Abd El Mohsen MM, Kuhnle G, Rechner AR, et al. Uptake and metabolism of epicatechin and its access to the brain after oral ingestion. *Free Radic Biol Med.* 2002;33:1693–1702.
- Rechner AR, Kuhnle G, Bremner P, et al. The metabolic fate of dietary polyphenols in humans. *Free Radic Biol Med.* 2002;33:220–235.
- Unno T, Tamemoto K, Yayabe F, et al. Urinary excretion of 5-(3',4'-dihydroxyphenyl)-gamma-valerolactone, a ring-fission metabolite of (-)-epicatechin, in rats and its in vitro antioxidant activity. *J Agric Food Chem.* 2003;51:6893–6898.

45. Holt RR, Lazarus SA, Sullards MC, et al. Procyanidin dimer B2 [epicatechin-(4beta-8)-epicatechin] in human plasma after the consumption of a flavanol-rich cocoa. *Am J Clin Nutr*. 2002;76:798–804.
46. Spencer JP, Schroeter H, Shenoy B, et al. Epicatechin is the primary bioavailable form of the procyanidin dimers B2 and B5 after transfer across the small intestine. *Biochem Biophys Res Commun*. 2001;285:588–593.
47. Zhu QY, Holt RR, Lazarus SA, et al. Stability of the flavan-3-ols epicatechin and catechin and related dimeric procyanidins derived from cocoa. *J Agric Food Chem*. 2002;50:1700–1705.
48. Steinberg FM, Holt RR, Schmitz HH, et al. Cocoa procyanidin chain length does not determine ability to protect LDL from oxidation when monomer units are controlled. *J Nutr Biochem*. 2002;13:645–652.
49. Urpi-Sarda M, Monagas M, Khan N, et al. Epicatechin, procyanidins, and phenolic microbial metabolites after cocoa intake in humans and rats. *Anal Bioanal Chem*. 2009;394:1545–1556.
50. Gonthier MP, Donovan JL, Texier O, et al. Metabolism of dietary procyanidins in rats. *Free Radic Biol Med*. 2003;35:837–844.
51. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med*. 1996;20:933–956.
52. Schroeder P, Klotz LO, Sies H. Amphiphilic properties of (–)-epicatechin and their significance for protection of cells against peroxynitrite. *Biochem Biophys Res Commun*. 2003;307:69–73.
53. Actis-Goretti L, Mackenzie GG, Oteiza PI, et al. Comparative study on the antioxidant capacity of wines and other plant-derived beverages. *Ann N Y Acad Sci*. 2002;957:279–283.
54. Erlejtman AG, Fraga CG, Oteiza PI. Procyanidins protect Caco-2 cells from bile acid- and oxidant-induced damage. *Free Radic Biol Med*. 2006;41:1247–1256.
55. Fraga CG, Martino VS, Ferraro GE, et al. Flavonoids as antioxidants evaluated by in vitro and in situ liver chemiluminescence. *Biochem Pharmacol*. 1987;36:717–720.
56. Guo Q, Zhao B, Li M, et al. Studies on protective mechanisms of four components of green tea polyphenols against lipid peroxidation in synaptosomes. *Biochim Biophys Acta*. 1996;1304:210–222.
57. Kurosawa T, Itoh F, Nozaki A, et al. Suppressive effect of cocoa powder on atherosclerosis in Kurosawa and Kusanagi-hypercholesterolemic rabbits. *J Atheroscler Thromb*. 2005;12:20–28.
58. Lotito SB, Actis-Goretti L, Renart ML, et al. Influence of oligomer chain length on the antioxidant activity of procyanidins. *Biochem Biophys Res Commun*. 2000;276:945–951.
59. Lotito SB, Fraga CG. (+)-Catechin prevents human plasma oxidation. *Free Radic Biol Med*. 1998;24:435–441.
60. Lotito SB, Fraga CG. Catechins delay lipid oxidation and alpha-tocopherol and beta-carotene depletion following ascorbate depletion in human plasma. *Proc Soc Exp Biol Med*. 2000;225:32–38.
61. Verstraeten SV, Hammerstone JF, Keen CL, et al. Antioxidant and membrane effects of procyanidin dimers and trimers isolated from peanut and cocoa. *J Agric Food Chem*. 2005;53:5041–5048.
62. Orozco TJ, Wang JF, Keen CL. Chronic consumption of a flavanol- and procyanidin-rich diet is associated with reduced levels of 8-hydroxy-2'-deoxyguanosine in rat testes. *J Nutr Biochem*. 2003;14:104–110.
63. Osakabe N, Natsume M, Adachi T, et al. Effects of cacao liquor polyphenols on the susceptibility of low-density lipoprotein to oxidation in hypercholesterolemic rabbits. *J Atheroscler Thromb*. 2000;7:164–168.
64. Videla LA, Fraga CG, Koch OR, et al. Chemiluminescence of the in situ rat liver after acute ethanol intoxication—effect of (+)-cyanidanol-3. *Biochem Pharmacol*. 1983;32:2822–2825.
65. Fraga CG. Plant polyphenols: how to translate their in vitro antioxidant actions to in vivo conditions. *IUBMB Life*. 2007;59:308–315.
66. Fraga CG. Cocoa, diabetes, and hypertension: should we eat more chocolate? *Am J Clin Nutr*. 2005;81:541–542.
67. Radi R. Nitric oxide, oxidants, and protein tyrosine nitration. *Proc Natl Acad Sci U S A*. 2004;101:4003–4008.
68. Ghafourifar P, Richter C. Nitric oxide synthase activity in mitochondria. *FEBS Lett*. 1997;418:291–296.
69. Giulivi C, Poderoso JJ, Boveris A. Production of nitric oxide by mitochondria. *J Biol Chem*. 1998;273:11038–11043.
70. Boveris A, Cadenas E. Mitochondrial production of superoxide anions and its relationship to the antimycin insensitive respiration. *FEBS Lett*. 1975;54:311–314.
71. Turrens JF, Boveris A. Generation of superoxide anion by the NADH dehydrogenase of bovine heart mitochondria. *Biochem J*. 1980;191:421–427.
72. Sarkis A, Roman RJ. Role of cytochrome P450 metabolites of arachidonic acid in hypertension. *Curr Drug Metab*. 2004;5:245–256.
73. Ward R, Cirkovic-Velichovia T, Ledequ F, et al. Neuroprotection by taurine and taurine analogues. *Adv Exp Med Biol*. 2006;583:299–306.
74. Jones SA, O'Donnell VB, Wood JD, et al. Expression of phagocyte NADPH oxidase components in human endothelial cells. *Am J Physiol*. 1996;271:H1626–H1634.
75. Touyz RM. Recent advances in intracellular signalling in hypertension. *Curr Opin Nephrol Hypertens*. 2003;12:165–174.
76. Park YC, Rimbach G, Saliou C, et al. Activity of monomeric, dimeric, and trimeric flavonoids on NO production, TNF-alpha secretion, and NF-kappaB-dependent gene expression in RAW 264.7 macrophages. *FEBS Lett*. 2000;465:93–97.
77. Jovanovic SD, Steenken S, Tosic M, et al. Flavonoids as antioxidants. *J Am Chem Soc*. 1994;116:4846–4851.
78. Oteiza PI, Erlejtman AG, Verstraeten SV, et al. Flavonoid-membrane interactions: a protective role of flavonoids at the membrane surface? *Clin Dev Immunol*. 2005;12:19–25.
79. Rosenkranz S, Knirel D, Dietrich H, et al. Inhibition of the PDGF receptor by red wine flavonoids provides a molecular explanation for the “French paradox”. *FASEB J*. 2002;16:1958–1960.
80. Tachibana H, Koga K, Fujimura Y, et al. A receptor for green tea polyphenol EGCG. *Nat Struct Mol Biol*. 2004;11:380–381.
81. Schewe T, Kuhn H, Sies H. Flavonoids of cocoa inhibit recombinant human 5-lipoxygenase. *J Nutr*. 2002;132:1825–1829.
82. Schewe T, Sadik C, Klotz LO, et al. Polyphenols of cocoa: inhibition of mammalian 15-lipoxygenase. *Biol Chem*. 2001;382:1687–1696.
83. Lee KW, Kundu JK, Kim SO, et al. Cocoa polyphenols inhibit phorbol ester-induced superoxide anion formation in cultured HL-60 cells and expression of cyclooxygenase-2 and activation of NF-kappaB and MAPKs in mouse skin in vivo. *J Nutr*. 2006;136:1150–1155.
84. Zhang WY, Liu HQ, Xie KQ, et al. Procyanidin dimer B2 [epicatechin-(4beta-8)-epicatechin] suppresses the expression of cyclooxygenase-2 in endotoxin-treated monocytic cells. *Biochem Biophys Res Commun*. 2006;345:508–515.
85. Oak MH, El Bedoui J, Anglard P, et al. Red wine polyphenolic compounds strongly inhibit pro-matrix metalloproteinase-2 expression and its activation in response to thrombin via direct inhibition of membrane type 1-matrix metalloproteinase in vascular smooth muscle cells. *Circulation*. 2004;110:1861–1867.
86. Ottaviani JJ, Carrasquedo F, Keen CL, et al. Influence of flavan-3-ols and procyanidins on UVC-mediated formation of 8-oxo-7,8-dihydro-2'-deoxyguanosine in isolated DNA. *Arch Biochem Biophys*. 2002;406:203–208.
87. Fraga CG, Oteiza PI. Flavanols and NF-κB activation: relevance for inflammation and associated diseases. Surh YJ, Dong Z, Cadenas E, et al, eds. *Dietary Modulation of Cell Signaling Pathways*. Boca Raton, FL: CRC Press; 2008:137–151.
88. Khan N, Afaq F, Saleem M, et al. Targeting multiple signaling pathways by green tea polyphenol (–)-epigallocatechin-3-gallate. *Cancer Res*. 2006;66:2500–2505.
89. Surh YJ. Cancer chemoprevention with dietary phytochemicals. *Nat Rev Cancer*. 2003;3:768–780.
90. Mackenzie GG, Oteiza PI. Modulation of transcription factor NF-kappaB in Hodgkin's lymphoma cell lines: effect of (–)-epicatechin. *Free Radic Res*. 2006;40:1086–1094.
91. Erlejtman AG, Jagers G, Fraga CG, et al. TNFalpha-induced NF-kappaB activation and cell oxidant production are modulated by hexameric procyanidins in Caco-2 cells. *Arch Biochem Biophys*. 2008;476:186–195.
92. Verstraeten SV, Mackenzie GG, Oteiza PI, et al. (–)-Epicatechin and related procyanidins modulate intracellular calcium and prevent oxidation in Jurkat T cells. *Free Radic Res*. 2008;42:864–872.
93. Shepherd J, Cobbe SM, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med*. 1995;333:1301–1307.
94. Stamler J, Wentworth D, Neaton JD. Is relationship between serum cholesterol and risk of premature death from coronary heart disease continuous and graded? Findings in 356,222 primary screeners of the

- Multiple Risk Factor Intervention Trial (MRFIT). *J Am Med Assoc.* 1986;256:2823–2828.
95. Borradaile NM, de Dreu LE, Wilcox LJ, et al. Soya phytoestrogens, genistein and daidzein, decrease apolipoprotein B secretion from HepG2 cells through multiple mechanisms. *Biochem J.* 2002;366:531–539.
 96. Ikeda I, Imasato Y, Sasaki E, et al. Tea catechins decrease micellar solubility and intestinal absorption of cholesterol in rats. *Biochim Biophys Acta.* 1992;1127:141–146.
 97. Pal S, Ho N, Santos C, et al. Red wine polyphenolics increase LDL receptor expression and activity and suppress the secretion of ApoB100 from human HepG2 cells. *J Nutr.* 2003;133:700–706.
 98. Wilcox LJ, Borradaile NM, de Dreu LE, et al. Secretion of hepatocyte apoB is inhibited by the flavonoids, naringenin and hesperetin, via reduced activity and expression of ACAT2 and MTP. *J Lipid Res.* 2001;42:725–734.
 99. Baba S, Natsume M, Yasuda A, et al. Plasma LDL and HDL cholesterol and oxidized LDL concentrations are altered in normo- and hypercholesterolemic humans after intake of different levels of cocoa powder. *J Nutr.* 2007;137:1436–1441.
 100. Fraga CG, Actis-Goretta L, Ottaviani JI, et al. Regular consumption of a flavanol-rich chocolate can improve oxidant stress in young soccer players. *Clin Dev Immunol.* 2005;12:11–17.
 101. Grassi D, Desideri G, Necozione S, et al. Blood pressure is reduced and insulin sensitivity increased in glucose-intolerant, hypertensive subjects after 15 days of consuming high-polyphenol dark chocolate. *J Nutr.* 2008;138:1671–1676.
 102. Baba S, Osakabe N, Kato Y, et al. Continuous intake of polyphenolic compounds containing cocoa powder reduces LDL oxidative susceptibility and has beneficial effects on plasma HDL-cholesterol concentrations in humans. *Am J Clin Nutr.* 2007;85:709–717.
 103. Mursu J, Voutilainen S, Nurmi T, et al. Dark chocolate consumption increases HDL cholesterol concentration and chocolate fatty acids may inhibit lipid peroxidation in healthy humans. *Free Radic Biol Med.* 2004;37:1351–1359.
 104. Steinberg D, Parthasarathy S, Carew TE, et al. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med.* 1989;320:915–924.
 105. Kondo K, Hirano R, Matsumoto A, et al. Inhibition of LDL oxidation by cocoa. *Lancet.* 1996;348:1514.
 106. Mathur S, Devaraj S, Grundy SM, et al. Cocoa products decrease low density lipoprotein oxidative susceptibility but do not affect biomarkers of inflammation in humans. *J Nutr.* 2002;132:3663–3667.
 107. Osakabe N, Baba S, Yasuda A, et al. Daily cocoa intake reduces the susceptibility of low-density lipoprotein to oxidation as demonstrated in healthy human volunteers. *Free Radic Res.* 2001;34:93–99.
 108. Rein D, Lotito S, Holt RR, et al. Epicatechin in human plasma: in vivo determination and effect of chocolate consumption on plasma oxidation status. *J Nutr.* 2000;130:S2109–S2114.
 109. Wan Y, Vinson JA, Etherton TD, et al. Effects of cocoa powder and dark chocolate on LDL oxidative susceptibility and prostaglandin concentrations in humans. *Am J Clin Nutr.* 2001;74:596–602.
 110. Renaud S, de Lorgeril M. Wine, alcohol, platelets, and the French paradox for coronary heart disease. *Lancet.* 1992;339:1523–1526.
 111. McCullough ML, Chevaux K, Jackson L, et al. Hypertension, the Kuna, and the epidemiology of flavanols. *J Cardiovasc Pharmacol.* 2006;47:S103–S109.
 112. Grassi D, Lippi C, Necozione S, et al. Short-term administration of dark chocolate is followed by a significant increase in insulin sensitivity and a decrease in blood pressure in healthy persons. *Am J Clin Nutr.* 2005;81:611–614.
 113. Taubert D, Berkels R, Roesen R, et al. Chocolate and blood pressure in elderly individuals with isolated systolic hypertension. *J Am Med Assoc.* 2003;290:1029–1030.
 114. Taubert D, Roesen R, Schomig E. Effect of cocoa and tea intake on blood pressure: a meta-analysis. *Arch Intern Med.* 2007;167:626–634.
 115. Faridi Z, Njike VY, Dutta S, et al. Acute dark chocolate and cocoa ingestion and endothelial function: a randomized controlled crossover trial. *Am J Clin Nutr.* 2008;88:58–63.
 116. Ralay Ranaivo H, Diebolt M, Andriantsitohaina R. Wine polyphenols induce hypotension, and decrease cardiac reactivity and infarct size in rats: involvement of nitric oxide. *Br J Pharmacol.* 2004;142:671–678.
 117. Bingham SA, Vorster H, Jerling JC, et al. Effect of black tea drinking on blood lipids, blood pressure and aspects of bowel habit. *Br J Nutr.* 1997;78:41–55.
 118. Negishi H, Xu JW, Ikeda K, et al. Black and green tea polyphenols attenuate blood pressure increases in stroke-prone spontaneously hypertensive rats. *J Nutr.* 2004;134:38–42.
 119. Stensvold I, Tverdal A, Solvoll K, et al. Tea consumption. Relationship to cholesterol, blood pressure, and coronary and total mortality. *Prev Med.* 1992;21:546–553.
 120. Yang YC, Lu FH, Wu JS, et al. The protective effect of habitual tea consumption on hypertension. *Arch Intern Med.* 2004;164:1534–1540.
 121. Muniyappa R, Hall G, Kolodziej TL, et al. Cocoa consumption for 2 wk enhances insulin-mediated vasodilatation without improving blood pressure or insulin resistance in essential hypertension. *Am J Clin Nutr.* 2008;88:1685–1696.
 122. Osterud B. A global view on the role of monocytes and platelets in atherogenesis. *Thromb Res.* 1997;85:1–22.
 123. Pearson DA, Holt RR, Rein D, et al. Flavanols and platelet reactivity. *Clin Dev Immunol.* 2005;12:1–9.
 124. Rein D, Paglieroni TG, Wun T, et al. Cocoa inhibits platelet activation and function. *Am J Clin Nutr.* 2000;72:30–35.
 125. Hermann F, Spieker LE, Ruschitzka F, et al. Dark chocolate improves endothelial and platelet function. *Heart.* 2006;92:119–120.
 126. Flammer AJ, Hermann F, Sudano I, et al. Dark chocolate improves coronary vasomotion and reduces platelet reactivity. *Circulation.* 2007;116:2376–2382.
 127. de Graaf JC, Banga JD, Moncada S, et al. Nitric oxide functions as an inhibitor of platelet adhesion under flow conditions. *Circulation.* 1992;85:2284–2290.
 128. Pignatelli P, Di Santo S, Buchetti B, et al. Polyphenols enhance platelet nitric oxide by inhibiting protein kinase C-dependent NADPH oxidase activation: effect on platelet recruitment. *FASEB J.* 2006;20:1082–1089.
 129. Pignatelli P, Pulcinelli FM, Celestini A, et al. The flavonoids quercetin and catechin synergistically inhibit platelet function by antagonizing the intracellular production of hydrogen peroxide. *Am J Clin Nutr.* 2000;72:1150–1155.
 130. Schramm DD, Wang JF, Holt RR, et al. Chocolate procyanidins decrease the leukotriene-prostacyclin ratio in humans and human aortic endothelial cells. *Am J Clin Nutr.* 2001;73:36–40.
 131. Navarro-Nunez L, Castillo J, Lozano ML, et al. Thromboxane A2 receptor antagonism by flavonoids: structure-activity relationships. *J Agric Food Chem.* 2009;57:1589–1594.
 132. Sadik CD, Sies H, Schewe T. Inhibition of 15-lipoxygenases by flavonoids: structure-activity relations and mode of action. *Biochem Pharmacol.* 2003;65:773–781.
 133. Vasquez-Martinez Y, Ohri RV, Kenyon V, et al. Structure-activity relationship studies of flavonoids as potent inhibitors of human platelet 12-hLO, reticulocyte 15-hLO-1, and prostate epithelial 15-hLO-2. *Bioorg Med Chem.* 2007;15:7408–7425.
 134. Chun OK, Chung SJ, Claycombe KJ, et al. Serum C-reactive protein concentrations are inversely associated with dietary flavonoid intake in U.S. adults. *J Nutr.* 2008;138:753–760.
 135. di Giuseppe R, Di Castelnuovo A, Centritto F, et al. Regular consumption of dark chocolate is associated with low serum concentrations of C-reactive protein in a healthy Italian population. *J Nutr.* 2008;138:1939–1945.
 136. Ramiro-Puig E, Castell M. Cocoa: antioxidant and immunomodulator. *Br J Nutr.* 2009;101:931–940.
 137. Selmi C, Cocchi CA, Lanfredini M, et al. Chocolate at heart: the anti-inflammatory impact of cocoa flavanols. *Mol Nutr Food Res.* 2008;52:1340–1348.
 138. Cho ES, Jang YJ, Kang NJ, et al. Cocoa procyanidins attenuate 4-hydroxynonenal-induced apoptosis of PC12 cells by directly inhibiting mitogen-activated protein kinase kinase 4 activity. *Free Radic Biol Med.* 2009;46:1319–1327.
 139. Kang NJ, Lee KW, Lee DE, et al. Cocoa procyanidins suppress transformation by inhibiting mitogen-activated protein kinase kinase. *J Biol Chem.* 2008;283:20664–20673.
 140. Jenny M, Santer E, Klein A, et al. Cacao extracts suppress tryptophan degradation of mitogen-stimulated peripheral blood mononuclear cells. *J Ethnopharmacol.* 2009;122:261–267.