

Oviductal fluid counterbalances the negative effect of high temperature on sperm
in an ectotherm model

Authors: Rossi N^{1,2}, Lopez Juri G^{1,2}, Chiaraviglio M^{1,2} and Cardozo G^{1,2}

Affiliation:

¹ Universidad Nacional de Córdoba, Facultad de Ciencias Exactas Físicas y Naturales. Laboratorio de Biología del Comportamiento; Córdoba, Argentina.

² Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Instituto de Diversidad y Ecología Animal (IDEA), Córdoba, Argentina.

Corresponding author: Nicola Rossi

Email address of corresponding author: nicola.rossi20@gmail.com

ORCID identifier: 0000-0003-0780-7117

Telephone number of corresponding author: +54 3512274631

Keywords: Sperm dynamic - Sperm motility - Temperature - Oviductal fluid - Global warming

Summary statement: Our work found that oviductal fluid enhances sperm dynamics and survival and mitigates the negative effects of increasing temperatures on sperm in an ectotherm model.

Abstract. Global warming is affecting biodiversity; however, the extent to which animal reproductive processes respond to predicted temperature increments remains largely unexplored. The thermal environment has a pronounced impact on metabolic rates of ectotherms; therefore, an interesting question to assess is whether temperature increase might affect specific reproductive mechanisms like sperm performance in ectotherms. Moreover, in many species, oviductal fluid (OF) is known to regulate and maintain sperm quality; however, the role of oviductal fluid in relation to the effects of high temperature on sperm remains unclear. Our aim was to experimentally test the effect of increased temperature on sperm velocity, swimming path and percentage of motility in neutral conditions at ejaculation (without OF) and in female's reproductive tract fluid (with OF), in a social ectotherm lizard model, *Tropidurus spinulosus*, which has specific thermal requirements for reproduction. Our results suggest that a rising temperature associated with global warming (+4°C) affects negatively sperm dynamics and survival. However, OF fluid ameliorated the harmful effects of high temperature. This is an important point, as this study is the first that ever tested the role of OF to preserve sperm from a warmer pre-fertilization environment. These results contribute to our understanding of how thermal environment changes might affect post-copulatory reproductive mechanisms.

Introduction

Increase of global surface temperatures, frequent heatwaves and extreme temperature events have been predicted by the Intergovernmental Panel on Climate Change (IPCC) for the end of the century (Allen et al., 2018; Stocker et al., 2013). Global warming is affecting biodiversity, however, there is insufficient data concerning physiological and performance impacts to extreme temperatures in natural populations (Cooper et al., 2020; Vicenzi et al., 2020). Specifically, the extent to which animal reproductive traits respond to the predicted temperature increments, remains largely unexplored (Bestion et al., 2015; Dunn and Winkler, 2010; Isaac, 2009; Kingsolver et al., 2013); and reproductive sensitivity to temperature variations is better known in endotherms (Hurley et al., 2018).

By contrast, limited attention has been given to ectotherm taxa, in which reproductive processes are more directly influenced by thermal environment changes (Carey, 2014; Dubey and Shine, 2011; Paaajmans et al., 2013). In the context of climate change, both thermoregulatory behaviour and thermal physiology can be targets of selection in ectotherms (Gilbert and Miles, 2017). However, evolutionary changes in thermoregulatory capability are unlikely to keep pace with current rates of environmental change. Therefore, since the thermal environments have a pronounced impact on body temperatures of ectotherms, temperature increase is hypothesized to impact specific reproductive mechanisms, like sperm performance.

The temperature of the different pre-fertilization environments to which the sperm cells are exposed can have many fundamental effects on sperm performance and function (Fenkes et al., 2017a; Sinclair et al., 2016). In fishes, increasing temperatures related to global warming affect sperm cells metabolism (Dadras et al., 2017), beat frequency of sperm flagella and make viability decrease leading to fewer motile cells (Mansour et al., 2002). In terrestrial ectotherms such as insect groups, heatwaves compromise male fertility by reducing sperm number and viability, thus lessening its competitive ability (Gasparini et al., 2018; Sales et al., 2018). Moreover, not only average sperm parameters but also their intramale variability, which has been identified as a sperm competition strategy to maximize their fertilization success (Blengini et al., 2014; Calhim et al., 2011; Helfenstein et al., 2010), might be affected by warmer conditions.

Mechanisms in the female reproductive tract (FRT), such as sperm storage and gamete interaction, are also affected by temperature increments (Sales et al., 2018). Some mechanisms mediated by female fluids offer females the opportunity to exert post-mating female choice in several taxa (Devigili et al., 2018; Firman et al., 2017; Fitzpatrick et al., 2020). Oviductal fluid (OF) may actively enhance the motility of sperm by increasing differentially the sperm velocity of some males; this phenomenon leads to the Cryptic Female Choice (CFC) (Cardozo and Pilastro, 2018; Gasparini and Pilastro, 2011). Consequently, sperm performance in the female medium could be affected by female factors evolved as a consequence of sexual conflict (Cardozo et al., 2020; Møller et al., 2009). However, a question that has not been addressed yet is whether the role of oviductal fluid gets modified due to the effects of high temperature. Rising global surface temperatures (1.1 to 6.4°C) by 2100 would increase the metabolic rates of ectotherms by 10-75%, which could negatively affect reproduction. In the light of this alarming projection (Huey et al., 2009), testing if there is an interaction between temperature increments and OF effect on sperm performance is a mandatory first step to understand evolutionary responses in a changing environment.

Reproductive processes may involve specific thermal requirements (e.g. preferred temperatures, a suitable thermal range and thermal stability) in ectotherms both in males and females (Fenkes et al., 2017b; López Juri et al., 2018a). Furthermore, the effects of temperature on sperm motility and velocity are related to the reproductive thermal ecology of each species (Tourmente et al., 2011a). Specifically, temperature of microenvironments plays a pivotal role in sperm performance in multiple instances, both before and after copulation (Dosemane and Bhagya, 2015). Therefore, assessing sperm performance when sperm is ejaculated, as well as under post-mating conditions, is fundamental to understand the different mechanistic bases of sperm sensitivity to temperature (Marshall, 2015).

To test the interaction between OF and sperm, effective and safe procedures to obtain both sperm and OF samples are required. In lizards, sperm collection has been addressed (López Juri et al., 2018); however, non-lethal and non-invasive OF collection protocols are not available. In birds and fishes, some techniques have been used successfully (Cardozo and Pilastro, 2018; Cramer et al., 2014; Devigili et al., 2018; Gasparini and Pilastro, 2011). In our study, we novelly adapted them to lizards providing a thorough and safe method for OF collection to test the impact of temperature on sperm-OF interaction under a climate change scenario.

Tropidurus spinulosus, the spiny lava lizard, is a suitable ectotherm model for this study because reproductive individuals have specific thermal requirements (López Juri et al., 2018a). Moreover, it shows intense social reproductive interactions and male-biased operative sex ratio (López Juri, 2019), which leads to sperm competition and CFC (Pizzari and Wedell, 2013). The species distribution in temperate areas makes it an interesting model because warming would be more marked in these regions (Allen et al., 2018; Stocker et al., 2013) and because species adapted to seasonality would be more sensitive to climate change (Tang et al., 2012).

Our aim was to experimentally test the effect of increased temperature associated to an extreme global warming on sperm velocity, swimming path and percentage of motile sperm under neutral conditions at ejaculation (without OF) and to evaluate the role of female's reproductive tract fluid (with OF) on the sperm subjected to high temperature in an ectotherm model. We predict that a significant temperature increment above the preferred temperature documented for reproductive individuals will impair sperm velocity and motility whereas OF interaction will enhance the spermiatic parameters.

Materials and methods

Capture and handling

Reproductive females (n=15) and males (n=27) of *T. spinulosus* were captured in Córdoba province, Argentina (31°23'33''W, 64°35'48''S) during two consecutive breeding seasons (2017-2018) (López Juri et al., 2018a).

As reproduction in *T. spinulosus* is seasonal, we evaluated reproductive structures (by ultrasound scanning Sonosite 180 Plus) to confirm the stage of the reproductive cycle of males and females. We used receptive females with enlarged vitellogenic follicles and reproductive males with enlarged testes as they possess sperm in their deferent ducts (López Juri et al., 2018b). The individuals were isolated for one week to ensure that no recent inseminated sperm was available in the cloaca of females, and that males could replenish sperm reserves (Gasparini et al., 2019). Lizards were kept individually under light (9-17hs, UVB 5.0) and at 28°C (mean environmental temperature from capture site in their natural habitat); larvae of *Tenebrio molitor* and water were provided *ad libitum*.

This research was approved by the Governmental Environmental Agency and the Ethical Committee of the Instituto de Diversidad y Ecología Animal CONICET-UNC (protocol number: 2/2017 and 12/2019).

Experimental design: temperature treatments in relation to IPCC projections

In brief, we collected semen and OF of individuals from wildlife. Semen of each male was tested under the concomitant effect of two factors: presence/absence of OF and two temperature levels. OF presence/absence will allow us to disentangle the effect of OF from sperm performance *per se*. In regards to temperature levels, IPCC projects climate warming within the range from 2°C to 4.5°C for the end of the century (Allen et al., 2018; Stocker et al., 2013). Moreover, in the absence of near-term mitigation to reduce emissions, the likelihood of 4 °C warming being reached during the present century has increased (Fawcett et al., 2015). Consequently, we tested the following temperature treatments: a) 34°C, which corresponds to the preferred temperature of males and receptive females during the breeding season (Tsel_females: 33.79±1.42°C; Tsel_males: 33.50±2.27 °C; López Juri et al, 2018a; Thermal preference is the body temperature selected in the absence of ecological constraints; see Gilbert and Miles, 2017); and b) 38°C, according to a +4°C increment in surface temperatures projected by the Representative Concentration Pathway 8.5 (RCP 8.5, Allen et al., 2018; Stocker et al., 2013). Thirty-eight degrees is close to the upper thermal range tolerated by *Tropidurus spinulosus*, however it is still a temperature they may voluntarily select to thermoregulate in a laboratory setup (López Juri et al., 2018).

OF collection

Before beginning OF collection, the cloaca was rinsed with sterile saline solution (0.9% NaCl) to eliminate any pasty urates and feces. Initially, a ventro-lateral stroking was performed to achieve relaxation of the musculature. Later the anal plate was massaged to produce distension of the cloacal musculature (Mengden et al., 1980). After that, the cloaca aperture was rubbed with a metal probe by performing circular movements. These movements favoured the relaxation of the cloacal musculature and therefore allowed us to locate the urinary papillae; the latter were gently lowered to expose the entrance to the oviduct where OF was then collected. There, 10 µl of the saline solution was injected and retrieved using a micropipette. This operation was repeated four times; a total 40 µl of saline solution was injected and approximately 25 µl of OF solution (OF + saline solution) was collected obtaining an OF dilution similar to that used in CFC studies (Cardozo and Pilastro, 2018; Gasparini et al., 2012). OF solution was examined under optical microscope to confirm the absence of recently inseminated sperm.

Sperm collection

The lizards were electrostimulated according to (López Juri et al., 2018b) to trigger the expulsion of semen into the cloacal ampulla. We added 10 µl of saline solution to the expelled semen: this mixture was immediately aspirated with a micropipette and then rinsed with into 10 µl of saline solution. This procedure was repeated until all samples were completed to a fixed volume of 20 µl of semen solution (semen + saline solution). A mean sperm number of $2.1 \pm 1.8 \times 10^6$ spermatozoa was obtained. Because the number of captured receptive females was nearly half of reproductive males, the OF solution from each female was usually tested with sperm samples of two males.

Animal welfare

After semen/OF collection, lizards were checked once a week for normal alertness, calm chemical sample of surrounding, unhurried locomotion, relaxed feeding and breathing (Warwick et al., 2013). After that, the specimens were released at the capture sites (GPS coordinates).

Experimental protocol

To simulate sperm ejaculation and post-mating conditions, sperm velocity was measured in two different solutions (Fig. S1): A) OF absence treatment (OF-): consisting of 8 µl (40% volume) of Ham's F-10 culture medium (Ham's F-10. Gibco, New York, USA), supplemented with 1% bovine serum albumin, and 12 µl (60% volume) of saline solution; B) OF presence treatment (OF+): consisting of 8 µl (40 % volume) of similarly supplemented Ham's F-10 culture medium and 12 µl (60% volume) of OF solution; this proportions were adjusted according to (Gasparini and Pilastro, 2011). Semen solution (2 µl) was added to the A and B solutions, in random order. The total volume (22µl) of each solution was then divided and used for both incubation temperature treatments: 34°C and 38°C. Thus, the sample series was composed of four treatments: 1) OF+ 34°C, 2) OF+ 38°C, 3) (OF- 34°C), and 4) (OF- 38°C). Samples were incubated in thermal baths (Thermo Scientific Precision™) for 15 minutes which are sufficient to observe relevant effects on motility (Awda et al., 2009; Williams and Ford, 2005) and then processed in a randomized fashion. The total number of series was 27.

Sperm parameters

Dynamic parameters were measured using a video microscopy system (Eclipse 50i Nikon phase contrast; camera Nikon Digital Sight DS-Fi2). Sperm tracks were followed for 3s (mean \pm SD = 21 \pm 3 cells/sample) and transformed to a matrix of Cartesian coordinates using ImageJ version 1.43u (NIH) and its plug-in MtrackJ v. 1.1.0 (Meijering et al., 2012). Sperm dynamic parameters were calculated using Spermtrack v. 4.2 (Universidad Nacional de Cordoba, Argentina): straight line velocity (VSL; $\mu\text{m/s}$), curvilinear velocity (VCL; $\mu\text{m/s}$), and swimming path estimated as linearity (LIN = VSL/VCL). The percentage of motile spermatozoa was estimated over approximately 50 sperm cells per analysis with random changes of the microscope fields. Variability of the motility parameters (VSL, VCL and LIN) was evaluated by calculating Coefficients of Variation (CV), which is a useful statistics to evaluate sperm variability associated with motility (Blengini et al., 2014; Immler et al., 2008).

Statistical analysis

We fitted a linear mixed effect models with ‘Female ID’, ‘Male ID’ and ‘Year’ as random effects and ‘Temperature’ and ‘OF treatment’ and their interaction as fixed factors (package lme4, Bates et al., 2015). Percentage of motile spermatozoa was tested with a generalized linear model with a binomial link function and the same random and fixed effects as the linear models. Since ‘Year’ was always non- significant, we removed it from all models. Normality and homoscedasticity were checked by inspecting residuals distribution. Shapiro-Wilks test was also performed when checking linear models. *A posteriori* Tuckey test was performed to determine differences among treatments. Statistical analyses were performed in R version 3.6.1 (R Core Team, 2019).

Data availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Results

Straight line velocity (VSL; $\mu\text{m/s}$) and curvilinear velocity (VCL; $\mu\text{m/s}$) presented a similar pattern both varying among treatments; (Figure 1, Table 1). Sperm velocity was the highest at 34°C in (OF+). The increment of 4°C caused a reduction in spermatozoa velocity in both (OF-) and (OF+). However, considering the interaction, the sperm at high temperature in (OF+) was significantly faster

than in (OF-). Variability in curvilinear sperm velocity (VCL) was highest when temperature increased and in (OF-).

Under (OF-) conditions sperm linearity (LIN) was high and was not significantly modified by temperature increment. When sperm was under (OF+) conditions at 34°C spermatozoa linearity was low, however, at 38°C spermatozoa linearity increased (but with marginal significance), reaching similar values to (OF-) conditions.

The percentage of motile spermatozoa was the highest at 34°C in (OF+). Temperature increment caused a decrease in sperm motility. However, at high temperature the percentage of motile sperm in (OF+) was significantly higher than in (OF-).

Discussion

We found that a temperature increment associated to an extreme global warming affects negatively sperm velocity and percentage of motile cells. However, OF enhance sperm performance and ameliorated the negative effects of high temperature on sperm. Additionally, our method of OF extraction, based on other taxa (Cardozo and Pilastro, 2018; Møller et al., 2009) was safe and efficient, and was applied in squamates for the first time. The technique did not harm females as they did not show abnormal behaviour after the procedure (Warwick et al., 2013).

The reduction in the velocity of the spermatozoa caused by the temperature increment might impact fertilization success of vitellogenic follicles since sperm velocity is known to be a major determinant of male fertility (Beausoleil et al., 2012; Birkhead and Møller, 1998; Gomendio and Roldan, 2004). The decrease in sperm velocity might also affect the competitive ability of males; i.e. males under sperm competition risk often vary in sperm quality (Blengini et al., 2016; Cardozo et al., 2020); however, if warmer thermal environments modify sperm velocity, the ability of some males to reach the fertilization site faster might be altered.

In other squamates and mammals, high temperatures also harm sperm dynamic, possibly through an increase in sperm metabolic rates that alters enzymes or the resistance of the lipid membrane to Reactive Oxygen Species (ROS) (Soren et al., 2018; Tourmente et al., 2011b). In ectotherm marine species, warming environments can either boost (Ho et al., 2013) or harm sperm velocity (Binet and

Doyle, 2013). In internally fertilizing ectotherms, sperm production (Zeh et al., 2012) and performance within the female tract might be negatively impacted, since sperm deteriorates rapidly after being exposed to heatwaves (Sales et al., 2018).

On the other hand, although OF has already been identified as an improver of sperm motility in some fishes and birds (Rosengrave et al., 2009; Sasanami et al., 2013), in our lizard model, the OF effect on the increment of sperm velocity was notable (on average 20%) compared to that observed in other species (approximately 10% in guppies, Gasparini and Pilastro 2011 and Cardozo and Pilastro, 2018; 10% in bovines, (Grippio et al., 1995); 12% in boars, (Coy et al., 2010); 10-17% in mice, (Oliveira et al., 1999). The ions and protein composition naturally present in the OF may stimulate spermatozoa ATP metabolism and, thus, enhance motility (Rosengrave et al., 2009); a similar mechanism involving the pH of secreted materials in the birds vagina might modify sperm motility (Sasanami et al., 2013).

Remarkably the negative effect of high temperature on sperm velocity seems to be less pronounced in OF than in neutral conditions. The compensation effect of OF suggests that postcopulatory mechanism mediated by FRT fluids might partially protect sperm in the oviductal tract from increasing temperatures even when the increment is above the T_{set_max} and near to T_{max} (in *Tropidurus spinulosus* for reproductive females: T_{set_max} 35.34 ± 1.40 and T_{max} 37.49 ± 1.81 ; for reproductive males: T_{set_max} = 35.07 ± 2.04 and T_{max} 37.53 ± 2.00 ; see Lopez Juri et al 2018). As in mammals, the compensation might be mediated by OF enzymes that inhibit sperm ROS action (Kobayashi et al., 2014) or help in restoring motility of spermatozoa under oxidative stress (Bilodeau et al., 2002). This might be important in species with prolonged mate search where sperm parameters seem to be modelled by the temperature of the habitat that males pass through (Tourmente et al., 2011a).

OF may also activate sperm through capacitation, which in reptiles has been poorly studied. Some works in crocodiles and lizards added to recent evidence that reptilian sperm may have the full range of phosphorylation-mediated cellular mechanisms and activation of mitochondria associated with capacitation, drastically increasing motility (Nixon et al., 2016). The influence of temperature on capacitation events has also been largely neglected, however the few evidence available show different impacts on motility depending on the taxa considered. In bovines, incubation gradients (20°C-40°C) affect the % of hyperactivated sperm but not sperm velocity, while in humans the opposite is true (Hammerstedt and Hay, 1980). In reptiles these processes should be studied further

to properly quantify the relative contribution to motility of the capacitation mediated by OF and/or the protective effect of OF towards high temperatures.

To our knowledge velocity as a predictor of fertilization has not been tested as exclusive in squamates, so it is possible that other post-copulatory traits, such as sperm viability, are influential for fertilization success. The percentage of motile sperm is unaffected by temperature in some bird and fish species (Bonato et al., 2012), whereas high temperatures may have deleterious effects on sperm of squamates (Tourmente et al., 2011a). Our results support the latter, but also highlight the importance of OF in maintaining spermatozoa alive despite the thermal stress. Therefore, this study is the first that has ever tested the role of OF to preserve not only sperm performance but also the number of motile spermatozoa from the harmful effect of high temperature from a climate change perspective.

Sperm linearity at the preferred temperature and in OF was low. In other taxa, non-linear movement would reflect sperm cells, redirecting themselves towards a chemoattractant, like progesterone, through transitional movements (Blengini et al., 2011). Here, since the receptive females were at a late stage of vitellogenesis, progesterone level in the collected OF was probably high (Ramírez-Pinilla et al., 2009) and may have caused the observed non-linear pattern. In addition, linearity increased with temperature, which has been interpreted as a strategy to reach the follicles or storage sites earlier at the expense of lower cell viability (Mehlis and Bakker, 2014).

Variability in curvilinear sperm velocity (VCL) was highest when temperature increased and in (OF-). In other lizards, sperm variability has been found in relation to sperm competition risk (Blengini et al., 2014). In this study, the observed inter-male variability suggests differential responses of sperm performance to temperature increments, although OF apparently acts as a levelling factor. Sperm variability could be related to sperm age (Gasparini et al., 2019) or male traits e.g. body size, as occurs in *Tropidurus spinulosus* (López Juri et al., 2020). Moreover, trade-offs between colour ornamentation variability (Rossi et al., 2019) and sperm resistance to oxidative stress activated by increased temperature (Tomášek et al., 2017) might also produce variation in sperm velocity.

In this paper we focused on short-term mechanisms following ejaculation when the sperm reach the swimming environment within the female reproductive tract. However, the OF-temperature interaction might be important for longer term postcopulatory mechanisms such as CFC mediated by female sperm storage that is well known in lizards (Siegel et al., 2011). For example, the strong

linearization of sperm trajectory that we observed might alter sperm competence to reach the oviductal crypts, thus modifying CFC and sperm precedence patterns (Magris et al., 2017). However, to test it, experiments that consider exposure to temperature for longer time should be addressed. Furthermore, submitting the animals to temperature treatment instead of testing the sole interaction between OF and sperm, may highlight thermoregulatory response mechanisms that could function as mitigation strategies. Indeed, although lizard may choose to perform certain activities even when superficial temperatures are high and body temperature in the upper preferred range (Gunderson and Leal, 2016), they may regulate their body temperature behaviourally, since most ectotherms do not have a physiological thermal-safety margin (Sunday et al., 2014). By hiding in crevices, lizard may avoid unsuitable temperatures (Cecchetto et al., 2019), thus buffering reproductive mechanisms from high temperatures.

In conclusion, our study assesses potential evolutionary responses of reproductive processes of ectotherms to global warming. Temperature increments may modify sperm dynamics and survival, possibly affecting male reproductive success. By contrast, oviductal fluid enhanced sperm performance and remarkably ameliorated the harmful effects of high temperature. These results contribute to our understanding of how global warming might affect reproductive postcopulatory mechanisms.

Acknowledgements

We are very grateful to Andrea Pilastro, Clelia Gasparini and Alessandro Devigili for improving the manuscript with their comments. We thank the Graduate School of Biological Sciences of the National University of Córdoba and also the Private Natural Reserve “Cascada Los Chorrillos”, Flor Serrana, Tanti, Córdoba. We thank two anonymous reviewers for their interesting suggestions, which were implemented in the manuscript.

Funding

This study was funded by Consejo Nacional de Investigaciones Científicas y Técnicas CONICET PIP 2011e2013; CONICET PIP 2015e2017; Fondo para la Investigación Científica y Tecnológica PICT-2011- 1599 Res 140/12; PICT- BICENTENARIO Nro. 2010-2782; CONICET PUE-2016 D2555/16 and Animal Behavior Society due to Student Research Grant 2019.

REFERENCES

- Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., Kainuma, M., Kala, J., Mahowald, N., Muluguetta, Y., Perez, R., Wairiu, M., and Zickfeld, K. (2018). Framing and Context. In *Global Warming of 1.5 OC: An IPCC Special Report on the Impacts of Global Warming of 1.5° C Above Pre-industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty* (Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield). In Press.
- Awda, B. J., Mackenzie-Bell, M., and Buhr, M. M. (2009). Reactive oxygen species and boar sperm function. *Biology of Reproduction*, *81*(3), 553–561.
- Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B., Grothendieck, G., Green, P., and Bolker, M. B. (2015). Package ‘lme4.’ *Convergence*, *12*(1), 2.
- Beausoleil, J.-M. J., Doucet, S. M., Heath, D. D., and Pitcher, T. E. (2012). Spawning coloration, female choice and sperm competition in the redbside dace, *Clinostomus elongatus*. *Animal Behaviour*, *83*(4), 969–977.
- Bestion, E., Teyssier, A., Richard, M., Clobert, J., and Cote, J. (2015). Live Fast, Die Young: Experimental Evidence of Population Extinction Risk due to Climate Change. *PLoS Biology*, *13*(10), 1–19. Scopus. <https://doi.org/10.1371/journal.pbio.1002281>
- Bilodeau, J.-F., Blanchette, S., Cormier, N., and Sirard, M.-A. (2002). Reactive oxygen species-mediated loss of bovine sperm motility in egg yolk Tris extender: Protection by pyruvate, metal chelators and bovine liver or oviductal fluid catalase. *Theriogenology*, *57*(3), 1105–1122.
- Binet, M., and Doyle, C. (2013). Effect of near-future seawater temperature rises on sea urchin sperm longevity. *Marine and Freshwater Research*, *64*(1), 1–9.
- Birkhead, T. R., and Møller, A. P. (1998). *Sperm competition and sexual selection*. Academic Press.

- Blengini, C., Naretto, S., Cardozo, G., Giojalas, L. C., and Chiaraviglio, M. (2016). Relationship between pre-and post-copulatory traits in *Salvator rufescens* (Squamata: Teiidae). *Biological Journal of the Linnean Society*, *119*(4), 932–942.
- Blengini, C., Sergio, N., Gabriela, C., Giojalas, L. C., and Margarita, C. (2014). Variability in sperm form and function in the context of sperm competition risk in two Tupinambis lizards. *Ecology and Evolution*, *4*(21), 4080–4092. <https://doi.org/10.1002/ece3.1262>
- Blengini, C., Teves, M. E., Unates, D. R., Guidobaldi, H. A., Gatica, L. V., and Giojalas, L. C. (2011). Human sperm pattern of movement during chemotactic re-orientation towards a progesterone source. *Asian Journal of Andrology*, *13*(5), 769.
- Bonato, M., Cornwallis, C. K., Malecki, I. A., Rybnik-Trzaskowska, P. K., and Cloete, S. W. (2012). The effect of temperature and pH on the motility and viability of ostrich sperm. *Animal Reproduction Science*, *133*(1–2), 123–128.
- Calhim, S., Double, M. C., Margraf, N., Birkhead, T. R., and Cockburn, A. (2011). Maintenance of sperm variation in a highly promiscuous wild bird. *PLoS One*, *6*(12), e28809.
- Cardozo, G., Devigili, A., Antonelli, P., and Pilastro, A. (2020). Female sperm storage mediates postcopulatory costs and benefits of ejaculate anticipatory plasticity in the guppy. *Journal of Evolutionary Biology*.
- Cardozo, G., and Pilastro, A. (2018). Female nutritional condition affects ovarian fluid quality in guppies. *Biology Letters*, *14*(5), 20180122.
- Carey, C. (2014). Climate change, extinction risks, and reproduction of terrestrial vertebrates. In *Reproductive sciences in animal conservation* (pp. 35–54). Springer.
- Cecchetto, N. R., Medina, S. M., Taussig, S., and Ibarquengoytía, N. R. (2019). The lizard abides: Cold hardiness and winter refuges of *Liolaemus pictus argentinus* in Patagonia, Argentina. *Canadian Journal of Zoology*, *97*(9), 773–782.
- Cooper, C. E., Hurley, L. L., Deviche, P., and Griffith, S. C. (2020). Physiological responses of wild zebra finches (*Taeniopygia guttata*) to heatwaves. *Journal of Experimental Biology*.
- Coy, P., Lloyd, R., Romar, R., Satake, N., Matas, C., Gadea, J., and Holt, W. V. (2010). Effects of porcine pre-ovulatory oviductal fluid on boar sperm function. *Theriogenology*, *74*(4), 632–642.

- Cramer, E. R., Laskemoen, T., Eroukhmanoff, F., Haas, F., Hermansen, J. S., Lifjeld, J. T., Rowe, M., Slatre, G.-P., and Johnsen, A. (2014). Testing a post-copulatory pre-zygotic reproductive barrier in a passerine species pair. *Behavioral Ecology and Sociobiology*, *68*(7), 1133–1144.
- Dadras, H., Sampels, S., Golpour, A., Dzyuba, V., Cosson, J., and Dzyuba, B. (2017). Analysis of common carp *Cyprinus carpio* sperm motility and lipid composition using different in vitro temperatures. *Animal Reproduction Science*, *180*, 37–43.
- Devigili, A., Fitzpatrick, J. L., Gasparini, C., Ramnarine, I. W., Pilastro, A., and Evans, J. P. (2018). Possible glimpses into early speciation: The effect of ovarian fluid on sperm velocity accords with post-copulatory isolation between two guppy populations. *Journal of Evolutionary Biology*, *31*(1), 66–74. Scopus. <https://doi.org/10.1111/jeb.13194>
- Dosemane, D., and Bhagya, M. (2015). In vitro Study of the Spermatozoa Motility in the Lizard *Eutropis carinata*. *International Journal of Zoological Research*, *11*(3), 89.
- Dubey, S., and Shine, R. (2011). Predicting the effects of climate change on reproductive fitness of an endangered montane lizard, *Eulamprus leuraensis* (Scincidae). *Climatic Change*, *107*(3–4), 531–547.
- Dunn, P. O., and Winkler, D. W. (2010). Effects of climate change on timing of breeding and reproductive success in birds. *Effects of Climate Change on Birds*, 113–128.
- Fawcett, A. A., Iyer, G. C., Clarke, L. E., Edmonds, J. A., Hultman, N. E., McJeon, H. C., Rogelj, J., Schuler, R., Alsalam, J., Asrar, G. R., and others. (2015). Can Paris pledges avert severe climate change? *Science*, *350*(6265), 1168–1169.
- Fenkes, M., Fitzpatrick, J. L., Ozolina, K., Shiels, H. A., and Nudds, R. L. (2017a). Sperm in hot water: Direct and indirect thermal challenges interact to impact on brown trout sperm quality. *Journal of Experimental Biology*, *220*(14), 2513–2520.
- Fenkes, M., Fitzpatrick, J. L., Ozolina, K., Shiels, H. A., and Nudds, R. L. (2017b). Sperm in hot water: Direct and indirect thermal challenges interact to impact on brown trout sperm quality. *Journal of Experimental Biology*, *220*(14), 2513–2520.
- Firman, R. C., Gasparini, C., Manier, M. K., and Pizzari, T. (2017). Postmating female control: 20 years of cryptic female choice. *Trends in Ecology and Evolution*, *32*(5), 368–382.

- Fitzpatrick, J. L., Willis, C., Devigili, A., Young, A., Carroll, M., Hunter, H. R., and Brison, D. R. (2020). Chemical signals from eggs facilitate cryptic female choice in humans. *Proceedings of the Royal Society B*, 287(1928), 20200805.
- Gasparini, C., Andreatta, G., and Pilastro, A. (2012). Ovarian fluid of receptive females enhances sperm velocity. *Naturwissenschaften*, 99(5), 417–420.
- Gasparini, C., Devigili, A., and Pilastro, A. (2019). Sexual selection and ageing: Interplay between pre-and post-copulatory traits senescence in the guppy. *Proceedings of the Royal Society B*, 286(1897), 20182873.
- Gasparini, C., Lu, C., Dingemanse, N. J., and Tuni, C. (2018). Paternal-effects in a terrestrial ectotherm are temperature dependent but no evidence for adaptive effects. *Functional Ecology*, 32(4), 1011–1021.
- Gasparini, C., and Pilastro, A. (2011). Cryptic female preference for genetically unrelated males is mediated by ovarian fluid in the guppy. *Proceedings of the Royal Society of London B: Biological Sciences*, 278(1717), 2495–2501.
- Gilbert, A. L., and Miles, D. B. (2017). Natural selection on thermal preference, critical thermal maxima and locomotor performance. *Proceedings of the Royal Society B: Biological Sciences*, 284(1860), 20170536.
- Gomendio, M., and Roldan, E. R. (2004). Implications of diversity in sperm size and function for sperm competition and fertility. *International Journal of Developmental Biology*, 52(5–6), 439–447.
- Grippio, A., Way, A., and Killian, G. (1995). Effect of bovine ampullary and isthmic oviductal fluid on motility, acrosome reaction and fertility of bull spermatozoa. *Reproduction*, 105(1), 57–64.
- Gunderson, A. R., and Leal, M. (2016). A conceptual framework for understanding thermal constraints on ectotherm activity with implications for predicting responses to global change. *Ecology Letters*, 19(2), 111–120.
- Hammerstedt, R. H., and Hay, S. R. (1980). Effect of incubation temperature on motility and cAMP content of bovine sperm. *Archives of Biochemistry and Biophysics*, 199(2), 427–437.

- Helfenstein, F., Losdat, S., Møller, A. P., Blount, J. D., and Richner, H. (2010). Sperm of colourful males are better protected against oxidative stress. *Ecology Letters*, *13*(2), 213–222.
- Ho, M., Price, C., King, C., Virtue, P., and Byrne, M. (2013). Effects of ocean warming and acidification on fertilization in the Antarctic echinoid *Sterechinus neumayeri* across a range of sperm concentrations. *Marine Environmental Research*, *90*, 136–141.
- Huey, R. B., Deutsch, C. A., Tewksbury, J. J., Vitt, L. J., Hertz, P. E., Pérez, H. J. Á., and Garland, T. (2009). Why tropical forest lizards are vulnerable to climate warming. *Proceedings of the Royal Society of London B: Biological Sciences*, *276*(1664), 1939–1948.
<https://doi.org/10.1098/rspb.2008.1957>
- Hurley, L. L., McDiarmid, C. S., Friesen, C. R., Griffith, S. C., and Rowe, M. (2018). Experimental heatwaves negatively impact sperm quality in the zebra finch. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1871), 20172547.
- Immler, S., Calhim, S., and Birkhead, T. R. (2008). Increased postcopulatory sexual selection reduces the intramale variation in sperm design. *Evolution: International Journal of Organic Evolution*, *62*(6), 1538–1543.
- Isaac, J. L. (2009). Effects of climate change on life history: Implications for extinction risk in mammals. *Endangered Species Research*, *7*(2), 115–123.
- Kingsolver, J. G., Diamond, S. E., and Buckley, L. B. (2013). Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology*, *27*(6), 1415–1423. Scopus. <https://doi.org/10.1111/1365-2435.12145>
- Kobayashi, M., Wada, M., Hori, T., and Kawakami, E. (2014). Superoxide dismutase activity in the oviductal and uterine fluid of the bitch and the effects of the enzyme on viability, motility and hyperactivation of canine sperm in Vitro. *Journal of Veterinary Medical Science*, *76*(5), 741–743. Scopus. <https://doi.org/10.1292/jvms.13-0545>
- López Juri, G. (2019). *Evolución fenotípica en relación a las estrategias reproductivas en lagartos del Dominio Chaqueño* [PhD Thesis]. Doctorado en Ciencias Biológicas, Universidad Nacional de Córdoba.

- López Juri, G., Chiaraviglio, M., and Cardozo, G. (2018a). Do female reproductive stage and phenotype influence thermal requirements in an oviparous lizard? *Journal of Thermal Biology*, *71*, 202–208.
- López Juri, G., Chiaraviglio, M., and Cardozo, G. (2018b). Electrostimulation is an effective and safe method for semen collection in medium-sized lizards. *Theriogenology*.
- López Juri, G., Rossi, N., Chiaraviglio, M., and Cardozo, G. (2020). Phenotypic diversity and its relationship to reproductive potential in changing social contexts in a lizard model. *Biological Journal of the Linnean Society*.
- Magris, M., Cardozo, G., Santi, F., Devigili, A., and Pilastro, A. (2017). Artificial insemination unveils a first-male fertilization advantage in the guppy. *Animal Behaviour*, *131*, 45–55.
- Mansour, N., Lahnsteiner, F., and Patzner, R. (2002). The spermatozoon of the African catfish: Fine structure, motility, viability and its behaviour in seminal vesicle secretion. *Journal of Fish Biology*, *60*(3), 545–560.
- Marshall, D. J. (2015). Environmentally induced (co) variance in sperm and offspring phenotypes as a source of epigenetic effects. *Journal of Experimental Biology*, *218*(1), 107–113.
- Mehlis, M., and Bakker, T. C. (2014). The influence of ambient water temperature on sperm performance and fertilization success in three-spined sticklebacks (*Gasterosteus aculeatus*). *Evolutionary Ecology*, *28*(4), 655–667.
- Meijering, E., Dzyubachyk, O., and Smal, I. (2012). Methods for cell and particle tracking. In *Methods in enzymology* (Vol. 504, pp. 183–200). Elsevier.
- Mengden, A., Platz, C. G., Hubbard, R., and Quinn, H. (1980). Semen collection, freezing and artificial insemination in snakes. *Contributions to Herpetology Reproductive Biology and Diseases of Captive Reptiles*. St. Louis: St. Louis University. p, 71–78.
- Møller, A., Mousseau, T., Rudolfson, G., Balbontin, J., Marzal, A., Hermosell, I., and De Lope, F. (2009). Senescent sperm performance in old male birds. *Journal of Evolutionary Biology*, *22*(2), 334–344.
- Nixon, B., Anderson, A. L., Smith, N. D., McLeod, R., and Johnston, S. D. (2016). The Australian saltwater crocodile (*Crocodylus porosus*) provides evidence that the capacitation of spermatozoa may extend beyond the mammalian lineage. *Proceedings of the Royal Society B*:

Biological Sciences, 283(1830), 20160495.

- Oliveira, R., Tomasi, L., Rovasio, R., and Giojalas, L. (1999). Increased velocity and induction of chemotactic response in mouse spermatozoa by follicular and oviductal fluids. *Reproduction*, 115(1), 23–27.
- Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock, C. C., and Thomas, M. B. (2013). Temperature variation makes ectotherms more sensitive to climate change. *Global Change Biology*, 19(8), 2373–2380.
- Pizzari, T., and Wedell, N. (2013). *The polyandry revolution*. The Royal Society.
- R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <https://www.R-project.org>
- Ramírez-Pinilla, M. P., Calderón-Espinosa, M. L., Flores-Villela, O., Muñoz-Alonso, A., and de la Cruz, F. R. M. (2009). Reproductive activity of three sympatric viviparous lizards at Omiltemi, Guerrero, Sierra Madre del Sur, Mexico. *Journal of Herpetology*, 409–420.
- Rosengrave, P., Taylor, H., Montgomerie, R., Metcalf, V., McBride, K., and Gemmell, N. J. (2009). Chemical composition of seminal and ovarian fluids of chinook salmon (*Oncorhynchus tshawytscha*) and their effects on sperm motility traits. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 152(1), 123–129. Scopus. <https://doi.org/10.1016/j.cbpa.2008.09.009>
- Rossi, N., Benitez-Vieyra, S., Cocucci, A., Chiaraviglio, M., and Cardozo, G. (2019). Sexual dichromatism and color diversity in the spiny lava lizard *Tropidurus spinulosus* using lizard visual modelling. *Scientific Reports*, 9(1), 1–10.
- Sales, K., Vasudeva, R., Dickinson, M. E., Godwin, J. L., Lumley, A. J., Michalczyk, \Lukasz, Hebberecht, L., Thomas, P., Franco, A., and Gage, M. J. (2018). Experimental heatwaves compromise sperm function and cause transgenerational damage in a model insect. *Nature Communications*, 9(1), 4771.
- Sasanami, T., Matsuzaki, M., Mizushima, S., and Hiyama, G. (2013). Sperm storage in the female reproductive tract in birds. *Journal of Reproduction and Development*, 59(4), 334–338.
- Siegel, D. S., Miralles, A., Chabarria, R. E., and Aldridge, R. D. (2011). Female reproductive anatomy: Cloaca, oviduct, and sperm storage. *Reproductive Biology and Phylogeny of*

Snakes. Science Publishers, Inc., Enfield.

- Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., Dong, Y., Harley, C. D., Marshall, D. J., Helmuth, B. S., and others. (2016). Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecology Letters*, *19*(11), 1372–1385.
- Soren, S., Singh, S. V., Borah, S., Kumar, A., and Dwivedi, D. (2018). Global Warming: Impact, Adaptation and Ameliorative Measures of Semen Quality under Tropical Climatic Conditions in Crossbred Bulls. *Journal of Animal Research*, *8*(5), 731–739.
- Stocker, T., Qin, D., Plattner, G., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, B., and Midgley, B. (2013). *IPCC, 2013: Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change.*
- Sunday, J. M., Bates, A. E., Kearney, M. R., Colwell, R. K., Dulvy, N. K., Longino, J. T., and Huey, R. B. (2014). Thermal-safety margins and the necessity of thermoregulatory behavior across latitude and elevation. *Proceedings of the National Academy of Sciences*, *111*(15), 5610–5615.
- Tang, X.-L., Yue, F., Yan, X.-F., Zhang, D.-J., Xin, Y., Wang, C., and Chen, Q. (2012). Effects of gestation temperature on offspring sex and maternal reproduction in a viviparous lizard (*Eremias multiocellata*) living at high altitude. *Journal of Thermal Biology*, *37*(6), 438–444.
- Tomášek, O., Albrechtová, J., Němcová, M., Opatová, P., and Albrecht, T. (2017). Trade-off between carotenoid-based sexual ornamentation and sperm resistance to oxidative challenge. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1847), 20162444.
- Tourmente, M., Giojalas, L. C., and Chiaraviglio, M. (2011). Sperm parameters associated with reproductive ecology in two snake species. *Herpetologica*, *67*(1), 58–70.
- Tourmente, M., Gomendio, M., and Roldan, E. R. (2011). Sperm competition and the evolution of sperm design in mammals. *BMC Evolutionary Biology*, *11*(1), 12.
- Vicenzi, N., Laspiur, A., Sassi, P. L., Massarelli, R., Krenz, J., and Ibarguengoytia, N. R. (2020). Impact of temperature on bite force and bite endurance in the Leopard Iguana (*Diplolaemus leopardinus*) in the Andes Mountains. *Journal of Experimental Biology*.

- Warwick, C., Arena, P., Lindley, S., Jessop, M., and Steedman, C. (2013). Assessing reptile welfare using behavioural criteria. *In Practice*, 35(3), 123–131.
- Williams, A. C., and Ford, W. C. L. (2005). Relationship between reactive oxygen species production and lipid peroxidation in human sperm suspensions and their association with sperm function. *Fertility and Sterility*, 83(4), 929–936.
- Zeh, J. A., Bonilla, M. M., Su, E. J., Padua, M. V., Anderson, R. V., Kaur, D., Yang, D.-S., and Zeh, D. W. (2012). Degrees of disruption: Projected temperature increase has catastrophic consequences for reproduction in a tropical ectotherm. *Global Change Biology*, 18(6), 1833–1842. <https://doi.org/10.1111/j.1365-2486.2012.02640.x>

Figure




Figure 1 Effects of temperature increment and oviductal fluid (OF) on sperm performance. VSL: straight line velocity, VCL: curvilinear velocity, LIN: linearity ($LIN = VSL/VCL$), Motility: percentage of motile spermatozoa (%). CV: Coefficients of Variation. Lines within boxes correspond to the median value of the sample, the lower and upper edge of the box correspond to quartiles 1 and 3 respectively, whiskers denote the minimum and maximum values and outliers are highlighted with points. Different letters indicate significant differences according to Tukey's test ($p < 0.05$). The total sample size was 27 independent replicates.

Table 1. Statistic parameters for dynamic spermatic variables in *Tropidurus spinulosus*

Sperm parameter	OF factor	Temperature factor	OF*Temperature
VSL	F=.22.01, p<0.0001	F =22, p<0.0001	F=1.83, p=0.17
VCL	F=105.409, p<0.0001	F =152.04, p<0.0001	F=29.83, p=<0.0001
LIN	F=1.32, p=0.25	F =8.66, p=0.0058	F=3.73, p=0.056
% Motile sperm	X ² =16.25, p<0.0001	X ² =16.25, p<0.0001	X ² =0.02, p=0.88
VSL_CV	F=1.04, p=0.31	F=0.42, p=0.54	F=1.61, p=0.20
VCL_CV	F=6.83, p=0.01	F=5.54, p=0.02	F=0.07, p=0.78
LIN_CV	F=0.01, p=0.91	F=1.27, p=0.26	F=5.09, p=0.02




Fig. S1. Schematic representation of the experimental protocol to assess the influence of oviductal fluid (OF) and temperature on sperm dynamics and viability in *Tropicurus spinulosus*