



XI CONGRESO ARGENTINO DE MICROBIOLOGÍA GENERAL

5 al 7 de Agosto de 2015
Córdoba, Argentina

SAMIGE

Asociación Civil de Microbiología General

ANALYSING BACTERIAL COMMUNITIES FROM MICROBIAL MATS AND SEDIMENTS LOCATED IN THE ATACAMA DESERT

A.B. Fernández¹, M.C. Rasuk¹, D. Kurth², M. Contreras², F. Novoa³, D. Poire¹, M.E. Farías¹.

¹Laboratorio de Investigaciones Microbiológicas de Lagunas Andinas, PROIMI-CONICET, Argentina. ²Centro de Ecología Aplicada, Santiago, Chile. ³Universidad Nacional de la Plata. ⁴Centro de Investigaciones Geológicas, UNLP-CONICET, Argentina.

anabfg82@gmail.com

The Atacama Desert has more than 100 basins with interior drainage and most of them contain salt flats. These ecosystems have extreme environmental conditions that allow the development of unique microbial communities. The objective of this was to study the bacterial diversity using independent culture tools of microbial mats and sediments from salt flats in the Atacama Desert. Some physicochemical conditions of the water surrounding these samples were analysed to discover if any physicochemical characteristic could be influencing in its taxonomic composition. Five samples were collected, three of them were microbial mats and two were sediments. The mat samples were taken from Laguna Llamara (samples named LL1 and LL2) and Laguna Cejar (Cej). Sediments were taken from Laguna Jachucoposa (Cop) and Laguna Pujsa (Puj) where microbial mats are not present. Total metagenomic DNA extraction was performed on each sample and the V4 hypervariable region of the bacterial 16S rRNA gene was amplified by pyrosequencing using the Ribosomal Database Project (RDP)-suggested universal primers. Diversity of the microbial community was assessed using the QIIME software package. Lakes that harbor microbial mats have a higher salinity and a lower dissolved oxygen concentration and proportion of organic matter and total phosphorous than lakes where mats are absent. All the samples have important concentrations of arsenic, with an extremely high amount in Puj. *Proteobacteria* and/or *Bacteroidetes* are the major phyla represented in all samples. Also, other phyla as *Spirochaetes*, *Chloroflexi* or *Verrucomicrobia* are found. However, cyanobacterial sequences are only observed in LL2 and Puj. On the other hand, we have found a higher diversity in sediment than in mat samples. The sediments samples contain phyla not observed in mat samples. 16S rRNA gene sequences classified within *Actinobacteria* and *Gracilibacteria* are only found in Puj and related to *Tenericutes*, *Gemmatimonadetes* and *Acidobacteria* are only observed in Cop. Finally, an important fraction of the sequences could not be classified at phylum level. The high diversity found in sediment samples may be explained by the physicochemical conditions in the environment. For example, they have a lower conductivity than mat samples. It is known hypersaline environments have a low diversity, where halophilic microorganisms are able to survive to these extreme conditions because they have specific strategies to balance the osmotic pressure. Besides, we found a low proportion or absence of *Cyanobacteria* in the ecosystems studied, suggesting the possibility that other groups may be playing an essential role as primary producers in these extreme environments. Additionally, the large proportion of 16S rRNA gene sequences that could not be affiliated to any known bacterial phyla suggesting that in these ecosystems there are potential novel representatives of bacterial phyla not yet described.

EXTREME-HALOPHILES: THEIR ROLE IN THE ARSENIC BIOGEOCHEMICAL CYCLE

M.C. Rasuk¹, O.F. Ordoñez¹, M. Soria¹, M.E. Farías¹.

¹Laboratorio de Investigaciones Microbiológicas de Lagunas Andinas, PROIMI-CONICET, Argentina.

cecirasuk@gmail.com

Biofilms, mats and microbialites dwell under extreme environmental conditions (high salinity, extreme aridity, pH and arsenic concentration) in the Argentinean Puna and the Atacama Desert. Microbial communities inhabiting those ecosystems are poorly known. Arsenic metabolism is proposed to be an ancient mechanism in microbial life. Besides, some bacteria and archaea are not only able to use detoxification processes to grow under high arsenic concentration, but also, some of them are able to exploit arsenic as a bioenergetic substrate in either anaerobic arsenate respiration or chemolithotrophic growth on arsenite. Only four aioAB coding for arsenite oxidase and two *arrA* coding for arsenate reductase sequences from haloarchaea were previously deposited in the NCBI Database, but have not been reported in the literature. The *arrA* arsenate reductases are reliable indicators of anaerobic As (V) respiration and catalyze the electron transfer to the As (V) terminal acceptor in dissimilatory arsenatereducing prokaryotes (DARPs). In this work, we are presenting our first steps in the study of the arsenic biogeochemical cycle in these ecosystems. Thus, the aim of this study was to isolate and to study the arsenic metabolism genes of the isolated extreme halophile microorganisms as well as to test the growth in minimal medium using different carbon sources. Mats and microbialites samples were taken from the water's edge of Laguna Tebenquiche, Laguna Brava (Salar de Atacama, Chile)