

Ecological characterization of wild *Helianthus annuus* and *Helianthus petiolaris* germplasm in Argentina

Monica Poverene^{1,2*}, Miguel Cantamutto^{1,3} and Gerald J. Seiler⁴

¹Department of Agronomy, Universidad Nacional del Sur, 8000 Bahía Blanca, Argentina,

²Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS-CONICET),

Bahía Blanca, Argentina, ³Centro Udl-IRTA, Lleida, Cataluña, Spain and ⁴USDA-ARS

Northern Crop Science Laboratory, PO Box 5677, Fargo, ND 58105, USA

Received 8 January 2008; Accepted 30 April 2008 – First published online 14 July 2008

Abstract

Helianthus annuus and *H. petiolaris* (Asteraceae) are wild sunflowers native to North America but have become naturalized in central Argentina covering an area of about 5 million hectares. Wild *H. annuus* has been recognized as invader species in several countries, but no research has been done to study the ecological determinants of their distribution. In a survey covering seven provinces between 31°58'–38°S and 60°33'–69°W, we described the ecology of the main wild populations. Wild *Helianthus* populations were located in three of the 18 ecological regions of Argentina, on five Mollisol and seven Entisol soil groups. The associated plant communities were comprised 60 species belonging to 16 families, all being frequent components of the native flora. Disease symptoms were seldom observed in wild populations, with *Alternaria helianthi* being the most commonly observed pathogen. Population size varied from less than 100 to more than 100,000 plants, covering from 100 to more than 60,000 m² with densities most frequently up to 3 plants/m², but reaching 80 plants/m² at certain sites. Intermediate plant phenotypes between wild species and cultivated sunflower were found in one-third of the populations providing evidence of intense gene flow. Hybrid swarms were found at three localities with population sizes between 100 and 10,000 individuals.

Keywords: community; density; diseases; habitat; populations; sunflower

Introduction

Helianthus annuus L. and *H. petiolaris* Nutt. (Asteraceae) are annual diploid species native to North America where the former has a wide distribution and the latter is restricted to the central region (Heiser *et al.*, 1969; Rogers *et al.*, 1982). Wild or common *H. annuus* tends to be weedy, always located in habitats that have been disturbed. The prairie sunflower, *H. petiolaris*, usually grows in sandy soils, but it is also found as an adventive weed elsewhere (Seiler and Rieseberg, 1997). Both species have several botanical forms and are systematically complex (Heiser,

1954, 1961; Seiler and Rieseberg, 1997; Jan and Seiler, 2007). *H. annuus* is the ancestral species of cultivated sunflower (Heiser, 1978; Burke *et al.*, 2002).

Both species are valuable germplasm resources with traits that have been transferred into cultivated sunflower, i.e. cytoplasmic male sterility (CMS) from *H. petiolaris* (Leclercq, 1969), but Rieseberg and Seiler (1990) provided evidence that CMS may have been derived from *H. annuus*, disease and pest resistance, oil quality and other traits for crop breeding (Seiler, 1992). These species are also crop weeds in North America (Geier *et al.*, 1996; Rosales Robles *et al.*, 2002; Deines *et al.*, 2004) and are beginning to invade summer crops in Argentina.

Sixty years after the first introduction of *H. annuus* and *H. petiolaris*, they have become naturalized in the central

* Corresponding author. E-mail: poverene@criba.edu.ar

area of Argentina (Covas, 1966; Bauer, 1991; Poverene *et al.*, 2002). At present, their distribution significantly overlaps that of the sunflower crop. As in the Northern Hemisphere, flowering time of both wild species and the cultivated sunflower coincide and they share pollinators, mainly honeybees, bumblebees and other wild bees, favouring gene flow (Burke *et al.*, 2002; Poverene *et al.*, 2004) and natural hybridization processes (Rieseberg *et al.*, 1998, 1999b).

Wild *H. annuus* has been recognized as an invader species in several countries (Berville *et al.*, 2005), but at present, *H. petiolaris* has been naturalized only in Argentina. The study of the invasive process of these annual species could help to understand and prevent analogous processes in other regions of the world. Cantamutto *et al.* (2008) studied the environmental conditions of these invader species distributions, but there is no available information about the ecology of these wild sunflower species. The objective of this research was to

describe the eco-geographic distribution of *H. annuus* and *H. petiolaris* in Argentina and to characterize the populations in their natural habitats.

Materials and methods

Agro-ecological descriptions of populations were made during a collection trip across seven central provinces of Argentina, during February 2007 (Fig. 1). Thirty previous explorations carried out between 2000 and 2006 provided data on population locations, habitat and soil type. Collected information included botanical name, collection site (province, district, location, latitude, longitude and altitude), environmental conditions (habitat) and community (dominance of co-occurring plant species), estimated population size, plant density, plant size and morphological variation. Also, the occurrence of prevalent sunflower diseases was recorded: downy




Fig. 1. Wild *Helianthus* populations (white diamonds) sampled in three ecological regions of central Argentina (black numbers): of central Argentina: 11, Espinal; 12, Pampa; 13, Shrubs of Plateau and Plains. Soils in the Pampa region are mainly Mollisols, whereas Entisols predominate in the other two regions. Provinces are Buenos Aires (BA), Cordoba (CO), Entre Rios (ER), La Pampa (LP), Mendoza (ME), San Juan (SJ) and San Luis (SL) (map from Burkart *et al.*, 1999; scale 1:15,000,000).

mildew (*Plasmopara halstedii*), rust (*Puccinia helianthi*), white rust (*Albugo tragopogonis*), *Alternaria helianthi*, *Verticillium dahliae*, *Phoma macdonaldii* and *Sclerotinia sclerotiorum* wilt.

The geographic coordinates were used to determine the agro-ecological regions (Burkart *et al.*, 1999) and soil taxa (INTA, 1990) corresponding to each population. The order, great group, suborder, area and predominant texture of each soil type were used to describe the habitat environment of both species (Bouma, 2003).

Data recorded for each population included occupied area, density and total number of individuals. The total area was the sum of one to five quadrants measured at each site that contained all the individuals. The density was estimated by ten samples within the quadrants, taken at regular intervals along the main transect across each population, with a 0.25 m² circle. The total number of plants was then calculated as a product of area × mean density. In the case of populations growing in continuous patches, the limit was established as the point where the distance between two patches was greater than the longest side of the quadrant.

Plant community density was recorded for each quadrant following a semi-quantitative method (Clay and Johnson, 2002). At each collection site, data were collected from ten points on a uniformly spaced grid coordinate system. At each grid point (a 2 m² circle), abundance was qualified as following: 0, absent; 1, less than 5 plants/m²; 2, 6–10 plants/m² and 3, more than 10 plants/m². The 20 most frequent species were characterized by life cycle, origin and status.

To estimate the number of plants potentially exposed to gene flow, a mean was obtained for population size based on the ranges observed in more than 50% of the populations. Cultivated plant number was computed taking into account the minimum acreage per cultivated field (30–60 ha) and the minimum number of plants per hectare (40,000) usually sown. Frequencies of gene flow between the three taxa were obtained from our previous research (Poverene *et al.*, 2004; Cantamutto *et al.*, 2007; Ureta *et al.*, 2008).

Results

H. annuus populations were widespread ranging from 31°58' to 37°31'S, and 60°33' and 68°14'W, at an altitude of 128–600 m.a.s.l. (Table 1). Plants grew in patches in disturbed habitats such as roadsides, ditches, fence rows and field margins in the sunflower production areas (Buenos Aires, La Pampa and Cordoba provinces). They were also patchily distributed along irrigation channels in the western provinces (Mendoza and San Juan) and growing along crags for several kilometres of the

Table 1. Frequency of selected populations and habitat characteristics of wild *Helianthus annuus* and *Helianthus petiolaris* collected in Argentina

Ecogeographic data	<i>H. annuus</i> (%)	<i>H. petiolaris</i> (%)	Both (mixed stands; %)
Altitude (m.a.s.l.)			
< 300	55	60	25
> 300	18	40	75
Not recorded	27	0	0
Population size (no. of plants)			
< 100	9	0	0
101–1000	0	30	50
1001–5000	55	30	25
5001–10,000	18	20	25
10,001–50,000	9	20	0
> 50,000	9	0	0
Surface area (m ²)			
100–1000	27	20	25
1001–10,000	27	70	75
10,001–50,000	18	0	0
> 50,000	27	10	0
Mean plant density (plant/m ²)			
< 1	27	40	50
1–3	64	50	50
> 3	9	10	0
Maximum plant density (plant/m ²)			
16–25	22	40	0
11–15	23	10	50
5–10	33	20	0
< 5	22	30	50
Plant height (cm)			
> 280	22	0	0
200–280	45	10	50
< 200	33	90	50
Soil texture			
Loam	9	0	0
Loamy sand	18	40	25
Sand	0	60	0
Sandy loam	64	0	75
Silt loam	9	0	0
Habitat			
Roadside, intersection	55	80	75
Riverside	9	0	0
Field margin	18	10	25
Within crop	0	10	0
Ditch	18	0	0
Volunteers			
Present	46	10	25
Absent	55	90	75
Intermediate plants			
Present	36	30	100
Absent	64	70	0

coastal rivers in eastern Entre Ríos province. Population size varied from a few dozen individuals to more than 100,000 plants, with mean densities varying between 0.25 and 6 plants/m². However, some populations

reached 72 and 80 plants/m² in Mendoza and Córdoba provinces, respectively. Most plants were very robust with heights over 2.80 m.

H. petiolaris ranged from 35°08' to 38°08'S, and 62°16' and 65°56'W, and up to 455 m in altitude (Table 1). Populations were very numerous in the eastern La Pampa and western Buenos Aires provinces, reaching 18,000 plants with a mean density of 0.25–6 plants/m², and up to 40 plants/m² in the former. Sparser populations were found in southern San Luis. In Córdoba, this species seemed to be confined to the southern extreme of the province.

Two perennial *Helianthus* populations were found in Mendoza, probably *Helianthus tuberosus* or *Helianthus × laetiflorus*, but were difficult to identify because they were just beginning to flower.

Volunteer plants from the cultivated sunflower crop were found among wild ones and many plants showed intermediate morphological traits. Variation was observed for leaf size, the presence of anthocyanin in stems and petioles, ray colour and white pubescent disc flowers in the centre of the head.

Three hybrid swarms were found, one in Buenos Aires and two in La Pampa. These swarms included wild-type plants of both species, intermediate plants and a number of volunteers from crop plants. In the largest swarm from La Pampa, the northern half comprised 2600 plants with about 15% being annuus-like and a plant density of 1.34 plants/m². The southern part comprised about 8000 plants, with 50% being annuus-like and the rest petiolaris-like. Many intermediate plants were observed in the central zone.

Wild species populations were found in three ecological regions: Pampa, Espinal and Shrubs of Plateau and Plains (Fig. 1). *H. petiolaris* and *H. annuus* populations were found on five Mollisol and seven Entisol groups (Table 2). Species distributions were significantly associated with soil subgroups according to Pearson's chi-squared test (Pearson's $\chi^2 < 0.001$, highly significant).

The plant communities associated with the wild *Helianthus* species comprised 60 species belonging to 16 families. Of these, 32 were found associated with both wild species. Most frequent species were *Sorghum halepensis*, *Cynodon dactylon*, *Eragrostis curvula* (Poaceae), *Chenopodium album*, *Salsola kali* (Chenopodiaceae) and *Centaurea solstitialis* (Asteraceae). Nineteen other species were found only in *H. annuus* communities, with *Melilotus albus* (Fabaceae) being the most common. Nine other species were only found in *H. petiolaris* communities, where *Cenchrus pauciflorus* (Poaceae) was the most frequent. Differences for the latter two species' association with wild sunflowers were significant ($P < 0.05$). Table 3 presents life cycle, origin and status of the 20 species most frequently found associated with wild sunflowers in the explored provinces, considered as the dominant community species.

Disease symptoms were observed only in 25% of the wild populations. *Alternaria* lesions on leaves were the most frequent, with *A. helianthi* being the most likely pathogen. *Puccinia helianthi* was often found on volunteer plants, but never on wild plants. Table 4 presents the observed diseases and the frequency of affected plants.

Table 2. Frequency of stable wild *Helianthus petiolaris* (PET), *Helianthus annuus* (ANN) populations and mixed stands (MIX) associated with 16 of the 65 soil taxa defined by INTA 1990 for the colonized provincial counties

Order	Group	Subgroup	Population frequency (%)		
			ANN	MIX	PET
Mollisols	Argiaquolls	Typic	4.65		
Mollisols	Argiudolls	Aquic	6.98		
Mollisols	Argiudolls	Typic	4.65		
Mollisols	Argiustolls	Typic	6.98	25.0	2.70
Mollisols	Hapludolls	Various	4.66		
Mollisols	Hapludolls	Entic	18.60		12.16
Mollisols	Hapludolls	Thapto-argidic			4.06
Mollisols	Hapludolls	Typic	6.98		4.06
Mollisols	Haplustolls	Entic	9.30	50.00	31.08
Mollisols	Haplustolls	Litic	4.65	25.00	2.70
Mollisols	Haplustolls	Various			12.16
Mollisols	Haplustolls	Udortentic	6.98		
Entisols	Torrifluvents	Typic	13.95		
Entisols	Torripsammets	Various	6.98		1.35
Entisols	Udipsammets	Typic	2.32		1.35
Entisols	Ustisfluvents	Typic	2.32		
Entisols	Ustipsammets	Typic			20.27
Entisols	Ustorthentst	Typic			8.11

Table 3. The 20 dominant community species most frequently associated with wild sunflower populations in central Argentina

Species	Family	Cycle	Origin	Status	Provinces
<i>Chenopodium album</i>	Chenopodiaceae	A	E	I, MS	BA, CO, LP, SL, ME, SJ
<i>Sorghum halepense</i>	Poaceae	P	E, A	W ^a	All
<i>Cynodon dactylon</i>	Poaceae	P	E, A	W ^a	All
<i>Centaurea solstitialis</i>	Asteraceae	A	E, A	W, SS	BA, CO, ER, LP, SL
<i>Salsola kali</i>	Chenopodiaceae	A	E, A	W ^a	BA, LP, ME, SL
<i>Eragrostis curvula</i>	Poaceae	P	E, N	SS	BA, CO, LP, SL, ME
<i>Melilotus albus</i>	Fabaceae	A	E, A	SS	All
<i>Portulaca oleracea</i>	Portulacaceae	A	E, N	MS	All
<i>Tagetes minuta</i>	Asteraceae	A	Na	SS	All
<i>Setaria verticillata</i>	Poaceae	A	E, A	MS	All
<i>Eleusine indica</i>	Poaceae	P	Na	SS	BA, CO, ER, LP
<i>Diplotaxis tenuifolia</i>	Cruciferae	P	E, A	W ^a	BA, CO, LP, SL, ME, SJ
<i>Amaranthus quitenensis</i>	Amarantaceae	A	Na	W ^a	BA, ER, ME, SJ
<i>Chenopodium multifidum</i>	Chenopodiaceae	P	Na	MS	All
<i>Medicago sativa</i>	Fabaceae	P	E, A	MS	BA, CO, ER, LP, ME, SJ
<i>Cenchrus pauciflorus</i>	Poaceae	A	Na	SS	BA, CO, ER, LP, SL, ME
<i>Heterotheca latifolia</i>	Asteraceae	A	E, A	W, SS	CO, LP, SL
<i>Onopordon acanthium</i>	Asteraceae	B	E, N	W ^a	BA, LP
<i>Polygonum aviculare</i>	Polygonaceae	A	E, A	MS	All
<i>Solanum elaeagnifolium</i>	Solanaceae	P	Na	MS	All

Life cycle: A, annual; P, perennial; B, biannual. Origin: Na, native; E, exotic; A, adventive; N, naturalized. Status: W, weed; I, invasive; MS, modified soils; SS, sandy soils. Provinces: BA, Buenos Aires; LP, La Pampa; SL, San Luis; CO, Cordoba; ME, Mendoza; SJ, San Juan; ER, Entre Rios; all, all the seven explored provinces.

^aAgricultural epidemic.

Discussion

Collection site data characterization

Regarding the number of plants, wild *H. annuus* populations in the centre of the country were only one magnitude of order lower than crop populations that are usually between 1 and 2.5 million plants (Ureta *et al.*, 2008). *H. petiolaris* was the most frequent species, but was geographically more restricted. This species often grows in field margins and seldom invades sunflower, corn or pasture crops. Most populations were found in roadsides and road intersections on disturbed sandy soils. Compared with previous collection trips, both species seemed to be more widespread, although population size and density are strongly dependent on climatic conditions, particularly moisture. Wild *Helianthus* populations were found in the agricultural regions where soybean, maize, sunflower and wheat are the predominate crops. Perennial *Helianthus* were only found in Mendoza; this kind of feral populations also occur in the Buenos Aires province, where they are usually established by rhizomes discarded from gardens (Sala *et al.*, 1990).

Gene flow

Morphologically intermediate plants found among volunteers indicate a frequent crop–wild gene exchange in

Buenos Aires and La Pampa provinces, where there is a large sunflower crop acreage, and Mendoza where there are areas devoted to sunflower seed production. Volunteers can considerably enhance sunflower crop–wild hybridization acting as a bridge for genetic transfer of crop traits into wild populations (Reagon and Snow, 2006). Persistent cultivar gene flow determines high levels of introgression and the replacement of wild populations by advanced generation hybrids (Linder *et al.*, 1998). Crop–*H. petiolaris* hybridization also occurs when they come into contact.

The magnitude of crop–wild gene flow in Argentina has been estimated through field observations and previous experiments (Poverene *et al.*, 2004; Ureta *et al.*, 2008 and unpublished data). Although the hybridization frequency was similar to that observed in North America (Arias and Rieseberg, 1994; Rieseberg *et al.*, 1999a), plants are so numerous that even at a low frequency of interspecific crosses, thousands of F1 hybrids are likely produced every year (Fig. 2).

Sunflower crop genes persist for several generations in wild populations (Whitton *et al.*, 1997; Linder *et al.*, 1998) and can modify wild populations depending on their fitness (Alexander *et al.*, 2001; Cummings *et al.*, 2002) and on the environments where they grow (Mercer *et al.*, 2007). The high number of morphologically intermediate crop–wild plants observed in two-thirds of the populations in central Argentina could be assessed to

Table 4. Observed diseases on wild *Helianthus* populations from central Argentina

Population	Disease	Frequency of infected plants (%)
1007 <i>Helianthus annuus</i>	Alternaria	70
1107 <i>H. annuus</i>	Alternaria	90
	Phoma black stem	90
1207 <i>H. annuus</i>	Alternaria	10
1307 <i>Helianthus petiolaris</i>	Alternaria	10
1407 <i>H. annuus</i>	Virus (SuCMoV) ^a	40
	Powdery mildew	10
1607 <i>H. annuus</i>	Alternaria	20
2107 <i>H. petiolaris</i>	Alternaria	40
2507 <i>H. annuus</i>	Alternaria	30
2607 <i>H. petiolaris</i> crop–wild hybrids	Alternaria	20
	Alternaria	50
	Phoma black stem	20
3507 <i>H. petiolaris</i> , crop–wild hybrid	Phoma black stem	10

^aField identification by leaf lesions.

gene flow. The consequences of the frequent hybridization process have yet to be fully evaluated.

When both wild species come into contact in North America, they can often form hybrid zones that have given rise to three other species via homoploid speciation (Rieseberg *et al.*, 1990, 1991; Rieseberg, 1991). Although both species have become established in Argentina rather recently, hybridization and introgression processes are taking place in this new environment.

Ecology

Wild species populations were found in three of the 18 ecological regions described in Argentina by Burkart *et al.* (1999). From the east, Pampa is a grass steppe without woody species, followed by Espinal, an intermediate savannah, with grasses and scarce xeric trees. The western Shrubs of Plateau and Plains is an arid steppe with

the predominance of shrubs and tough grasses. Both wild sunflower species extend along a SE–NW boundary that coincides with the limit between Pampa and Espinal regions. The subhumid region called Pampa is a cultivated area that corresponds to grasslands ploughed within last 140 years, while in the semi-arid region called Espinal agriculture is much more recent.

Soil taxonomy as an indicator of ecosystem processes can predict potential plant species suitable habitats (Mann *et al.*, 1999). Mollisols, Alfisols and Entisols orders cover only 18% of the world's temperate areas, but predominate in the centre of origin of the genus *Helianthus*, where together they cover 48% of the US surface. The soils of the Central Great Plains of North America, the common distribution area for the two annual species *H. annuus* and *H. petiolaris* (Rogers *et al.*, 1982) belong to these orders (USDA, 1999). In Argentina, both species are also associated with Mollisols and Entisols. The 14 soil subgroups where the *H. annuus*




Fig. 2. Gene flow frequencies among cultivated and wild sunflowers in Argentina and number (*n*) of plants estimated as a range from data in Table 1.a: from Ureta *et al.* (2008); b: from Poverene *et al.* (2004); c: from Cantamutto *et al.* (2007). Gene flow values were estimated in natural conditions, except one which came from a planned field experiment (dotted arrow).

populations were found cover 9.9 million hectares, while the 11 subgroups associated with *H. petiolaris* cover 13.1 million hectares (INTA, 1990). In the central area, where the sunflower crop has moved to in the last 10 years, there is a high probability of observing new wild sunflower populations because of the favourable macro-habitat conditions.

Except for *Eragrostis curvula*, the most frequent plant community species associated with wild *Helianthus* are weeds and related to disturbed soils (Marzocca, 1994). *E. curvula*, 'weeping lovegrass', has become established in sandy soils subjected to wind erosion. *Melilotus albus* associated with wild *H. annuus* is a salt tolerant forage species and has become established in humid soils with medium to high salt levels; *Cenchrus pauciflorus* associated with *H. petiolaris* is a noxious weed, very common in sandy soils of the western central region (Marzocca, 1994). Among species in Table 3, six are considered noxious weeds of agriculture in Argentina. All the cited species are frequent components of the flora in central Argentina and none of them was indicative of a specific ecosystem. This reinforces the hypothesis that abiotic factors, particularly disturbance determine the wild *Helianthus* colonization (Cantamutto et al., 2007) and that the two wild sunflowers will expand their distribution when the habitat and opportunity arise. Most populations were free from diseases and confirmed that wild sunflower species are potential gene reservoirs for fungus and virus resistance.

Wild *H. annuus* and *H. petiolaris* form large populations distributed over an area of about 5 million hectares in central Argentina. Since their establishment 60 years ago they have continuously increased their area, behaving as an invasive species providing evidence that they will continue spreading. These species offer opportunities for research covering various scopes. First, both wild *Helianthus* constitute germplasm reservoirs of biotic and abiotic gene resistance for crop improvement. Second, wild populations subjected to gene flow may acquire crop traits (i.e. herbicide tolerance) that modify their fitness enhancing invasiveness or weediness, changing ecological relationships in their environment. Finally, hybrid zones allow comparative studies with the centre of origin regarding processes of parallel adaptation and speciation.

Acknowledgements

This research was supported by grant UNS-PGI 24A106 and special funds from Postgraduate Studies Department of the Universidad Nacional del Sur for the visit of Dr Gerald J. Seiler.

References

- Alexander HM, Cummings CL, Kahn L and Snow AA (2001) Seed size variation and predation of seeds produced by wild and crop-wild sunflowers. *American Journal of Botany* 88: 623–627.
- Arias DM and Rieseberg LH (1994) Gene flow between cultivated and wild sunflowers. *Theoretical and Applied Genetics* 89: 655–660.
- Bauer HA (1991) Cuarenta años en el mejoramiento del girasol (*Helianthus annuus* L.) en Argentina 1947–1987. *Helia* 14: 63–68.
- Berville A, Müller MH, Poinso B and Serieys H (2005) Ferality: risks of gene flow between sunflower and other *Helianthus* species. In: Gressel J (ed.) *Crop Ferality and Volunteerism: A Threat to Food Security in the Transgenic Era?* Boca Raton, FL: CRC Press, pp. 209–230.
- Bouma J (2003) Back to the old paradigms of soil classification. In: Eswaran H, Rice T, Ahrens R and Stewart BA (eds) *Soil Classification, a Global Desk Reference*. Boca Raton, FL: CRC Press, pp. 51–56.
- Burkart R, Bárbaro NO, Sánchez RO and Gómez DA (1999) *Eco-regiones de la Argentina*. Administración de Parques Nacionales, Secretaría de Recursos Naturales y Desarrollo Sostenible, Presidencia de la Nación, Argentina
- Burke JM, Tang S, Knapp SJ and Rieseberg LH (2002) Genetic analysis of sunflower domestication. *Genetics* 161: 1257–1267.
- Cantamutto M, Ureta S, Gutiérrez A, Presotto A and Poverene M (2007) Zonas híbridas entre especies silvestres de *Helianthus* en Argentina. 36º Congreso Argentino de Genética. *Basic and Applied Genetics (Argentina)* 18(Suppl II): 192.
- Cantamutto M, Poverene M and Peinemann N (2008) Multi-scale analysis of two annual *Helianthus* species naturalization in Argentina. *Agriculture, Ecosystems and Environment* 123: 69–74.
- Clay S and Johnson G (2002) Scouting for weeds. *Crop Management*, doi:10.1094/CM-2002-1206-01-MA.
- Covas G (1966) Antófitas nuevas para la flora pampeana. *Apuntes para la Flora de La Pampa (Argentina)* 22: 88.
- Cummings CL, Alexander HM, Snow AA, Rieseberg LH, Kim MJ and Culley TM (2002) Fecundity selection in a sunflower crop-wild study: can ecological data predict crop allele changes? *Ecological Applications* 12: 1661–1671.
- Deines SR, Dille JA, Blinka EL, Regehr DL and Staggenborg SA (2004) Common sunflower (*Helianthus annuus*) and shattercane (*Sorghum bicolor*) interference in corn. *Weed Science* 52: 976–983.
- Geier PW, Maddux LD, Moshier LJ and Stahlman PW (1996) Common sunflower (*Helianthus annuus*) interference in soybean (*Glycine max*). *Weed Technology* 10: 317–321.
- Heiser CB Jr (1954) Variation and sub-speciation in the common sunflower *Helianthus annuus*. *American Midland Naturalist* 51: 287–305.
- Heiser CB Jr (1961) Morphological and cytological variation in *Helianthus petiolaris* with notes on related species. *Evolution* 15: 247–258.
- Heiser CB Jr (1978) Taxonomy of *Helianthus* and origin of domesticated sunflower. In: Carter JF (ed.) *Sunflower Science and Technology, Agronomy series 19*. Madison, WI: American Society of Agronomy, Inc, pp. 31–53.
- Heiser CB Jr, Smith DM, Clevenger SB and Martin Jr WC (1969) The North American Sunflowers (*Helianthus*). *Memories of the Torrey Botanical Club* 22: 1–37.

- INTA – Instituto Nacional de Tecnología Agropecuaria (1990) *Atlas de Suelos de la República Argentina*. Buenos Aires: Tomos I y II.
- Jan CC and Seiler GJ (2007) Sunflower. In: Singh RJ (ed.) *Genetic Resources, Chromosome Engineering, and Crop Improvement*. Vol 4 Oilseed crops. Boca Raton, FL: CRC Press, pp. 103–165.
- Leclercq P (1969) Une sterilite cytoplasmique chez le tournesol. *Annales d'Amélioration des Plantes* 19: 99–106.
- Linder CR, Taha I, Seiler GJ, Snow AA and Rieseberg LH (1998) Long-term introgression of crop genes into wild sunflower populations. *Theoretical and Applied Genetics* 96: 339–347.
- Mann LK, King AW, Dale VH, et al., (1999) The role of soil classification in geographic information system modelling of habitat pattern; Threatened calcareous ecosystems. *Ecosystems* 2: 524–538.
- Marzocca A (1994) *Guía descriptiva de Malezas en el Cono Sur*. Buenos Aires, Argentina: INTA.
- Mercer KL, Andow DA, Wyse DL and Shaw RG (2007) Stress and domestication traits increase the relative fitness of crop-wild hybrids in sunflower. *Ecology Letters* 10: 383–393.
- Poverene MM, Cantamutto MA, Carrera AD, et al. (2002) El girasol silvestre (*Helianthus* spp.) en la Argentina: caracterización para la liberación de cultivares transgénicos. *Revista de Investigaciones Agropecuarias (Argentina)* 31: 97–116.
- Poverene M, Carrera A, Ureta S and Cantamutto M (2004) Wild *Helianthus* species and wild-sunflower hybridization in Argentina. *Helia* 27: 133–142.
- Reagon M and Snow AA (2006) Cultivated *Helianthus annuus* (Asteraceae) volunteers as a genetic 'bridge' to weedy sunflower populations in North America. *American Journal of Botany* 93: 127–133.
- Rieseberg LH (1991) Homoploid reticulate evolution in *Helianthus* (Asteraceae): evidence from ribosomal genes. *American Journal of Botany* 78: 1218–1237.
- Rieseberg LH and Seiler GJ (1990) Molecular evidence and the origin and development of the domesticated sunflower (*Helianthus annuus*). *Economic Botany* 44(Suppl 3): 79–91.
- Rieseberg LH, Carter R and Zona S (1990) Molecular tests of the hypothesized hybrid origin of two diploid *Helianthus* species (Asteraceae). *Evolution* 44: 1498–1511.
- Rieseberg LH, Beckstrom-Sternberg SM, Liston A and Arias DM (1991) Phylogenetic and systematic inferences from chloroplast DNA and isozyme variation in *Helianthus* sect *Helianthus* (Asteraceae). *Systematic Botany* 16: 50–76.
- Rieseberg LH, Baird S and Desrochers A (1998) Patterns of mating in wild sunflower hybrid zones. *Evolution* 52: 713–726.
- Rieseberg LH, Kim MJ and Seiler GJ (1999a) Introgression between the cultivated sunflower and a sympatric relative *Helianthus petiolaris* (Asteraceae). *International Journal of Plant Science* 160: 102–108.
- Rieseberg LH, Whitton J and Gardner K (1999b) Hybrid zones and the genetic architecture of a barrier to gene flow between two wild sunflower species. *Genetics* 152: 713–727.
- Rogers CE, Thompson TE and Seiler GJ (1982) *Sunflower Species of the United States*. Fargo, ND: National Sunflower Association.
- Rosales Robles E, Salinas Garcia JR, Sánchez de la Cruz R and Esqueda Esquivel V (2002) Interference and control of wild sunflower (*Helianthus annuus* L.) in spring wheat (*Triticum aestivum* L.) in northeastern México. *Cereal Research Communications* 30: 439–446.
- Sala CA, Echarte AM and Rodriguez R (1990) Una nueva especie de *Helianthus* (Compositae) para la flora argentina. *Darwiniana (Argentina)* 30: 293–296.
- Seiler GJ (1992) Utilization of wild sunflower species for the improvement of cultivated sunflower. *Field Crops Research* 30: 195–230.
- Seiler GJ and Rieseberg LH (1997) Systematics, origin, and germplasm resources of the wild and domesticated sunflower. In: Schneiter AA (ed.) *Sunflower Technology and Production, Agronomy Series 35*. Madison, WI: American Society of Agronomy Inc, pp. 21–65.
- USDA – United States Department of Agriculture (1999) *Soil Taxonomy. A basic system of soil classification for making and interpreting soils surveys*. Agriculture Handbook No. 436.
- Ureta MS, Carrera AD, Cantamutto MA and Poverene MM (2008) Gene flow among wild and cultivated sunflower *Helianthus annuus* in Argentina. *Agriculture, Ecosystems and Environment* 123: 343–349.
- Whitton J, Wolf DE, Arias DM, Snow AA and Rieseberg LH (1997) The persistence of cultivar alleles in wild populations of sunflowers five generations after hybridization. *Theoretical and Applied Genetics* 95: 33–40.