

# Evaluation of new pasture legumes options in the Pampa-Patagonia transitional region of Argentina

November 16, 2022

Renzi, J.P.<sup>1,2</sup>; Chantre, G.R.<sup>2,3</sup>; Reinoso, O.<sup>1</sup>; Quintana, M.<sup>1,3</sup>; Grand, A.<sup>1</sup>; Cantamutto, M.A.<sup>1,2,3</sup>

## ABSTRACT

Biodiversity in forage legumes adapted to semiarid-arid conditions is scarce. In the Pampa-Patagonia transitional region of Argentina (PTRA), the annual rainfall is 440 mm with annual evapotranspiration of 1690 mm. Only *Vicia villosa* Roth shows agronomic adaptation to such an environment. Dry matter production of 19 selected legume species was compared to *V. villosa* (cv. Patagonia INTA) in the first year of establishment during 2015-to-2019. *Medicago truncatula* Gaertn. and *Trifolium subterraneum* L. (cv. Goulburn) showed high winter dry matter production, similar to *V. villosa*, but poor biomass production in spring. Spring dry matter contributed  $\approx 80\%$  of the total biomass per year. Preliminary results showed that *Onobrychis viciifolia* Scop. overcame the rest of the accessions evaluated, except for *V. villosa*. *O. viciifolia* could be considered a complementary species to *V. villosa*, a candidate for initiating a process of breeding as a new pasture legume for semiarid-arid environments. This is the first record of the potential use as a pasture of *O. viciifolia* in the PTRA.

**Keywords:** semiarid-arid regions; vetch; sainfoin.

## RESUMEN

La biodiversidad de leguminosas forrajeras adaptadas a condiciones semiáridas-áridas es escasa. En la región de transición Pampa-Patagonia de Argentina (PTRA), la precipitación anual es de 440 mm con una evapotranspiración anual de 1690 mm. Solo *Vicia villosa* Roth muestra adaptación agronómica a dicho ambiente. Se comparó la producción de materia seca de 19 especies de leguminosas seleccionadas con *V. villosa* (cv. Patagonia INTA) en el primer año de establecimiento durante 2015-2019. *Medicago truncatula* Gaertn. y *Trifolium subterraneum* L. (cv. Goulburn) mostraron una alta producción de materia seca invernal, similar a la de *V. villosa*, pero una pobre producción de biomasa en primavera. La materia seca de primavera aportó  $\approx 80\%$  de la biomasa total por año. Los resultados preliminares mostraron que *Onobrychis viciifolia* Scop. superó al resto de las accesiones evaluadas, excepto a *V. villosa*. *O. viciifolia* podría considerarse una especie complementaria a *V. villosa*, candidata a iniciar un proceso de mejoramiento como nueva leguminosa de pastoreo para ambientes semiáridos-áridos. Este es el primer registro del uso potencial como forrajera de *O. viciifolia* en la PTRA.

**Palabras clave:** regiones semiáridas-áridas, vicia, esparceta.

<sup>1</sup>Instituto Nacional de Tecnología Agropecuaria (INTA), Estación Experimental Agropecuaria (EEA) Hilario Ascasubi, Ruta 3 km 794 (8142) Hilario Ascasubi, Buenos Aires, Argentina. Correo electrónico: renzipugni.juan@inta.gob.ar

<sup>2</sup>Universidad Nacional del Sur, Departamento de Agronomía, San Andrés 800, (8000) Bahía Blanca, Argentina.

<sup>3</sup>Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS-CONICET), Camino de la Carrindanga km 7 (8000) Bahía Blanca, Buenos Aires, Argentina.

## INTRODUCTION

The semiarid-arid Pampa-Patagonia transitional region of Argentina (PTRA) encompasses nearly 18 million hectares. The predominant farming systems combine cereal cropping with cattle production. Annual precipitation (PP) varies between 370 to 590mm, with 67±13% of the precipitation distributed from autumn until spring, but high evapotranspiration (PET) especially during the summer months. During the spring-summer period, the aridity index (AI=PP/PET) is less than 0.2, corresponding to the arid climate type (UNEP 1993). The mean annual temperature is about 14.8°C with an average minimum temperature during the winter of 7.9°C and a maximum of 22.4°C during summer. As in other semiarid-arid regions of the world, the soils of the PTRA present high risk of eolic erosion, compaction, as well as low fertility (Burkart *et al.*, 1999; Silenzi *et al.*, 2012). Additionally, this region receives low inputs of fertilization and control of weeds, insects, and diseases.

The diversity of native herbaceous legumes is low and perennial grasses dominate the grasslands of the PTRA (Busso, 1997). Naturalised populations of woolly burr medic (*Medicago minima* L.) show a notorious adaptation in about 10 million hectares (Fresnillo, 1990). However, this species shows fast senescence, low fodder production and very low drought tolerance (Covas, 1978). Moreover, traditional perennial pasture legumes, such as lucerne (*Medicago sativa* L.), white clover (*Trifolium repens* L.), and red clover (*Trifolium pratense* L.) show lack of adaptation to dry conditions and are not used in the PTRA (Iurman *et al.*, 2008).

Scientific information about the benefits of legumes in cereal rotations is extensive. The Australian farming systems, known as 'ley farming' and 'phase farming', are based on legume crops. Australian 'ley farming' practice, which stands for the inclusion of an annual pasture legume in a cereal crop rotation, is a well-known solution to revert environmental and economic problems derived from monoculture practices (Nichols *et al.*, 2007, 2012). Traditionally, the most common species used for this purpose were subterranean clover (*Trifolium subterraneum* L.) and annual medics (*Medicago* spp.) (Loi *et al.*, 2005).

The western Mediterranean-Asiatic region is a major area of legume biodiversity which includes a wide range of annual and perennial species, currently grown under semiarid-arid environments. Among the diverse range of adapted pasture legume species, vetches (*Vicia* spp.) and vetchlings (*Lathyrus* spp.) are considered the most agronomically important species. The *Onobrychis*, *Hedysarum* and *Ornithopus* genera have great potential as future pasture legumes (Hayot Carbonero *et al.*, 2011; Ates *et al.*, 2013).

Despite all the scientific advances generated for the conditions in Australia and the Mediterranean basin, no studies have yet been conducted in the PTRA. In this region, legume adaptation also presents some difficulties mainly associated with drought and high rainfall variability. *Vicia villosa* Roth is an annual legume species, well adapted to different semiarid-arid temperate areas of the world. It is considered as a reference legume in the PTRA due to its confirmed adaptation and a large number of references for its use and management (Renzi *et al.*, 2017, 2019).

The current work aims to explore opportunities for developing new pasture legumes to complement *V. villosa* (Renzi and Cantamutto, 2013). A key requirement for farmers to adopt a new pasture is that it should present an acceptable biomass production during the first year of establishment. The objec-

tive of the present work was to assess the suitability of different legume species compared to *V. villosa* during the first year of establishment in the PTRA.

## MATERIALS AND METHODS

### Field experiment

Legume germplasm was provided by the Gene Bank of the Research Station Hilario Ascasubi (62°37'W, 39°23'S), located in the south of Buenos Aires province. Twenty-seven legume accessions were cultivated in the area of influence of the Research Station from 2015-2019. The selected legumes include 5 species of the genus *Medicago*, thirteen of *Trifolium*, and one of *Onobrychis*. All species showed antecedents of autumnal sowing and use as forage. Most of them are annual species, except for *T. fragiferum*, *T. hybridum*, *T. medium*, *T. montanum*, *T. ochroleucon*, *T. pannonicum*, and *O. viciifolia* (Kaiser, 2015). Within the evaluated accessions, the cultivars (cv) indicate a greater breeding effort compared to the rest of the accessions (table 1).

A field experiment was performed each year from 2015-2019 with similar sowing dates (10<sup>th</sup> of April 2015, 27<sup>th</sup> of April 2016, 11<sup>th</sup> of April 2017, 22<sup>th</sup> of March 2018, 5<sup>th</sup> of April 2019). The accessions evaluated, taxonomic names, common names, references for cultivars and sowing rates are listed in table 1.

The experiment was arranged in a randomized block design (n=3). The experimental units were 4x1.4m. A small plot cone seeder or row manual seeder was used for sowing at 20cm row spacing. During the growing season, observations of flowering time were recorded (2015, 2016, and 2019). Growing degree days (GDD; threshold temperature = 0°C) for the duration between sowing and flowering were estimated. A single cut per year of the forage dry matter (DM) production was measured at the end of winter (mid-September) and in late spring (November) by cutting plant shoots at ground level in 0.16 m<sup>2</sup> samples. The aboveground biomass harvested was oven-dried at 65°C for 72 hours.

The legume seed was inoculated with a commercially available rhizobial inoculum. Seed germination testing was previously carried out according to ISTA rules (International Seed Testing Association, 2019) to adjust the sowing rates (table 1). Fertilizers were avoided to exclude chemical interactions with bacterial activity. The seedbed was disked lightly and then levelled in the late summer before sowing the plots. Small grains were cultivated before establishing the field experiments with legumes.

The soil surface layer (0-20 cm) of the experimental sites were *entic haplustoll*, sandy loam, slightly alkaline (pH = 7.9±0.2), with a low organic matter content (1.7±0.5%) and a high P content (15.6±5.9 ppm P Bray & Kurtz). Weather data from each year were gathered from a meteorological station located less than 800 m from the experimental site (<http://inta.gob.ar/documentos/informes-meteorologicos>).

### Statistical analysis

A linear mixed model (LMM) was performed to test if the accessions differed in the winter and spring DM production, and GDD. We considered the accession as fixed factors and the year as a random factor. The blocks were nested within the year. The LMM was performed with R version 3.6.3 using Infostat software (Di Rienzo *et al.*, 2019) interface to R. The coeffi-

| Species                            | Common name        | Accession or cultivar        | Seed        | Sowing rate** | Inoculant group*** | Year                    |
|------------------------------------|--------------------|------------------------------|-------------|---------------|--------------------|-------------------------|
|                                    |                    |                              | weight (mg) |               |                    |                         |
| <i>Medicago lupulina</i> L.        | Black medic        | SVKVEF-004                   | 1.57±0.23   | 10            | AM                 | 15'; 19'                |
| <i>M. minima</i> (L.) Bart.        | Woolly burr medic  | CZENIJ-104                   | 1.20±0.04   | 10            | AM                 | 15'                     |
| <i>M. rigidula</i> L.              | Tifton medic       | SRBIST-104                   | 4.45±0.13   | 10            | AM                 | 15'                     |
| <i>M. polymorpha</i> L.            | Burr medic         | cv. Scimitar                 | 2.80±0.17   | 10            | AM                 | 15'; 16'; 17'; 18'; 19' |
| <i>M. truncatula</i> Gaertn.       | Barrel medic       | cv. Chetah                   | 4.37±0.06   | 10            | AM                 | 15'; 16'; 17'; 18'; 19' |
|                                    |                    | cv. Jester                   | 3.47±0.06   | 10            | AM                 | 15'; 16'; 17'; 18'; 19' |
| <i>Onobrychis viciifolia</i> Scop. | Sainfoin           | SRBIST-059                   | 20.8±1.02*  | 42            | Special#           | 15'; 16'; 17'; 18'; 19' |
|                                    |                    | Advanced Line INTA (AD-INTA) | 22.9±1.76*  | 40            | Special#           | 16'; 17'; 18'; 19'      |
| <i>Trifolium ochroleucon</i> Huds. | Sulphur clover     | CZENIJ-112                   | 2.17±0.15   | 12            | B                  | 15'                     |
| <i>T. arvense</i> L.               | Rabbit-foot clover | SRBIST-033                   | 0.47±0.06   | 6             | B                  | 15'                     |
| <i>T. fragiferum</i> L.            | Strawberry clover  | CZENIJ-111                   | 1.77±0.21   | 12            | B                  | 15'                     |
| <i>T. hybridum</i> L.              | Alsike clover      | SRBIST-032                   | 0.48±0.13   | 15            | B                  | 15'                     |
|                                    |                    | cv. Taipan                   | 1.10±0.10   | 6             | C                  | 15'; 16'                |
|                                    |                    | cv. Viper                    | 1.07±0.12   | 5             | C                  | 15'; 16'; 17'; 19'      |
| <i>T. michelianum</i> Savi.        | Balansa clover     | cv. Frontier                 | 0.90±0.05   | 5             | C                  | 16'; 17'; 18'; 19'      |
|                                    |                    | SRBIST-008                   | 2.09±0.11   | 12            | B                  | 15'                     |
| <i>T. medium</i> L.                | Zigzag clover      | SRBIST-008                   | 2.09±0.11   | 12            | B                  | 15'                     |
| <i>T. montanum</i> L.              | Mountain clover    | SVKVEF-016                   | 0.68±0.06   | 10            | B                  | 15'                     |
| <i>T. retusum</i> L.               | Teasel clover      | SRBIST-106                   | 0.13±0.06   | 5             | B                  | 15'                     |
| <i>T. striatum</i> L.              | Knotted clover     | SRBIST-101                   | 2.13±0.15   | 10            | B                  | 15'                     |
| <i>T. pannonicum</i> L.            | Hungarian clover   | SRBIST-068                   | 2.90±0.07   | 10            | B                  | 15'                     |
| <i>T. resupratum</i> L.            | Persian clover     | cv. Nitro                    | 0.57±0.06   | 10            | C                  | 15'; 16'; 17'; 18'      |
|                                    |                    | cv. Goulburn                 | 4.90±0.36   | 15            | C                  | 15'; 16'; 17'; 18'      |
|                                    |                    | cv. Riverina                 | 4.97±0.15   | 17            | C                  | 15'; 16'                |
|                                    |                    | cv. Urana                    | 4.83±0.29   | 15            | C                  | 15'; 16'; 17'; 18'      |
|                                    |                    | cv. Bindoon                  | 4.77±0.21   | 15            | C                  | 15'; 16'; 17'; 18'      |
| <i>T. vesiculosum</i> Savi.        | Arrowleaf clover   | cv. Zulú II                  | 1.37±0.06   | 8             | C                  | 15'; 16'; 17'; 18'; 19' |
| <i>Vicia villosa</i> Roth          | Hairy vetch        | cv. Patagonia INTA           | 31.2±0.74   | 20            | E                  | 15'; 16'; 17'; 18'; 19' |

Table 1. Sample size and serology results. Analysis of the 3'UTR of *SLC11A1* bovine gene

\*unhulled seed

\*\* kg of clean, germinable seed ha<sup>-1</sup>

\*\*\*Inoculant group: AM: *Sinorhizobium medicae*, B and C: *Rhizobium leguminosarum* bv. *trifolii*, E: *R. leguminosarum* bv. *viciae*, and Special: *Rhizobium* sp.

# Nodulaid®

cient of variation (CV%) for each accession was calculated only in those that were evaluated for more than one year (table 1).

The relationship between spring DM and GDD was plotted using GraphPad Prism Software version 8.0 (GraphPad, San Diego, California, USA).

## RESULTS

Pampa-Patagonia transitional region of Argentina displays strong climate variability with favourable periods for plant growth during autumn and spring. The annual precipitation from 2015-2017 was similar to the long-term mean (>400 mm), while

<400 mm year<sup>-1</sup> was registered in 2018 and 2019 (table 2).

*Medicago rigidula*, *Trifolium arvense*, *T. montanum*, *T. retusum* and *T. striatum* L. failed to establish (death of seedlings) during 2015. Therefore, they were not evaluated in subsequent years. Similarly, *M. minima*, *T. ochroleucon*, *T. fragiferum*, *T. hybridum*, *T. medium* and *T. pannonicum* showed very low winter DM yield during 2015 (figure 1a), thus they were excluded from further evaluations.

Both winter and spring DM mean values are presented in figure 1a, showing a wide range of oscillation (0 to 2.1 t ha<sup>-1</sup> and 0 to >6 t ha<sup>-1</sup> during winter and spring seasons, respectively).

| Month     | Precipitation (mm) |      |      |      |      | PET (mm) | Temp. (°C) |
|-----------|--------------------|------|------|------|------|----------|------------|
|           | 2015               | 2016 | 2017 | 2018 | 2019 |          |            |
| January   | 69                 | 133  | 4    | 5    | 19   | 270±21   | 23.5±1.3   |
| February  | 113                | 74   | 44   | 35   | 5    | 202±14   | 22.7±1.2   |
| March     | 20                 | 30   | 96   | 33   | 119  | 198±18   | 19.1±0.9   |
| April     | 73                 | 26   | 58   | 31   | 3    | 97±26    | 15.7±1.9   |
| May       | 7                  | 67   | 27   | 44   | 38   | 44±19    | 11.5±1.2   |
| June      | 1                  | 15   | 25   | 11   | 30   | 42±16    | 8.4±1.1    |
| July      | 45                 | 11   | 22   | 44   | 2    | 54±36    | 7.5±0.9    |
| August    | 34                 | 23   | 86   | 2    | 4    | 76±9     | 9.7±1.0    |
| September | 15                 | 10   | 7    | 24   | 18   | 103±14   | 12.1±1.0   |
| October   | 58                 | 55   | 8    | 36   | 44   | 131±22   | 14.2±1.1   |
| November  | 64                 | 37   | 61   | 59   | 72   | 194±27   | 18.1±1.6   |
| December  | 116                | 15   | 1    | 35   | 30   | 240±21   | 21.4±0.9   |
| Annual    | 614                | 495  | 439  | 359  | 384  | 1652±72  | 15.4±0.6   |
| Historic  |                    |      | 446  |      |      | 1695     | 14.9       |

Table 2. Monthly precipitation (mm), average monthly potential evapotranspiration (PET, mm) and temperatures (°C) during the experiments in comparison to the historic average (1966-2018).

Interaction between accessions per year was observed both in winter ( $F_{46,127} = 4.9$ ;  $P < 0.001$ ) and spring DM ( $F_{46,127} = 12.6$ ;  $P < 0.001$ ). Figure 1b and c show the relationship between the coefficient of variation and winter and spring DM among accessions as measured of stability between years.

*V. villosa* (cv. Patagonia INTA) had a higher winter DM followed by *M. truncatula* (cv. Jester and Chetah) and *T. sub-*

*terraneum* (cv. Goulburn) (figure 1b). Spring DM production contributed approximately 80% of the total DM per year under PTRAs conditions. The best performance in spring DM were *V. villosa* (cv. Patagonia INTA), *O. viciifolia* (AD-INTA>ARHA-059), and *T. vesiculosum* (cv. Zulu II) (figure 1c). However, the latter showed low stability ( $> CV\%$ ) due to reducing the spring DM yield when annual rainfall was  $< 400$  mm (figure 1c).





Figure 1. Winter and spring dry matter (DM) yield in different legume accessions (a), and relation to the coefficient of variance (%) (b and c). Least significant differences (LSD) ( $p < 0.05$ ) were used to indicate differences among accessions.



**Figure 2.** Spring dry matter (DM) yields as a function of growing degree days (above 0°C) necessary to reach flowering. Least significant difference (LSD) ( $p < 0.05$ ) was used to indicate differences among accessions.

No significant relationship between flowering time (GDD) and spring DM was observed, although mid-maturing species, as the case of *V. villosa*, *T. vesiculosum*, and *O. viciifolia* ( $\approx 2000^\circ\text{Cd}$ ) showed higher spring DM production, compared to early-maturing ( $< 2000^\circ\text{Cd}$ ) and late-maturing accessions ( $> 2000^\circ\text{Cd}$ ) (figure 2).

## DISCUSSION

This study explored opportunities for developing new pasture legumes to complement the *V. villosa* (Renzi *et al.*, 2019). *M. truncatula* and *T. subterraneum* L. (cv. Goulburn) showed high winter dry matter production, similar to *V. villosa*, but a poor biomass production in spring. Spring dry matter contributed  $\approx 80\%$  of the total biomass per year. We found evidence that *O. viciifolia* performs well for spring DM production and moderate stability (CV%) during the establishment (figure 1c). *O. viciifolia* is a drought tolerant pasture legume with many positive agronomic, environmental, nutritional and nutraceutical attributes, and currently, there has been a resurgence of interest in it (Mora-Ortiz and Smith, 2018). Cultivation as a pasture legume is increasing in semiarid-arid environments, like the western United States (Hybner, 2013; Sintim *et al.*, 2015) and northeast Spain (Delgado *et al.*, 2002; Cirujeda *et al.*, 2019). This situation supports the possibility of its development as a new pasture legume under PTRAs conditions.

While in some legume accessions the DM production was very low, these experiments were valid for accession comparison but not for generalizing about the species performance as a whole. However, like in the dryland of West Asia and North Africa, it would seem that vetches would appear to have greater potential use than annual medicines (Ates *et al.*, 2013; Cicek *et al.*, 2020). *V. villosa* was the best species in DM production due to its breeding and development locally, and *O. viciifolia* could be a new option for PTRAs.

A common outcome in this type of study is the need for earlier accessions for semiarid-arid Mediterranean climates,

based on the fact that delaying flowering is counterproductive because it is exposed to terminal drought (Thomson *et al.*, 1997; Berger *et al.*, 2002). Moreover, in Mediterranean climates most rainfall occurs at the same time as the low evaporative demand, improving the rainfall-use efficiency and pasture growth (Cobon *et al.*, 2019). Our results show that mid-maturing accessions present the best performance under PTRAs conditions. The mid-maturing accessions would perform better than the early flowering types in low and unpredictable rainfall environments because they show a greater ability to modify growth according to the occurrence of rainfall events. The good adaptation of *V. villosa* does not correspond to a greater tolerance to drought periods, being higher in *O. viciifolia* (Schvachsa, 2018). In *V. villosa*, the shallow root system with good lateral branches as well as the mid-maturity cycle could take advantage of the low and occasional rain events (Renzi and Cantamutto, 2013). *V. villosa* is well adapted to PTRAs farming systems, possibly due to different factors, such as an autumn–spring adjusted growing season, cold hardiness, a long vegetative phase that maximizes productivity, phenotypic plasticity with the ability to modify growth according to the occurrence of rainfall events, and excellent nodulation (Renzi *et al.*, 2017). *O. viciifolia* shows good behaviour both in the spring DM production and tolerance to drought, with a higher survival rate under the evaluated conditions.

Based on a derivation from current experiments, *O. viciifolia* is included as a crop for persistence evaluation and on-farm weed control trials. We observed that *O. viciifolia* with a range of  $0.9 \pm 0.3$  to  $5.8 \pm 2.0$  t DM ha<sup>-1</sup> year<sup>-1</sup>, depending on rainfall occurrence (232-to-575 mm year<sup>-1</sup>), could persist during more than three years under PTRAs (unpublished data). As *O. viciifolia* has shown adaptation to dryland conditions, it is imperative to evaluate different genotypes for further use in breeding programs for developing cultivars. Also, annual pasture legumes, like *V. villosa*, could be sown in mixtures with deep-rooted *O. viciifolia* to improve the forage seasonal distribution, water use, and increase the resilience of pastures to environmental

stresses (Dear 2003; Cicek *et al.*, 2020). This study can be useful for a similar semiarid-arid environment considering also the predicted climate change scenarios.

## CONCLUSIONS

This study explored the dry matter production of 19 selected legume species during the first year of establishment. Besides the well-known hairy vetch, we found evidence to show that *O. viciifolia* (sainfoin) demonstrated good performance for spring DM production and drought tolerance. This is the first exploratory study of the potential use of sainfoin as a pasture in the PTRA.

## DECLARATION OF COMPETING INTEREST

The authors declare that they have no competing interests

## ACKNOWLEDGEMENTS

The study reported in this manuscript was supported by the Federal Institute of Agricultural Technology-INTA (PE-I142, PE-I086, and Convenio MAIBA-INTA). Appreciation is extended to J. Castillo and M. Bruna, agricultural specialists, for assistance in establishing the field experiments.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in zenodo at <https://doi.org/10.5281/zenodo.5363431>.

## REFERENCES

- ATES, S.; FEINDEL, D.; EL MONEIM, A.; RYAN, J. 2013. Annual forage legumes in dryland agricultural systems of the West Asia and North Africa Regions: research achievements and future perspective. *Grass and Forage Science*, 69, 17-31. doi: 10.1111/gfs.12074
- BERGER, J.D.; ROBERTSON, L.D.; COCKS, P.S. 2002. Agricultural potential of Mediterranean grain and forage legumes: Key differences between and within *Vicia* species in terms of phenology, yield, and agronomy give insight into plant adaptation to semi-arid environments. *Genetic Resources and Crop Evolution*, 49, 313-325.
- BURKART, R.; BÁRBARO, N.O.; SANCHEZ, R.O.; GÓMEZ, D.A. 1999. *Eco-Regiones De La Argentina*. Administración de Parques Nacionales. Secretaría de Recursos Naturales y Desarrollo Sostenible, Presidencia de la Nación Argentina.
- BUSSO, C.A. 1997. Towards an increased and sustainable production in semi-arid rangelands of central Argentina: Two decades of research. *Journal of Arid Environments*, 36, 197-210.
- CICEK, H.; ATES, S.; OZCAN, G.; TEZEL, M.; KLING, J. G.; LOUHAICHI, M.; KELES, G. 2020. Effect of nurse crops and seeding rate on the persistence, productivity and nutritive value of sainfoin in a cereal-based production system. *Grass and Forage Science*, 1-10; doi:10.1111/gfs.12467
- CIRUJEDA, A.; MARÍ, A.I.; MURILLO, S.; AIBAR, J.; PARDO, G.; ORIOL SOLÉ-SENAN, X. 2019. May the Inclusion of a Legume Crop Change Weed Composition in Cereal Fields? Example of Sainfoin in Aragon (Spain). *Agronomy* 9, 134; doi:10.3390/agronomy9030134
- COBON, D.H.; KOUADIO, L.; MUSHTAQ, S.; JARVIS, C.; CARTER, J.; STONE, G.; DAVIS, P. 2019. Evaluating the shifts in rainfall and pasture-growth variabilities across the pastoral zone of Australia during 1910–2010. *Crop and Pasture Science*, 70, 634-647.
- COVAS, G. 1978. Forrajeras indígenas. *Ciencia e Investigación*, 34, 209-213.
- DEAR, B.S.; MOORE, G.A.; HUGHES, S.J. 2003. Adaptation and potential contribution of temperate perennial legumes to the southern Australian wheatbelt: a review. *Australian Journal of Experimental Agriculture*, 43, 1-18. doi:10.1071/EA01202
- DELGADO, I.; ANDRÉS, C.; SIN, E.; OCHOA, M.J. 2002. Estado actual del cultivo de la esparceta (*Onobrychis viciifolia* Scop.). Encuesta realizada a agricultores productores de semilla. *Pastos* 12, 235-247.
- DI RIENZO, J.; CASANOVES, F.; BALZARINI, M.G.; GONZALEZ, L.; TABLADA, M.; ROBLEDI, C. 2019. Infostat Versión. 2019. Grupo InfoStat, FCA, Universidad Nacional de Córdoba: Córdoba, Argentina.
- FRESNILLO, D.E. 1990. Estrategias ecológicas de *Medicago minima* (L.) Grufb. var. *minima* y *Erodium cicutarium* (L.) L'Herit., dos anuales de valor forrajero en el Caldenal (Ecological strategies of *Medicago minima* (L.) Grufb. var. *minima* and *Erodium cicutarium* (L.) L'Herit, two annuals of forage value in the Caldenal.) M.Sc Thesis, Universidad Nacional del Sur, 132 p.
- HAYOT CARBONERO, C.; MUELLER-HARVEY, I.; BROWN, T.A.; SMITH, L. 2011. Sainfoin (*Onobrychis viciifolia*): a beneficial forage legume. *Plant Genetic Resources: Characterization and Utilization* 9, 70-85.
- HYBNER, R.M. 2013. *Onobrychis viciifolia* Scop.: An Introduced Legume for Conservation Use in Montana and Wyoming. *Plant Materials Technical Note* N.º MT-91.
- INTERNATIONAL SEED TESTING ASSOCIATION. 2019. International rules for seed testing. Zürich, ISTA. 300 p. doi:10.15258/istarules.2019.F
- IURMAN, D.; MARINISSEN, J.; CASTOLDI, F.; LARREGUY, V.; AGAMENNONI, R.; PEREZ PIZARRO, J.; MOSCIANO, M.; TOSI, C. 2008. Sistemas agropecuarios representativos de Villarino y Patagones. Análisis y propuestas. Informe EEA. H. Ascasubi 110 p.
- KAISER, H. 2015. Adaptación, crecimiento y producción de Fabáceas forrajeras en el sur de Buenos Aires. Ing. Agr. thesis, Bahía Blanca, Buenos Aires, Argentina: Universidad Nacional del Sur, 40 p.
- LOI, A.; HOWIESON, J.G.; NUTT, B.J.; CARR, S.J. 2005. A second generation of annual pasture legumes and their potential for inclusion in Mediterranean-type farming systems. *Australian Journal of Experimental Agriculture*, 45, 289-299.
- MORA-ORTIZ, M.; SMITH, M.J. 2018. *Onobrychis viciifolia*; a comprehensive literature review of its history, etymology, taxonomy, genetics, agronomy and botany. *Plant Genetic Resources*. doi:10.1017/S1479262118000230
- NICHOLS, P.G.H.; LOI, A.; NUTT, B.J.; EVANS, P.M.; CRAIG, A.D.; PENGELLY, B.C.; DEAR, B.S.; LLOYD, D.L.; REVELL, C.K.; NAIR, R.M.; EWING, M.A.; HOWIESON, J.G.; AURICHT, G.A.; HOWIE, J.H.; SANDRAL, G.A.; CARR, S.J.; DE KONING, C.T.; HACKNEY, B.F.; CROCKER, G.J.; SNOWBALL, R.; HUGHES, S.J.; HALL, E.J.; FOSTER, K.J.; SKINNER, P.W.; BARBETTI, M.J.; YOU, M.P. 2007. New annual and short-lived perennial pasture legumes for Australian agriculture – 15 years of revolution. *Field Crops Research*, 104, 10-23. doi:10.1016/j.fcr.2007.03.016
- NICHOLS, P.G.H.; REVELL, C.K.; HUMPHRIES, A.W.; HOWIE, J.H.; HALL, E.J.; SANDRAL, G.A.; GHAMKHAR, K.; HARRIS, C.A. 2012. Temperate pasture legumes in Australia - their history, current use, and future prospects. *Crop and Pasture Science*, 63: 691-725.
- RENZI, J.P.; CANTAMUTTO, M.A. 2013. *Vicias: Bases agronómicas para el manejo en la Región Pampeana*. Ediciones INTA. Buenos Aires, Argentina. 299 p.
- RENZI, J.P.; CHANTRE, G.R.; CANTAMUTTO, M.A. 2017. Self-regeneration of hairy vetch (*Vicia villosa* Roth) as affected by seedling density and soil tillage method in a semi-arid agroecosystem. *Grass and Forage Science*, 72, 535-544.
- RENZI, J.P.; CHANTRE, G.R.; GONZÁLEZ-ANDÚJAR, J.L.; CANTAMUTTO, M.A. 2019. Development and validation of a simulation model for hairy vetch (*Vicia villosa* Roth) self-regeneration under different crop rotations. *Field Crops Research*, 235, 79-86.
- SCHVACHSA, A.E. 2018. Tolerancia a la sequía y producción de *Onobrychis viciifolia* Scop. en el Partido de Villarino (Buenos Aires). Ing. Agr. thesis, Universidad Nacional del Sur, Bahía Blanca, Buenos Aires, Argentina. 31 p.
- SILENZI, J.C.; ECHEVERRIA, N.E.; VALLEJOS, A.G.; BOUZA, M.E.; DE LUCIA, M.P. 2012. Wind erosion risk in the southwest of Buenos Aires province, Argentina and its relationship to the productivity index. *Aeolian Research*, 3, 419-425.
- SINTIM, H.Y.; ADJESIWOR, A.T.; ZHELJAZKOV, V.D.; ISLAM, M.A.; OBOUR, A.K. 2015. Nitrogen Application in Sainfoin under Rain-Fed Conditions in Wyoming: Productivity and Cost Implications. *Agronomy Journal*, 108, 294-300. doi:10.2134/agronj2015.0317
- THOMSON, B.D.; SIDDIQUE, K.; BARR, M.D.; WILSON, L.M. 1997. Grain legume in low rainfall Mediterranean type environments. 1. Phenology and seed yield. *Field Crop Research*, 54, 173-187.
- UNEP. 1993. *World Atlas of Desertification*. The United Nations Environment Programme (UNEP), London.