

EGU21-3543

<https://doi.org/10.5194/egusphere-egu21-3543>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



## Petrotectonic implications of metabasites of the Eastern Andean Metamorphic Complex at Lago O´ Higgins-San Martin, southern Patagonia

**Diego Rojo**<sup>1,2</sup>, Mauricio Calderón<sup>3</sup>, Matias Ghiglione<sup>2</sup>, Rodrigo Javier Suárez<sup>2</sup>, Paulo Quezada<sup>4</sup>, Francisco Hervé<sup>3,5</sup>, Marly Babinski<sup>6</sup>, and C. Mark Fanning<sup>7</sup>

<sup>1</sup>Arturo Prat University, Arturo Prat, Facultad de Ingeniería, Iquique, Chile (d.rojomartel@gmail.com)

<sup>2</sup>Instituto de Estudios Andinos IDEAN (Universidad de Buenos Aires - CONICET), Buenos Aires, Argentina

<sup>3</sup>Carrera de Geología, Facultad de Ingeniería, Universidad Andrés Bello, Sazié 2119, Santiago, Chile

<sup>4</sup>Instituto LAMIR, Universidad de Federal do Paraná, Centro Politécnico, Jardim das Américas, Curitiba 81531-980, PR, Brazil

<sup>5</sup>Departamento de Geología, Universidad de Chile, Plaza Ercilla 803, Santiago, Chile

<sup>6</sup>Instituto de Geociências, Universidade de São Paulo, São Paulo, Brazil

<sup>7</sup>Research School of Earth Sciences, The Australian National University, Canberra, Australia

The Eastern Andean Metamorphic Complex (EAMC) in southwestern Patagonia (4°-52°S) is a 450 km long belt mainly composed by low-grade metasedimentary rocks of Upper Devonian-lower Carboniferous, and Permian-lower Triassic ages. Previous works have suggested a passive margin environment for the deposition of the protolith. The EAMC comprise scarce interleaved tectonic slices of marbles, metabasites, and exceptional serpentinite bodies. At Lago O´ Higgins-San Martin (48°30'S-49°00'S) the metasedimentary successions are tectonically juxtaposed with lenses of pillowed metabasalts and greenschists having OIB, N-MORB, BABB and IAT geochemical affinities. The Nd-isotopic composition of metabasalts is characterized by  $\epsilon\text{Nd}_{(t=350 \text{ Ma})}$  of +6 and +7. The metabasalts show no signal of crustal contamination, instead, the mantle source was probably modified by subduction components. New and already published provenance data based on mineralogy, geochemistry and zircon geochronology indicate that the quartz-rich protolith of metasediments were deposited during late Devonian-early Carboniferous times (youngest single zircon ages around of latest Devonian-earliest Carboniferous times) sourced from igneous and/or sedimentary rocks located in the interior of Gondwana, as the Deseado Massif, for instance. Noticeable, the detrital age patterns of all samples reveal a prominent population of late Neoproterozoic zircons, probably directly derived from igneous and/or metaigneous rocks of the Brasiliano/Pan-African orogen or from reworked material from variably metamorphosed sedimentary units that crops out at the same latitudes in the extra-Andean region of Patagonia. We propose that the protolith of metabasites formed part of the upper part of an oceanic-like lithosphere generated in a marginal basin above a supra-subduction zone, where plume-related oceanic island volcanoes were generated. The closure of the marginal basin, probably in mid-Carboniferous times, or soon after. The oceanic lithosphere was likely underthrust within an east-to-northeast-dipping subduction zone, where ophiolitic rocks and metasedimentary

sequences were tectonically interleaved at the base of an accretionary wedge.