

RELATIONSHIP BETWEEN CHANGES IN CORE BODY TEMPERATURE IN LAMBS AND POST-SLAUGHTER MUSCLE GLYCOGEN CONTENT AND DARK-CUTTING

Pighin, D. G.^{1,3*}, Brown, W.², Ferguson, D. M.⁴, Fisher, A. D.⁵ and Warner, R. D.²

¹ Instituto Tecnología de Alimentos, Instituto Nacional de Tecnología Agropecuaria - INTA, De Los Reseros y Las Cabañas, Morón, Argentina

² CSIRO Animal, Food and Health Sciences, 677 Sneydes Road, Werribee, Vic., 3030, Australia

³ Consejo Nacional de Investigaciones Científica y Técnicas - CONICET, Av. Rivadavia 1917, Buenos Aires, Argentina

⁴ Livestock Welfare, CSIRO Livestock Industries, Locked Bag 1, Armidale, Australia

⁵ Faculty of Veterinary Science and Animal Welfare Science Centre, The University of Melbourne, Werribee, Australia

* Corresponding author (phone: +54-11-4621-0446; fax: +54-11-4621-2012; e-mail: dpighin@cnia.inta.gov.ar)

Abstract

Pre-slaughter stress may decrease muscle glycogen content, a key element for a suitable low ultimate pH and prevention of dark-cutting meat. Body temperature monitoring is a tool used in research on animal stress, as an indicator of stress events. Possible relationships between body temperature of sheep and *post-mortem* muscle glycogen were investigated in this study. Body temperature was measured with intravaginal loggers inserted into each animal at 3 days pre-slaughter, to record body temperature every 3 minutes over a period of 3 days. Blood samples were collected from each animal at exanguination for measurement of glucose and lactic acid concentrations. The muscle content of glycogen and lactic acid were determined in samples of *Longissimus* muscle collected at the level of the 13th rib, at 1 h post-slaughter. A plot of body temperature vs time showed a rise in body temperature from all animals during events such as mustering, loading onto the truck, unloading at the abattoir, during pre-slaughter handling and at slaughter. Pearson's correlation coefficients were determined between 1) the main temperature increments occurring between farm and slaughter; and 2) post-slaughter muscle glycogen and lactate levels. A significant negative correlation was detected between elevation in core body

temperature due to physical stress of sheep and muscle glycogen levels at slaughter. A low correlation was detected between body temperature and blood glucose or lactate concentrations. Further research should examine the relationship between core body temperature and meat quality in order to better understand the complex relationship between animal stress and meat quality.

Keywords—Glycogen, Temperature, Stress, Dark-Cutting, Ultimate pH

I. INTRODUCTION

It is well known that muscle glycogen concentration at the time of slaughter is a key factor involved in the *post mortem* anaerobic glycolysis of muscle cells (Lawrie, 1958). Adequate muscle glycogen content will lead to a suitable pH decline in the conversion of muscle into meat, which in turn is related to the overall acceptability of meat (Tarrant, 1989a). In contrast, low levels of muscle glycogen will limit pH decline, leading to high ultimate pH of meat, also known as dark-cutting. Dark-cutting meat generally has less desirable flavour, tenderness, shelf-life and overall acceptability (Shorthose, 1989; Purchas & Aungsupakorn, 1993). The storage of muscle glycogen and the rate of *post mortem* pH decline also depend on several factors including diet, exercise, muscle fibre composition, stress and muscle temperature (Tarrant, 1989b; Zhua, Ruusunenb, Gusellac, Zhoua & Puolanne, 2011).

Pre-slaughter stress may decrease muscle glycogen content through activation of the nervous system, mediated by catecholamines (Kuchel, 1991), and/or the endocrine system, mediated by glucocorticoids (Dantzer, 1994). A number of biochemical, physiological and behavioural measurements have been previously used to measure potential effects of animal stress (Moberg, 2000). However, most of these measures are invasive or not well established, requiring expert staff and being time consuming.

In recent years, body temperature has started to be examined more closely as an indicator of animal stress. It is known that an animal's temperature is a constant balance between heat gain and

heat loss. To maintain body temperature at a constant point, several key mechanisms are involved, known collectively as the thermoregulatory system (Robertshaw, 1985). Moreover, core body temperature can also be dramatically increased with muscular activity, nervous and hormonal factors (such as sympathetic nervous activity), catecholamines, and thyroid hormones (King, 2004). At present, the capacity to continuously log body temperature of freely behaving animals has become easy, non invasive and cost effective (Bluett, Fisher & Waugh, 2000; Lea, Niemeyer, Reed, Fisher & Ferguson, 2008). Moreover, an interesting relationship between body temperature and the level of stress, either physical and/or psychological, at several stages of animal handling has been also shown (Ferguson, 2003; Takakazu, Oka, & Tetsuro, 2001). As temperature logging is a method that significantly lacks stress associated with animal handling, it may be a useful tool in animal stress physiology research.

The aim of the present study was to investigate the possible relationships between the body temperature of sheep and *post mortem* muscle glycogen concentration, in order to predict dark-cutting meat.

II. MATERIALS AND METHODS

Animals

Forty cross-bred female lambs (first-cross: Merino x Border Leicester -BL-, and second-cross: Merino x BL x Poll Dorset), approximately 6 months old were used in the study. Animals were raised in the fields of the experimental farm at DPI Rutherglen (VIC, Australia) and were fed with a pasture diet (lucerne, cereal regrowth and cereal stubble). The experiment was conducted in May 2009, when daily temperatures ranged in average between 0.8 °C and 15.0 °C.

Animal handling and experimental procedures were approved by the Agricultural Research and Extension Animal Ethics Committee of Victorian Department of Primary Industries, Australia.

Temperature logging

Temperature data loggers (Dallas Thermocron iButton, DS1921 H, Maxim Integrated Products, Sunnyvale, CA, USA) were programmed to start logging on Sunday at 2.00 PM and to end logging on Tuesday at 7.00 PM, when slaughter was planned to be completed, with individual temperatures recorded every 3 minutes. Loggers were coupled to the plastic wings of progesterone-free ovine intravaginal progesterone releasing devices (CIDR Interag, Hamilton, New Zealand) as described by Lea, Niemeyer, Reed, Fisher, & Ferguson (2008). The devices were washed in disinfectant solution and were inserted intravaginally in each animal two days before starting to log, using an applicator lubricated in the same disinfectant solution.

Temperature monitoring included a period of 16 h of grazing on the farm, 20 min of mustering prior to transport, 6 h of yarding, 1 h of pre-transport weighing, 13 min of truck loading, 6 h of transport, 5 min of truck unloading, about 16 h at the abattoir's awaiting slaughter in lairage pens, and finally, the assembly prior to slaughter (5-10 min). Temperature loggers were removed at about 5 min post-slaughter, while the carcass was hanging upside-down. One data logger was not able to be located and retrieved from a carcass and was recorded as missing. Data was downloaded to a PC using the iButton receptor connected to a 1-Wire adapter (Dallas Thermocron, DS1402D-DR8 iButton receptor and DS9490R USB 1-Wire adapter, Maxim Integrated Products, Sunnyvale, CA, USA). Data was displayed and analysed using the software provided by Dallas Thermocron (Maxim Integrated Products, Sunnyvale, CA, USA).

Samples and pH measurement

Blood samples were collected from each animal at exanguination into tubes containing heparin as anticoagulant and immediately placed on ice. Samples were centrifuged (3000 x g, 10 min, 4 °C) within 5 h post-slaughter, plasma was separated and then stored at -20 °C until analysis for glucose and lactic acid concentrations.

Muscle samples were removed from *m. longissimus lumborum* (LL) at 1 h post-slaughter. Samples of LL were immediately frozen in liquid nitrogen until processing for glycogen and lactic acid levels. The ultimate pH of the *m. semitendinosus* (ST) and LL was measured at 24 h *post*

mortem, using a Micrometer pH Vision Model 6007 (Jenco Instruments, San Diego, CA) with a direct pH probe (Ionode Model No. IJ42).

Laboratory Methods

Plasma glucose and lactic acid concentrations were measured using enzymatic commercial kits (Sigma-Aldrich Pty, MO and Randox Labs Ltd., UK, respectively).

Muscle glycogen concentration was determined according to Dreiling, Brown, Casale & Kelly (1987). Briefly, muscle homogenates were incubated in amyloglucosidase solution during 90 min to convert glycogen to glucose. The concentration of glucose was determined spectrophotometrically after 30 min incubation in GOD/POD solution. Muscle L-lactic acid level was determined in the same muscle homogenates as described by Noll (1985). Glycolytic potential was calculated from muscle glycogen and lactic acid levels according to Monin & Sellier (1985).

Statistical and Data Analysis

Basal temperature from each animal was calculated as the mean of the four lowest temperature values recorded in order to obtain a more representative lineal body temperature nadir. Peak-maximum temperatures were used to evaluate changes in core body temperature associated to animal handling. Thus, main peak increments from each animal were calculated as the difference between the maximum temperature and the corresponding basal temperature and were expressed as percentage of individual basal temperature (%). Results obtained are expressed as mean \pm standard deviation (SD). Pearson's linear regression coefficient was calculated in order to determine correlations between the different parameters assayed.

III. RESULTS AND DISCUSSION

The mean core body temperature of sheep recorded from yarding, day 1, to the time of slaughter,

day 3, is shown in Fig. 1. Main temperature increments are marked with arrows. As can be seen, core body temperature reflected the presumed stress associated with some key events. In accordance with previous observations (Bluett, Fisher, & Waugh, 2000), results obtained in the present study successfully show the application of intravaginal loggers to measure core body temperature.

Mustering, truck loading and pen change were the most representative events reflected in core body temperature increases. All these procedures involved physical stress, and thereby, physical activity. The calculated temperature increments, expressed as percentages, corresponding to these peaks are shown in Table 1. As can be seen, peak increments belonging to mustering and truck loading were the greatest contributors to the total peak increment observed during the experimental period.

Zimerman, Grigioni, Taddeo, & Domingo (2011) have recently reported a significant increase in blood cortisol concentration in castrated kids exposed to physical exercise (moved at 3 km/h) during 30 min prior to slaughter, comparable to cortisol concentration found in animals exposed to psychological stress. The authors have proposed that fear to human presence during the assay may be responsible for this finding. However, in the present study, it would be possible that the temperature increase recorded was consequence of both physical exercise and psychological stress. Both mechanisms would lead to the consumption of stored energy in the muscle.

Results of biochemical parameters assayed in blood and muscle collected at slaughter are shown in Table 2. Blood glucose was detected slightly increased in comparison with normal resting values previously published by other authors (Ross & Kitts, 1969), possibly reflecting stress and fear of the animals immediately before slaughter. Nevertheless, lactic acid concentrations were detected within values previously published (Cottrell, McDonagh, Dunshea & Warner, 2008; Pighin, Cunzolo, Zimerman, Domingo, Pazos, & Grigioni, 2012; Ponnampalam, Warner & Dunshea, 2012). This finding does not agree with recent studies which suggested that excitation or fear sharply increases the blood concentrations of lactic acid in pigs (Edwards *et al.*, 2010) and cattle (Warner, Ferguson, Cottrell & Knee, 2007). A possible explanation could be the different stress susceptibility of animal

species and the level of stress involved in the different studies. Nevertheless, taking into account the metabolite levels, it is possible to believe that relatively low stress took place during the slaughter period in the present study. On the other hand, the SD observed also suggests an important animal variability associated with stress susceptibility.

Glycogen and lactic acid levels in the LL were within a normal range (Cottrell, Mc Donagh, Dunshea, & Warner, 2008), which was reflected in ultimate pH of both ST and LL muscles (Table 2). Consequently, no dark-cutting meat was observed in the present study. Muscle glycogen content at 1 h post-slaughter ranged within levels corresponding to well-fed animals (Apple, Dikeman, Minton, McMurphy, Fedde, Leith & Unruh, 1995). This finding would agree with the concept that some loss of glycogen *in vivo* -above the threshold of 40-57 $\mu\text{mol/g}$ - can be accommodated without deleterious effects on meat quality (Howard, 1963; Tarrant, 1989).

The Pearson's correlation coefficients between blood and muscle biochemical parameters, ultimate pH for the ST and LL and major temperature peak and total increment of temperature are shown in Table 3. As shown, the highest correlation coefficient obtained was between temperature Peak 1 and the LL muscle glycogen level or glycolytic potential ($r = -0.455, -0.419$ respectively; $P < 0.01$ for both). There was also a significant correlation between total temperature increase and muscle glycogen or glycolytic potential ($P < 0.05$ for both).

In the present study, none of the carcasses were defined as dark-cutting ($\text{pHu} > 6.0$), and therefore it would be expected that the relationship between ultimate pH and temperature would not be strong. Nevertheless, ultimate pH of ST muscle showed a significant negative correlation with the total temperature increase. Since the ST muscle is more involved with body movements than the LL, this finding could be explained by means of the different fibre composition of these muscles.

Core body temperature depends on the heat load, which in turn depends on the environment's heat and the temperature associated to food oxidation (Youseft, 1985). Since regulation of body temperature in mammals is inextricably linked to hydration state (Simon, Pierau, & Taylor, 1986), climatic conditions, exercise and diet would be key issues to take into account in temperature

monitoring. Recently, McKinley, Weissenborn, & Mathai (2009) have noted it that rehydration plays a major role in core body temperature and regulation ().

Bearing in mind that dark-cutting meat would involve greater impact of stressors on body biochemistry and, consequently, on body temperature, it is suggested that the inverse correlation found between body temperature peaks and muscle glycogen storage levels, means that temperature monitoring may be a useful tool to predict the probability of altered meat quality. Further studies should also focus attention on the effect of climatic and dietary conditions on stress-mediated body temperature increases.

IV. CONCLUSION

A significant negative correlation was found between changes in core body temperature due to physical stress of sheep and muscle glycogen levels at slaughter. There was little relationship between muscle ultimate pH and changes in body temperature but this is most likely because there was no dark-cutting high pH meat in the lamb carcasses.

Thus, changes in body temperature pre-slaughter in lambs show an interesting potential for enabling prediction of muscle glycogen levels and dark-cutting post-slaughter, and to propose changes in protocols of animal handling that contribute to avoid such meat quality alteration.

ACKNOWLEDGEMENT

The financial support from INTA (Argentina) and DPI (Australia) is gratefully acknowledged. Authors express their gratitude to Matthew Kerr, Athula Naththarampatha and Greg Seymour for their interest and technical assistance.

REFERENCES

- Aalhus, J. L. & Price, M. A. (1991). Endurance-exercised growing sheep: I. Post mortem and histological changes in skeletal muscles. *Meat Science*, 29, 43-56.
- Apple, J. K., Dikeman, M. E., Minton, J. E., McMurphy, R. M., Fedde, M. R., Leith, D. E and Unruh, J. A. (1995). Effects of restraint and isolation stress and epidural blockade on endocrine and blood metabolite status, muscle glycogen metabolism, and incidence of dark-cutting longissimus muscle of sheep. *Journal of Animal Science*, 73: 2295-2307.
- Bluett, S. J., Fisher, A. D., & Waugh, C. D. (2000). Heat challenge of diary cows in the Waikato: a comparison of spring and summer. *Proceedings of the New Zealand Society of Animal Production*, 60, 226-229.
- Cottrell, J. J., Mc Donagh, M. B., Dunshea, F. R., & Warner, R. D. (2008). Inhibition of nitric oxide release pre-slaughter increases *post-mortem* glycolysis and improves tenderness in ovine muscles. *Meat Science*, 80, 511–521.
- Dantzer, R. (1994). Animal welfare methodology and criteria. *Revue Scientifique et Technique del Office International des Epizooties*, 13, 291–302.
- Dreiling, C. E., Brown, D. E., Casale, L., & Kelly, L. (1987). Muscle glycogen: Comparison of iodine binding and enzyme digestion assays and application to meat samples. *Meat Science*, 20, 167–177.
- Edwards, L. N., Grandin, T., Engle, T. E., Porter, S. P., Ritter, M. J., Sosnicki, A. A., & Anderson, D. B. (2010). Use of exsanguinations blood lactate to assess the quality of pre-slaughter pig

handling. *Meat Science*, 86: 384-390.

Ferguson, D. M. (2003). Regulation of post mortem glycolysis in ruminant muscle. PhD Thesis. The University of New England.

Howard, A. (1963) The relation between physiological stress and meat quality. In 'Carcass composition and appraisal of meat animals'. (Ed. DE Tribe) pp. 11–21. (CSIRO: Melbourne).

King, J. (2004). Thermoregulation: Physiological responses and adaptations to exercise in hot and cold environments. *Journal of Hyperplasia Research*, 4 (3).

Kuchel, O. (1991). Stress and catecholamines. In G. Jasmin & M. Cantin (Eds.), *Stress revisited 1. Neuroendocrinology of stress* (pp. 80–103). Basel, Switz: Karger.

Lawrie, R. A. (1958). Physiological stress in relation to dark cutting beef. *Journal of Science in Food and Agriculture*, 9(11), 721-727.

Lea, J. M., Niemeyer, D. D. O., Reed, M. T., Fisher, A. D., & Ferguson, D. M. (2008). Development and validation of a simple technique for logging body temperature in free-ranging cattle. *Australian Journal of Experimental Agriculture*, 48, 741-745.

McKinley, M. J., Weissenborn, F. & Mathai, M. L. (2009) Drinking-induced thermoregulatory panting in rehydrated sheep: influences of oropharyngeal/esophageal signals, core temperature, and thirst satiety. *Am J Physiol Regul Integr Comp Physiol* 296:R1881-R1888.

Moberg, G. P. (2000). Biological response to stress: implications for animal welfare. In: G. P.

Moberg & J. A. Mench (Eds.). The biology of animal stress - Basic principles and implications for animal welfare (pp 1-22). Oxon, UK: CABI Publishing.

Monin, G. & Sellier, P. (1985). Pork of low technological quality with a normal rate of muscle pH fall in the immediate post-mortem period: the case of the Hampshire breed. *Meat Science*, 13, 49-63.

Noll, F. (1985). L+ lactate determination. In H. U. Bergmeyer (Ed.), *Methods of enzymatic analysis* (pp. 583–588). Weinheim, Germany: VCH Verlagsgesellschaft.

Pighin, D. G., Cunzolo, S. A., Zimmerman, M., Domingo, E., Pazos, A. A., & Grigioni, G. (2012). *Peri mortem* muscle biochemistry in an animal model of acute stress. *Proceedings of 58th International Congress of Meat Science and Technology*, August 2012, Montreal, Canada (*in press*).

Ponnampalam, E. N., Warner, R. D., & Dunshea, F. R. (2012). Basal and hormone metabolism in lambs varies with breed and diet quality. *Domestic Animal Endocrinology*, 42, 94-102.

Purchas, R. W., & Aungsupakorn, R. (1993). Further investigations into the relationship between ultimate pH and tenderness for beef samples from bulls and steers. *Meat Science*, 34, 163-178.

Robertshaw, D. (1985). Heat loss of cattle. In *Volume 1: Stress Physiology in Livestock. Basic Principles*, pp. 55-66. Florida: CRC Press.

Ross, J. P. & Kitts, W. D. (1969). Concentration of certain blood metabolites in obese pregnant and non pregnant ewes. *Can. J. Animal Sci.*, 49, 91-95.

- Simon, E., Pierau, K. F., & Taylor, D. C. M. (1986). Central and peripheral thermal control of effectors in homeothermic temperature regulation. *Physiol Rev*, 66, 235-300.
- Shorthose, W. R. (1989) Dark-cutting in beef and sheep carcasses under the different environments in Australia. In 'Dark-cutting in cattle and sheep'. (Eds SU Fabiansson, WR Shorthose, RD Warner) pp. 68-73. *Australian Meat and Livestock Research and Development Corporation: Sydney*.
- Takakazu, O., Oka, K., & Tetsuro, H. (2001). Mechanisms and Mediators of Psychological Stress-Induced rise in core temperature. *Psychosomatic Medicine*, 63, 476-486.
- Tarrant, P. V. (1989a). Animal behaviour and environment in the dark-cutting condition in beef- A review. *Irish Journal of Food Science and Technology*, 13: 1-21.
- Tarrant, P. V. (1989b). Animal behaviour and environment in the dark-cutting condition. In "Dark-cutting in cattle and sheep". (Eds S.U. Fabiansson, W. R. Shorthose, R. D. Warner) pp. 8-18. *Australian Meat and Livestock Research and Development Corporation: Sydney*.
- Warner, R.D., Ferguson, D. M., Cottrell, J. J., Knee, B. W. (2007). Acute stress induced by the pre-slaughter use of electric prodders causes tougher beef meat. *Australian Journal of Experimental Agriculture*, 47: 782-788.
- Yousef, M.K. (1985). Thermoneutral zone. In *Stress physiology in livestock – Volume I Basic Principles*. Ed. M.K. Yousef, 67-75. CRC Press: Florida.

Zhua, X., Ruusunenb, M., Gusellac, M., Zhoua, G. & Puolanne, E. (2011). High *post-mortem* temperature combined with rapid glycolysis induces phosphorylase denaturation and produces pale and exudative characteristics in broiler Pectoralis major muscles. *Meat Science*, 89 (2): 181-188.

Zimmerman, M., Grigioni, G., Taddeo, H. & Domingo, E. (2011). Physiological stress responses and meat quality traits of kids subjected to different pre-slaughter stressors. *Small Ruminant Research*, 100 (2-3), 137-142.

