

A Survey on Occurrence of *Cladosporium fulvum* Identifies Race 0 and Race 2 in Tomato-Growing Areas of Argentina

Rocío Medina, Centro de Investigaciones en Fitopatologías (CIDEFI), La Plata (1900), Argentina; and Centro de Investigación y Desarrollo en Fermentaciones Industriales (CINDEFI), La Plata (1900), Argentina; **Silvina M. Y. López**, Instituto de Fisiología Vegetal (INFIVE), La Plata (1900), Argentina; **Mario E. E. Franco, Cristina Rollan, and Blanca L. Ronco**, CIDEFI, La Plata (1900), Argentina; **Mario C. N. Saparrat**, INFIVE, La Plata (1900), Argentina; **Pierre J. G. M. De Wit**, Wageningen University, Laboratory of Phytopathology, Wageningen, The Netherlands; and **Pedro A. Balatti**, CIDEFI, La Plata (1900), Argentina; and Comisión de Investigaciones Científicas de la Provincia de Buenos Aires, Argentina

Abstract

Medina, R., López, S. M. Y., Franco, M. E. E., Rollan, C., Ronco, B. L., Saparrat, M. C. N., De Wit, P. J. G. M., and Balatti, P. A. 2015. A survey on occurrence of *Cladosporium fulvum* identifies race 0 and race 2 in tomato-growing areas of argentina. Plant Dis. 99:1732-1737.

The presence of *Cladosporium fulvum* (syn. *Passalora fulva*), causal agent of tomato leaf mold, was confirmed in the two main greenhouse-production areas for tomato in Argentina. Using both morphological characters and internal transcribed spacer sequencing, we confirmed the presence of physiological races of this pathogen. A diagnostic multiplex polymerase chain reaction (PCR) was also developed, using primers derived from *C. fulvum* avirulence (*Avr*) genes. In all, 20 isolates of *Cladosporium* spp. were obtained as monospore cultures and 12 were

identified as *C. fulvum*. By this method, we showed that, of these 12 isolates, 5 were race 0 (carrying functional *Avr2*, *Avr4*, *Avr4E*, and *Avr9* genes) and 7 were race 2 (lacking the *Avr2* gene). Race identity was confirmed by testing their virulence on a set of tomato differentials carrying different *Cf* resistance genes. All *Avr* genes could be amplified in single or multiplex PCR using DNA isolated from in vitro grown monospore cultures but only three *Avr* could be amplified when genomic DNA was isolated from *C. fulvum*-infected necrotic leaf tissue.

Tomato (*Solanum lycopersicum* L.) can be heavily infected by leaf mold (Cooke 1883), a disease caused by *Cladosporium fulvum* (Braun et al. 2003) (syn. *Passalora fulva*). This nonobligate biotrophic fungus infects tomato plants under conditions of high relative humidity and temperatures around 20°C (Thomma et al. 2005). Conidia settle on the lower side of the leaf, germinate, penetrate through open stomata, and colonize the intercellular space (De Wit 1977; Lazarovits and Higgins 1976). Within a week, on the upper side of the leaves, pale green or yellowish diffuse spots appear and gradually grow and become brownish. On the lower side, the fungus starts to sporulate (Blancard 1992; Thomma et al. 2005) and, in a few weeks, the spots become necrotic.

Leaf mold disease complies with the typical gene-for-gene relationship (De Wit 1992; Flor 1971). In the intercellular space of tomato leaves, the fungus secretes effector proteins that function as virulence factors on tomato plants lacking matching *C. fulvum* (*Cf*) resistance genes but function as avirulence (*Avr*) factors on tomato plants carrying the matching *Cf* genes (Dixon et al. 1996, 1998; Jones et al. 1994; Parniske et al. 1997; Takken et al. 1998; Thomas et al. 1997). Of the *Avr* genes of *C. fulvum*, four had been cloned and sequenced at the start of this study: *Avr2* (Luderer et al. 2002), *Avr4* (Joosten et al. 1994), *Avr4E* (Westerink et al. 2004), and *Avr9* (Van Kan et al. 1991). Four additional extracellular proteins (Ecp)

have been identified that are also considered to be virulence factors and are recognized by matching *Cf-Ecp* genes (Bolton et al. 2008; Joosten et al. 1994; Laugé et al. 2000; Van den Ackerveken et al. 1993; Van Esse et al. 2006).

In the past, races of *C. fulvum* were identified using a set of tomato differentials carrying different *Cf* resistance genes. (Bailey 1947; Kooistra 1964). Recent studies showed that new races of *C. fulvum* often appeared within a short period of time, especially after the introduction of a single *Cf* gene in new tomato cultivars (Iida et al. 2010). In Japan, only cultivars carrying the *Cf-2* resistance gene have been cultivated for many years. However, since 2006, new varieties of tomato carrying the *Cf-9* gene were released. Soon, the new race 2.9 of *C. fulvum* appeared (Iida et al. 2010), which could overcome both the *Cf-2* and the *Cf-9* gene (Enya et al. 2009; Satou et al. 2005; Yamada and Abiko 2002).

In Argentina, leaf mold is a rather new disease, mostly affecting tomato produced in greenhouses in the provinces Corrientes and Buenos Aires. All commercial hybrids that have been grown in these provinces are susceptible to the disease (Rollan et al. 2013). Not much is known about physiological specialization of *C. fulvum* population in Argentina. Recently, Rollan and colleagues identified two races of *C. fulvum* (Rollan et al. 2013).

Molecular biology has provided new tools that can be used for diagnostic purposes; they are accurate and not highly influenced by the environment. Such tools include intersimple sequence repeat markers that are used to study diversity among and within pathogen populations (Stenglein and Balatti 2006) and specific markers that allow the identification of species or races (Theerakulpisut et al. 2008).

Recently, Stergiopoulos and colleagues (2007) developed a set of primers based on the sequences of cloned *Avr* genes of *C. fulvum*. They were used to study the allelic variation of *Avr* genes in a collection of isolates of *C. fulvum*. Yan et al. (2007) developed a real-time polymerase chain reaction (PCR) assay aimed at detecting *C. fulvum* on tomato leaves with primers derived from three fungus-specific sequences. In this study, using morphological as well as molecular tools, we isolated and identified *C. fulvum* from the two main greenhouse production areas for tomato in Argentina. In addition, we developed a diagnostic multiplex

Corresponding author: P. A. Balatti; E-mail: pbalatti@gmail.com

Accession numbers: ITS sequences = JQ768317 to JQ768327, KC132842, KC132843, KM488552 to KM488554, KM488556, KM488557; AVR sequences = KC132845, KM609217 to KM609220, KC132827 to KC132831, KM609221 to KM609226, KC132832 to KC132835, KC166871, KC132836 to KC132840, and KP234076 to KP234087.

*The e-Xtra logo stands for “electronic extra” and indicates that two supplementary figures and one supplementary table are published online.

Accepted for publication 9 April 2015.

PCR to identify races of the fungus using primers derived from four known *C. fulvum* Avr sequences.

Materials and Methods

Isolation, subculture, and storage of monospore isolates from diseased tomato leaves. From six different locations in the two main tomato production areas of Argentina, located in the provinces of Corrientes and Buenos Aires, diseased tomato leaf samples showing typical tomato leaf mold symptoms, presumably caused by *C. fulvum*, were collected. The samples were obtained from tomato (*S. lycopersicum* L) cultivars 'Elpida' (Enza Zaden), 'Tarija', 'Potosí' (BHN Research), 'Arcos' (Wisdom Seeds), 'Compak', 'Keitor' (Syngenta Seeds S.A.), 'Cherry Colly', and 'Colibri' (Clausen) (Rollan et al. 2013) (Table 1).

Conidia were collected from sporulating lesions and were transferred with a needle to petri plates filled with water agar (Biopack) medium. Germinated conidia produced mycelium that was transferred and grown on 2% potato dextrose agar (PDA; Britania). Cultures of fungal isolates were maintained on PDA at 4°C. Isolates were deposited in the culture collection of the Centro de Investigaciones de Fitopatología, Facultad de Ciencias Agrarias y Forestales, Universidad Nacional de La Plata, La Plata, Pcia de Buenos Aires, Argentina.

Determination of virulence spectrum of monospore isolates on a differential set of tomato cultivars carrying different Cf genes. The virulence spectrum of the monospore isolates was determined by inoculating them on a set of differentials of 'Money Maker' (MM) tomato carrying no (*Cf-0*) or different resistance genes (*Cf-2*, *Cf-4*, *Cf-5*, or *Cf-9*). Seed were placed in trays containing sterilized perlite-soil mix substrate and were cultivated in greenhouses under controlled conditions of growth at 28 ± 2°C. Plantlets were transplanted to 2-liter plastic pots containing the same soil medium. Plants were inoculated with a conidial suspension of 1 × 10⁵ conidia ml⁻¹ essentially as described before (Rollan et al. 2013). Twenty days after inoculation, susceptible genotypes developed pale green to yellow spots on the abaxial leaf surface and pale brown to olivaceous brown sporulation on the adaxial side.

PCR amplification of internal transcribed spacer. Genomic DNA of fungi was isolated using the cetyltrimethylammonium bromide (Sigma-Aldrich) method (Galván et al. 2003) and quantified by comparing with a λ *Hind*III marker (Invitrogen) after electrophoresis on 0.7% agarose gels that were stained with ethidium bromide.

Genomic DNA was mixed with primers internal transcribed spacer (ITS)-4 (5'-TCCTCCGCTTATTGATATGC-3') and ITS-5 (5'-GGAAGTAAAGTCGTAACAAGG-3') (White et al. 1990) to amplify the ITS region. Reactions contained 1× amplification buffer (Inbio Highway), 1.5 mM MgCl₂ (Inbio Highway), 0.15 μM each primer, 200 μM each deoxynucleoside triphosphate (Inbio Highway), 25 ng of DNA template, and 1 U of *Taq* DNA polymerase (Inbio Highway) in a 15-μl volume. Reactions were performed in a PTC-1152 Mini Cycler (MJ Research) programmed as follows: an initial step at 94°C for 4 min; followed by 33 cycles of a denaturing step at 94°C for 45 s, annealing at 56°C for 45 s, and extension at 72°C for 1 min; with a final extension at 72°C for 5 min. Control experiments lacked template DNA. Amplified fragments were visualized and compared with a 100- to 1,000-bp marker (Inbio Highway) on 1% agarose gels stained with ethidium bromide and documented by means of an image analyzer (Gene Genus analyzer; Syngene).

PCR amplification of Avr2, Avr4, Avr4E, and Avr9 genes. The primers used in the amplification reactions were essentially those described by Stergiopoulos et al. (2007). We slightly modified them because the universal MP13 sequence was eliminated (AVR2f: 5'-CATCAGCA TATCCTCTTCCATCC-3', AVR2r: 5'-CAGTACGTTCAAAGCA GATAAGG-3', AVR4f: 5'-ACGGTAGGTCTGTACACGAGCC-3', AVR4r: 5'-ACCGAACTGGGTATGGAATG-3', AVR4Ef: 5'-GCC CGGTATATCGCTGTGC-3', AVR4Er: 5'-CGGAACCCCTGGCT GAGA-3', AVR9f: 5'-AATACAACCTTGAAACAGCTAGG-3', and AVR9r: 5'-GGACTCTACGGGCTTGG-3') (Table 2). The PCR mixture contained 1× amplification buffer (Inbio Highway), 1.5 mM MgCl₂ (Inbio Highway), 0.2 mM dNTPs (Inbio Highway, Argentina), 0.4 μM forward and reverse primer, 1 U of T-Plus *Taq* DNA polymerase (Inbio Highway), and 30 ng of genomic template DNA. PCR was performed in a 15-μl volume with a thermocycler (Model Multigene

gradient; Labnet International, Inc.) programmed as follows: an initial denaturing step at 94°C for 5 min; then, 40 cycles of 30 s at 94°C; an annealing step of 1 min at 63, 65, 53.5, or 62.5°C for Avr2, Avr4, Avr4E, and Avr9, respectively; and 1 min of extension at 72°C. There was a final extension of 7 min at 72°C. Amplified PCR products were separated by electrophoresis in 1% agarose gels that were stained with ethidium bromide. A 100- to 1,000-bp DNA ladder marker (Inbio Highway) was used to estimate the size of the amplicons. Gels were observed and documented in a GeneGenius analyzer (Syngene) and the bands were processed by means of GeneSnaps, GeneTools, and GeneDirectory software (Syngene).

Multiplex PCR amplification of Avr genes. Multiplex PCR was performed in a 15-μl volume with a thermocycler programmed as previously described, except for the annealing temperature that was changed to 58.4°C. Reactions were performed with genomic DNA isolated either from monospore cultures (40 to 60 ng) or necrotic diseased leaf tissue (100 ng), 1.5 mM MgCl₂ (Inbio Highway), 0.20 μM each pair of primers (corresponding to Avr2, Avr4, Avr4E, and Avr9), 0.2 mM dNTPs (Inbio Highway), and 1 U of T-Plus DNA polymerase (Inbio Highway) in 1× reaction buffer (Inbio Highway). PCR products were resolved by electrophoresis in 1% ultrapure agarose (Invitrogen) gels that were stained with ethidium bromide and exposed to UV light with a transilluminator. Gels were documented in a GeneGenius analyzer (Syngene) and the bands were processed by means of GeneSnaps, GeneTools and GeneDirectory software (Syngene).

The amplified bands generated in multiplex PCR were eluted from the gels as described by Sambrook et al. (1989), precipitated, and sequenced by Macrogen Inc.

Sequencing of ITS and Avr genes. The amplified DNA fragments were precipitated by adding 10% of a 3-M NaAc solution and one volume of isopropyl alcohol. The DNA was sequenced by the dideoxy termination method (Sanger et al. 1977) using the BigDye Terminator Cycle Sequencing Ready Reaction kit and the automated ABI Prism 3730 DNA sequencer (Applied Biosystems, Macrogen). The sequences of the ITS and the Avr genes were deposited in the National Center for Biotechnology Information (NCBI) GenBank.

Table 1. Place of isolation and identity of the isolates used in this study

Species, isolate, cultivar ^a	Site of isolation ^b
<i>Cladosporium cladosporioides</i>	
CFOT7 (this article), Tarija	Los Hornos
AAS17 (this article), Arcos	Arana
AAS2 (this article), Arcos	Arana
HD10 (this article), ND	Los Hornos
CFOPC (this article), Potosí	Olmos
CFP14 (this article), ND	Abasto
AAS16 (this article), Arcos	Arana
<i>C. sphaerospermum</i>	
HD8 (this article), ND	Los Hornos
<i>C. fulvum</i>	
ALH (this article), Elpida	Los Hornos
ELH (this article), Elpida	Los Hornos
EMP (this article), Elpida	Los Hornos
CoA (this article), Colibrí	Arana
CH6 (this article), Elpida	Corrientes
EOP (this article), Elpida	Olmos
ColA (this article), Colibrí	Arana
CK813 (this article), Keitor	Corrientes
ELS (this article), Elpida	Etcheverry
ComA (this article), Compak	Arana
EAV (this article), Elpida	Arana
CHEAV (this article), Cherry-Colly	Arana
<i>Ensifer fredii</i>	
HH103 (Cleyet-Marel 1987), Soybean	Vietnam
<i>Humicolopsis cephalosporioides</i>	
EU (Allegrucci et al. 2009), Nothofagus	Tierra del Fuego

^a ND = not determined.

^b Los Hornos Pcia de Buenos Aires, Arana Pcia de Buenos Aires, Olmos Pcia de Buenos Aires, Abasto Pcia de Buenos Aires, Corrientes = Pcia de Corrientes, Etcheverry Pcia de Buenos Aires, and Tierra del Fuego = forest soil.

Results

Isolation and identification of *C. fulvum* as the causal agent of the disease. Twenty monosporic cultures were obtained from diseased leaves and analyzed. Eight isolates developed 40-mm colonies in 15 days on 2% PDA. Among them, one produced small spherical conidia (ball like, 3 to 4 μm) and the other seven small ovoid conidia (measuring about 3 to 7 by 2 to 4 μm). They were identified as *C. sphaerospermum* and *C. cladosporioides*, respectively (Bensch et al. 2010; Crous et al. 2006, 2007, 2009; Schubert et al. 2007; Zalar et al. 2007). The 12 remaining isolates grew at a slower rate and developed 40-mm colonies in 30 days. They produced one-celled, large, pale green, ovoid conidia in accordance with previous reports (Crous 2009; Rollan et al. 2013) and were identified as *C. fulvum* (Table 1). The morphological identification was confirmed by analysis of the ITS region sequences.

All 12 isolates of *C. fulvum* were inoculated on a differential set of tomato cultivars. Isolates ELH, EOP, ELS, ComA, and EAV infected only MM-*Cf*-0 tomato, which indicated that they should be designated race 0, because they contain functional *Avr2*, *Avr4*, *Avr4E* and *Avr9* genes (Boukema 1981). The remaining seven isolates of *C. fulvum* infected MM-*Cf*-0 and MM-*Cf*-2 tomato, which indicated that they lack a functional *Avr2* gene and should be designated race 2 (data not shown).

These results were confirmed by PCR with primers derived from *C. fulvum* *Avr* genes and genomic DNA from *C. fulvum* as template. DNA fragments of 570, 806, 640, and 710 bp were amplified (Fig. 1), each representing the expected size of *Avr2*, *Avr4*, *Avr4E*, and *Avr9*, respectively. These bands were sequenced and a BLAST analysis indicated that they corresponded to *Avr2*, *Avr4*, *Avr4E*, and *Avr9*.

Multiplex PCR as a diagnostic tool. Determining the virulence spectrum of *C. fulvum* on a set of differentials containing different *Cf* genes is time consuming and expensive. Therefore, we developed a molecular diagnostic tool aimed at allowing researchers to identify the causal agents. Because the annealing temperature of the reactions designed to amplify *Avr2*, *Avr4*, *Avr4E*, and *Avr9* (Stergiopoulos et al. 2007) varied within a narrow temperature range, it was postulated that it would be possible to amplify all *Avr* genes in one reaction. The method would be more powerful if the multiplex assay could be performed directly on diseased tissue, avoiding fungal isolation. PCR containing genomic DNA of *C. fulvum* race "0" (isolate ELH) and all the primers homologous to *Avr2*, *Avr4*, *Avr4E*, and *Avr9* were run within a temperature range of 52 to 65°C. All four *Avr* genes were successfully amplified at an annealing temperature of 58°C (Fig. 2A). Such results were obtained if the reactions contained 40 to 60 ng of template DNA from *C. fulvum* and 20 ng of each primer.

In order to confirm that the PCR are specific, reactions containing template DNA of *C. fulvum* mixed with increasing concentrations of *C. cladosporioides* DNA (20, 30, and 40 ng) or DNA isolated from unrelated fungi or bacteria were performed. *Avr* genes were amplified only in reactions containing template DNA from *C. fulvum* (Fig. 2B and C).

The sizes of the fragments amplified with the multiplex PCR were identical to those obtained in individual PCR and to those described by Stergiopoulos et al. (2007) and Peteira et al. (2011). The identity of the *Avr* genes was further confirmed by analyzing the sequence of the amplicons, which showed that they were homologous to *Avr2*, *Avr4*, *Avr4E*, and *Avr9*.

To develop the diagnostic tool further, the multiplex PCR was attempted on DNA extracted from infected tissue of tomato leaves inoculated with race 0 of *C. fulvum*. In four amplification reactions aimed at amplifying each *Avr* gene using template DNA from infected leaf tissue, we successfully amplified the *Avr2*, *Avr4*, *Avr4E*, and *Avr9* genes (Fig. 3A). However, when genomic DNA was extracted from infected leaf tissue, the multiplex reaction only amplified *Avr2*, *Avr4*, and *Avr4E*. *Avr9* could only be amplified in reactions containing the *Avr9*-specific pair of primers (Fig. 3B). Also, in this case, the amplified bands were sequenced and all appeared to represent correct *Avr* gene sequences.

As described above, among all the *C. fulvum* isolates analyzed, only two different races were found. However, the *Avr* gene sequences were polymorphic (Fig. 4; Supplementary Fig. S2). *Avr4* was polymorphic at the 5' end compared with other isolates from Argentina but also compared with those sequences available at the NCBI database that corresponded to *C. fulvum* representatives isolated in Europe. Regarding *Avr4E*, the sequences isolated from Argentina were the same but they were polymorphic compared with those available at the NCBI database that correspond to an isolate from another country (Fig. 4; Supplementary Fig. S2).

Discussion

C. fulvum occurs worldwide and causes large economic losses in new tomato-growing areas. This might be due to the fact that, in these new areas, the populations of the causal agent are not known and tomato cultivars lack *Cf* resistance genes. Therefore, whenever the environmental conditions are optimal, outbreaks of the disease may occur (Butler and Jones 1949). Tomato leaf mold is a new disease that affects greenhouse-grown tomato in Argentina. Therefore, in order to start to develop strategies to control the disease, we isolated 20 monosporic cultures from plants with typical symptoms of leaf mold from the two main production areas.

Twelve isolates were identified as *C. fulvum* using morphological characteristics as well as ITS (Schoch et al. 2012) and *Avr* sequences (Stergiopoulos et al. 2007). The four *Avr* genes could be amplified using primers homologous to cloned *Avr* genes of *C. fulvum* (Stergiopoulos et al. 2007) and enabled us to discriminate between races. We found that five isolates of *C. fulvum* were race 0 and seven isolates were race 2, as previously reported by Rollan et al. (2013). These findings suggest that tomato hybrids used in these two main areas in the province of Corrientes and Buenos Aires lack *Cf* resistance genes (except *Cf*-2). Therefore, in the future, hybrids carrying at least two *Cf* resistance genes out of the five known (*Cf*-2, *Cf*-4, *Cf*-4E, *Cf*-5, and *Cf*-9) should be used to manage the disease, provided that races 0 and 2 are the only ones present in Argentina.




Fig. 1. Amplification of avirulence genes *Avr2* (570 bp), *Avr4* (806 bp), *Avr4E* (640 bp), and *Avr9* (710 bp) in polymerase chain reaction containing genomic DNA isolated from different isolates of *Cladosporium fulvum* collected from tomato hybrid Elpida. Lanes 1 and 22: 100- to 1,000-bp marker; lanes 2, 5, 8, and 11: isolate ELH; lanes 3, 6, 9, and 12: isolate EMP; lanes 4, 7, 10, and 13: isolate ALH; lanes 14, 16, 18, and 20: isolate CoA; lanes 15, 17, 19 and 21: isolate CH6.

The tomato–*C. fulvum* interaction complies with the gene-for-gene hypothesis proposed by Flor (1956). As with other pathosystems, a classical way of identifying the virulence spectrum of isolates of *C. fulvum* is to perform virulence assays on a set of differential tomato cultivars carrying different *Cf* genes. Such experiments are laborious and time consuming. Several molecular studies described the identification, cloning, and characterization of both *Avr* and *Cf* genes,




Fig. 2. **A**, Multiplex polymerase chain reaction (PCR) of *Avr2* (570 bp), *Avr4* (806 bp), *Avr4E* (640 bp), and *Avr9* (710 bp) using different amounts of primers and annealing temperatures. Genomic DNA (40 ng) was used for isolate ELH. Lane 1, 100- to 1,000-bp marker; lanes 2 to 4, 20 ng of each primer; lanes 5 to 11, 12 ng of each primer. **B**, Multiplex PCR of *Avr2* (570 bp), *Avr4* (806 bp), *Avr4E* (640 bp), and *Avr9* (710 bp) in the presence of increasing amounts of template DNA from *Cladosporium fulvum* and competitive DNA from other organisms. Lane 1, 100- to 1,000-bp marker; lanes 2 to 4 reactions with 40 ng of genomic DNA of ALH, EMP, and ELH, respectively; lanes 5 to 8, reactions with 60, 45, 30, and 15 ng of genomic DNA of isolate ELH, respectively; lanes 9 to 11, reactions with 40 ng of genomic DNA of isolate ELH mixed with 40, 30, and 20 ng of genomic DNA of *C. cladosporioides*, respectively. **C**, Multiplex PCR of *Avr2* (570 bp), *Avr4* (806 bp), *Avr4E* (640 bp), and *Avr9* (710 bp). Lane 1, 100- to 1,000-bp marker; lanes 2 to 4, reactions with 40 ng of template DNA from *Ensifer fredii* HH103, *Humicolopsis cephalosporioides*, and *C. cladosporioides*, respectively; lanes 5 to 7, reactions with 40 ng of template DNA from isolates ALH, EMP, and ELH, respectively.

which provided researchers with tools for management of the disease (Dixon et al. 1996, 1998; Jones et al. 1994; Joosten and De Wit 1999; Stergiopoulos et al. 2007; Yan et al. 2008). Therefore, we developed a diagnostic tool for identification of *C. fulvum* as well as identifying the race.

We successfully amplified *Avr* genes *Avr2*, *Avr4*, *Avr4E*, and *Avr9* in single PCR; all of the reactions included the temperature of the annealing step within a narrow window, which suggested that amplifications might be performed in a single multiplex PCR. Such a reaction amplified all four *Avr* genes from *C. fulvum* template DNA but not from bacteria (*Ensifer fredii*) or fungi, including *Humicolopsis cephalosporioides* or *Cladosporium* spp. Also, the presence of contaminating DNA did not outcompete *C. fulvum* DNA, confirming that the reaction is specific: the DNA of saprophytes or opportunistic pathogens that might be accompanying *C. fulvum* in lesions does not interfere with the reaction.

The multiplex PCR was specific and particularly accurate when template DNA was isolated from cultures of *C. fulvum* grown in vitro. However, when the reaction was run with genomic DNA isolated from infected leaf tissue, only *Avr2*, *Avr4*, and *Avr4E* were amplified. We do not have an explanation for the failure to amplify *Avr9* under the conditions described, because the gene was amplified in reactions containing only *Avr9* primers. One possibility is that, because the *Avr9* sequence is short and located in a repeat-rich region, the corresponding primers might anneal less efficiently. Therefore, both single and multiple PCR effectively detected the presence or absence of *Avr* genes. However, multiplex PCR can only be successfully applied to identify the virulence spectrum of a race, when changes in virulence are correlated with the loss of an *Avr* gene, as can sometimes be the case for *C. fulvum*. Thus far, overcoming resistance genes such as *Cf-9* has always been correlated with loss of the *Avr9* gene, while overcoming *Cf-2* and *Cf-4E* has often, though not




Fig. 3. **A**, Amplification of *Avr2* (570 bp), *Avr4* (806 bp), *Avr4E* (640 bp), and *Avr9* (710 bp) in reactions containing genomic DNA isolated from necrotic leaf tissue infected by ELH. Lane 1, 100- to 1,000-bp marker; lane 2, *Avr2*; lane 3, *Avr4*; lane 4, *Avr4E*; lane 5, *Avr9*. **B**, Lane 1, 100- to 1,000-bp marker; lane 2, multiplex polymerase chain reaction using 100 ng of genomic DNA isolated from necrotic leaf tissue infected by ELH.




Fig. 4. Relative quantity of polymorphisms found in each *Avr* sequence as related to the tomato hybrid of isolation and number polymorphisms accumulated in each *Avr*.

always, been correlated with loss of the corresponding *Avr2* and *Avr4E* genes, respectively (Luderer et al. 2002; Van den Ackerveken et al. 1992; Westerink et al. 2004). Such a change is not based on loss of *Avr* gene but on nucleotide polymorphisms in the *Avr* genes leading to amino acid changes. In the *Avr* protein, the amplified bands need to be sequenced. This has been the case with races overcoming both the *Cf-4* (Joosten et al. 1994, 1997) and *Cf-2* and *Cf-4E* genes (Luderer et al. 2002; Westerink et al. 2004).

Multiplex PCR combined with sequencing of the multiplied bands can be an alternative for testing the virulence spectrum of races on a differential set of tomato lines carrying different *Cf* genes. However, new polymorphisms in *Avr* genes that have not been reported before may evolve that could overcome matching *Cf* genes. In such instances, the multiplex PCR will not be sufficient. The *Avr* genes will need to be tested for ability to cause a hypersensitive response (HR) when coexpressed in plants carrying the complementary *Cf* gene (van der Hoorn et al. 2000).

Although the areas sampled were distant, only two races of *C. fulvum* were found. This likely reflects the fact that few *Cf* genes are present in the hybrid cultivars that are used in Argentina. The analysis of the sequences of *Avr2*, *Avr4*, *Avr4E*, and *Avr9* in race 0 showed that they were almost identical to published sequences, except for a few polymorphisms that likely do not affect the biological activity of *Avr* proteins, because HR reactions occurred in cultivars carrying the corresponding *Cf* gene. The virulence spectrum of a given race of *C. fulvum* might be determined by polymorphisms in *Avr* genes, including nucleotide substitutions, frame-shift mutations, and the complete loss of an *Avr* gene. This means that, after having obtained the amplicon of an *Avr* gene, it still needs to be sequenced to see whether it may contain nucleotide polymorphisms that result in an *Avr* protein that can escape recognition by a matching *Cf* resistance gene. Moreover, the sequence of an *Avr* gene might help to identify the origin of a particular isolate within production areas and might allow researchers to follow its evolutionary history. Stergiopoulos and colleagues (2007) studied the allelic variation of effectors within isolates of *C. fulvum* and found that *Avr* genes seem to be under continuous selection pressure imposed by *Cf* genes present in tomato cultivars grown in production areas. In addition, neutral mutational shifts can occur that do not affect the biological activity of the encoded proteins. The presence of polymorphisms within the *Avr* gene sequences suggested that races might be under an evolutionary process, which might be important considering the reduced number of isolates analyzed.

In conclusion, two races of *C. fulvum*, race 0 and race 2, were identified within 12 isolates of *C. fulvum* obtained from the two most important greenhouse-grown tomato production areas in Argentina. The multiplex PCR method described successfully amplified four *Avr* genes from a mixture of genomic fungal DNA and also from genomic DNA isolated from infected leaf tissue although, in this case, two reactions are needed. Such methodology, much quicker than a virulence assay, is an important complementary diagnostic tool for identification. Introduction of *Cf-4*, *Cf-5*, and *Cf-9* resistance genes in the new tomato hybrids can protect them against the prevailing race 0 and 2 of *C. fulvum*. These results suggest that combining two or even three *Cf* genes into one hybrid should reduce the impact of the disease (Hörger et al. 2012).

Table 2. List of primers homologous to the 5' and 3' end of *Avr2*, *Avr4*, *Avr4E*, and *Avr9*

Primer	Sequences
AVR2f	5'-CATCAGCATATCCTCTTCCATCC-3'
AVR2r	5'-CAGTACGTTCAAAGCAGATAAGG-3'
AVR4f	5'-ACGGTAGGTCTGTACACGAGCC-3'
AVR4r	5'-ACCGAACTGGGTATGGAAATG-3'
AVR4Ef	5'-GCCCGGTATATCGCTGTGC-3'
AVR4Er	5'-CGGAACCCCTGGCTGAGA-3'
AVR9f	5'-AATACAACCTGAAACAGCTAGG-3'
AVR9r	5'-GGACTCTACGGGGCTTGG-3'

Acknowledgments

R. Medina worked on a scholarship from Centro de Investigación y Desarrollo en Fermentaciones Industriales, La Plata, Universidad Nacional de La Plata (UNLP). S. López received a scholarship from the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina. M. Saparrat is a researcher for the CONICET, Argentina. P. A. Balatti is a researcher for Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CICBA), Argentina. This research was partially supported by grants from CICBA, UNLP (Proyecto del Programa de Incentivos Secretaría de Políticas Universitarias, Ministerio de Educación de la Nación). This work was supported by the Agencia Nacional, PICT2012-2760.

Literature Cited

- Allegrucci, N., Cabello, M. N., and Arambarri, A. M. 2009. Diversity of saprotrophic anamorphic ascomycetes from native forests in Argentina: An updated review. *Darwiniana* 47:108-124.
- Bailey, D. L. 1947. The development through breeding of greenhouse tomato varieties resistant to leaf mold. *Ont. Hortic. Exp. Stn. Rep.* 1945-1946:56-60.
- Bensch, K., Groenewald, J. Z., Dijksterhuis, J., Starink-Willems, M., Andersen, B., Summerell, B. A., Shin, H. D., Dugan, F. M., Schreers, H. J., Braun, U., and Crous, P. W. 2010. Species and ecological diversity within the *Cladosporium cladosporioides* complex (*Davidiellaceae*, *Capnodiales*). *Stud. Mycol.* 67:1-94.
- Blancard, D. 1992. Enfermedades del tomate observar, identificar y luchar. Mundipress, Madrid.
- Bolton, M. D., Van Esse, H. P., Vossen, J. H., de Jonge, R., Stergiopoulos, I., Stulemeijer, I. J. E., Van den Berg, G. C. M., Borras-Hidalgo, O., Dekker, H. L., de Koster, C. G., De Wit, P. J. G. M., Joosten, M. H. A. J., and Thomma, B. P. H. J. 2008. The novel *Cladosporium fulvum* lysin motif effector Ecp6 is a virulence factor with orthologues in other fungal species. *Mol. Microbiol.* 69:119-136.
- Boukema, I. W. 1981. Races of *Cladosporium fulvum* Cooke (*Fulvia fulva*) and genes for resistance in the tomato (*Lycopersicon Mill.*). Genetics and breeding of tomato. Pages 287-292 in: Proc. Meet. Eucarpia Tomato Work. Group, Avignon, France. J. Philouze, ed.
- Braun, U., Crous, P. W., Dugan, F. M., Groenewald, J. Z., and de Hoog, G. S. 2003. Phylogeny and taxonomy of *Cladosporium*-like hyphomycetes, including *Davidiella* gen. nov., the teleomorph of *Cladosporium* s. str. *Mycol. Prog.* 2:3-18.
- Butler, E. J., and Jones, S. G. 1949. Plant Pathology. MacMillan and Co., London.
- Cleyet-Marel, J. C. 1987. Dynamics des Populations de *Rhizobium* et de *Bradyrhizobium* dans le Sol et la Rhizosphère. Thesis, University Claude Bernard, Lyon, France.
- Cooke, M. C. 1883. New American fungi. *Grevillea* 12:22-33.
- Crous, P. W. 2009. Taxonomy and phylogeny of the genus *Mycosphaerella* and its anamorphs. *Fungal Divers.* 38:1-24.
- Crous, P. W., Braun, U., Schubert, K., and Groenewald, J. Z. 2007. Delimiting *Cladosporium* from morphologically similar genera. *Stud. Mycol.* 58:33-56.
- Crous, P. W., Schöch, C. L., Hyde, K. D., Wood, A. R., Gueidan, C., de Hoog, G. S., and Groenewald, J. Z. 2009. Phylogenetic lineages in the *Capnodiales*. *Stud. Mycol.* 64:17-47.
- Crous, P. W., Schreers, H. J., Groenewald, J. Z., Braun, U., and Schubert, K. 2006. *Metulocladosporiella* gen. nov. for the causal organism of *Cladosporium* speckle disease of banana. *Mycol. Res.* 110:264-275.
- De Wit, P. J. G. M. 1977. A light and scanning electron microscopic study of infection of tomato plants by virulent and avirulent races of *Cladosporium fulvum*. *Neth. J. Plant Pathol.* 83:109-122.
- De Wit, P. J. G. M. 1992. Molecular characterization of gene-for-gene systems in plant-fungus interactions and the application of avirulence genes in control of plant pathogens. *Annu. Rev. Phytopathol.* 30:391-418.
- Dixon, M. S., Hatzixanthis, K., Jones, D. A., Harrison, K., and Jones, J. D. G. 1998. The tomato *Cf-5* disease resistance gene and six homologs show pronounced allelic variation in leucine rich repeat copy number. *Plant Cell* 10:1915-1925.
- Dixon, M. S., Jones, D. A., Keddie, J. S., Thomas, C. M., Harrison, K., and Jones, J. D. G. 1996. The tomato *Cf-2* disease resistance locus comprises two functional genes encoding leucine rich repeat proteins. *Cell* 84:451-459.
- Enya, J., Ikeda, K., Takeuchi, T., Horikoshi, N., Higashi, T., Sakai, T., Iida, Y., Nishi, K., and Kubota, M. 2009. The first occurrence of leaf mold of tomato caused by races 4.9 and 4.9.11 of *Passalora fulva* (syn. *Fulvia fulva*) in Japan. *J. Gen. Plant Pathol.* 75:76-79.
- Flor, H. H. 1956. The complementary genetic systems in flax and flax rust. *Adv. Genet.* 8:29-54.
- Flor, H. H. 1971. Current status of the gene-for-gene concept. *Annu. Rev. Phytopathol.* 9:275-296.
- Galván, M. Z., Bornet, B., Balatti, P., and Branchard, M. 2003. Inter simple sequence repeat (ISSR) markers as a tool for the assessment of both genetic diversity and gene pool origin of common bean (*Phaseolus vulgaris*). *Euphytica* 132:297-301.
- Hörger, A. C., Ilyas, M., Wolfgang, S., Tellier, A., van der Hoorn, R. A. L., and Rose, L. E. 2012. Balancing selection at the tomato RCR3 guardee gene family maintains variation in strength of pathogen defense. *PLoS Genet.* 8: e1002813.

- Iida, Y., Iwadate, Y., Kubota, M., and Terami, F. 2010. Occurrence of a new race 2.9 of leaf mold of tomato in Japan. *J. Gen. Plant Pathol.* 76:84-86.
- Jones, D. A., Thomas, C. M., Hammond-Kosack, K. E., Balint-Kurti, P. J., and Jones, J. D. G. 1994. Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science* 266:789-793.
- Joosten, M. H. A. J., Cozijnsen, T. J., and De Wit, P. J. G. M. 1994. Host resistance to a fungal tomato pathogen lost by a single base-pair change in an avirulence gene. *Nature* 367:384-386.
- Joosten, M. H. A. J., and De Wit, P. J. G. M. 1999. The tomato–*Cladosporium fulvum* interaction: A versatile experimental system to study plant-pathogen interactions. *Annu. Rev. Phytopathol.* 37:335-367.
- Joosten, M. H. A. J., Vogelsang, R., Cozijnsen, T. J., Verberne, M. C., and De Wit, P. J. G. M. 1997. The biotrophic fungus *Cladosporium fulvum* circumvents *Cf-4*-mediated resistance by producing unstable AVR4 elicitors. *Plant Cell* 9: 367-379.
- Kooistra, E. 1964. Recent experiences of breeding leaf mould resistant tomatoes. *Euphytica* 13:108-109.
- Laugé, R., Goodwin, P. H., De Wit, P. J. G. M., and Joosten, M. H. 2000. Specific HR-associated recognition of secreted proteins from *Cladosporium fulvum* occurs in both host and non-host plants. *Plant J.* 23:735-745.
- Lazarovits, G., and Higgins, V. J. 1976. Histological comparison of *Cladosporium fulvum* race 1 on immune, resistant, and susceptible tomato varieties. *Can. J. Bot.* 54:224-234.
- Luderer, R., Takken, F. L. W., De Wit, P. J. G. M., and Joosten, M. H. A. J. 2002. *Cladosporium fulvum* overcomes *Cf-2*-mediated resistance by producing truncated AVR2 elicitor proteins. *Mol. Microbiol.* 45:875-884.
- Parniske, M., Hammond-Kosack, K. E., Golstein, C., Thomas, C. M., Jones, D. A., Harrison, K., Wulff, B. B., and Jones, J. D. 1997. Novel disease resistance specificities result from sequence exchange between tandemly repeated genes at the *Cf-4/9* locus of tomato. *Cell* 91:821-832.
- Peteira, B., Bernal Cabrera, A., Martínez, B., and Miranda, I. 2011. Caracterización molecular de aislamientos de *Cladosporium fulvum* Cooke provenientes de tomate en condiciones de cultivo protegido. *Rev. Prot. Veg.* 26:5-15.
- Rollan, C., Protto, V., Medina, R., Lopez, S. Y. M., Vera Bahima, J., Ronco, L., Saparrat, M., and Balatti, P. A. 2013. Identification of Races 0 and 2 of *Cladosporium fulvum* (syn. *Passalora fulva*) on tomato in the Cinturón Hortícola de La Plata, Argentina. *Plant Dis.* 97:992.1.
- Sambrook, J., Fritsch, E. F., and Maniatis, T. 1989. Molecular Cloning: A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanger, F., Nicklen, S., and Coulson, A. R. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74:5463-5467.
- Satou, M., Shinozaki, T., Nishi, K., and Kubota, M. 2005. Leaf mold of tomato caused by races 4 and 4.11 of *Passalora fulva* in Japan. *J. Gen. Plant Pathol.* 71:436-437.
- Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W., and Fungal Barcoding Consortium. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proc. Natl. Acad. Sci. USA* 109:6241-6246.
- Schubert, K., Groenewald, J. Z., Braun, U., Dijksterhuis, J., Starink, M., Hill, C. F., Zalar, P., de Hoog, G. S., and Crous, P. W. 2007. Biodiversity in the *Cladosporium herbarum* complex (*Davallaceae*, *Capnodiales*), with standardisation of methods for *Cladosporium* taxonomy and diagnostics. *Stud. Mycol.* 58:105-156.
- Stenglein, S. A., and Balatti, P. A. 2006. Genetic diversity of *Phaeoisariopsis griseola* in Argentina as revealed by pathogenic and molecular markers. *Physiol. Mol. Plant Pathol.* 68:158-167.
- Stergiopoulos, I., De Kock, M. J. D., Lindhout, P., and De Wit, P. J. G. M. 2007. Allelic variation in the effector genes of the tomato pathogen *Cladosporium fulvum* reveals different modes of adaptive evolution. *Mol. Plant-Microbe Interact.* 20:1271-1283.
- Takken, F. L. W., Schipper, D., Nijkamp, H. J. J., and Hille, J. 1998. Identification and *Ds*-tagged isolation of a new gene at the *Cf-4* locus of tomato involved in disease resistance to *Cladosporium fulvum* race 5. *Plant J.* 14:401-411.
- Theerakulpisut, P., Kanawapee, N., Maensiri, D., Bunnag, S., and Chantanathai, P. 2008. Development of species-specific SCAR markers for identification of three medicinal species of *Phyllanthus*. *J. Syst. Evol.* 46:614-621.
- Thomas, C. M., Jones, D. A., Parniske, M., Harrison, K., Balint-Kurti, P. J., Hatzixanthis, K., and Jones, J. D. 1997. Characterization of the tomato *Cf-4* gene for resistance to *Cladosporium fulvum* identifies sequences that determine recognitional specificity in *Cf-4* and *Cf-9*. *Plant Cell* 9:2209-2224.
- Thomma, B. P. H. J., Van Esse, H. P., Crous, P. W., and De Wit, P. J. G. M. 2005. *Cladosporium fulvum* (syn. *Passalora fulva*), a highly specialized plant pathogen as a model for functional study. *Mol. Plant Pathol.* 6:379-393.
- Van den Ackerveken, G. F. J. M., Van Kan, J. A. L., and De Wit, P. J. G. M. 1992. Molecular analysis of the avirulence gene *Avr9* of the fungal tomato pathogen *Cladosporium fulvum* fully supports the gene-for-gene hypothesis. *Plant J.* 2:359-366.
- Van den Ackerveken, G. F. J. M., Van Kan, J. A. L., Joosten, M. H. A. J., Muysers, J. M., Verbakel, H. M., and De Wit, P. J. G. M. 1993. Characterization of two putative pathogenicity genes of the fungal tomato pathogen *Cladosporium fulvum*. *Mol. Plant-Microbe Interact.* 6:210-215.
- van der Hoorn, R. A. L., Laurent, F., Roth, R., and De Wit, P. J. G. M. 2000. Agroinfiltration is a versatile tool that facilitates comparative analysis of *Avr9/Cf-9*-induced and *Avr4/Cf-4*-induced necrosis. *Mol. Plant-Microbe Interact.* 13:439-446.
- Van Esse, H. P., Thomma, B. P., Van't Kloostewi, J. W., and De Wit, P. J. G. M. 2006. Affinity-tags are removed from *Cladosporium fulvum* effector proteins expressed in the tomato leaf apoplast. *J. Exp. Bot.* 57:599-608.
- Van Kan, J. A. L., Van den Ackerveken, G. F. J. M., and de Wit, P. J. G. M. 1991. Cloning and characterization of cDNA of avirulence gene *avr9* of the fungal pathogen *Cladosporium fulvum*, causal agent of tomato leaf mold. *Mol. Plant-Microbe Interact.* 4:52-59.
- Westerink, N., Brandwagt, B. F., De Wit, P. J. G. M., and Joosten, M. H. A. J. 2004. *Cladosporium fulvum* circumvents the second functional resistance gene homologue at the *Cf-4* locus (*Hcr9-4E*) by secretion of a stable *Avr4E* isoform. *Mol. Microbiol.* 54:533-545.
- White, T. J., Bruns, T., Lee, S., and Taylor, J. W. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. Pages 315-322 in: PCR Protocols: A Guide to Methods and Applications. M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White, eds. Academic Press Inc., New York.
- Yamada, K., and Abiko, K. 2002. Race composition of *Fulvia fulva* in Japan during 1997-1998 (in Japanese with English summary). *Jpn. J. Phytopathol.* 68:36-38.
- Yan, L., Zhang, C., Ding, L., and Ma, Z. 2008. Development of a real-time PCR assay for the detection of *Cladosporium fulvum* in tomato leaves. *J. Appl. Microbiol.* 104:1417-1424.
- Zalar, P., de Hoog, G. S., Schroers, H.-J., Crous, P. W., Groenewald, J. Z., and Gunde-Cimerman, N. 2007. *Stud. Mycol.* 58:157-183.