

ORIGINAL ARTICLE / ARTÍCULO ORIGINAL

HELMINTHIC ASSEMBLAGES OF *SERIOLELLA POROSA* GUICHENOT 1848 (PISCES:
CENTROLOPHIDAE) FROM SAN MATÍAS GULF (ARGENTINA)

ENSAMBLES HELMÍNTICOS DE *SERIOLELLA POROSA* GUICHENOT 1848 (PISCES:
CENTROLOPHIDAE) DEL GOLFO SAN MATÍAS (ARGENTINA)

Silvia Guagliardo¹, Carla Schwerdt¹, Noelia Galeano¹, Raúl González² & Rubén D. Tanzola¹

¹Laboratorio de Patología de Organismos Acuáticos- Depto. Biología, Bioquímica y Farmacia, Universidad Nacional del Sur, San Juan 670 (8000) Bahía Blanca, Argentina. ²Escuela Superior de Ciencias Marinas, Universidad Nacional del Comahue, sede Atlántica, San Martín 247, San Antonio Oeste, Río Negro, Argentina.
E-mail: rtanzola@uns.edu.ar

Suggested citation Guagliardo S, Schwerdt C, Galeano N, González R & Tanzola, RD. 2014. Helminthic assemblages of *Seriolella porosa* Guichenot 1848 (Pisces: Centrolophidae) from San Matías gulf (Argentina). *Neotropical Helminthology*, vol. 8, n°2, jul-dec, pp. 291-303.

Abstract

No previous studies were carried out on aspects of the helminth assemblages of *Seriolella porosa*. The present study aims to analyze the metazoan parasite of *S. porosa* and establish the relationship between the parasitic variations and both fish size and sex. A total of 101 specimens of *S. porosa* from San Matías Gulf, Argentina were examined between September and October 2006. Thirteen species were reported with a total of 3366 parasites. *Lecithocladium cristatum* was the most prevalent, dominant and abundant species. The average species richness was 3.2 ± 1.20 . The evenness showed an inverse correlation with the total number of parasites and did not reveal differences between host sexes. None of the parasite species showed interspecific associations with each other. The average richness of the infracommunity was significantly less than the component community. The helminth community of the silver warehou shows little diversity with many satellite species and only two core species (*Anisakis* sp. and *L. cristatum*). Further studies, using parasites as stock indicators, would indicate whether the San Matías population represents a reproductive aggregation that penetrates the gulf seasonally in winter or whether it is a permanent stock isolated from the continental shelf.

Keywords: Fish - helminthic assemblages - parasite ecology - *Seriolella porosa*.

Resumen

Hasta el momento no se llevaron a cabo estudios sobre aspectos de los ensambles de helmintos de *Seriolella porosa*. El presente estudio se propone analizar el ensamble de metazoos parásitos de *S. porosa* y establecer las interrelaciones entre variaciones en la estructura parasitaria y el sexo y tamaño del hospedador. Se examinó un total de 101 especímenes de *S. porosa* capturados entre septiembre y octubre de 2006 en el Golfo San Matías, Argentina. Trece especies y un total de 3366 parásitos fueron registrados. La especie más prevalente, dominante y abundante fue *Lecithocladium cristatum*. La riqueza específica fue de 3.2 ± 1.20 . La equitatividad mostró una correlación inversa con el número total de parásitos y no reveló diferencias entre sexos. Ninguna de las especies parásitas mostró asociación interespecífica con otras especies del ensamble. La riqueza promedio de la infracomunidad fue significativamente menor que aquella del componente comunitario. La comunidad de helmintos del savorín mostró poca diversidad con varias especies satélites y solo dos especies centrales (core) (*Anisakis* sp. and *L. cristatum*). El presente podría constituir la base para identificar stocks poblacionales, empleando parásitos, para demostrar si la población del Golfo San Matías es un conjunto reproductivo que penetra estacionalmente al Golfo en invierno, o si se trata de un stock propio del Golfo y separado física y reproductivamente de los savorines de la Plataforma Continental Argentina.

Palabras clave: Ecología parasitaria, ensambles helmínticos, peces, *Seriolella porosa*,

INTRODUCTION

Knowledge of helminth fish parasite communities from commercial fisheries reveals basic information as well as applicable data for studies on population stocks (Timi, 2007; Braicovich & Timi, 2008), the health of ecosystems (Palm & Rückert, 2009) and the sanitary state of the fish (Valero *et al.*, 2006; Jakob & Palm, 2006).

In view of recent studies, the parasitic fauna of fish appears to be governed by the trophic habits and availability of the hosts, as well as the range of depth and migration processes (Rhode, 1984; Kennedy *et al.*, 1986; Poulin, 1995; Klimpel *et al.*, 2003, 2006a, b). This implies that parasites are important indicators of the habitat and trophic status of fish in marine ecosystems (Klimpel *et al.*, 2003). Many studies consider that the host size is a factor that influences the formation of parasitic assemblages (Thoney, 1993; Sardella *et al.*, 1995; Balboa & George-Nascimento, 1998), being a more influential parameter than host sex (Luque *et al.*, 1996; Knoff *et al.*, 1997; González & Acuña, 2000;

Ramos Alves & Luque, 2001; Iannacone & Alvarino, 2008). *Seriolella porosa* Guichenot 1848 (silver warehou) is an important fish resource of the Argentine Sea (Sánchez & Bezzi, 2004). Its distribution extends from 35° to 55° S (Garciaarena & Perrotta, 2002). Commercial landings of this resource underwent a growing trend over the last 20 years as "by catch" in the hake fishery (Cousseau & Perrota, 1998). It was described by Perier & Di Giacomo (2002) as "a seasonal fish resource which uses the area as a spawning ground and nursery" in the San Matías Gulf fishery. Up to the present only two studies have been made on the parasitological status of this fish in the South Atlantic Ocean (Guagliardo *et al.*, 2009, 2010).

It would be expected that *S. porosa* carries several parasite species to upper levels of the food chain because it occupies an intermediate level in the food web. Since there are no records of aspects of the helminthic assemblages of the silver warehou, the aims of this study were to analyze them at the component and infracommunity levels and to evaluate the relationship between the parasitic burden and fish size and sex.

MATERIALS AND METHODS

A total of 101 specimens of *S. porosa* (68 males, 32 females and 1 undetermined) were examined between September and October 2006. The area of capture was the northern sector of San Matías Gulf (41°00'S, 64°00'W), Argentina. The fish were caught by commercial fleet trawlers from San Antonio Oeste port. Host size (cm), weight (g) and sex were determined. The gonad/somatic index was estimated as $GSI = (\text{gonad weight} / \text{total weight}) * 100$. Parasites were extracted from the gills, body cavity and digestive lumen. They were subsequently relaxed in hot saline solution, fixed in 5% formalin and stored in 70% alcohol. Part of the helminth sample was stained with Langeron's hydrochloric carmine cleared in clove oil and mounted in Canada balsam. Nematodes and acanthocephalans were made transparent in hot alcohol-glycerine. Parasite prevalence, abundance and the intensity of infection were calculated following Bush *et al.* (1997). Specimens of digenean trematodes were deposited in the collection of the Museum of La Plata (Argentina) under the following numbers of collection: *Lecithocladium cristatum* (Voucher MLP 5935), *Elytrophalloides oatesi* (Voucher MLP 5936), *Gonocerca* *cf.* *phycidis* (Voucher MLP 7937) & *Aponurus laguncula* (Voucher MLP 5938).

The community structure was analyzed at the component community and infracommunity levels. Species with prevalences $\geq 10\%$ were considered as component species in the infracommunities (Bush *et al.*, 1990).

The variance/mean ratio (coefficient of dispersion) was used to determine the distribution patterns at the level of infrapopulations. In order to quantify the influence of parasite species in the community, the specific importance ($IE = (\text{relative prevalence} + \text{relative abundance}) * 100$) was estimated according to Bursey *et al.* (2001). At the infracommunity level, the richness, mean diversity and evenness when richness was higher than four species (Brillouin based evenness index) were calculated (Mouillot *et al.*,

2005). The qualitative association between pairs of parasite species was estimated by using the correlation coefficient (Combes, 1983). At the component community level, the relative dominance (number of specimens of one species in respect of the total number of specimens of all species in the infracommunity) of each parasite species was calculated according to Morales & Pino de Morales (1987). The dominance frequencies and the co-dominances were also calculated. The ecto/endoparasite ratio was determined as the number of ectoparasite species/number of endoparasite species.

The species richness, the total number of parasites, the Simpson index of diversity (Pielou, 1975; Krebs, 1989), the Shannon-Wiener index of diversity and evenness were calculated after Magurran (1988). The correlations between the total number of parasites, richness, diversity and evenness in relation to host weight and total length were tested using Spearman's rank correlation coefficient (Siegel & Castellan, 1995). Evenness was also correlated with the total number of parasites. This latter parameter was compared with host sex using the nonparametric Mann-Whitney U test (Siegel & Castellan, 1995). The strategy of colonization, allogenic or autogenic species, was assessed according to Bush *et al.* (2001).

RESULTS

The mean total body length and weight of fish were 39.60 ± 4.68 and 514.55 ± 264.25 respectively. The mean GSI was 0.052 (0.009 – 0.230) in females. Most of the females (75%) had not reached the first sexual maturity length (< 45 cm). A total of 3366 helminths, mostly endoparasites, were recorded and a total of thirteen taxa were identified. The ectoparasite versus endoparasite relationship was low ($R = 0.18$). The helminth assemblage was highly dominated by digenean (2309 or 68.6% of the specimens). All component community species (prevalence $\geq 10\%$) had an aggregated pattern of distribution. Table 1 shows that *Lecithocladium cristatum* was not only dominant but also the

Table 1. Prevalence, abundance, mean intensity and specific importance of the parasitic helminths of *Seriotelella porosa* from the San Matías Gulf (Argentina). 95% confidence interval. SD: standard deviation.

Species	Developmental stage	Site	Prevalence (%) ± CI	Abundance ± SD	Mean intensity ± SD	Specific importance	Frequency of Dominance single	Frequency of Dominance 3 species	Previous records from South Atlantic Ocean (*)
Nematoda									
<i>Anisakis</i> sp.	LIII	Body cavity	75.25 ± 8.5	7.22 ± 10.96	9.59 ± 11.71	797.25	25	1	1,2,3,4,5,6,7,8,9,10
			1.98 ± 8.68	0.02 ± 0.14	1 ± 0	21.98	0	0	5,6,10,11,12,13
<i>Pseudoterranova</i> sp.	LIII	Body cavity	2.97 ± 10.57	0.03 ± 0.17	1 ± 0	32.97	1	0	1,2,3,4,5,6,7,8,10,14,15,16
			2.97 ± 10.57	0.05 ± 0.30	1.67 ± 0.58	52.97	0	0	---
<i>Hysterothylacium</i> sp.	LIV	Body cavity	2.97 ± 10.57	0.03 ± 0.17	1 ± 0	32.97	1	0	1,2,3,4,5,6,7,8,10,14,15,16
			2.97 ± 10.57	0.05 ± 0.30	1.67 ± 0.58	52.97	0	0	---
Acanthocephala									
<i>Philometra</i> cf. <i>seriotelellae</i>	Adult	Ovary	2.97 ± 10.57	0.05 ± 0.30	1.67 ± 0.58	52.97	0	0	---
			2.97 ± 10.57	0.03 ± 0.17	18 ± 0	32.97	0	0	1,7,8,10,17,18
<i>Corynosoma cetaceum</i>	Juvenile	Body cavity	2.97 ± 10.57	0.03 ± 0.17	18 ± 0	32.97	0	0	1,7,8,10,17,18
			1.98 ± 8.68	0.04 ± 0.31	2 ± 1.41	41.98	0	0	1,3,4,6,7,8,13,17,18,20,22,23
Cestoda									
<i>Corynosoma australe</i>	Juvenile	Body cavity	1.98 ± 8.68	0.04 ± 0.31	2 ± 1.41	41.98	0	0	1,3,4,6,7,8,13,17,18,20,22,23
<i>Nybelinia</i> sp.	Plerocercus	Body cavity	0.99 ± 1.95	0.01 ± 0.10	1	10.99	0	0	3,6,7,8

Table 1. Continuation.

Table 1. Continuation.

Species	Developmental stage	Site	Prevalence (%) ± CI	Abundance ± SD	Mean intensity ± SD	Specific importance	Frequency of Dominance single	Frequency of Dominance 3 species	Previous records from South Atlantic Ocean (*)
Monogenea									
<i>Neorubaea seriolellae</i>	Adult	Gills	63.4 ±9.49	2.65 ±3.98	4.19 ±4.32	328.4	6	2	---
<i>Eurysorchis australe</i>	Adult	Gills	28.7 ±8.91	0.42 ±0.80	1.44 ±0.87	70.7	2	0	---
Digenea									
<i>Lecithocladium cristatum</i>	Adult	Stomach	80.19 ±7.85	17.90 ±24.63	22.32 ±25.74	1870.2	56	2	7.24
<i>Aponurus laguncula</i>	Adult	Stomach	19.8 ±7.85	0.40 ±1.08	2 ±1.65	59.8	0	1	7,10,24,25,26
<i>Ellytrophaloides oatesi</i>	Adult	Stomach	31.68 ±9.16	4.55 ±13.88	14.38 ±21.81	486.68	8	0	7,11,12,15, 24,25,27,28,29,30
<i>Gonocerca. cf. phycidis</i>	Adult	Stomach	0 ±1.95	0.01 ±0.10	1	0.09	0	0	15,24,28,29,31

(*) References: 1- Guagliardo (2003), 2- Tanzola & Guagliardo (2004), 3- Timi *et al.* (2005), 4- Timi & Poulin (2003), 5- Sardella *et al.* (1998), 6- Sardella & Timi (2004), 7- Braicovich (2008), 8- Braicovich & Timi (2008), 9- Guagliardo *et al.* (2009), 10- Timi *et al.* (2008), 11- Gaevsckaya *et al.* (1990), 12- Sardella & Timi (1996), 13- Cremona & Sardella (1997), 14- Navone *et al.* (1998), 15- Mackenzie & Longshaw (1995), 16- Incorvaia & Diaz de Astarloa (1998), 17- Timi & Lanfranchi (2009), 18- Sardella *et al.* (2005), 19- Tanzola & Guagliardo (2000), 20- Timi *et al.* (2001), 21- Zdzitowiecki (1989), 22- Tanzola *et al.* (2005), 24- Guagliardo *et al.* (2010), 25- Szidat (1950), 26- Szidat (1961), 27- Brickle *et al.* (2006), 28- Suriano & Sutton (1981), 29- Szidat (1955), 30- Szidat & Graefe (1968), 31- Laskowski & Zdzitowiecki (2005).

Table 2. Community descriptors of *Seriolella porosa* in San Matías Gulf (Argentina).

Component community		Infracommunity	
Total number of species	13	Mean species per host (\pm SD)	3.12 \pm 1.20
Total number of helminths	3366	Mean helminth per host (\pm SD)	33.3 \pm 27.6
Diversity indices		Mean diversity (Brillouin) (\pm SD, rank)	
- Shannon-Wiener	1.28	0.23 \pm 0.13 [0.00-0.56]	
- Simpson	0.64		
Evenness (Shannon-Wiener)	0.50	Mean evenness (Brillouin) (\pm SD, rank)	
		0.40 \pm 0.19 [0.127-0.96]	
Dominance (Berger-Parker)		Percentage of infracommunities dominated by the dominant species	
0.53		55.45 %	

Table 3. Relationships (Spearman correlation) between infracommunities descriptors and the size and weight of *Seriolella porosa* from Golfo San Matías (Argentina). r_s : Spearman's coefficient; critical p-value = 0.05.

Infracommunity descriptor	Size		Total weight	
	r_s	p	r_s	p
Total number of parasites	0.31	0.002**	0.18	0.062
Richness	0.23	0.022*	0.18	0.074
Diversity (Brillouin)	0.16	0.098	0.35	0.00**
Evenness (Brillouin)	-0.13	0.21	0.04	0.73

most abundant and prevalent species and it was also of greater relative influence in the community (IE= 1870.2).

Only one fish was free of parasites, five fishes (4.95 %) harboured a single species while 95 hosts (94.05 %) showed multiple infestations, with a richness range of two to six parasite species. The average infracommunity species richness was 3.2 ± 1.20 . Co-dominances were not frequent. All parasite species were autogenic in their colonization strategy, completing their life-cycles in aquatic hosts within the same ecosystem.

Both the infracommunity and the component community descriptors are shown in Table 2.

The richness and total number of parasites showed significant correlation with the size of the hosts, while only diversity was correlated with body weight (Table 3). The evenness showed an inverse correlation with the total number of parasites.

No significant differences were observed between males and females ($U= 1057$, $p= 0.82$) in the total parasite burden and none of the species showed any strong interspecific associations.

DISCUSSION

Two species of *Seriolella* reported from the Southwestern Atlantic are: *S. porosa* Guichenot 1848 and *S. caerulea* Guichenot 1848. Although there is a latitudinal overlap between these two species, they are rarely found together as they live at different levels in the water column (Cousseau & Díaz de Astarloa, 1993). In the San Matías Gulf, *S. porosa* is the species that has undergone rising and sustained catches, so it is an important fishery resource in its distribution area (Garciaarena & Perrota, 2002; Perier & Di Giacomo, 2002).

This study is the first comprehensive approach on silver warehou helminth assemblages. Our

records barely coincide with any previous studies of *Seriolella* spp.; in *S. violacea* Guichenot, 1848 from the Peruvian coast there were no species in common with the present survey. The monogenean *Paraeurysorchis sarmientoi* (Tantalean, 1974) Tantalean, Martinez & Escalante, 1985 and the cestode *Neobothriocephalus aspinosus* Mateo & Bullock, 1966 are the only two species reported from this South Pacific host (Iannacone 2003). Additionally, *S. porosa* showed no similarities with the endoparasites of *S. brama* (Günther, 1860) from New Zealand although the monogenean species were the same (Rhode *et al.*, 1980). None of the digeneans in this work were recorded for *S. punctata* in Tasmania (Bray & Cribbs, 2003).

According to Cortés & Muñoz (2008) infracommunity studies constitute the first step to any parasitological analysis within or between species of hosts through their parasitological descriptors. In the present study, all component species of *S. porosa* follow the typical aggregated pattern of distribution. As stated by several authors, aggregation is common behavior in parasitic life (Crofton, 1971; Esch & Fernández, 1993). A wide range of biological processes generates this type of distribution (Morales & Pino de Morales, 1987). According to Janovy (1992) the structures of parasite populations and their frequency distributions would be the result of mostly ecological processes. The prey of the fishes could be considered as carriers of these infracommunities, in agreement with the concept of "instant infracommunities" (Bush *et al.*, 1993), perhaps transmitting many parasites with the intake of few intermediate hosts and vice versa, thus determining the distribution pattern of parasites.

In our study ectoparasites species were scarce. This fact agrees with the parasitism present in teleosts from the South Atlantic coast (Sardella & Timi, 1996; Tanzola *et al.*, 1997; Tanzola & Guagliardo, 2000; Sardella & Timi, 2004; Timi *et al.*, 2005; Timi & Lanfranchi, 2009). In the north Bonaerense region of the Argentine Sea, the dominance of larval parasites, in terms of

parasite prevalence and abundance, seems to be the rule (Timi & Lanfranchi, 2009). The fact that we have reported a dominance of adult stages at higher latitude in our study, probably reflects a particular biogeographic region where the ecology component governs the parasitic assemblages of the silver warehou in a different way. It was noted that the mean richness of the infracommunities was significantly less than the total number of species from the component community. This situation agrees with the idea that the infracommunities rarely include all species present at the component community level (Poulin, 1998).

Seven species, with prevalences lower than 10% were not included in the component community analysis (*Pseudoterranova* sp., *Hysterothylacium*, sp., *Philometra* cf. *seriolellae*, *Corynosoma australe*, *C. cetaceum*, *Nybelinia* sp. and *Gonocerca* cf. *phycidis*).

The average richness of the infracommunities (3.12 ± 1.20) represents a common value in other host/parasite systems, ranging from two to five species per host (Cordeiro & Luque, 2004; Mouillot *et al.*, 2005; Jacob & Palm, 2006; Rodríguez-González & Vidal-Martínez, 2008; Timi & Lanfranchi, 2009). In comparison, it should be noted that the average richness of *S. porosa* was slightly less than the value recorded for *S. violacea* in Chilean waters (4.60 ± 1.54) (Mouillot *et al.* 2005). In this study, the most recruited species with prevalence > 50% were *L. cristatum*, *N. seriolellae* and *Anisakis* sp.

The richness was positively correlated with the size of the hosts. In larger fish more food is consumed, so consequently an increase in the exposure to the infective stages of the parasites is expected (Poulin, 1995).

When both the Shannon-Wiener and Simpson indexes were compared in respect of the Brillouin values, the mean values for the infracommunities were lower than the component community.

Until the present time, the most common mean value of helminth diversity in marine fish

infracommunities is less than two. Based on previous records (Ramos Alves *et al.*, 2002; Cordeiro & Luque, 2004; Sardella & Timi, 2004; Timi & Lanfranchi, 2009; Palm & Rückert, 2009) the parasite assemblage of silver warehou shows little diversity, with many satellite species and only two central ones (*Anisakis* sp. and *L. cristatum*).

The evenness is a measure of the unequal representation of species in a hypothetical community in which all species are equally common (Krebs 1989). In this work, the evenness was 0.50, while the mean value of evenness in the infracommunities was lower (0.40) showing an unequal distribution of the specific abundance. These values can perhaps be explained by the high dominance of *L. cristatum*. Evenness was not correlated with either size or weight of the host but showed a highly significant inverse correlation with the total number of parasites. We have noted that the greatest number of helminths were *L. cristatum* and *Anisakis* sp., both of them with the highest dominances.

None of the species showed any interspecific association with each other (Table IV). A common pattern in many host species is the absence of sexual differences in the prevalence and/or parasitic intensities (Ramos Alves & Luque, 2001; Knoff *et al.*, 1997; Luque *et al.*, 1996; Iannacone, 2003). Accordingly, no differences in the total number of parasites were reported here, discarding selective parasitism or differential trophic behavior between sexes.

In respect of the colonizing ability in the Argentine Sea of cestodes, nematodes and acanthocephalans recorded in the silver warehou, they may be considered as generalists, as they have previously been recorded in several hosts in the South Atlantic. It is likely that some infect *S. porosa* accidentally, especially those with low prevalences, due to the trophic behavior of this host (Forciniti & Pérez Macri, 1992; Sánchez & Prenski, 1996). However it should be noted that both, monogeneans and digeneans, have demonstrated selective colonization strategies in the Atlantic Ocean.

Previous studies on the use of parasites as biological tags of South Atlantic fish stocks have been able to discriminate populations of *Merluccius hubbsi* Marini, 1933 (Sardella & Timi, 2004), *Scomber japonicus* (Houttuyn, 1782) (Cremonte & Sardella, 1997); *Engraulis anchoita* Hubbs & Marini, 1935 (Timi, 2003), *Cynoscion guatucupa* Cuvier 1830 (Timi *et al.*, 2005), *Eleginops maclovinus* (Valenciennes, 1839) Dollo, 1904 (Brickle & MacKenzie, 2007) *Percophis brasiliensis* Quoy & Gaimard, 1824 (Braicovich & Timi, 2008) and *Pinguipes brasilianus* Cuvier, 1829 (Timi *et al.*, 2008). The only information about seasonal migration in the area of distribution of the silver warehou in the Argentine Sea is in the San Matías Gulf (Perier & Di Giacomo, 2002). Further studies, using parasites as stock indicators, might show whether the San Matías Gulf population represents a reproductive aggregation that enters the gulf in winter, looking for better environmental conditions or whether it is a permanent stock isolated from the continental shelf.

ACKNOWLEDGEMENTS

Authors are grateful to Mrs. Rosemary Scoffield for her assistance with the English language and Mr. Paul Osovnikar from the *Instituto de Biología Marina y Pesquera Ate. Storni*, province of Río Negro, Argentina, for his collaboration in obtaining the samples of *S. porosa*.

BIBLIOGRAPHIC REFERENCES

- Balboa, L & George-Nascimento, M. 1998. *Variaciones ontogenéticas y entre años en las infracomunidades de parásitos metazoos de dos peces marinos de Chile*. Revista Chilena de Historia Natural, vol. 71, pp. 27 - 37.
- Braicovich, PE, González, RA & Tanzola, RD. 2005. *First record of Corynosoma australe (Acanthocephala, Polymorphidae) parasitizing seahorse*

- Hippocampus sp. (Pisces, Syngnathidae) in Patagonia. *Acta Parasitologica*, vol. 50, pp.145 - 149.
- Braicovich, PE & Timi, JT. 2008. Parasites as biological tags for stock discrimination of the Brazilian flathead *Percophis brasiliensis* in the south-west Atlantic. *Journal of Fish Biology*, vol. 73, pp. 557 - 571.
- Braicovich, PE. 2008. *Diversidad y ecología parasitarias a de Percophys brasiliensis Quoid et Gaimard, 1824, (Osteichthyes: Percophidae) (Pez Palo) en el Mar Argentino*, PhD Thesis, Universidad Nacional de Mar del Plata, 214 pp.
- Bray, RA & Cribbs, TH. 2003. *New species of Opechona Looss, 1907 and Cephalolepidapedon Yamaguti, 1970 (Digenea: Lepocreadiidae) from fishes off northern Tasmania*. *Papers and Proceedings of the Royal Society of Tasman*, vol. 137, pp. 1 - 5.
- Brickle, P, Mackenzie, K & Pike, A. 2006. *Variations in the parasite fauna of the Patagonian toothfish (Dissostichus eleginoides Smitt, 1898), with length, season, and depth of habitat around the Falkland Islands*. *Journal of Parasitology*, vol. 92, pp. 282 - 291.
- Brickle, P & Mackenzie, K. 2007. *Parasites as biological tags for Eleginops maclovinus (Teleostei: Eleginopidae) around the Falkland Islands*. *Journal of Helminthology*, vol. 81, pp. 147-153.
- Burse, CR, Goldberg, SR & Parmelee, JR. 2001. *Gastrointestinal helminths of 51 species of anurans from Reserva Cuzco Amazónico, Peru*. *Comparative Parasitology*, vol. 68, pp. 21-35.
- Bush, AO, Aho, JM & Kennedy, CR. 1990. *Ecological versus phylogenetic determinants of helminth parasite community richness*. *Evolutionary Ecology*, vol. 4, pp. 1- 20.
- Bush, AO, Heard, RW & Overstreet, RM. 1993. *Intermediate hosts as source communities*. *Canadian Journal of Zoology*, vol. 71, pp. 1358 - 1363.
- Bush, AO, Lafferty, KD, Lotz, JM & Shostak, AW. 1997. *Parasitology meets ecology on its own terms: Margolis et al. revisited*. *Journal of Parasitology*, vol. 83, pp. 575 - 583.
- Bush, AO, Fernández, JC, Esch, GW & Seed, JR. 2001. *Parasitism: The diversity and ecology of animal parasites*. Cambridge University Press, U.K., 566 pp.
- Combes, C. 1983. *Aplicación a l'ecología parasitaire des indices d'association fondés sur le caractère présence-absence*. *Vie et Milieu*, vol. 33, pp. 203 - 212.
- Cordeiro, AS & Luque, JL. 2004. *Community ecology of the metazoan parasites of atlantic moonfish, Selene setapinnis (Osteichthyes: Carangidae) from the coastal zone of the state of Rio de Janeiro, Brazil*. *Brazilian Journal of Biology*, vol. 64, pp.399-406.
- Cortés, Y & Muñoz, G. 2008. *Infracomunidades de parásitos eumetazoos del bagre de mar Aphos porosus (Valenciennes, 1837) (Actinopterygii: Batrachoidiformes) en Chile central*. *Revista de Biología Marina y Oceanografía*, vol. 43, pp.255-263.
- Cousseau, MB & Díaz de Astarloa, JM. 1993. *El género Brevoortia en la costa Atlántica sudamericana*. *Frente Marítimo*, vol.14, pp.49-57.
- Cousseau, MB & Perrotta, RG. 1998. *Peces marinos de Argentina. Biología, distribución, pesca*. Publicaciones Especiales INIDEP, Mar del Plata, Argentina, 163 pp.
- Cremonte, F & Sardella, NH. 1997. *The parasite Fauna of Scomber japonicus Houttuyn, 1782 (Pisces: Scombridae) in two zones of the Argentine Sea*. *Fisheries Research*, vol. 31, pp. 1 - 9.
- Crofton, HD. 1971. *A quantitative approach to parasitism*. *Parasitology*, vol. 62, pp. 179 - 193.
- Esch, GW & Fernández, JC. 1993. *A functional biology of parasitism*. Chapman & Hall (Eds.), London, 337 pp.
- Forcitini, L & Pérez Macri, G. 1992. *Contribución al crecimiento del sistema digestivo y la alimentación del savorín (Seriollella porosa)*. *Frente Marítimo*, vol.12, pp.129 - 137.

- Gaevskaya, AV, Rodyuk, GN & Parukhin, AM. 1990. Peculiarities and formation of parasitofauna of the Patagonian toothfish *Dissostichus eleginoides*. *Biologiyal Morya*, vol.4, pp.23-28.
- Garciaarena, AD & Perrotta, RG. 2002. *Características biológicas y de la pesca del savorín Seriolella porosa (Pisces: Centrolophidae) del Mar Argentino*. *Revista del Instituto Nacional de Investigación y Desarrollo Pesquero. Informe Técnico*, vol.4, pp.1-19.
- González, T & Acuña, E. 2000. *Influence of host size and sex on the endohelminth infracommunities of the red rockfish *Sebastes capensis* off northern Chile*. *Journal of Parasitology*, vol. 86, pp.854 - 857.
- Guagliardo, SE. 2003. *Estudio sistemático, biológico y ecológico de los cestodes tripanorhincos (Eucestoda: Trypanorhyncha) parásitos de peces marinos en el área de Bahía Blanca*. PhD Thesis, Universidad Nacional de Sur, Bahía Blanca, Argentina, 298 pp.
- Guagliardo, SE, De Salvo, N, Schwerdt, CB, Galeano, NA & Tanzola, RD. 2009. *Anisákidos del savorín, Seriolella porosa (Pisces, Centrolophidae). Análisis de la interacción parásito-hospedador*. *BioScriba*, vol. 2, pp.106-114.
- Guagliardo, SE, Schwerdt, CB, Martorelli, SR, Galeano, NA & Tanzola, RD. 2010. *Digenean trematodes of Seriolella porosa (Pisces, Centrolophidae) in San Matías Gulf, Argentina*. *Acta Parasitologica* vol. 55, pp.29-38.
- Iannacone, J. 2003. *Tres metazoos parásitos de la cojinoba Seriolella violacea Guichenot (Pisces, Centrolophidae), Callao, Perú*. *Revista Brasileira de Zoología* vol. 20, pp.257-260.
- Iannacone, J & Alvarino, L. 2008. *Influencia del tamaño y sexo de *Peprilus medius* (Peters) (Stromateidae: Perciformes) capturados en chorrillos, Lima, Perú, sobre su comunidad parasitaria*. *Neotropical Helminthology*, vol. 2, pp.62-70.
- Incorvaia, IS & Díaz de Astarloa, JM. 1998. *Estudio preliminar de las larvas (Nematoda: Ascaridida) parásito de Paralichthys orbygnyanus (Valenciennes, 1839) y Paralichthys patagonicus (Pisces: Pleuronectiformes)*. *Boletín Chileno de Parasitología*, vol.53, pp.38-42.
- Jakob, E & Palm, H. 2006. *Parasites of commercially important fish species from the southern Java coast, Indonesia, including the distribution pattern of trypanorhynch cestodes*. *Verhandlungen der Gesellschaft für Ichthyologie*, vol. 5, pp. 165-191.
- Janovy, J, Clopton, RE & Percival, TJ. 1992. *The roles of ecological and evolutionary influences in providing structure to parasite species assemblages*. *Journal of Parasitology*, vol. 78, pp. 630 - 640.
- Kennedy, CR, Bush A & Aho, JM. 1986. *Patterns in helminth communities: why are birds and fish so different?* *Parasitology*, vol. 93, pp. 205 - 215.
- Klimpel, S, Seehagen, A & Palm, H. 2003. *Metazoan parasites and feeding behaviour of four small-sized fish species from the central North Sea*. *Parasitology Research* vol. 91, pp. 290 - 297.
- Klimpel, S, Palm, H, Busch, MW, Kellermanns, E & Ruckert, S. 2006a. *Fish parasites in the Arctic deep-sea: poor diversity in pelagic fish species vs. heavy parasite load in a demersal fish*. *Deep-Sea Research Pt I*, vol. 53, pp.1167 - 181.
- Klimpel, S, Ruckert, S, Piatkowski, U, Palm, H & Hanel, R. 2006b. *Diet and metazoan parasites of silver scabbard fish *Lepidopus caudatus* from the Great Meteor Seamount (North Atlantic)*. *Marine Ecology Progress Series*, vol.315, pp.249-257.
- Knoff, M, Luque, JL & Amato, JF. 1997. *Community ecology of the metazoan parasites of grey mullets, *Mugil platanus* (Osteichthyes: Mugilidae) from the littoral of the state of Rio de Janeiro, Brazil*. *Revista Brasileira de Biología*, vol. 57, pp. 441- 454.
- Krebs, CJ. 1989. *Ecological Methodology*. Harper Colins Publishers, New York, 654 pp.
- Laskowski, Z & Zdzitowiecki, K. 2005. *The*

- helminth fauna of some notothenioid fishes collected from the shelf of Argentine Islands, West Antarctica. Polish Polar Research, vol. 26, pp.315-324.
- Luque, JL, Amato, JF & Takemoto, RM. 1996. Comparative analysis of the communities of metazoan parasites of *Orthopristis ruber* and *Haemulon steindachneri* (Osteichthyes: Haemiuridae) from the southeastern Brazilian litoral: I. Structure and influence of the size and sex of hosts. Revista Brasileira de Biología, vol. 56, pp.279-292.
- Magurran, AE. 1988. *Ecological diversity and its measurement*. Princeton University Press, Princeton, USA, 167 pp.
- Mackenzie, K & Longshaw, M. 1995. Parasites of the hakes *Merluccius australis* and *M. hubbsi* in the waters around the Falk land Islands, southern Chile, and Argentina, with an assessment of their potential value as biological tags. Canadian Journal of Fisheries and Aquatic Sciences, vol. 52, pp. 213-224.
- Morales, G & Pino de Morales, LA. 1987. *Parasitología cuantitativa*. Fundación Fondo Editorial Acta Científica Venezolana, Caracas, Venezuela, 132 pp.
- Mouillot, D, George-Nascimento, M & Poulin, R. 2005. Richness, structure and functioning in metazoan parasite communities. Oikos, vol.109, pp.447-460.
- Navone, GT, Sardella, NH & Timi, JT. 1998. Occurrence of larvae and adults of *Hysterothylacium aduncum* (Rudolphi, 1802) (Nematoda: Anisakidae) in fishes and invertebrates in the Argentine Sea. Parasite, vol. 5, pp. 127-136.
- Palm, H & Rückert, S. 2009. A new approach to visualize ecosystem health by using parasites. Parasitology Research, vol. 105, pp. 539-553.
- Perier, M & Di Giácomo, E. 2002. *El savorín Seriolella porosa como un recurso estacional en el Golfo San Matías, República Argentina*. Revista del Instituto Nacional de Investigación y Desarrollo Pesquero, vol.15, 15-26.
- Pielou, EC. 1975. *Ecological diversity*. New York, J. Willey & Sons, 165 pp.
- Poulin, R. 1995. *Phylogeny, ecology, and the richness of parasite communities in vertebrates*. Ecological Monographs, vol. 65, pp. 283 - 302.
- Poulin, R. 1998. *Evolutionary ecology of Parasites*. Chapman & Hall, London, 212 pp.
- Ramos Alves, D & Luque, JL. 2001. Aspectos quantitativos das infrapopulações de metazoários parasitos de *Micropogonias furnieri* (Osteichthyes: Sciaenidae) do litoral do estado do Rio de Janeiro, Brasil. Parasitología al Día, vol. 25, pp. 30-35.
- Ramos Alves, D, Luque, JL & Paraguassú, AR. 2002. *Community Ecology of the Metazoan Parasites of Pink Cusk-eel, Genypterus brasiliensis* (Osteichthyes: Ophidiidae), from the Coastal Zone of the State of Rio de Janeiro, Brazil. Memórias do Instituto Oswaldo Cruz, vol. 97, pp. 683 - 689.
- Rodríguez-González, A & Vidal-Martínez, VM. 2008. *The helminth communities of tonguefish (Symphurus plagiusa) of the Campeche coast, México*. Revista Mexicana de Biodiversidad, vol. 79, pp. 159 - 173.
- Rohde, K. 1984. *Ecology of marine parasites*. Helgoland Marine Research, vol. 37, pp. 5 - 33.
- Rohde, K, Roubal, F & Hewitt, G. 1980. *Ectoparasitic Monogenea, Digenea and Copepoda from the gills of some marine fishes of New Caledonia and New Zealand*. New Zealand Journal of Marine and Freshwater Research, vol. 14, pp.1-13.
- Sánchez, RP & Bezzi, SI. 2004. *El Mar Argentino y sus recursos pesqueros*. Tomo 4. *Los peces marinos de interés pesquero. Caracterización biológica y evaluación del estado de explotación*. Publicaciones especiales Revista del Instituto Nacional de Investigación y Desarrollo Pesquero, Mar del Plata, 359 pp.
- Sánchez, F & Prenski, B. 1996. *Ecología trófica de peces demersales en el Golfo San Jorge*. Revista del Instituto Nacional de Investigación y Desarrollo Pesquero, vol. 10, pp.57-72.

- Sardella, NH, Etchegoin, JA & Martorelli, SR. 1995. *Las comunidades parasitarias de Micropogonias furnieri (corvina) en Argentina*. Boletín del Instituto Oceanográfico de Venezuela, 34, pp. 41-47.
- Sardella, NH & Timi, JT. 1996. *Parasite communities of Merluccius hubbsi from the Argentinean-Uruguayan Common Fishing Zone*. Fisheries Research, 27, pp. 81 - 88.
- Sardella, NH & Timi, JT. 2004. *Parasites of Argentine hake in the Argentine Sea: population and infracommunity structure as evidence for host stock discrimination*. Journal of Fish Biology, vol. 65, pp. 1472 - 1488.
- Sardella, NH, Avendaño, MF & Timi, JT. 1998. *Parasite communities of Genypterus blacodes and G. brasiliensis (Pisces: Ophidiidae) from Argentina*. Helminthologia, vol. 35, pp. 209 - 218.
- Sardella, NH, Mattiucci, S, Timi, J, Bastida, RO, Rodríguez, DH & Nascetti, G. 2005. *Corynosoma australe Johnston, 1937 and C. cetaceum Johnston & Best, 1942 (Acanthocephala: Polymorphidae) from marine mammals and fishes in Argentinian waters: allozyme markers and taxonomic status*. Systematic Parasitology, vol. 61, pp. 143-156.
- Siegel, S & Castellan, NJ. 1995. *Estadística no Paramétrica aplicada a las ciencias de la conducta*. Editorial Trillas, México, 437 pp.
- Suriano, DM & Sutton, CA. 1981. *Contribución al conocimiento de la fauna parasitológica Argentina. VII. Digeneos de peces de la plataforma del mar Argentino*. Revista del Museo de La Plata, Nueva Serie, Sección Zoología, vol. 12, pp. 261-271.
- Szidat, L. 1950. *Los parásitos del róbalo (Eleginops maclovinus Cuv & Val)*. Actas Primer Congreso Nacional de Pesquerías Marítimas e Industriales. Mar del Plata, Argentina, vol. 2, pp. 235-270.
- Szidat, L. 1955. *La fauna de parásitos de Merluccius hubbsi como carácter auxiliar para la solución de problemas sistemáticos y zoogeográficos del género Merluccius L.* Revista del Museo Argentino de Ciencias Naturales Bernardino Rivadavia Zool, vol. 4, pp. 1-54.
- Szidat, L. 1961. *Versuch einer Zoogeographie des Sud-Atlantik mit Hilfe von Leitparasiten der Meeresfische*. Parasitologische Schriftreihe, vol. 13, pp. 1-98.
- Szidat, L & Graefe, G. 1968. *The parasites of Parachaenichthys charcoti an Antarctic fish, in relation to problems of zoogeography* Symposium on Antarctic Oceanography (Santiago). International Council of Science Unions, Special Committee in Antarctic Research 169 - 170.
- Tanzola, RD, Guagliardo, SE, Brízzola, SM & Arias, MV. 1997. *Helminth fauna of Porichthys porosissimus (Pisces: Batrachoidiformes) in the estuary of Bahía Blanca Argentina*. Helminthologia, vol. 34, pp. 221-227.
- Tanzola, RD & Guagliardo, SE. 2000. *Helminth fauna of the Argentine conger, Conger orbignyanus (Pisces: Anguilliformes)*. Helminthologia, vol. 37, pp. 229-232.
- Tanzola, RD & Guagliardo, SE. 2004. *Nematodes anisákidos presentes en peces del área de Bahía Blanca y el riesgo potencial de anisakidosis humana*. Revista de la Asociación Médica Bahía Blanca, vol. 14, pp. 67-73.
- Thoney, DA. 1993. *Community ecology of the parasites of adult spot, Leiostomus xanthurus, and Atlantic croaker, Micropogonias undulatus (Sciaenidae) in the Cape Hatteras region*. Journal of Fish Biology, vol. 43, pp. 781-804.
- Timi, JT. 2003. *Parasites of Argentine anchovy in the Southwest Atlantic: latitudinal patterns and their use for discrimination of host populations*. Journal of Fish Biology, vol. 63, pp. 90-107.
- Timi, JT. 2007. *Parasites as biological tags for stock discrimination in marine fish from South American Atlantic waters*. Journal of Helminthology, vol. 81, pp. 107-111.
- Timi, JT & Poulin, R. 2003. *Parasite community*

- structure within and across host populations of a marine pelagic fish: how repeatable is it?* International Journal for Parasitology vol.33, pp.1353-1362.
- Timi, JT, Luque, JL & Sardella, NH. 2005. *Parasites of Cynoscion guatucupa along South American Atlantic coasts: evidence for stock discrimination.* Journal of Fish Biology, vol. 67, pp. 1603 - 1618.
- Timi, JT, Sardella, NH & Navone, GT. 2001. *Parasitic nematodes of Engraulis anchoita Hubbs et Marini, 1935 (Pisces: Engrauliidae) off the Argentine and Uruguayan coasts, South West Atlantic.* Acta Parasitologica, vol. 46, pp.186 - 193.
- Timi, JT, Lanfranchi, AL, Etchegoin, JA & Cremonte, F. 2008. *Parasites of the brazilian sandperch, Pinguipes brasiliensis. A tool for stock discrimination in the Argentine Sea.* Journal of Fish Biology, vol. 72, pp.1332 - 1342.
- Timi, JT & Lanfranchi, AL. 2009. *The importance of the compound community on the parasite infracommunity structure in a small benthic fish.* Parasitology Research, vol. 104, pp. 295 - 302.
- Valero, A, Paniagua, MI, Hierro, I, Díaz, V, Valderrama, MJ, Benitez, R & Adroher, FJ. 2006. *Anisakid parasites of two forkbeards (Phycis blennoides and Phycis phycis) from the Mediterranean coasts of Andalucía (Southern Spain).* Parasitology International, vol. 55, pp. 1-5.
- Zdzitowiecki, K. 1989. *New data on the morphology and distribution of two acanthocephalans, Andracantha baylisi (Zdzitowiecki, 1986) comb. n. and Corynosoma australe Johnston, 1937.* Acta Parasitologica, vol. 34, pp.167-172.

Received June 11, 2014.
Accepted August 14, 2014.