



Original article

Effect of water activity and temperature on the growth of *Eurotium* species isolated from animal feeds



Mariana Greco ^{a,d,*}, Alejandro Pardo ^{a,d}, Graciela Pose ^{b,d}, Andrea Patriarca ^{c,d}

^a Laboratorio de Micología Molecular, Departamento de Ciencia y Tecnología, Universidad Nacional de Quilmes (UNQ), Roque Sáenz Peña 352, B1876BXD Bernal, Buenos Aires, Argentina

^b Universidad Nacional de Río Negro (UNRN), Escuela de Producción, Tecnología y Medio Ambiente, Mitre 331, 8336 Villa Regina, Río Negro, Argentina

^c Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Química Orgánica, Laboratorio de Microbiología de Alimentos, Ciudad Universitaria, Intendente Güiraldes 2160, Pabellón II, 3 Piso, C1428EGA Buenos Aires, Argentina

^d Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina

ARTICLE INFO

Article history:

Received 11 October 2016

Accepted 20 April 2017

Available online 12 November 2017

Keywords:

Eurotium
Water activity
Temperature
Growth
Animal feeds

ABSTRACT

Background: Xerophilic fungi represent a serious problem due to their ability to grow at low water activities causing the spoiling of low and intermediate moisture foods, stored goods and animal feeds, with the consequent economic losses.

Aims: The combined effect of water activity and temperature of four *Eurotium* species isolated from animal feeds was investigated.

Methods: *Eurotium amstelodami*, *Eurotium chevalieri*, *Eurotium repens* and *Eurotium rubrum* were grown at 5, 15, 25, 37 and 45 °C on malt extract agar adjusted with glycerol in the range 0.710–0.993 of water activities.

Results: The cardinal model proposed by Rosso and Robinson (2001) was applied to fit growth data, with the variable water activity at fixed temperatures, obtaining three cardinal water activities ($a_{w\min}$, $a_{w\max}$, $a_{w\text{opt}}$) and the specific growth rate at the optimum a_w (μ_{opt}). A probabilistic model was also applied to define the interface between growth and no-growth. The cardinal model provided an adequate estimation of the optimal a_w to grow and the maximum growth rate. The probabilistic model showed a good performance to fit growth/no-growth cases in the predicted range.

Conclusions: The results presented here could be applied to predict *Eurotium* species growth in animal feeds.

© 2017 Asociación Española de Micología. Published by Elsevier España, S.L.U. All rights reserved.

Efecto de la actividad del agua y la temperatura en el crecimiento de especies de *Eurotium* aisladas de alimentos para animales

RESUMEN

Palabras clave:

Eurotium
Actividad del agua
Temperatura
Crecimiento
Piensos para animales

Antecedentes: Los hongos xerófilos son un problema importante debido a su capacidad de crecer a bajas actividades del agua, lo que causa el deterioro de alimentos a humedades bajas e intermedias, de materias primas almacenadas y de piensos para animales, con las consecuentes pérdidas económicas.

Objetivos: Se llevó a cabo un estudio sobre el efecto de los factores ambientales (temperatura y actividad del agua) sobre el crecimiento de cuatro especies pertenecientes al género *Eurotium* aisladas de piensos para animales.

Métodos: Se estudió el crecimiento de *Eurotium amstelodami*, *Eurotium chevalieri*, *Eurotium repens* y *Eurotium rubrum* a valores de actividad de agua en el rango 0,710–0,993 en el medio de cultivo agar extracto de malta modificado con glicerol, y valores de temperatura de 5, 15, 25, 37 y 45 °C.

Resultados: El modelo cardinal propuesto por Rosso y Robinson (2001) se aplicó para realizar el ajuste de datos con la actividad del agua como variable a una temperatura fija; se obtuvieron tres valores cardinales

* Corresponding author.

E-mail address: mariana.greco@gmail.com (M. Greco).

de actividad del agua ($a_{w\min}$, $a_{w\max}$, $a_{w\text{opt}}$) y la tasa de crecimiento específico en el valor óptimo de a_w (μ_{opt}). También se aplicó un modelo probabilístico para definir la interfase entre crecimiento y no crecimiento. El modelo cardinal presentó una adecuada estimación del $a_{w\text{opt}}$ y la máxima velocidad de crecimiento. El modelo probabilístico fue adecuado para el ajuste de los casos de crecimiento/falta de crecimiento en el rango previsto.

Conclusiones: Los resultados presentados en este artículo pueden aplicarse para pronosticar el crecimiento de especies de *Eurotium* en piensos para animales.

© 2017 Asociación Española de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Animal feeds are made from different ingredients such as corn, wheat, rice, soybean, sorghum, barley and other grain mill products, animal proteins, vitamins, minerals, etc. Appropriate amounts of all nutrients are required to maintain animals healthy and get faster at maximum weight. For feedstuff stability, maximum moisture contents are critical; below 0.6 water activity (a_w) spoilage by microorganisms would not be expected.²⁷ Feedstuffs are produced by commercial feed mills, as well as by home mixture, being low water activity a characteristic of these feeds (average 0.537). However, improper production or inadequate storage conditions allow the growth and proliferation of xerophilic fungi, resulting in a negative impact on nutritional and organoleptic properties of feeds and mycotoxin production.^{4,23} Thus, feedstuff should be discarded, with the subsequent economic losses.

According to the recent changes in nomenclature, the genus *Eurotium* was transferred to the anamorphic *Aspergillus* genus,^{31,41} although a proposal has been recently submitted to Taxon by Pitt and Taylor³⁶ in order to keep the name *Eurotium*, advocating the recognition of the diversity in the morphological and physiological phenotypes among these fungi. The genus belongs to the phylum Ascomycota, characterized by the production of ascospores resistant to high temperatures inside bright yellow cleistothecia.

Eurotium species are universally known as spoiling fungi, and are responsible for substantial economic losses of food commodities, leather goods and textiles. Most of its species are of particular interest to food and feed mycology because of their xerophilic physiology; many isolates are able to grow at water activities below 0.75 and some have been reported to grow at values as low as 0.64 a_w .¹⁰ Twenty *Eurotium* species are known approximately; among the most common ones, *Eurotium amstelodami*, *Eurotium chevalieri*, *Eurotium repens*, and *Eurotium rubrum* are usually associated with stored goods.³⁵ The growth and development of these fungi cause changes in the organoleptic characteristics and the nutritional quality of the raw materials and commodities, affecting the shelf life of processed products. Furthermore, these species have been reported to produce secondary metabolites.^{1,10,24} Some of these compounds, such as echinulin, physcion and flavoglaucin, have shown toxicity to animals.²⁴ Although most of the studies on feedstuffs mycobiota have been focused on the presence of the micotoxigenic genera *Aspergillus*, *Fusarium* and *Penicillium*,^{11,12,18,22,29,32,34,38} *Eurotium* species have also been reported as frequent contaminants.^{7,43} In Argentina, *Eurotium* species have been determined as major components of the mycobiota in poultry,^{18,23} chinchilla, rabbit and rainbow trout feeds.^{22,25} The interaction between environmental factors and food characteristics might provide the required conditions for the growth and development of filamentous fungi, followed by potential toxic metabolite production. In order to predict and avoid undesirable fungal growth it is necessary to determine the influence of those environmental factors, such as water activity and temperature.

The aim of the present work was to evaluate the effect of a_w and temperature on the germination time and growth of four xerophilic *Eurotium* species isolated from animal feeds. Mathematical modelling tools were tested to describe and characterize the ecophysiology of the xerophilic fungi on synthetic media.

Materials and methods

Fungal isolates

Four species belonging to the genus *Eurotium* were used in the present study: *E. amstelodami*, isolated from rabbit feed, and *E. chevalieri*, *E. repens* and *E. rubrum*, isolated from chinchilla feeds. The identity of the four isolates had been previously confirmed by scanning electron microscopy of ascospores and DNA sequencing of two independent DNA loci (ITS and beta-tubulin).²⁴

Media

The basal medium used was malt extract agar (MEA), consisting of 2% malt extract (Oxoid, LP0039, UK), 0.1% peptone (Britania, Buenos Aires, Argentina), 2% glucose (Oxoid, LP0071) and 2% agar (Oxoid, LP0011), at nine different water activities (a_w). Media were adjusted by substituting equal parts of water by glycerol (w/w). Different water contents were prepared in the range 0.710–0.993 (0.710, 0.750, 0.790, 0.830, 0.870, 0.914, 0.952, 0.989 and 0.993). Glycerol concentrations were calculated according to Chirife et al.¹³ and Ross.³⁸ All media were sterilized by steam treatment at 121 °C for 15 min. The a_w was measured with an AquaLab CX-2 water activity metre (Decagon Devices, Inc., USA).

Preparation of the inoculum

Fungi were grown on Czapek yeast extract agar with 20% sucrose (CY20S) for seven days at 25 °C to obtain heavily sporulating cultures. After incubation, conidia from CY20S plates were removed and suspended in 5 ml of sterile water/glycerol solutions with their a_w previously modified to the required value, and containing 0.05% of Tween 80 (Anedra, Research AG, Buenos Aires, Argentina). The number of conidia from this stock suspension was determined using a Neubauer chamber, and then the final concentration was adjusted to 10⁵ conidia/ml.

Inoculation and incubation

Petri plates (55 mm diameter) with approximately 10 ml of medium were centrally inoculated with a calibrated loop containing 1 µl of the conidia suspension (average 100 conidia). Petri plates with the same a_w values were enclosed in polyethylene bags and incubated at 5, 15, 25, 37 and 45 °C for a maximum of 90 days. To minimize the transfer of water to or from the media, plates with glycerol solutions adjusted to the corresponding a_w , were placed

inside the plastic bags and changed weekly. Non-inoculated control plates were included inside each bag and measured at the end of the experiment to detect any significant deviation of the initial a_w , and no variation higher than 0.001 was detected. Each treatment ($a_w T$) was carried out in four replicates for each *Eurotium* species, either for germination and growth rate measurements. To obtain enough growth/no-growth data for the probabilistic models, four additional plates were inoculated and incubated under each set of conditions, and the combined data of 8 replicates were used for each species at each $a_w T$ treatment.

Germination measurement

To determine the germination, Petri plates were examined twice a day using a stereomicroscope ($\times 40$). The criterion to decide that germination had occurred was the production of a germination tube with a similar length to the diameter of the conidia in at least 50% of the conidia.²⁶ Germination times were recorded in hours and reported in days as the mean of four replicates for each species under each $a_w T$ treatment. Plates were examined until 90 days.

Growth measurement

The radial mycelial growth rate was determined by periodical measurement of two perpendicular diameters of the colonies. Growing colonies were measured daily until the end of the experiment (90 days) or until the colony reached the edge of the Petri plate.

Statistical methods

Growth rates (mm/day) for each a_w and temperature combination were calculated as the linear regression from the linear phase of the growth curve. Secondary models were used to describe the influence of water activity and temperature on fungal growth. The Rosso cardinal model^{39,40} at a fixed temperature was applied to describe the effect of water activity on the radial growth rate. As the *Eurotium* strains isolated from animal feeds used in the present study did not grow at many of the temperatures tested, a secondary model was applied at fixed temperatures of 15 and 25 °C, at which enough growth data were available.

The model is described by the following equation:

$$\mu_{\max} = \frac{\mu_{\text{opt}} (a_w - a_{w\max})(a_w - a_{w\min})^2}{(a_{w\text{opt}} - a_{w\min}) \{ (a_{w\text{opt}} - a_{w\min})(a_w - a_{w\text{opt}})(a_{w\text{opt}} - a_{w\max})(a_{w\text{opt}} + a_{w\min} - 2a_w) \}}$$

The equation was fitted using the Statistica v 10.0 (StatSoft Inc.) non-linear estimation procedure. The goodness of fit of the model was evaluated by the root mean square error (RMSE). Homogeneity of the variance of the dependent variable was tested by Levene's test using Statistica v 10.0.

To treat the data of each of the four fungal species, a linear logistic regression analysis was applied to determine the growth/no-growth boundaries under the different a_w and temperature values evaluated. Data from 8 replicates per species were expressed as values of 1 or 0 corresponding to growth or no-growth, respectively. The resulting data were fitted to a logistic regression equation³⁷ with a full second order logistic regression model¹⁸ that includes the linear term for time:

$$\text{Logit } P = \ln(P/(1-P)) = b_0 + b_1 a_w + b_2 T + b_{11} a_w^2 + b_{22} T^2 + b_{12} a_w T + t$$

where P is the probability of growth (range 0–1), b_i are the coefficients to be estimated, t (days) is the incubation time, a_w is the water activity of the medium, and T is the incubation

temperature in °C. The equation was fitted using Statistica 10.0 linear logistic regression procedure. The predicted growth/no-growth interfaces for $P=0.1$, 0.5 and 0.9 were calculated and plotted using Microsoft Excel 2003 Solver and MatLab (MathWorks, Version 7.0.1R14SP1).

Results

Germination data

Influence of a_w on the germination time for each species is shown in Fig. 1. For *E. amstelodami*, germination occurred in the range 0.830–0.993 a_w at 15 °C and 0.790–0.993 a_w at 25 °C. Germination at 37 °C was only observed at 0.989 a_w (2 days). No germination was registered at any of the tested a_w values at 5 and 45 °C. The lowest germination time was observed at 0.952 a_w and 25 °C (0.96 days) and the highest was at 14 days, occurring at 15 °C and 0.830 a_w .

Germination of *E. chevalieri* occurred in a wider range of a_w (0.750–0.993), both at 15 and 25 °C. Moreover, at 37 °C and 0.989 a_w or 0.993 a_w , germination followed by mycelial growth was also registered after 0.88 and 0.92 days, respectively. No germination was observed at the other a_w values at 37 °C or at any of the a_w values tested between 5 and 45 °C. The shortest time required for germination was 0.79 days, registered at 25 °C and 0.989 a_w , while the longest for this species (27.3 days) was observed at 15 °C and 0.750 a_w .

E. repens presented a quite different behaviour from the above-mentioned species. Germination only occurred at the highest a_w values (range 0.914–0.993), both at 15 and 25 °C. In the range 0.914–0.952 a_w , germination was also observed at 5 °C, being this species the only one capable of growing at this temperature. No germination was registered at any of the tested a_w values at 37 and 45 °C. The fastest germination for *E. repens* occurred at 25 °C and 0.989 a_w (0.79 days), while it took 60.4 days to germinate at 5 °C and 0.914 a_w ; this was the longest germination time observed during the whole incubation period at all the evaluated conditions.

Unlike the other *Eurotium* species, *E. rubrum* presented an upper limit of a_w , since it did not germinate above 0.952 a_w at any of the tested temperatures. The lower limit for germination was 0.750 a_w , at 15 and 25 °C. No germination was observed at any other assayed

temperature (5, 37, and 45 °C). The shortest germination time was 1.57 days at 0.952 a_w and 25 °C, and the longest was 41.8 days at 0.750 a_w and 15 °C.

Growth data

Growth of *E. amstelodami* was observed in the whole range of conditions in which germination occurred (0.790–0.993 a_w at 25 °C, 0.830–0.993 a_w at 15 °C). The minimum growth rate was registered at 15 °C and 0.830 a_w (1.1 mm/day). Nevertheless, it grew slowly at 37 °C and 0.989 a_w (0.18 mm/day). The optimal a_w levels for growing were in the range 0.914–0.952 a_w at 15 and 25 °C. The maximum growth rate was 8.3 mm/day at 25 °C and 0.952 a_w (Fig. 2).

E. chevalieri growth also occurred at all the conditions which allowed germination (0.750–0.993, both at 15 and 25 °C). It was also capable of growing at 37 °C and 0.989 and 0.993 a_w , but with low growth rates (0.6 and 0.5 mm/day, respectively). The optimal conditions for growing were 0.952 a_w and 25 °C, with a maximum




Fig. 1. Influence of a_w on the germination time at 15 and 25 °C for *Eurotium amstelodami*, *E. chevalieri*, *E. repens* and *E. rubrum*.




Fig. 2. Growth rates (mm/day) as a function of a_w at the different temperatures for *E. amstelodami*, *E. chevalieri*, *E. repens* and *E. rubrum*. Fitted curves of the Rosso cardinal model at 25 °C (solid line) and 15 °C (dashed line) are shown for *E. amstelodami*, *E. chevalieri*, and *E. rubrum*.

growth rate of 9.6 mm/day. At 15 °C, 0.952 was also the optimum a_w level, at which the growth rate was 3.3 mm/day (Fig. 2).

Regarding *E. repens*, the a_w range for growing was again the same as for the germination. At temperatures of 15 and 25 °C, the growth only occurred in the range 0.914–0.993 a_w . Unlike

the species previously discussed, *E. repens* could grow at 5 °C and 0.914 a_w and 0.952 a_w levels, although at low rates (0.43 and 0.51 mm/day, respectively). The optimal a_w value to grow was 0.952 at 5, 15 and 25 °C, with a maximum growth rate of 13.5 mm/day (25 °C and 0.952 a_w) (Fig. 2).

Table 1

Cardinal values of water activity (minimum, maximum and optimum) and growth rate (optimum) predicted by secondary cardinal models, experimental values and RMSE obtained for *Eurotium* species from animal feeds.

Species	Parameter	T=25 °C			T=15 °C		
		Estimated value ± SE	Experimental value	RMSE	Estimated value ± SE	Experimental value	RMSE
<i>E. amstelodami</i>	μ_{opt}	8.44 ± 0.14	8.26	0.299	2.51 ± 0.05	2.57	0.126
	$a_{w\text{max}}$	1.00 ± 0.00	0.993		1.01 ± 0.00	0.993	
	$a_{w\text{min}}$	0.71 ± 0.01	0.790		0.77 ± 0.01	0.830	
	$a_{w\text{opt}}$	0.95 ± 0.00	0.952		0.92 ± 0.00	0.914	
<i>E. chevalieri</i>	μ_{opt}	8.89 ± 0.21	9.58	0.529	2.92 ± 0.12	3.36	0.317
	$a_{w\text{max}}$	0.99 ± 0.00	0.993		0.99 ± 0.00	0.993	
	$a_{w\text{min}}$	0.71 ± 0.01	0.750		0.74 ± 0.01	0.750	
	$a_{w\text{opt}}$	0.95 ± 0.00	0.952		0.95 ± 0.00	0.952	
<i>E. rubrum</i>	μ_{opt}	7.47 ± 0.20	8.17	0.581	3.67 ± 0.04	3.58	0.118
	$a_{w\text{max}}$	0.99 ± 0.00	0.952		0.99 ± 0.00	0.952	
	$a_{w\text{min}}$	0.71 ± 0.01	0.750		0.70 ± 0.00	0.750	
	$a_{w\text{opt}}$	0.90 ± 0.00	0.914		0.90 ± 0.00	0.914	
<i>E. repens</i>	μ_{opt}	—	13.53	—	—	6.35	—
	$a_{w\text{max}}$	—	0.993		—	0.993	
	$a_{w\text{min}}$	—	0.914		—	0.914	
	$a_{w\text{opt}}$	—	0.952		—	0.952	

For *E. rubrum*, growth was registered both at 15 and 25 °C in the range 0.750–0.952 a_w . The optimal a_w values were in the range 0.870–0.914 a_w at 15 °C (3.6 mm/day at both a_w levels) and 0.914 a_w at 25 °C, conditions at which the maximum growth rate was registered (8.2 mm/day) (Fig. 2). The minimum growth rate for this species was 0.62 mm/day and it occurred at 15 °C and 0.750 a_w .

Secondary model

Several existing secondary kinetic models were evaluated (data not shown) to describe fungal growth response in the assayed environmental conditions (Davey, Miles, Gibson, Parra and polynomial models),¹⁹ obtaining the best fit to the experimental data when applying the Rosso cardinal model at fixed temperatures of 15 and 25 °C. In this model it is common to substitute $a_{w\text{max}}$ for 1,⁴² since a_w cannot exceed this value, and most bacteria and several fungal species have no upper limit to grow. However, as this was not the case with xerophilic fungi, this parameter was left to be estimated by the model. The cardinal values of environmental factors obtained (minimum, maximum and optimum) by the secondary cardinal model for each species are shown in Table 1, while graphics of the adjusted models (solid and dashed lines) and experimental data (points) are presented in Fig. 2. The root mean square error (RMSE) was calculated to evaluate the performance of the predictive model for each studied species (Table 1). This parameter determines the average deviation between the observed values and the predicted ones. According to RMSE values, the model was adequate to fit the experimental growth rate data for *E. amstelodami* and *E. chevalieri*, at both temperatures, although better estimations were obtained at 15 °C than at 25 °C for all the tested species. The highest RMSE and, consequently, the poorest fit comparatively among species, was obtained for *E. rubrum* at 25 °C. It was not possible to apply the Rosso cardinal model to *E. repens* due to insufficient growth data.

For *E. amstelodami*, the cardinal parameters estimated by the model were in agreement with the experimental data, except for the minimum a_w for growing, which was lower than the experimental value at both temperatures (Table 1). The same was observed for *E. chevalieri* at 25 °C, while at 15 °C the predictions were in good agreement with the measured values. For *E. rubrum*, however, cardinal values estimated by the model differed from the experimental ones, except for the optimum a_w to grow ($a_{w\text{opt}}$) and the maximum growth rate (μ_{opt}) at 15 °C.

Table 2
Second order logistic regression models for *Eurotium* species.

Species	Parameter	Estimated value ± SE
<i>E. amstelodami</i>	Intercept	377 ± 17
	t	-0.073 ± 0.005
	a_w	-682 ± 39
	T	-6.4 ± 0.2
	a_w^2	346 ± 20
	T^2	0.109 ± 0.001
<i>E. chevalieri</i>	$a_w T$	1.9 ± 0.2
	Intercept	74 ± 5
	t	-0.068 ± 0.003
	a_w	-99 ± 14
	T	-0.80 ± 0.08
	a_w^2	95 ± 9
<i>E. rubrum</i>	T^2	0.127 ± 0.005
	$a_w T$	-5.8 ± 0.3
	Intercept	395 ± 15
	t	-0.066 ± 0.003
	a_w	-807 ± 29
	T	-5.3 ± 0.1
<i>E. repens</i>	a_w^2	457 ± 16
	T^2	0.102 ± 0.001
	$a_w T$	1.3 ± 0.1
	Intercept	2809 ± 97
	t	-0.104 ± 0.008
	a_w	-5951 ± 432

p value < 0.001.

Probability model

Table 2 shows second order logistic regression models. Models were generated including linear terms, quadratic (except the quadratic term of time), and interactions terms, for each of these four species. As all the estimated parameters were significant ($p < 0.001$), no backward elimination of terms was done. Figs. 3–6 show the predicted growth/no-growth boundaries and growth/no-growth cases at probabilities 0.1, 0.5 and 0.9, for time periods of 7, 30 and 90 days. A high percentage of agreement between the experimental data and model fit can be observed. All cases contained in the area $P > 90\%$ have shown 100% growth (all replicates showed growth at the reported time); 59% of the cases in the area $10 < P < 90$ showed 100% growth, and there were no-growth cases in the $P < 10\%$ region.




Fig. 3. Predicted growth/no-growth boundaries for *E. amstelodami* at: (a) 7, (b) 30, and (c) 90 days, with respect to a_w and temperature at probabilities of 0.1 (dotted line), 0.5 (dashed line) and 0.9 (solid line). (●) 100% observed growth; (○) no growth.

Discussion

According to the data obtained in this study for *E. amstelodami*, minimum values of water activity and temperature at which germination and growth occurred were $0.790\ a_w$ at $25\ ^\circ\text{C}$, and $0.830\ a_w$ at $15\ ^\circ\text{C}$, showing the highest growth rate at $0.952\ a_w$ and $25\ ^\circ\text{C}$. Other authors reported an optimal temperature to grow at $33\text{--}35\ ^\circ\text{C}$,¹⁶ and a maximum of $43\text{--}46\ ^\circ\text{C}$.⁹ El Halouat and Debevere¹⁷ reported that the germination of *E. amstelodami* isolated from prunes in synthetic medium under modified atmosphere occurred at temperatures as high as $40\ ^\circ\text{C}$ at 0.83 , 0.92 and $0.95\ a_w$. However, in our study, growth at $37\ ^\circ\text{C}$ was only detected at $0.989\ a_w$, and no growth was observed at $45\ ^\circ\text{C}$. Armolik and Dickson⁵ reported growth of

E. amstelodami from stored grain at $25\ ^\circ\text{C}$ and $0.75\ a_w$, a lower a_w value than that observed in the present work. In addition, Abellana et al.² reported growth of *Eurotium* species from bakery products, including *E. amstelodami*, on flow wheat-sucrose agar at $0.775\ a_w$ and temperatures above $20\ ^\circ\text{C}$. It seems that the a_w and temperature ranges of *E. amstelodami* from food origin is wider than for the animal feed strain used in the present work.

Regarding *E. chevalieri*, the minimum a_w value at which germination and growth occurred was 0.750 , both at 25 and $15\ ^\circ\text{C}$; maximum growth was registered at $0.952\ a_w$ and $25\ ^\circ\text{C}$. Information from the literature indicates that the optimum temperature for growing is in the range $30\text{--}35\ ^\circ\text{C}$ ¹⁶ with a maximum at $40\text{--}43\ ^\circ\text{C}$.⁹ In the present study the growth at $37\ ^\circ\text{C}$ only occurred at the two




Fig. 4. Predicted growth/no-growth boundaries for *E. chevalieri* at: (a) 7, (b) 30, and (c) 90 days, with respect to a_w and temperature at probabilities of 0.1 (dotted line), 0.5 (dashed line) and 0.9 (solid line). (●) 100% observed growth; (○) 10–90% growth.

highest a_w levels, and no growth was registered at 45 °C. Ayerst⁶ reported the minimum growth at 0.71 a_w and 42 °C, while Abellana et al.³ determined that an *E. chevalieri* strain from bakery products was able to grow at 0.75 a_w at 30 °C. However, both Abellana et al.³ and Marín et al.³⁰ found that *E. chevalieri* could not grow on sponge cake analogue medium at 0.75 a_w and 25 °C; meanwhile a strain of *E. chevalieri* isolated from milk jam was able to grow at 0.74 a_w at this temperature.¹² Even though literature data differ on the growing extreme conditions, the strain isolated from animal feeds was able to grow in a more restricted a_w and temperature range than most of the reported strains from foods.

E. repens showed the highest growth rate at 0.952 a_w and 25 °C, while 0.91 was the minimum a_w value at which germination and

growth was observed, both at 15 and 25 °C. This was the only species able to grow at 5 °C, but unlike the two previous mentioned ones, it did not grow at 37 °C. According to González et al.²¹ and Panasenko,³³ *E. repens* grows in the range of temperatures from 4–5 to 38–40 °C, with the optimum temperature in the range 25–27 °C. Several authors have reported 0.72 a_w as the minimum value at which germination occurs at temperatures of 20–25 °C.^{5,28} Meanwhile, Gock et al.²⁰ reported a minimum a_w for germination and growth of 0.74 at 25 °C for an isolate from dried prunes. According to Dagnas et al.,¹⁴ *E. repens* isolated from bakery products showed estimated values of optimum temperature and a_w of 29 °C and 0.91, respectively, and could grow in the temperature range 0–35 °C. More recently, Dagnas et al.¹⁵ estimated a minimum a_w of 0.74 for this species. The narrow range of a_w levels at which the animal feed




Fig. 5. Predicted growth/no-growth boundaries for *E. repens* at: (a) 7, (b) 30, and (c) 90 days, with respect to a_w and temperature at probabilities of 0.1 (dotted line), 0.5 (dashed line) and 0.9 (solid line). (●) 100% observed growth; (○) 10–90% growth; (○) no growth.

strain was able to grow differs from most reports in the literature and from the other *Eurotium* species studied.

Minimum a_w values of 0.70–0.72 have been reported in the germination of *E. rubrum* at 25 °C,^{5,20} with a minimum growth temperature of 5 °C and an optimum of 25–27 °C.³³ Gock et al.²⁰ reported minimum growth parameters of 0.74 a_w and 25 °C on synthetic medium for isolates from prunes. The *E. rubrum* isolate from animal feeds used in the present study could not grow below 0.750 a_w at any of the tested temperatures. The highest growth rate for this strain was observed at 0.914 a_w at 25 °C, which is similar to that reported by Wheeler et al.⁴⁴ for dried fish strains (0.91–0.94).

These wide discrepancies could be due to geographical variabilities among isolates, nutritional differences of the substrates used

to assess growth,³⁴ and climate adaptation. However, in general, isolates from food products were able to germinate and grow in a wider range of a_w and temperature than those from animal feeds evaluated in the present work. Thus, it is relevant to understand the ecophysiology of these particular strains in order to prevent fungal contamination of animal feeds and reduce the costs associated with their spoilage.

In general terms, the cardinal model was adequate for the estimation of the $a_{w\text{opt}}$ and μ_{opt} , the latter with the only exception of *E. rubrum* at 25 °C. However, limiting conditions were poorly predicted by this model, especially the $a_{w\text{min}}$ for growing, which was underestimated in all three *Eurotium* species at both temperatures, except for *E. chevalieri* at 15 °C. Regarding the growing-upper a_w limit, the model could appropriately estimate it when this




Fig. 6. Predicted growth/no-growth boundaries for *E. rubrum* at: (a) 7, (b) 30, and (c) 90 days, with respect to a_w and temperature at probabilities of 0.1 (dotted line), 0.5 (dashed line) and 0.9 (solid line). (●) 100% observed growth; (○) 10–90% growth; (○) no growth.

parameter was close to 1, but it failed to provide a suitable estimation when an a_w limit for growing existed, as the case with *E. rubrum*.

Since for spoiling xerophilic fungi predicting the restricting environmental conditions are more relevant than the optimal ones for growing, probabilistic models were also developed for each *Eurotium* species with the aim to predict the growth/no-growth boundaries. As no-growth data sets are useful for the construction of these models, both a_w and T terms were included in the probability equation. The model showed an adequate fit to the experimental data and could accurately predict growth/no-growth cases in the range under study (Figs. 3–6). Validation with literature data is difficult due to the high variability observed between the strains isolated from different sources and geographical origins.

Additionally, in some cases, the range of environmental conditions studied does not include minimum, maximum or optimum values to grow. Comparison of the minimum a_w predicted by probability models for the different *Eurotium* species with previously published data shows certain agreement. For *E. amstelodami*, minimum predicted a_w was 0.82 at 15 °C and 0.76 at 25 °C, values in agreement with Abellana et al.^{2,3} for strains from bakery products ($a_{w\min}$ 0.80–0.825 at 15 °C, and 0.75–0.80 at 25 °C). The minimum a_w reported by Char et al.¹² for *E. chevalieri* on milk jam (<0.74) shows a better agreement with the one predicted by the model than those reported by Abellana et al.^{2,3} (0.825–0.875 at 15 °C, and 0.775–0.80 at 25 °C), as well as the minimum a_w (<0.74 reported by Gock et al.²⁰ for *E. rubrum* in malt extract agar. The predicted cardinal for *E. repens* did not show agreement with literature

data,^{14,20} as it was expected by the difference observed between this strain and most food strains reported elsewhere.

The data reported in the present study are relevant as a contribution to the understanding of the ecophysiological behaviour of these four *Eurotium* species isolated from animal feeds. The results presented here can be applied, after validation in the respective substrates, to predict *Eurotium* spp. growth in feeds and similar matrices. This information will be useful to develop control strategies to prevent fungal spoilage of feeds.

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was supported by UNQ, UNRN, CONICET, and ANPCyT. G. Pose, A. Patriarca and A. Pardo are members of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). Engineer G. Pacheco is acknowledged for his assistance.

References

1. Abarca ML, Bragulat MR, Castellá G, Accensi F, Cabañes FJ. Hongs productores de micotoxinas emergentes. *Rev Iberoam Micol.* 2000;17:S63–8.
2. Abellana M, Benedi J, Sanchis V, Ramos AJ. Water activity and temperature effects on germination and growth of *Eurotium amstelodami*, *E. chevalieri* and *E. herbariorum* isolates from bakery products. *J Appl Microbiol.* 1999;87:371–80.
3. Abellana M, Magrí X, Sanchis V, Ramos AJ. Water activity and temperature effects on growth of *Eurotium amstelodami*, *E. chevalieri* and *E. herbariorum* on a sponge cake analogue. *Int J Food Microbiol.* 1999;52:97–103.
4. Alam S, Shah HU, Afzal M, Magan N. Influence of calcium propionate, water activity and storage time on mold incidence and aflatoxins production in broiler starter feed. *Anim Feed Sci Technol.* 2014;188:137–44.
5. Armolik N, Dickson JG. Minimum humidity requirements for germination of conidia associated with storage of grain. *Phytopathology.* 1956;46:462–5.
6. Ayerst G. The effects of moisture and temperature on growth and spore germination in some fungi. *J Stored Prod Res.* 1969;5:127–41.
7. Barbosa TS, Pereyra CM, Soleiro CA, Dias EO, Oliveira AA, Keller KM, et al. Mycobiota and mycotoxins present in finished fish feeds from farms in the Rio de Janeiro State, Brazil. *Int Aquat Res.* 2013;5:1–9.
8. Battey AS, Duffy S, Schaffner DW. Modeling yeast spoilage in cold-filled ready-to-drink beverages with *Saccharomyces cerevisiae*, *Zygosaccharomyces bailii*, and *Candida lipolytica*. *Appl Environ Microb.* 2002;68:1901–6.
9. Blaser P. Taxonomische und physiologische Untersuchungen über die Gatung *Eurotium* Link ex Fries. *Syndowia.* 1975;28:1–49.
10. Butinar L, Zalar P, Frisvad J, Gunde-Cimerman N. The genus *Eurotium* – members of indigenous fungal community in hypersaline waters of salterns. *FEMS Microbiol Ecol.* 2005;51:155–66.
11. Cegielska-Radziejewska R, Stuper K, Szablewski T. Micoflora and mycotoxins contamination in poultry feed mixtures from western Poland. *Ann Agric Environ Med.* 2013;20:30–5.
12. Char C, Guerrero S, González L, Alzamora SM. Growth response of *Eurotium chevalieri*, *Aspergillus fumigatus* and *Penicillium brevicompactum* in Argentine milk jam. *Food Sci Technol Int.* 2005;11:297–305.
13. Chirife J, Ferro Fontán C, Benmergui EA. The prediction of water activity in aqueous solutions in connection with intermediate moisture foods. IV. a_w prediction in aqueous non-electrolyte solutions. *J Food Technol.* 1980;70:15–59.
14. Dagnas S, Onno B, Membré JM. Modeling growth of three bakery product spoilage molds as a function of water activity, temperature and pH. *Int J Food Microbiol.* 2014;186:95–104.
15. Dagnas S, Gougli M, Onno B, Koutsoumanis KP, Membré JM. Quantifying the effect of water activity and storage temperature on single spore lag times of three moulds isolated from spoiled bakery products. *Int J Food Microbiol.* 2017;240:75–84.
16. Domsch KH, Gams W, Anderson TH. Compendium of soil fungi 2. London: Academic Press; 1980.
17. El Halouat A, Debevere JM. Effect of water activity, modified atmosphere packaging and storage temperature on spore germination of moulds isolated from prunes. *Int J Food Microbiol.* 1997;35:41–8.
18. Fraga ME, Curvello F, Gatti LR, Cavagliere KR, Dalcerio AM, da Rocha Rosa CA. Potential aflatoxin and ochratoxin A production by *Aspergillus* species in poultry feed processing. *Vet Res Commun.* 2007;31:343–53.
19. García D, Ramos AJ, Sanchis V, Marín S. Predicting mycotoxins in foods: a review. *Food Microbiol.* 2009;26:757–69.
20. Gock MA, Hocking AD, Pitt JI, Poulos PG. Influence of temperature, water activity and pH on growth of some xerophilic fungi. *Int J Food Microbiol.* 2003;2481:11–9.
21. González HHL, Resnik SL, Vaamonde G. Influence of temperature on growth rate and lag phase of fungi isolated from Argentine corn. *Int J Food Microbiol.* 1988;6:179–83.
22. Greco MV, Pardo AG, Ludemann V, Martino PE, Pose GN. Mycoflora and natural incidence of selected mycotoxins in rabbit and chinchilla feeds. *Sci World J.* 2012;1:1–6.
23. Greco MV, Franchi ML, Rico Golba SL, Pardo AG, Pose GN. Mycotoxins and mycotoxicogenic fungi in poultry feed for food-producing animals. *Sci World J.* 2014;1:9.
24. Greco M, Kempainen M, Pose G, Pardo A. Taxonomic characterization and secondary metabolite profiling of *Aspergillus* section *Aspergillus* contaminating feeds and feedstuffs. *Toxins.* 2015;7:3515–37.
25. Greco M, Pardo A, Pose G. Mycotoxicogenic fungi and natural co-occurrence of mycotoxins in rainbow trout (*Oncorhynchus mykiss*) feeds. *Toxins.* 2015;7:4595–609.
26. Hocking AD, Miscamble BF. Water relations of some zygomycetes isolated from food. *Mycol Res.* 1995;99:1113–5.
27. Lowe JA, Kershaw SJ. Water activity-moisture content relationship as a predictive indicator for control of spoilage in commercial pet diet components. *Anim Feed Sci Technol.* 1995;56:187–94.
28. Magan N, Lacey J. Effect of temperature and pH on the water relations of field and storage fungi. *Trans Br Mycol Soc.* 1984;82:71–81.
29. Magnoli C, Dalcerio AM, Chiacchiera SM, Mazzola R, Saenz MA. Enumeration and identification of *Aspergillus* group and *Penicillium* species in poultry feeds from Argentina. *Mycopathologia.* 1998;142:27–32.
30. Marín S, Abellana M, Rubiat M, Sanchis V, Ramos AJ. Efficacy of sorbates on the control of the growth of *Eurotium* species in bakery products with near neutral pH. *Int J Food Microbiol.* 2003;87:251–8.
31. McNeill J, Barrie FF, Buck WR, editors. International Code of Nomenclature for algae, fungi, and plants (Melbourne Code). Königstein (Regnumvegetabile no. 154): KoeltzScientificBooks; 2012.
32. Monge MP, Magnoli CE, Chiacchiera SM. Survey of *Aspergillus* and *Fusarium* species and their mycotoxins in raw materials and poultry feeds from Córdoba, Argentina. *Mycotoxin Res.* 2012;28:111–2.
33. Panasenko VT. Ecology of microfungi. *Bot Rev.* 1967;33:189–215.
34. Pardo E, Ramos AJ, Sanchis V, Marín S. Modelling of effects of water activity and temperature on germination and growth of ochratoxigenic isolates of *Aspergillus ochraceus* on a green coffee-based medium. *Int J Food Microbiol.* 2005;98:1–9.
35. Pitt JI, Hocking AD. Fungi and food spoilage. 3rd ed. Berlin, Germany: Springer; 2009.
36. Pitt JI, Taylor JW. (2441) Proposal to conserve the name *Aspergillus* (Fungi: Eurotiidae: Trichocomaceae) with a conserved type to maintain also the name *Eurotium*. *Taxon.* 2016;65:631–2.
37. Ratkowsky D, Ross T. Modelling the bacterial growth/no growth interface. *Lett Appl Microbiol.* 1995;20:29–33.
38. Ross K. Estimation of water activity in intermediate moisture foods. *Food Technol.* 1975;29:26–30.
39. Rosso L, Lobry JR, Flandrois JP. An unexpected correlation between cardinal temperatures of microbial growth highlighted by a new model. *J Theoret Biol.* 1993;162:447–63.
40. Rosso L, Robinson TP. A cardinal model to describe the effect of water activity on the growth of moulds. *Int J Food Microbiol.* 2001;63:265–73.
41. Samson RA, Visagie CM, Houbraken J, Hong SB, Hubka V, Klaassen CHW, et al. Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Stud Mycol.* 2014;78:141–73.
42. Sautour M, Dantigny P, Divies C, Bensoussan M. A temperature-type model for describing the relationship between fungal growth and water activity. *Int J Food Microbiol.* 2001;67:63–9.
43. Vesonder RF, Lambert R, Wicklow DT, Biehl ML. *Eurotium* spp. and echinulin in feed refused by swine. *Appl Environ Microb.* 1988;54:830–1.
44. Wheeler KA, Hocking AD, Pitt JI. Water relations of some *Aspergillus* species isolated from dried fish. *Trans Br Mycol Soc.* 1988;91:631–7.