

Seasonal Oviposition Activity of *Aedes aegypti* (Diptera: Culicidae) in San Miguel de Tucumán, Northwestern Argentina

Author(s): Giselle A. Rodríguez, Cecilia A. Veggiani Aybar, Guillermo L. Claps and Mercedes S. Lizarralde de Grosso

Source: Florida Entomologist, 98(4):1241-1243.

Published By: Florida Entomological Society

URL: <http://www.bioone.org/doi/full/10.1653/024.098.0435>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Seasonal oviposition activity of *Aedes aegypti* (Diptera: Culicidae) in San Miguel de Tucumán, northwestern Argentina

Giselle A. Rodríguez^{1,2,*}, Cecilia A. Veggiani Aybar^{1,2,*}, Guillermo L. Claps¹, and Mercedes S. Lizarralde de Grosso^{1,2}

Aedes aegypti (L.) (Diptera: Culicidae) is known as an efficient vector of dengue, chikungunya, and yellow fever viruses (PAHO 2011). This domestic urban mosquito shares an important connection with humans because the immature stages develop in water storage containers that are used in both domestic and peri-domestic habitats (Juliano & Lounibos 2005). In 1986, the re-infestation of Argentina by *Ae. aegypti* occurred in the northern provinces and later attained its current distribution, which includes the north, center and south of the country with the southernmost record from Patagonia (Almirón 2002; Grech et al. 2012; Diez et al. 2014). In 1991, the vector was found in some urban areas of Tucumán, and in 2009, a dengue epidemic resulted in the first autochthonous cases in the province (Augier 2000; Ramirez et al. 2013).

In tropical and subtropical regions, the seasonal distribution of mosquito vectors typically follows precipitation patterns, whereas in temperate regions, population size is regulated by the duration and intensity of winter, precipitation levels, and temperature (Rossi & Almirón 2004). Studies in Argentina have shown that oviposition patterns of *Ae. aegypti* correlate with temperature, humidity, and precipitation (Domínguez et al. 2000; Micieli & Campos 2003; Vezzani et al. 2004; Stein et al. 2005; Micieli et al. 2006); for Tucumán, this relationship has not been established. The aim of our study therefore was to observe the relative abundance and seasonal patterns of oviposition by *Ae. aegypti* in San Miguel de Tucumán.

The study was conducted from Sep 2012 to Jun 2013 in 5 neighborhoods: Oeste II (26°48'S, 65°15'W), El Bosque (26°48'S, 65°13'W), Ciudadela (26°49'S, W 65°13'W), Norte (26°49'S, 65°11'W), and Modelo (26°48'S, 65°14'W). All neighborhoods were located in San Miguel de Tucumán, Tucumán Province (Fig. 1). Oviposition by *Ae. aegypti* was monitored weekly by placing 8 ovitraps at each sampling site ($n = 40$ traps). The abundance and percentage of total eggs oviposited (for all sites for that sampling week) was determined for each sampling site and these data examined for correlation(s) with temperature, precipitation level, and other climatic variables.

We collected 9,476 eggs in San Miguel de Tucumán with the greatest abundance at Ciudadela followed by El Bosque, Oeste II, Modelo, and Norte. The main peak of egg abundance was in summer (Dec-Jan-Mar) and was followed by secondary peaks of smaller scale in spring (Nov) and autumn (Apr). The first eggs were observed in ovitraps in late

Oct 2012 and the last in mid-May 2013. Oviposition activity decreased towards autumn (May) and ceased with mean daily temperatures of approximately 15 °C; no eggs were collected in winter (Jun). Pearson's correlation coefficient revealed significant associations between oviposition activity and temperature ($r = 0.77$; $P < 0.008$) and precipitation level ($r = 0.76$; $P < 0.010$) in Ciudadela, and with temperature ($r = 0.63$; $P < 0.049$) in Oeste II (Fig. 2).

Although *Ae. aegypti* has previously been reported from Tucumán (Augier 1998), this is the first report of the oviposition patterns for this species in an epidemic area of the province. Other studies in Argentina have reported oviposition patterns similar to those observed in our study. For example, Micieli & Campos (2003) detected oviposition activity in Salta Province throughout the year, with peaks of abundance in Mar that correlated with precipitation levels. Stein et al. (2005) observed oviposition activity for *Ae. aegypti* from Oct to Jun in Chaco, which also correlated with the timing and amounts of precipitation. In Córdoba and Buenos Aires, oviposition by *Ae. aegypti* was observed from Oct/Nov to May/Jun, with peaks of activity associated with precipitation levels and temperature (Campos & Maciá 1996; Domínguez et al. 2000; Vezzani et al. 2004; Micieli et al. 2006).

Human activity could be a factor that affects the abundance of *Ae. aegypti* in Ciudadela and El Bosque. It is known that water-filled containers, improperly maintained swimming pools, unmanaged vegetation, and socio-economic factors favor proliferation of *Ae. aegypti*. Carbajo et al. (2006) reported the ready adaptation of this vector species to urban areas. Gürtler et al. (2009) and Fisher & Schweigmann (2010) noted the effects of vegetation, shade conditions, improperly maintained swimming pools, and housing and waste management in areas with high human densities as factors that maintain high infestation levels of mosquito vectors, including *Ae. aegypti*.

Our results show the presence of oviposition activity by *Ae. aegypti* in Tucumán and provide information that may be useful to future surveillance efforts targeted at *Ae. aegypti*. In this regard, it is important also to monitor changes in human density and activity in endemic areas, especially during times of high *Ae. aegypti* density when the potential for disease transmission may lead to a repetition of the 2009 dengue outbreak in Tucumán.

We are grateful to the families Rodríguez, Angeletti, Cano, and Veggiani, to Alvaro Galbán and Andrea Saracho, residents of the houses

¹Instituto Superior de Entomología "Dr. Abraham Willink", Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, Miguel Lillo 205, CP 4000, San Miguel de Tucumán, Tucumán, Argentina

²Consejo Nacional de Investigaciones Científicas y Técnicas, Crisóstomo Álvarez 722, CP 4000, San Miguel de Tucumán, Tucumán, Argentina

*Corresponding authors; E-mail G. A. R.: gisellerodriguez@csnat.unt.edu.ar; E-mail C. A. V. A.: ceciliaveggianiaybar@csnat.unt.edu.ar



Fig. 1. Aerial photograph of the study area in San Miguel de Tucumán, northwestern Argentina.




Fig. 2. Seasonal patterns of oviposition activity by *Aedes aegypti*, including temperature and precipitation in (A) Ciudadela and temperature in (B) Oeste II.

where ovitraps were placed. We thank Andrés Angeletti, Maximiliano Rodríguez, Miguel Pomares, and Gabriel Herrera for their support in the field work. Also thanks to Sofia Nanni and Lucia Krapovickas for the help in translating the manuscript.

Summary

Aedes aegypti (L.) (Diptera: Culicidae) has recolonized many areas of Argentina. The purpose of this study was to determine the seasonal patterns of oviposition activity for this mosquito vector and the association between this factor and climate in an urban area (Tucumán Province) with a history of dengue transmission. We collected 9,476 eggs by ovitraps in San Miguel de Tucumán with the highest abundance of eggs observed in the 2 neighborhoods Ciudadela and El Bosque. Peak oviposition activity by *Ae. aegypti* occurred in the summer. The number of eggs laid was positively correlated with temperature and precipitation levels.

Key Words: emerging tropical disease; arbovirus; ovitrap; *Aedes (Stegomyia) aegypti*

Sumario

Aedes aegypti (L.) (Diptera: Culicidae) ha recolonizando diversas áreas de la Argentina. El objetivo de este estudio fue determinar los patrones estacionales de la actividad de oviposición para este mosquito vector, y la asociación entre este factor y el clima en un área urbana (Provincia de Tucumán) con antecedentes de transmisión de dengue. 9.476 huevos fueron recolectados con ovitrampas en San Miguel de Tucumán, con la mayor abundancia de huevos en los barrios Ciudadela y el Bosque. Picos de actividad de oviposición de *Ae. aegypti* ocurrieron en el verano. El número de huevos puestos fue correlacionado positivamente con la temperatura y los niveles de precipitación.

Palabras Clave: enfermedades tropicales emergentes; arbovirus; ovitrampas; *Aedes (Stegomyia) aegypti*

References Cited

- Almirón W. 2002. Culicidae (Diptera) de la provincia de Córdoba, pp. 97-106. In Salomón O, [ed.], Actualizaciones en artropodología sanitaria Argentina, Enfermedades transmisibles, Publicación monográfica 2. Fundación Mundo Sano, Buenos Aires, Argentina.
- Augier L. 1998. Presencia de *Aedes aegypti* (Diptera: Culicidae) en Tucumán, Argentina. Revista de la Sociedad Entomológica Argentina 57: 66.
- Augier L. 2000. El Dengue, 14 pp. In Abdala SL, Fogliatta AS, Tereschuk ML, [eds.], Serie Monográfica y Didáctica, Facultad de Ciencias Naturales e IML, UNT, Tucumán, Argentina.
- Campos R, Maciá A. 1996. Observaciones biológicas de una población natural de *Aedes aegypti* (Diptera: Culicidae) en la provincia de Buenos Aires, Argentina. Revista de la Sociedad Entomológica Argentina 55: 67-72.
- Carbajo A, Curto S, Schweigmann N. 2006. Spatial distribution pattern of oviposition in the mosquito *Aedes aegypti* in relation to urbanization in Buenos Aires: southern fringe bionomics of an introduced vector. Medical and Veterinary Entomology 20: 209-218.
- Diez F, Bresler V, Quirán E, Rossi G. 2014. Niveles de infestación y nuevos registros de *Aedes aegypti* (Diptera: Culicidae) en la provincia de La Pampa, Argentina. Revista Chilena de Entomología 39: 67-72.
- Domínguez M, Ludueña-Almeida F, Almirón W. 2000. Dinámica poblacional de *Aedes aegypti* (Diptera: Culicidae) en Córdoba capital. Revista de la Sociedad Entomológica Argentina 59: 42-50.
- Fisher S, Schweigmann N. 2010. Seasonal occurrence of immature mosquitoes in swimming pools in Buenos Aires, Argentina. Journal of the American Mosquito Control Association 26: 95-98.
- Grech M, Visintín A, Laurito M, Estallo E, Lorenzo P, Rocca I, Korin M, Goya F, Ludueña-Almeida F, Almirón W. 2012. New records of mosquito species (Diptera: Culicidae) from Neuquén and La Rioja Provinces, Argentina. Revista de Saúde Pública 46: 387-389.
- Gürtler R, Garelli F, Coto H. 2009. Effects of a five-year citywide intervention program to control *Aedes aegypti* and prevent dengue outbreaks in northern Argentina. PLoS Neglected Tropical Diseases 3: e427.
- Juliano S, Lounibos L. 2005. Ecology of invasive mosquitoes: effects on resident species and human health. Ecology Letters 8: 558-574.
- Mieli M, Campos R. 2003. Oviposition activity and seasonal pattern of a population of *Aedes (Stegomyia) aegypti* (L.) (Diptera: Culicidae) in subtropical Argentina. Memórias do Instituto Oswaldo Cruz 98: 659-663.
- Mieli M, García J, Achinelly M, Martí G. 2006. Dinámica poblacional de los estadios inmaduros del vector del Dengue *Aedes aegypti* (Diptera: Culicidae): un estudio longitudinal (1996–2000). Revista de Biología Tropical 54: 979-983.
- PAHO. 2011. Preparedness and Response for Chikungunya Virus. Introduction in the Americas. http://www.paho.org/hq/index.php?option=com_docman&task=doc_download&Itemid=&gid=29607&lang=en (last accessed 3 Sep 2015).
- Ramírez F, Cuezzo R, Sarrouf E, Hilal FA, Alcaraz G, Nuñez Mrad R, Morales A, Enría D, Fabbri C, Caillou S, Echenique H, Flores I, Calli Flores R. 2013. Seroprevalencia de anticuerpos contra dengue en niños de uno a nueve años luego de una epidemia, Aguilares, Tucumán, 2009. Revista Argentina de Salud Pública 4: 18-22.
- Rossi G, Almirón W. 2004. Clave ilustrada para la identificación de larvas de mosquitos de interés sanitario encontradas en criaderos artificiales en la Argentina. Serie enfermedades transmisibles, Publicación monográfica 5. Fundación Mundo Sano, Buenos Aires, Argentina.
- Stein M, Oria G, Almirón W, Willener J. 2005. Fluctuación estacional de *Aedes aegypti* en Chaco, Argentina. Revista de Saúde Pública 39: 559-564.
- Vezzani D, Velázquez S, Schweigmann N. 2004. Seasonal pattern of abundance of *Aedes aegypti* (Diptera: Culicidae) in Buenos Aires City, Argentina. Memórias do Instituto Oswaldo Cruz 99: 351-356.