



Original article

Communities of anamorphic fungi on green leaves and leaf litter of native forests of *Scutia buxifolia* and *Celtis tala*: Composition, diversity, seasonality and substrate specificity



Natalia Allegrucci^{a,*}, Ana María Bucszinszky^a, Marcelo Arturi^b, Marta Noemí Cabello^{a,c}

^a Instituto de Botánica Spegazzini, FCNyM, UNLP, 53 # 477, 1900 La Plata, Provincia de Buenos Aires, Argentina

^b Laboratorio de Investigación en Sistemas Ecológicos y Ambientales (LISEA), Facultad de Ciencias Agrarias y Forestales, UNLP, diagonal 133, n° 469, 1900 La Plata, Provincia de Buenos Aires, Argentina

^c Comisión de Investigaciones Científicas, Provincia de Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 7 June 2013

Accepted 5 November 2013

Available online 21 March 2014

Keywords:

Microfungi

Xeric forest

Leaf litter

Decomposition

ABSTRACT

Background: Xeric forests dominated by two tree species, *Scutia buxifolia* (Rhamnaceae) and *Celtis tala* (Ulmaceae), are temperate, semi-deciduous wooded communities that represent the most abundant woodlands on the eastern plains of Buenos Aires Province, Argentina. The district of Magdalena has one of the most well-preserved native-forest areas, with an environmental heterogeneity that gives rise to the wide variability in the vegetation present.

Aims: The aim of this study was to analyze the species composition, diversity, seasonal variations, and substrate specificity of anamorphic fungi (Ascomycota) on the green leaves and in the leaf litter of native forests dominated by *Scutia buxifolia* and *Celtis tala* from Magdalena, Buenos Aires, Argentina.

Methods: In order to obtain the mycobiota of decomposition, seasonal samples of green leaves and leaf litter from both types of trees were collected over a two-year period. In the laboratory, the leaves were placed in a moist chamber and incubated at room temperature.

Results: A total of 100 species of anamorphic Ascomycota were identified in both forests. No significant variations were observed in the richness, diversity, or evenness of the fungal communities of the green leaves and leaf litter of both forests between seasons.

Conclusions: The species that characterized the fungal communities in the leaves of each of the trees were found to be different. The type of substrate had a stronger influence in determining the composition of the fungal community in both types of forests.

© 2013 Revista Iberoamericana de Micología. Published by Elsevier España, S.L.U. All rights reserved.

Comunidades de hongos anamórficos en hojas verdes y hojarasca de bosques nativos de *Scutia buxifolia* y *Celtis tala*: composición, diversidad, estacionalidad y especificidad del sustrato

RESUMEN

Antecedentes: Los bosques xerófilos dominados por las especies arbóreas *Scutia buxifolia* (Rhamnaceae) y *Celtis tala* (Ulmaceae) conforman comunidades boscosas semicaducas y templadas que constituyen los bosques nativos más abundantes de la provincia de Buenos Aires, Argentina. En el distrito de Magdalena constituyen una de las comunidades naturales de este tipo mejor conservada, con una heterogeneidad ambiental que da lugar a la gran variabilidad en la composición de la vegetación presente.

Objetivos: El objetivo del presente trabajo fue analizar la distribución de especies, la diversidad, las variaciones estacionales y la especificidad por el sustrato de hongos anamórficos (Ascomycota) presentes en las hojas verdes y en la hojarasca de los bosques nativos xerófilos dominados por las especies arbóreas *Scutia buxifolia* y *Celtis tala* en el distrito de Magdalena, Provincia de Buenos Aires, Argentina.

Métodos: Para obtener la micobiota descomponedora se tomaron muestras estacionales de hojas verdes y hojarasca de los 2 tipos de árboles durante un período de 2 años. En el laboratorio, las hojas se colocaron en cámara húmeda y se incubaron a temperatura ambiente.

Palabras clave:

Microhongos

Bosque xerófilo

Hojarasca

Descomposición

* Corresponding author.

E-mail address: nataliaallegrucci@yahoo.com (N. Allegrucci).

Resultados: Se identificaron 100 especies de anamorfos de la división Ascomycota en las hojas de ambos tipos de árboles. No se observaron diferencias en cuanto a la diversidad, equidad y riqueza de especies entre ambos tipos de hojas.

Conclusiones: Las especies que componen la comunidad fúngica de las hojas de *Scutia buxifolia* y *Celtis tala* es diferente. El tipo de sustrato que representan las hojas de cada especie arbórea condiciona la composición de la comunidad de anamorfos de Ascomycota por especie de hoja.

© 2013 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

In a forest, dead leaves constitute the main source of nutrients for the fauna and microorganisms with about 80% being degraded by microorganisms, mainly fungi.³¹ As the dead leaves are decomposed, the associated fungal communities undergo sequential alterations in their composition and substrate-degradation potential as a result of changes in the availability of nutrients, moisture, pH, and oxygen tension.

The litter and humus layers that overlie the mineral horizons of soils harbor various types of fungal communities. The litter is a transitional phase between the living plant community and the soil; thus, the mycotic composition of litter is closely linked to that of the plant community and contains many plant-inhabiting species as well as a distinctive assemblage of fungal taxa that are largely restricted to decaying plant material *per se*.¹⁴

The species composition and density of propagules are clearly affected by the sequence of changes occurring through the fragmentation, decomposition, and humidification of the plant detritus. Stable organic compounds such as waxes, lignins, and phenols—that may persist for hundreds of years—become incorporated in the mineral soil horizons and are, in turn, colonized by characteristic groups of fungal species.⁵

The phyllosphere fungi, that is the first group to colonize, have the advantage of gaining access to readily available organic compounds in freshly fallen leaves, before subsequent colonizing fungi alter the litter fall.³³ A sterilization of the leaves and consequent exclusion of the phyllosphere fungi was found to reduce the decomposition rate of the leaves⁴²; this change might reduce the substrate use by succeeding fungal decomposers,³² thus suggesting the potential relevance of prior colonization by phyllosphere fungi to the initial and subsequent stages of decomposition and fungal succession of a decaying leaf fall.

Many studies have been published on the ecology of fungi in forest litter and the identification of specific fungal communities.³⁶ Other investigations have focused on a possible nutritional specialization among saprotrophs.^{15,20,23,25–28,30,34,35,40,41} Several mycological studies have assessed the soil fungi^{6,10,11,13} and the mycorrhizae^{17–19} present in the native forests of *Scutia buxifolia* Reissek (coronillo) and *Celtis tala* Gillies ex Planch. (tala) in Magdalena (Buenos Aires province, Argentina). However, there are no studies of fungal communities present on green leaves and litter in both types of forest.

The aim of this investigation was to describe the composition, diversity, seasonal variations, and substrate specificity of the anamorphic saprotrophic fungi (Ascomycota) on the green leaves and in the leaf litter of native forests dominated by *S. buxifolia* and *C. tala* from Magdalena, Buenos Aires, Argentina (Biosphere Reserve, MAB-UNESCO).

Materials and methods

Study area

The study area (35°11' S, 57°17' W) is located in the Magdalena district, 20 km southeast of Magdalena town in the province of Buenos Aires, Argentina. Xeric forests dominated by two tree

species, *S. buxifolia* (Rhamnaceae) and *C. tala* (Ulmaceae), are temperate, semi-deciduous wooded communities that represent the most abundant woodlands on the eastern plains of Buenos Aires province, ranging from the banks of the Paraná River in the north-west to the suburbs of Mar Chiquita town in the southeast. The district of Magdalena has one of the most well-preserved native-forest areas, with an environmental heterogeneity that gives rise to the wide variability in the vegetation present. Wooded ranges run in parallel to La Plata River, where the forest grows on highly calcareous substrates derived from the sea's incursions and withdrawals during the Quaternary Period.³⁷

The climate of the region includes an annual mean temperature of 15.9 °C and annual mean precipitation of 885 mm, with peaks of rainfall during autumn and spring. The characteristics of each tree species provide contrasting features in each forest: *C. tala* is a deciduous species, whose senescent leaves begin to drop at the end of the summer, reaching the peak of litter fall in the autumn. These forests contain an abundant development of herbaceous vegetation. In contrast, *S. buxifolia* is an evergreen, with periods of leaf fall through replacement in both the autumn and spring. Thus, during these seasons, with their greater contribution of leaf litter, the forest has an appearance different from that of the *C. tala* forest, featuring constant shade and sparse herbaceous vegetation.⁴ Considered together, the leaves of these two species of trees contribute more than 90% of the total amount of leaf fall.

Sampling

Two areas were studied: one of pure *S. buxifolia* forest and the other of pure *C. tala* forest. A pure forest was defined as one consisting of 90% of the dominant tree species.³ Samples were collected seasonally during a two-year period: in autumn (May 2004, 2005), in winter (August 2004, 2005), in spring (October 2004, 2005), and in summer (February 2005, 2006).

In order to obtain the mycobiota of decomposition, 150 leaves were collected directly from the plant and another 150 leaves from the litter layer of each forest during each season. In the laboratory, the leaves were placed in a moist chamber (10 leaves in a 90-mm Petri dish) and incubated at room temperature.¹² The presence of each taxon on each leaf was recorded by means of the direct observation of both leaf surfaces. The taxonomic determination and preservation of the fungal species recorded were performed by isolation in plates with agar-containing media (malt-extract, potato-dextrose, corn-meal, or *S. buxifolia* and *C. tala* leaf litter extract agar).

Determination of substrate specificity

In order to evaluate whether the characteristics of the substrate consisting in the leaves of a given species of tree *per se* (here, the leaf litter from *S. buxifolia* or *C. tala*) was responsible for the species composition of the fungal communities rather than an effect at the forest level of the dominant species of tree present, 150 dead leaves of *S. buxifolia* in a pure *C. tala* forest and 150 dead leaves of *C. tala* in a pure *S. buxifolia* forest were collected seasonally. In the laboratory,

the leaves were placed in the moist chambers described above and incubated at room temperature. Weekly observations were made and the taxa present on each leaf recorded.

Data analysis

A leaf was considered as a natural sampling unit. After direct observation of both leaf surfaces, the taxa present on each leaf were recorded and their relative frequency determined as follows: (number of leaves where one species was recorded/total number of leaves analyzed) \times 100. These data were used to calculate the Shannon diversity index (H'), species richness (S), and species evenness (E).^{6,29} The differences in the values of these indices between seasons were evaluated through the analysis of variance (ANOVA). A principal-components analysis (PCA)⁹ was used to evaluate differences in composition between the fungal communities on the green leaves and on the leaf litter for the two forests. In order to determine whether there were differences between the fungal community developing on the *C. tala* leaves collected in the *S. buxifolia* forest and the one developing on the *S. buxifolia* leaves collected in the *C. tala* forest and also between the fungal communities found in the homogeneous leaf litter of both forests (*i.e.*, the *S. buxifolia* leaf litter in the *S. buxifolia* forest and the *C. tala* leaf litter in the *C. tala* forest), we applied a PCA using the data of species frequencies from the different sampling occasions.

Results

A total of 100 species of anamorphic Ascomycota were identified by traditional taxonomy on the green leaves and in the leaf litter from the *S. buxifolia* and *C. tala* forests (Table 1), with 25 of these species being recorded in the leaf litter of both forests.

The species identified belonged to the following orders: Hypocreales (27 taxa), Pleosporales (15 taxa), Helotiales (11 taxa), Xylariales (3 taxa), Eurotiales (6 taxa), Capnodiales (1 taxon), Chaetosphaeriales (1 taxon), Erysiphales (1 taxon), Orbiliales (2 taxa), and 33 species of anamorphs of uncertain position within the Ascomycota.

Cultures of these isolates were maintained in selective medium, being then deposited in the culture collection of the Instituto Spegazzini, UNLP, La Plata, Argentina (LPSC) (Table 1).

A total of 69 and 57 taxa were recorded in the *S. buxifolia* and *C. tala* forest respectively on the green leaves and in the leaf litter (Table 1).

In the *S. buxifolia*–forest data, axis 1 and axis 2 of the PCA (Fig. 1) explained a total of 57.78% of the total variability (at 41.34% and 16.43%, respectively). The first axis discriminated between the green-leaf and the leaf-litter data, with the former being located in the negative end and the latter in the positive end of the axis.

In the *C. tala*–forest data, the PCA diagram (Fig. 2) showed that both axis 1 (31.98%) and axis 2 (17.34%) together explained 49.32% of the cumulative variance. The second axis separated the samples of green leaf and leaf litter since species characteristic of green leaves such as *Tubercularia* sp., *Fusarium* sp., and *Acremonium* sp. could be observed, and their frequency were higher than that seen in leaf litter.

The PCA (Fig. 3) discriminated the *S. buxifolia* samples (of both the green and the dead leaves) from the *C. tala* samples. The first two axes explained 54.82% of the total variation, with the first component being the one that better separated the two forest types. Toward the positive end, the species on the *S. buxifolia* dead leaves were characterized by *Ciliochorella buxifoliae*, *Idriella tropicalis*, *Dactylaria obtriangularia*, *Circinotrichum chathamense*, *Gyrothrix verticiclada*, *Gyrothrix podosperma*, *Cylindrocladium colhounii*, *Chalara hughesii*, *Cir-*

cinotrichum maculiforme, and *Pestalotiopsis guepinii*, whereas toward the negative end, the taxa on the green and dead leaves of *C. tala* featured *Dematiocladium celtidis*, *Volutella ciliata*, *Tubercularia* sp., *Acremonium exigenum*, *Fusarium* sp., *Alternaria* sp. 1, and *Cylindrocarpon orthosporum*.

The greatest difference in fungal composition was observed among the leaf-litter samples from both species of forest, whose data tended to be located at opposite ends of the axes. In the *C. tala* forest, the data from the green leaves and those from the dead leaves were less separated from each other than were those two data sets from the *S. buxifolia* forest. Moreover, the data from the green leaves of both species tended to be located in the intermediate positions of axis 1, indicating a smaller difference in composition than that observed among the samples of leaf litter.

The richness (S), evenness (E), and diversity (Shannon–Wiener $H' \log 2$) of both forest types did not show significant differences between seasons ($P > 0.05$, Tables 2A, B and 3C, D). The values of these indices, however, were higher for the leaf litter than for the green leaves in both the years 2004 and 2005.

The variations in the diversity index were related to the variability of species richness. The evenness remained constant, indicating that changes in the species composition were present, but not in the distribution of the species in the samples.

The differences in the frequencies observed between the fungal species registered in samples of *S. buxifolia* and *C. tala* leaf litter collected from the opposite forest type and those recorded in the two leaf-litter samples removed from the same respective forest type were compared by means of PCA. The first component absorbed 35.24% of the total variation (Fig. 3) and was an informative discriminant of samples of *S. buxifolia* and *C. tala*. This discrimination allowed the separation of two groups in all instances. The first was located at the negative end of the axis, consisted in the *C. tala* green leaves and leaf litter along with *C. tala* leaf litter in a forest of *S. buxifolia*, and was characterized throughout by the following species: *D. celtidis*, *V. ciliata*, *Fusarium* sp., *Tubercularia* sp., *Alternaria* sp. 1, and *Acremonium exiguum*; the second was found at the positive end of the axis, consisted in the *S. buxifolia* green leaves and leaf litter along with *S. buxifolia* leaf litter in a forest of *C. tala*, and featured throughout *C. buxifoliae*, *D. obtriangularia*, *I. tropicalis*, *C. chathamense*, *G. verticiclada*, *G. podosperma*, *C. hughesii*, *Circinotrichum papakuriae*, and *C. colhounii*.

Discussion

The species that characterized the fungal communities of the forest of *S. buxifolia* were different from those of the forest of *C. tala*. Three taxa were recorded as new species in the leaf litter of *S. buxifolia*: *Thozetella buxifolia*,¹ *C. buxifoliae*, and *Koorchaloma scutiae*,² along with one new genus in the *C. tala* litter, represented by the species *D. celtidis*.⁸ According to Cai et al.,⁷ differences like these may indicate that the habitat and/or substrate could play a decisive role in the composition of these species, either through stimulating or inhibiting the growth of specific fungal species. The presence of different fungal communities is the result of the differing initial colonization mechanisms on the part of the species involved, and of the ability of those species both to explore the substrate and to compete for available resources.

The values of biodiversity indices were higher for the leaf litter than for the green leaves in both years. This result agrees with previous observations,²² where it was found that senescent leaves were significantly more densely colonized by endophytes when compared with young leaves. It was suggested that the increased density of colonization of older leaves was due to repeated reinfection of the leaf over time, probably from air-borne inoculum.

Table 1
Frequency of fungal species isolated from green leaves (gl) and leaf litter (ll) from the forest of *S. buxifolia* and *C. tala* during the 2004–2005 period. Species are ordered alphabetically within each order of Ascomycota.

Orders of Ascomycota	Species of anamorphs of Ascomycota	<i>S. buxifolia</i>		<i>C. tala</i>	
		gl	ll	gl	ll
Capnodiales	<i>Cladosporium cladosporioides</i> (Fresen.) G.A. de Vries 1952 (LPSC 788)	27.67	22.42	16.08	24.67
Chaetosphaeriales	<i>Thozetella buxifolia</i> Allegr., Cazau, Cabello & Aramb. 2004 (LPS 47406)	0	0.917	0	0
Erysiphales	<i>Oidium</i> sp.	0	0	0.167	0
Eurotiales	<i>Aspergillus niger</i> Tiegh. 1867	0	0	0	4.583
	<i>Aspergillus</i> sp.	0.167	0	0	0
	<i>Purpureocillium lilacinum</i> (Thom) Luangsa-ard, Pywel-Jones & Samson, 2011 (LPSC 862)	0	0	0.083	0
	<i>Penicillium</i> sp. 1	0	0	0.167	0.583
	<i>Penicillium</i> sp. 2	1.333	0	0	0
	<i>Penicillium</i> sp. 3	0	0.333	0	0
Helotiales	<i>Articulospora</i> sp.	0.083	0	0	0
	<i>Dactylaria obtriangularia</i> Matsush. 1975 (LPSC 849)	0.167	17.83	0	0.083
	<i>Dactylaria</i> sp. (LPSC 865)	0	0.5	0	0.083
	<i>Dactylella</i> sp.	0	0	0.083	1.25
	<i>Idriella lunata</i> P.E. Nelson & S. Wilh. 1956 (LPSC 852)	0.083	1.333	0	0.333
	<i>Idriella setiformis</i> R.F. Castañeda & G.R.W. Arnold 1985	0	3.417	0	0
	<i>Idriella tropicalis</i> Lunghini & Rambelli 1978 (LPSC 848)	0.583	19.92	0	0
	<i>Idriella variabilis</i> Matsush. 1971	0	0	0	4.75
	<i>Idriella</i> sp. 1	0	0.083	0	0
	<i>Idriella</i> sp. 2	0	0	0	2.917
	<i>Mirandina corticola</i> G. Arnaud ex Matsush. 1975	0	1.167	0	0.5
Hypocreales	<i>Acremonium fusidioides</i> (Nicot) W. Gams 1971 (LPSC 854)	0	0	0	2.583
	<i>Acremonium</i> sp.	0	0.25	11.83	9.75
	<i>Acrostalagmus luteoalbus</i> (Link) Zare, W. Gams & Schroers 2004 (LPSC 858)	0.083	0.083	6	1.75
	<i>Calcarisporium</i> sp.	0	0.083	0.083	0.083
	<i>Clonostachys compactiuscula</i> (Sacc.) D. Hawksw. & W. Gams 1975	0	0	0	0.917
	<i>Clonostachys cylindrospora</i> Höhn. 1952	2.5	0.833	0	0
	<i>Clonostachys rogersoniana</i> Schroers 2001 (LPSC 921)	0	0.167	0	5.75
	<i>Clonostachys setosa</i> (Vittal) Schroers 2001 (LPSC 855)	0	0.083	0	8.417
	<i>Cylindrocarpon orthosporum</i> (Sacc.) Wollenw. 1916	0.083	1	0.75	22
	<i>Cylindrocarpon</i> sp.	0	0	0	0.333
	<i>Cylindrocladium colhounii</i> Peeraly 1973 (LPSC 789)	1.5	13.67	0	0
	<i>Cylindrocladium</i> sp.	0	0	0.583	0.583
	<i>Dematiocladium celtidis</i> Allegr., Aramb., Cazau & Crous 2005 (LPSC 812)	0	0	11.08	22.25
	<i>Fusarium incarnatum</i> (Desm.) Sacc. 1886	0	0	0.083	0
	<i>Fusarium oxysporum</i> Schltld. 1824 (LPSC 870)	0.083	0.667	0	0.25
	<i>Fusarium solani</i> (Mart.) Sacc. 1881 (LPSC 868)	0	0	0.917	0.25
	<i>Fusarium sulphureum</i> Schltld. 1824 (LPSC 859)	0	0	0	0.083
	<i>Fusarium</i> sp. 1	0	0	18.58	5.5
	<i>Fusarium</i> sp. 2	1.083	0	0	0
	<i>Gliocladium penicillioides</i> Corda 1840	0.083	0	0	0
	<i>Kionochaeta ramifera</i> (Matsush.) P.M. Kirk & B. Sutton 1985 (LPSC 864)	0	3.833	0	0.333
	<i>Metarhizium anisopliae</i> (Metschn.) Sorokin 1883	0	0	0	1.083
	<i>Myrothecium lachatrae</i> Sacc. 1882 (LPSC 860)	0.083	0.5	0.5	16.5
	<i>Stilbella</i> sp.	0	0	0	0.083
	<i>Trichothecium roseum</i> (Pers.) Link 1809 (LPSC 1021)	0.083	0	0	0
	<i>Tubercularia</i> sp.	0.75	0	28.08	3.583
	<i>Volutella ciliata</i> (Alb. & Schwein.) Fr. 1832 (LPSC 807)	0.333	0.25	0.833	27.67
Orbiliiales	<i>Arthrobotrys conoides</i> Drechsler 1937 (LPSC 1012)	0.417	0	0.417	0.417
	<i>Arthrobotrys</i> sp. 2	0	0.083	0	0
Pleosporales	<i>Alternaria</i> sp. 1 (LPSC 846)	11.58	4.583	17.17	23.5
	<i>Alternaria</i> sp. 2	0	0	0	0.5
	<i>Alternaria</i> sp. 3	0.5	0	0	0
	<i>Alternaria</i> sp. 4	0	0	0	0.167
	<i>Cochliobolus spicifer</i> R.R. Nelson 1964	0	0	0	0.333
	<i>Curvularia lunata</i> (Wakker) Boedijn 1933 (LPSC 873)	0	0	0	0.333
	<i>Drechslera</i> sp.	0	0	0	0.167
	<i>Epicoccum nigrum</i> Link 1815 (LPSC 861)	1.333	0	0	1.75
	<i>Helicosporium</i> sp.	0	0	0.083	0
	<i>Helicosporium tenuis</i> Speg. 1910 (LPSC 1013)	0	0.417	0	0
	<i>Helicosporium griseum</i> Berk. & M.A. Curtis 1874	0	0.167	0	0
	<i>Periconia byssoides</i> Pers. 1801 (LPSC 809)	2.083	0.25	4.25	6.75

Table 1 (Continued)

Orders of Ascomycota	Species of anamorphs of Ascomycota	<i>S. buxifolia</i>		<i>C. tala</i>	
		gl	ll	gl	ll
Other anamorph Ascomycetes	<i>Sporidesmium adscendens</i> Berk. 1840	0	0.167	0	0
	<i>Stemphylium</i> sp.	0.5	0	0	0
	<i>Stemphylium vesicarium</i> (Wallr.) E.G. Simmons 1969	0	0	0.667	0.083
	<i>Camposporium</i> sp.	0	0.25	0	0
	<i>Chaetospermum</i> sp.	0	0.083	0	0
	<i>Chalara hughesii</i> Nag Raj & W.B. Kendr. 1974 (LPSC 869)	0.083	11.17	0	0
	<i>Chalara minima</i> W.P. Wu 2004	0.083	1.417	0	0
	<i>Chalara</i> sp.	0	1.167	0	1.167
	<i>Ciliochorella buxifoliae</i> Alleg. Eliades & Aramb. 2011 (LPSC 847)	5.25	67	0	0
	<i>Circinotrichum chathamense</i> McKenzie 1993	0.667	13.5	0	0
	<i>Circinotrichum maculiforme</i> Nees 1816	0	8.417	0	0
	<i>Circinotrichum olivaceum</i> (Speg.) Piroz. 1962	0	0	0	0.167
	<i>Circinotrichum papakurae</i> S. Hughes & Piroz	0	7.167	0	0
	<i>Dactylaria fusiformis</i> (Matsush.) de Hoog & Arx 1974	0	0.083	0	0.25
	<i>Fusariella atrovirens</i> (Berk.) Sacc. 1884	0	0	0.333	3.083
	<i>Gyrothrix citricola</i> Piroz. 1962	0.083	0	0	0.25
	<i>Gyrothrix flagella</i> (Cooke & Ellis) Piroz. 1962	0	2.917	0	0
	<i>Gyrothrix grisea</i> Piroz. 1962	0	0	0	0.167
	<i>Gyrothrix podosperma</i> (Cord.) S. Hughes. 1844	0.5	12.17	0	0.917
	<i>Gyrothrix verticillata</i> (Goid.) S. Hughes & Piroz. 1971	0.167	11.75	0	0
	<i>Gyrothrix verticillata</i> (Goid.) S. Hughes & Piroz. 1971	0	0	0	4.417
	<i>Koorchaloma scutiae</i> Alleg. Eliades & Aramb. 2011 (LPSC 791)	1.083	1.417	0	0
	<i>Matsushimomyces</i> sp.	0	1.167	0	0
	<i>Mycoleptodiscus</i> sp.	0.5	0.583	0	0
	<i>Mycoleptodiscus terrestris</i> (Gerd.) Ostaz. 1968	0	0.583	0	0
	<i>Nigrospora</i> sp.	0.083	0	0	0
	<i>Oidiiodendron</i> sp.	0.167	0	0	0
	<i>Olpitrichum macrosporum</i> (Farl. ex Sacc.) Sumst. 1911 (LPSC 863)	2.333	0.083	0	0
	<i>Polyscytalum pustulans</i> (M.N. Owen & Wakef.) M.B. Ellis 1976	0.75	0.333	0	9.667
	<i>Septonema</i> sp.	0.083	0.167	0	0
	<i>Septopatella</i> sp.	1.083	0	0	0
	<i>Solosympodiella</i> sp.	0	0.167	0	0
<i>Spermospora</i> sp.	0	0.25	0	0	
<i>Sporula harunganae</i> (Hansf.) M.B. Ellis 1968	0	0	0	0.083	
<i>Torula herbarum</i> (Pers.) Link 1809	0	0	0	0.583	
<i>Zygosporium oscheoides</i> Mont. 1842	0	0.25	0	0.083	
Xylariales	<i>Beltraniella</i> sp.	0	0.25	0	0
	<i>Beltraniella portoricensis</i> (F. Stevens) Piroz. & S.D. Patil 1970	0	2.083	0	0
	<i>Pestalotiopsis guepinii</i> (Desm.) Steyaert 1949 (LPSC 786)	18.83	6.5	0	0

LPSC: number Culture Collection of Instituto Spegazzini.

The assemblage of plant-associated species usually intergrades with the assemblages of soil species at the humus–soil interface.¹⁴ The leaf litter presented species that had reached that stage from the original green leaves. These species thus came from propagules that had already been present on the green leaves, together with other species that were able to colonize and decompose the leaf litter.²⁴ Accordingly, in *S. buxifolia*, 26 out of the 69 species (38%),

and in *C. tala* 19 out of the 57 species (33%) recorded, had already been present on the green leaves. Species such as *Cladosporium cladosporioides*, *Alternaria* sp. 1, and *Periconia byssoides* were among the most frequent, having been recorded in all the green-leaf and leaf-litter samples of both forests. In this study, we found no significant seasonal variations in species richness or diversity between the two forests studied.

Table 2

Values of richness (*S*), evenness (*E*), and Shannon–Wiener diversity (*H'* log 2) during the seasons (mean of three samples ± SD) in *Scutia buxifolia* green leaves and leaf litter (A and B).

Index	2004				2005			
	Autumn	Winter	Spring	Summer	Autumn	Winter	Spring	Summer
<i>(A) Green leaves</i>								
<i>S</i>	6a ± 4.6	8a ± 4.5	7a ± 5.8	10a ± 5	2a ± 1	1a ± 1	3a ± 2.3	3a ± 1.5
<i>E</i>	0.73a ± 0.05	0.73a ± 0.07	0.52a ± 0.45	0.65a ± 0.17	0.31a ± 0.27	0.33a ± 0.58	0.68a ± 0.13	0.65a ± 0.56
<i>H'</i> log 2	1.68a ± 1	2.16a ± 0.82	1.71a ± 1.5	2.14a ± 1.03	0.4a ± 0.39	0.33a ± 0.58	1.05a ± 0.65	1.17a ± 1.04
<i>(B) Leaf litter</i>								
<i>S</i>	20a ± 2.08	19a ± 6.03	16a ± 2.08	17a ± 3.06	16a ± 5.03	15a ± 3.79	11a ± 9.64	11a ± 1.73
<i>E</i>	0.85b ± 0.01	0.82ab ± 0.03	0.76a ± 0.06	0.8ab ± 0.03	0.85a ± 0.04	0.84a ± 0.02	0.59a ± 0.51	0.72a ± 0.04
<i>H'</i> log 2	3.68a ± 0.13	3.44a ± 0.49	3.03a ± 0.37	3.28a ± 0.32	3.32a ± 0.25	3.24a ± 0.22	2.36a ± 2.04	2.48a ± 0.3

Same letter indicates that the values do not differ significantly ($P > 0.05$) between the seasons.




Fig. 1. *S. buxifolia* forest. Principal Component Analysis (PCA) ordination of green leaves and leaf litter samples (Abbreviations: ▲ gl/04 = green leaves 2004, ▼ gl/05 = green leaves 2005, ■ ll/04 = leaf litter 2004, ◆ ll/05 = leaf litter 2005).

Litter degradation by the soil microbiota is dependent on the substrate composition in terms of organic as well as inorganic compounds and/or the availability of nutrients.^{16,21,38} With respect to nutrient requirements, the levels in organic-matter needed for its fungal degradation are also a function of the litter species with the habitat *cum* substrate, with the litter being also influenced by conditions such as the soil, the vegetation, and the climate.³⁹ The PCA applied to the characteristic of substrate specificity separated the samples into two groups: those from *S. buxifolia* and those from *C. tala*, thus indicating that the type of substrate was a more influential determinant of the composition

of the fungal community present than either the differences in the environmental conditions between the two forest types or the availability of fungal species within the environment. Rambelli et al.³⁶ assumed that in environments where conditions are optimal for microfungal development (e.g., in terms of humidity and temperature), saprotrophic specialization (e.g., in terms of primarily to nutritional characteristics and secondarily to the chemistry of the substrata).

In conclusion, the nature and amount of nutrients along with the physical characteristics of the substrate, the availability of water, the humidity, and the temperature determine the success, the




Fig. 2. *C. tala* forest. Principal Component Analysis (PCA) ordination of green leaves and leaf litter samples (Abbreviations: ■ gl/04 = green leaves 2004, ▲ gl/05 = green leaves 2005, ○ ll/04 = leaf litter 2004, □ ll/05 = leaf litter 2005).




Fig. 3. Principal Component Analysis (PCA) ordination of green leaves and leaf litter samples of *S. buxifolia* and *C. tala* forest (Abbreviations: ■ S.b. gl/04 = *S. buxifolia* green leaves 2004; ▲ S.b. ll/04 = *S. buxifolia* leaf litter 2004; ◆ S.b. ll/05 = *S. buxifolia* leaf litter 2005; ⊠ S.b. gl/05 = *S. buxifolia* green leaves 2005; ◇ C.t. gl/04 = *C. tala* green leaves 2004; ○ C.t. ll/04 = *C. tala* leaf litter 2004; □ C.t. gl/05 = *C. tala* green leaves 2005; ▽ C.t. ll/05 = *C. tala* leaf litter 2005; ● S.b. × C.t. = leaf litter of *S. buxifolia* collected from a *C. tala* forest; ♦ C.t. × S.b. = leaf litter of *C. tala* collected from a *S. buxifolia* forest).

Table 3

Values of richness (*S*), evenness (*E*), and Shannon–Wiener diversity (*H'* log₂) during the seasons (mean of three samples ± SD) in *Celtis tala* green leaves and leaf litter (C and D).

Index	2004				2005			
	Autumn	Winter	Spring	Summer	Autumn	Winter	Spring	Summer
<i>(C) Green leaves</i>								
<i>S</i>	8ab ± 4.5	10b ± 3.06	3a ± 1.53	8ab ± 2	7a ± 3.06	a	3a ± 3	8a ± 2
<i>E</i>	0.76a ± 0.08	0.79a ± 0.04	0.73a ± 0.24	0.74a ± 0.12	0.68a ± 0.12	a	0.55a ± 0.49	0.74a ± 0.12
<i>H'</i> log ₂	2.17b ± 0.5	2.55b ± 0.5	1.09a ± 0.1	2.22b ± 0.6	1.94a ± 0.8	a	1.12a ± 1	2.22a ± 0.6
<i>(D) Leaf litter</i>								
<i>S</i>	13a ± 2.52	13a ± 2.08	10a ± 1.15	11a ± 2.52	13b ± 2.9	6a ± 1	a	21c ± 4.93
<i>E</i>	0.87b ± 0.06	0.81ab ± 0.03	0.77ab ± 0.03	0.73a ± 0.11	0.83a ± 0.02	0.77a ± 0.09	a	0.85a ± 0.02
<i>H'</i> log ₂	3.17b ± 0.07	3.02ab ± 0.17	2.51a ± 0.15	2.57ab ± 0.61	3.1b ± 0.3	2a ± 0.35	a	3.69b ± 0.37

Same letter indicates that the values do not differ significantly ($P > 0.05$) between the seasons.

^a The lack of data of green leaves in winter and of leaf litter in spring in 2005 corresponds to the lack of leaves in those seasons.

degree of colonization, and the ultimate survival of anamorphic individuals that will constitute the eventual fungal-species composition of the community.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

This research is part of 11/N651 UNLP project. This work was performed with the financial support of CIC and CONICET. The authors thank Dr. Donald F. Haggerty, a retired career investigator and native English speaker, for editing the final version of the manuscript

References

- Allegrucci N, Arambarri AM, Cazau C, Cabello MN. *Thozetella buxifolia* a new hyphomycete from Argentina. *Mycotaxon*. 2004;90 Suppl. 2:275–9.
- Allegrucci N, Eliades LA, Cabello MN, Arambarri AM. New species of *Koorchaloma* and *Cliochoarella* from xeric forests in Argentina. *Mycotaxon*. 2011;115:175–81.
- Arturi MF (Tesis de Ph.D.) Regeneración de *Celtis tala* en el Noreste de la Provincia de Buenos Aires. Argentina: Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata; 1997.
- Arturi MF, Barrera MD, Brown AD. Caída y masa de hojarasca en bosques de *Celtis tala* Gill ex Planch y *Scutia buxifolia* Reiss del este de la provincia de Buenos Aires, Argentina. *Revista de la Facultad de Agronomía, La Plata*. 1996;101:151–8.
- Baath E. Microfungi in a clear-cut pine forest soil in central Sweden. *Can J Bot*. 1980;59:1331–7.
- Cabello MN, Arambarri AM. Diversity in soil fungi from undisturbed and disturbed *Celtis tala* and *Scutia buxifolia* forests in the eastern Buenos Aires province (Argentina). *Microbiol Res*. 2002;157:115–25.
- Cai LK, Frang JI, Hyde KD. Variation between freshwater and terrestrial fungal communities on decaying bamboo culms. *Antonie van Leeuwenhoek*. 2006;89:293–301.
- Crous PW, Allegrucci N, Arambarri AM, Cazau C, Groenewald JZ, Wingfield MJ. *Dematiocladium celtidis* gen. sp. nov. (Nectriaceae, Hypocreales) a new genus from *Celtis* leaf litter in Argentina. *Mycol Res*. 2005;109 Suppl. 7:833–40.
- Digby PGN, Kempton RA. Multivariate analysis of ecological systems. In: Chapman, Hall, editors. *Fungal ecology*. UK: London Press; 1987.
- Eliades LA, Bucsinszky AM, Cabello MN. Microbiota alcalofílica y alcalinotolerante en suelos de bosques xéricos en una localidad de la provincia de Buenos Aires, Argentina. *Bol Micol*. 2004;19:41–7.
- Eliades LA, Cabello MN, Voget CE. Contribution to the study of alkalophilic and alcali-tolerant Ascomycota from Argentina. *Darwiniana*. 2006;44 Suppl. 1:64–73.

12. Elíades LA, Voget CE, Arambarri AM, Cabello MN. Fungal communities on decaying saltgrass (*Distichlis spicata*) in Buenos Aires province (Argentina). *Sydowia*. 2007;59 Suppl. 2:227–34.
13. Elíades LA (Tesis de Ph.D.) Estudio de la micobiota alcalofílica y alcalinotolerante del suelo de los bosques de *Celtis tala* Gill (ex Planch) y *Scutia buxifolia* Reiss en el Partido de Magdalena, Provincia de Buenos Aires. Argentina: Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata; 2008.
14. Gerald F, Bills MC, Powell M, Thorn G. Saprobic soil fungi. In: Muller GM, Bills GF, Foster M, editors. Biodiversity of fungi: inventory and monitoring methods. USA: Elsevier Academia Press; 2004. p. 271–302.
15. Gusmão LFP, Grandi RAP, Milanez AL. Hyphomycetes from leaf litter of *Miconia cabussu* in the Brazilian Atlantic rain forest. *Mycotaxon*. 2001;79:201–13.
16. Hobbie SE. Interactions between litter lignin and soil nitrogen availability during leaf litter decomposition in a Hawaiian montane forest. *Ecosystems*. 2000;3:484–94.
17. Irrazabal GB (Tesis de Ph.D.) Estudio de las comunidades de hongos formadores de micorrizas arbusculares en bosques nativos de *Celtis tala*. Su influencia en el desarrollo temprano de las plántulas de tala en condiciones de invernadero. Argentina: Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata; 2007.
18. Irrazabal GB, Cabello MN, Velásquez MS. Infectividad y diversidad de hongos micorrízicos arbusculares de la rizósfera de los talares de Magdalena (Buenos Aires, Argentina). *Bol Micol*. 2004;19:49–57.
19. Irrazabal GB, Schalamuk S, Velásquez MS, Cabello MN. Especies de hongos formadores de micorrizas arbusculares. Nuevas citas para la República Argentina. *Bol Soc Argent Bot*. 2005;40:17–22.
20. Kabi OM, Rambelli A. Influencia del substrato sulla morfologia di *Beltrania rhombica* Penzig. *Micologia Italiana*. 1990;2 Suppl. 2:33–6.
21. Kassai K, Morinaga T, Horikoshi T. Fungal succession in the early decomposition process of pine cones on the floor of *Pinus densiflora* forests. *Mycoscience*. 1995;36:325–34.
22. Kumaresan V, Suryanarayanan TS. Endophyte assemblages in young, mature and senescent leaves of *Rhizophora apiculata*: evidence for the role of endophytes in mangrove litter degradation. *Fungal Divers*. 2002;9:81–91.
23. Laessle T, Lodge DJ. Three host-specific *Xylaria* species. *Mycologia*. 1994;86:436–46.
24. Lee OHK, Hyde KD. Phylloplane fungi in Hong Kong mangroves: evaluation of study methods. *Mycologia*. 2002;94 Suppl. 4:596–606.
25. Lodge DJ. Factors related to diversity of decomposer fungi in tropical forests. *Biodivers Conserv*. 1997;6:681–8.
26. Lodge DJ, Cantrell S. Fungal communities in wet tropical forests: variation in time and space. *Can J Bot*. 1995;73 Suppl. 1:S1391–8.
27. Lodge DJ, Fisher PJ, Sutton BC. Endophytic fungi of *Manilkara bidentata* leaves in Puerto Rico. *Mycologia*. 1996;88:733–8.
28. Lodge DJ, Laessle T. Host-specificity in *Camillea verruculospora*. *Mycologist*. 1995;9:152–3.
29. Magurran AE. Ecological diversity and its measurement. London: Croom Helm; 1988.
30. Mulas B, Rambelli A. Contribution to the study of the microfungi in the saprotrophic specialization in tropical forest litter. *Giornale Botanico Italiano*. 1995;129:1225–32.
31. Osono T. Phyllosphere fungi on leaf litter of *Fagus crenata*: occurrence, colonization, and succession. *Can J Bot*. 2002;80:460–9.
32. Osono T. Effect of prior decomposition of beech leaf litter by phyllosphere fungi on substrate utilization by fungal decomposers. *Mycoscience*. 2003;44:41–5.
33. Osono T. Colonization and succession of fungi during decomposition of *Swida controversa* leaf litter. *Mycologia*. 2005;97:589–97.
34. Pirozynski KA. Microfungi of Tanzania. I. Miscellaneous fungi on oil palm. II. New hyphomycetes. *Mycol Pap*. 1972;129:1–64.
35. Polishook JD, Bills GF, Lodge DJ. Microfungi from decaying leaves of two rain forest trees in Puerto Rico. *J Ind Microbiol*. 1996;17:284–94.
36. Rambelli A, Mulas B, Pasqualetti M. Comparative studies on microfungi in tropical ecosystems in Ivory Coast forest litter: behaviour on different substrata. *Mycol Res*. 2004;108:325–36.
37. Sánchez RO, Ferrer JA, Duymovich OA, Hurtado MA. Estudio pedológico integral de los Partidos de Magdalena y Brandsen (Provincia de Buenos Aires). *Anales del LEMIT, serie II N° 310*. Ministerio de Obras Públicas de la Provincia de Buenos Aires; 1976. p. 1–123.
38. Saparrat MCN, Arambarri AM, Balatti PA. Growth response and extracellular enzyme activity of *Ulocladium botrytis* LPSC 813 cultured on carboxymethylcellulose under a pH-range. *Biol Fertil Soils*. 2007;4:383–6.
39. Sariyildiz T, Anderson JM. Interactions between litter quality, decomposition and soil fertility: a laboratory study. *Soil Biol Biochem*. 2003;35:391–9.
40. Subramanian CV, Vittal BPR. Studies on litter fungi. II. Fungal colonization of *Atlantia monophylla* Corr. Leaves and litter. *Nova Hedwigia*. 1979;63:361–9.
41. Subramanian CV, Vittal BPR. Studies on litter fungi. IV. Fungal colonization of *Gymnosporia emarginata* leaves and litter. *Trans Mycol Soc Jpn*. 1980;21:339–44.
42. Tanner EVJ. The decomposition of leaf litter in Jamaican montane rain forest. *J Ecol*. 1981;69:263–75.