

Atmospheric Concentrations of New Persistent Organic Pollutants and Emerging Chemicals of Concern in the Group of Latin America and Caribbean (GRULAC) Region

Cassandra Rauert,[†] Tom Harner,^{*,†} Jasmin K. Schuster,[†] Anita Eng,[†] Gilberto Fillmann,[‡] Luisa Eugenia Castillo,[§] Oscar Fentanes,^{||} Martín Villa Ibarra,[⊥] Karina S. B. Miglioranza,[#] Isabel Moreno Rivadeneira,[▽] Karla Pozo,[○] and Beatriz Helena Aristizábal Zuluaga[◆]

[†]Air Quality Processes Research Section, Environment and Climate Change Canada, Toronto, Ontario M3H 5T4, Canada

[‡]Universidade Federal do Rio Grande, Instituto de Oceanografia, Rio Grande - RS, Brazil

[§]Central American Institute for Studies on Toxic Substances (IRET), Universidad Nacional, Heredia, Costa Rica

^{||}CGCSA/INECC, Ciudad de Mexico, Mexico

[⊥]Instituto Tecnológico Superior de Cájeme, Cájeme, Sonora México

[#]Universidad Nacional Mar del Plata, IIMyC-CONICET, Mar del Plata, Argentina


[▽]Laboratorio de Física de La Atmósfera, Instituto de Investigaciones Física, UMSA, La Paz, Bolivia

[○]Facultad de Ingeniería y Tecnología, Universidad San Sebastián, Lientur 1457, Concepción 4080871, Chile

[◆]Universidad Nacional de Colombia, Manizales, Colombia

S Supporting Information

ABSTRACT: A special initiative was run by the Global Atmospheric Passive Sampling (GAPS) Network to provide atmospheric data on a range of emerging chemicals of concern and candidate and new persistent organic pollutants in the Group of Latin America and Caribbean (GRULAC) region. Regional-scale data for a range of flame retardants (FRs) including polybrominated diphenyl ethers (PBDEs), organophosphate esters (OPEs), and a range of alternative FRs (novel FRs) are reported over 2 years of sampling with low detection frequencies of the novel FRs. Atmospheric concentrations of the OPEs were an order of magnitude higher than all other FRs, with similar profiles at all sites. Regional-scale background concentrations of the poly- and perfluoroalkyl substances (PFAS), including the neutral PFAS (n-PFAS) and perfluoroalkyl acids (PFAAs), and the volatile methyl siloxanes (VMS) are also reported. Ethyl perfluorooctane sulfonamide (EtFOSA) was detected at highly elevated concentrations in Brazil and Colombia, in line with the use of the pesticide sulfluramid in this region. Similar concentrations of the perfluoroalkyl sulfonates (PFAS) were detected throughout the GRULAC region regardless of location type, and the VMS concentrations in air increased with the population density of sampling locations. This is the first report of atmospheric concentrations of the PFAAs and VMS from this region.



INTRODUCTION

The capacity for the Global Monitoring Plan (GMP) to evaluate the effectiveness of the United Nations Environment Programme's (UNEP) Stockholm Convention on persistent organic pollutants (POPs) is limited by the lack of monitoring information on POPs from developing regions. The GMP aims to provide comparable monitoring data from all United Nations (UN) regions;¹ however, there is a substantial gap from developing regions such as the Group of Latin American and Caribbean countries (GRULAC).^{2–4} This limits the information provided to risk assessments on chemicals of concern; the ability to evaluate regional and global environmental trans-

port^{2,5} and prevents global spatial and temporal trends being developed, thus limiting the ability to evaluate the effectiveness of restrictions.⁶

Passive air sampling is a particularly useful methodology for atmospheric monitoring in developing regions^{7,8} because remote locations can be accessed with no electricity requirements, and they are cost-effective and easy to deploy. As such, a

Received: February 21, 2018

Revised: May 11, 2018

Accepted: May 30, 2018

Published: May 30, 2018



55 special initiative was implemented in 2012 by the Global
56 Atmospheric Passive Sampling (GAPS) network, with support
57 from UNEP, to address the lack of information on emerging
58 contaminants and candidate, and new POPs in the GRULAC
59 region. The GAPS network is a global passive sampling
60 monitoring network that contributes unique atmospheric data
61 on legacy and new and emerging POPs to the GMP.⁹ The
62 GAPS network is run by a central laboratory at Environment
63 and Climate Change Canada (ECCC) and contributes to
64 international regulatory programs such as the Stockholm
65 Convention and also contributes to domestic Canadian risk
66 assessments and management programs under the Chemicals
67 Management Plan (CMP).

68 The core GAPS sampling campaign includes a few sites in
69 the GRULAC region,^{9,10} and in this special initiative, additional
70 sites were added to provide a more-comprehensive monitoring
71 reach. The first regional-scale monitoring data for polychlori-
72 nated dioxins and furans in the region has been reported from
73 this initiative.¹¹ The second phase of this study has focused on
74 emerging contaminants and candidate and new POPs. The first
75 atmospheric data for organophosphate ester (OPE) flame
76 retardants (FRs) in the GRULAC region has since been
77 reported.² With the conclusion of the second phase of this
78 special initiative, a wide range of new and emerging POPs have
79 been monitored, including the polybrominated diphenyl ethers
80 (PBDEs), OPEs, a range of other brominated and chlorinated
81 alternative FRs (novel FRs), a range of poly- and perfluoroalkyl
82 substances (PFAS), and the volatile methyl siloxanes (VMS).

83 PBDEs have been used historically as FRs in a range of
84 consumer products including foam, electronic, and textile
85 products. They are produced in three different commercial
86 formulations: the PentaBDE, OctaBDE, and DecaBDE
87 formulations, which contain different contributions of the
88 PBDE congeners. Due to concerns over their persistency and
89 toxicity, all three technical formulations have now been added
90 to the Stockholm Convention for reduction with restricted
91 uses.¹² With the restrictions imposed on the use of PBDEs, the
92 use of alternative FRs has increased,^{13,14} including the OPEs
93 and a range of other brominated and chlorinated novel FRs.
94 These alternative FRs are also detected regularly in the
95 environment^{13,15} and, due to concerns over its persistency and
96 toxicity, hexabromocyclododecane (HBCD) was listed in
97 Annex A of the Stockholm Convention in 2013.¹⁶ The OPEs
98 have numerous applications other than as FRs, including as
99 plasticizers, in hydraulic fluid and antifoaming agents.¹⁴ OPE
100 atmospheric concentrations are often reported in levels that are
101 at least an order of magnitude higher than the PBDEs.^{10,14}
102 There are concerns over the persistency of the OPEs, and in
103 Canada, tris(2-chloroethyl) phosphate (TCEP) is prohibited in
104 foam products intended for children under the age of three.¹⁷
105 These alternative FRs are detected regularly in remote
106 locations, thus showing their potential for long-range
107 atmospheric transport.^{10,18–20}

108 PFAS are surfactants with applications including as stain
109 repellants in fabric and grease-proof paper, in fluoropolymer
110 manufacture and in aqueous film-forming foams.²¹ The
111 perfluoroalkyl acids (PFAAs) have raised concerns due to
112 their persistency and toxicity, and perfluorooctanesulfonate
113 (PFOS) and its salts and precursor compound (perfluorooctane
114 sulfonyl fluoride) were added to the Stockholm Convention in
115 2009 for restriction of production and use, with specific
116 exemptions.²² Perfluorohexanesulfonate (PFHxS) and perfluor-
117 ooctanoic acid (PFOA) are currently under review for addition

to the Stockholm Convention.²³ Also of concern are the neutral
PFAS (n-PFAS), which are known to degrade in the
environment to the toxic PFAAs.²¹ In the GRULAC region,
Brazil has registered the use of PFOS, its salts, and PFOSF in
the Register of Acceptable purposes for the use as an
intermediate in the production of the pesticide sulfluramid.²⁴
The active ingredient of sulfluramid is ethyl perfluorooctane
sulfonamide (EtFOSA), a known precursor for PFOS.²⁵

The VMS are used in the manufacture of silicone polymers,
as surfactants and lubricants, and also in a wide range of
personal-care products.²⁶ They are classified as high-produc-
tion-volume chemicals, and highly elevated concentrations (as
compared with the FRs and PFAS) are detected regularly in the
environment.^{27,28} Concerns arise over the toxicity of the VMS
to aquatic organisms,²⁹ and more monitoring is required to
understand their distribution in the environment. The PFAS
and VMS are also regularly detected in remote locations, again
demonstrating the persistency and potential for the long-range
transport of these chemicals.^{28,30–32}

Previous atmospheric monitoring studies of these chemicals
of concern tend to focus on North America and Europe, and
more monitoring information is needed from developing
regions to provide a complete picture of global distribution
and transport. This special initiative provides unique regional
monitoring data on emerging chemicals of concern and new
and candidate POPs from the GRULAC region, and this is the
first reporting of atmospheric concentrations of the PFAAs and
VMS from across this region.

■ MATERIALS AND METHODS

Sample Deployment. This special initiative included 9
sites covering 7 countries: Mexico ($n = 2$), Costa Rica ($n = 1$),
Colombia ($n = 1$), Brazil ($n = 2$), Bolivia ($n = 1$), Argentina (n
 $= 1$), and Chile ($n = 1$). The majority of sites were classified as
background sites ($n = 5$), with 3 urban and 1 agricultural site
included. Site classifications follow the guidelines under the
guidance document for the global monitoring plan.³³ Rio
Gallegos falls under the category of suburban following these
guidelines and, in this study, is referred to as an urban site.
Further site details are provided in Table S1, and Figure 1 maps
the location of each sampling site. At each site polyurethane
foam passive air samplers (PUF-PAS) were deployed for
sampling periods of 3 months, providing 4 sampling quarters
throughout a sampling year, referred to as Q1, Q2, Q3, and Q4.
The PUF-PAS were deployed during 2014 and 2015, and the
deployment details are listed in Tables S2–S5. Sorbent-
impregnated PUF passive air samplers (SIP-PAS) were also
deployed for one sampling quarter in 2015. The SIP-PAS have
a greater uptake capacity than the PUF-PAS,³⁴ and for more-
volatile compounds that form equilibrium quickly with the
PUF-PAS, the SIP-PAS will continue to sample in the linear
sampling phase, providing time-averaged concentrations of
these analytes. The PUF-PAS and SIP-PAS at Tapanti (Costa
Rica) were deployed for a full year (June 2015–June 2016) due
to restrictions accessing the site. The SIP-PAS at Chacaltaya
(Bolivia) was deployed in 2016 (not 2015) but was still
deployed in Q2 (April–July). Deployment details of the SIP-
PAS are listed in Table S6.

PUF-PAS and SIP-PAS were precleaned, stored in glass jars,
and shipped to the sampling sites following previously
published methods.^{11,27} The samples were deployed in
double-domed sampling chambers (TE-200, Tisch Environ-
mental, Cleves, OH), and the deployment and collection



Figure 1. Location of the nine sites included in the GRULAC study. The location type is listed below the site name. BA, background; AG, agricultural; and UR, urban.

analytes are listed in Table S7. The PUF-PAS were extracted and processed following previously reported methods.¹⁰ The sample extracts were analyzed for the PBDEs and novel FRs with gas chromatography–tandem quadrupole mass spectrometry (GC–MS/MS) and analyzed for the OPEs and the HBCD diastereomers using ultra-performance liquid chromatography–tandem quadrupole mass spectrometry (UPLC–MS/MS). Analysis details have been previously reported in Rauert et al.¹⁰

The SIP-PAS were extracted and analyzed for the n-PFAS, PFAAs, and VMS. Target analytes are listed in Table S8. The SIP-PAS were extracted and processed following previously reported methods.²⁸ The sample extracts were analyzed for the n-PFAS and VMS with GC–MS and analyzed for the PFAAs with UPLC–MS/MS. Analysis details have been reported previously in Rauert et al.²⁸

Quality Assurance and Quality Control. Air concentrations (pg/m^3) were derived from the mass of the target analyte detected on the sample (pg) divided by an effective air sampling volume. The air volume was determined using the GAPS template.³⁵ The OPEs and novel FRs were assumed to stay in the linear sampling phase during deployment, and air sampling volumes were calculated as the number of days the sample was deployed multiplied by a sampling rate of $4 \text{ m}^3/\text{day}$, as described in Rauert et al.¹⁰ Field blanks were sent to each site and consisted of an additional precleaned PUF-PAS in 2014 and 2015 and a SIP-PAS in 2015. The field blanks were collected following the same protocol as the sample deployment without actual deployment. Further details on field blanks and collection are provided in Schuster et al.¹¹ The field blanks were extracted and analyzed following the same procedures as the samples. Concentrations (ng/PAS) of target analytes detected in each field blank are listed in Tables S9–S15. Instrument detection limits (IDLs) were calculated as the concentration of a peak that would produce a signal-to-noise

180 followed procedures previously published.¹¹ After collection,
181 the samples were shipped back to ECCC for analysis.

182 **Extraction and Analysis.** The PUF-PAS were extracted
183 and analyzed for the PBDEs, novel FRs, and OPEs. Target




Figure 2. Box-and-whisker plots of target analyte concentrations (pg/m^3) in 2014 (red) and 2015 (blue) in the GRULAC region. The line indicates the median concentration; boxes correspond to the interquartile range with the upper and lower edges designating the 75th and 25th percentiles, respectively. The whiskers represent the maximum and minimum concentrations. The diamond indicates the mean concentration and the orange line indicates the MDL. $\Sigma_{18}\text{OPEs}$, sum of all OPEs; $\Sigma_{14}\text{PBDEs}$, sum of all PBDEs; $\Sigma_{15}\text{novel FRs}$, sum of all novel FRs; $\Sigma_2\text{FTOH}$, sum of 8:2 FTOH and 10:2 FTOH; $\Sigma_4\text{FOSA/Es}$, sum of all FOSA and FOSEs; $\Sigma_3\text{PFASs}$, sum of PFBS, PFHxS, and PFOS; and $\Sigma_4\text{VMS}$, sum of D3, D4, D5, and D6. Where an analyte was not detected, $1/2$ MDL was used.




Figure 3. Map of mean concentrations (pg/m^3) of TCEP, TCPP, TPhP, TBEP, and TDCPP in the GRULAC region in 2014 and 2015. The three urban locations are marked with an asterisk. NS indicates that a sample was not deployed in that year. <MDL indicates that all analytes were below detection limits in that year.

ratio of 3:1. Method detection limits (MDL) were calculated as the mean mass in the field blanks plus three times the standard deviation. For the OPEs, PBDEs, and novel FRs, MDLs were calculated as the mean concentration in field blanks, to keep data sets comparable with the global FR data from Rauert et al.¹⁰ Where a target analyte was detected regularly in field blanks, samples were blank corrected by subtracting the mean mass (in nanograms) from all the field blanks; hence, it is possible for reported concentrations to be below the MDLs. Limits of quantitation (LOQ) were calculated as the concentration of a peak that would produce a signal-to-noise ratio of 10:1 and inserted for MDLs where an analyte was not detected in the field blanks. For calculating average values, $1/2$ MDL or $1/2$ LOQ were inserted when an analyte was not detected. To convert MDLs and LOQs to pg/m^3 , the average air sampling volume (m^3) was applied.

All analytes, except for the novel FRs, were surrogate recovery corrected and the mean recoveries are listed in Table S16. As described in Rauert et al.¹⁰ an enhancement in the surrogate recoveries was observed for the PBDEs and hence they were not used to recovery correct the novel FRs. To improve this methodology, future analyses in the GAPS network will use mass labeled versions of the novel FRs (and the PBDEs) and adopt surrogate recovery correction for the novel FRs. However, for these samples, this option was not available, and we note that this increases the uncertainty with the analysis of the novel FRs. Statistical analyses (two-sample t tests assuming unequal variances) were conducted using Microsoft Excel 2010.

RESULTS AND DISCUSSION

PUF-PAS. The PUF-PAS were analyzed for the three FR classes. Concentrations of OPEs, PBDEs, and novel FRs from the 2014 deployment have been reported previously,^{2,10} and this is the first reporting of atmospheric concentrations of these analyte classes from 2015. Concentrations from each sample (2014 and 2015) are listed in Tables S17–S22. Figure 2 depicts box and whisker plots of concentrations of all the analyte

classes monitored (including the PFAS and VMS monitored with SIP-PAS), showing the variation in atmospheric concentrations between the different classes.

Organophosphate Esters. Total OPE concentrations (Σ_{18} OPEs) were similar between both sampling years (Figure 2), with Σ_{18} OPE concentrations in each sample ranging 90–1850 pg/m^3 in 2014 and 60–2200 pg/m^3 in 2015, and were not significantly different between the two years ($p > 0.05$). The urban site of Concepción (Chile) was the exception, where elevated concentrations were detected in 2015 of 730–7050 pg/m^3 (Figure 3). The other two urban sites of São Luis (Brazil) and Rio Gallegos (Argentina) had concentrations in line with the background sites in both sampling years.

In this GRULAC study five OPEs were detected the most frequently and in the highest concentrations (TCEP, tris(chloroisopropyl) phosphate (TCPP), triphenyl phosphate (TPhP), tris(2-butoxyethyl) phosphate (TBEP), and tris(1,3-dichloro-2-propyl) phosphate (TDCPP)). Figure S1 presents box and whisker plots of the concentrations of the five dominating OPEs in 2014 and 2015. The mean profiles of the five dominating OPEs at each site are mapped in Figure 3.

The profiles of the OPEs were similar at most sites in both 2014 and 2015, although variations were observed at Concepción and Celestún (Mexico). At Concepción, elevated concentrations of TBEP were detected in 2014; however, in 2015, the profile was similar to the other sites, although it is noted only two PUF-PAS were deployed in 2014. At Celestún (Mexico), tri-*n*-butyl phosphate (TnBP) was detected at elevated concentrations in Q1 of 2015 but was not detected at this site in 2014. The other OPEs at Celestún were in line with concentrations detected at the other background sites. As discussed later, one challenge of passive air sampling studies is deployment at a location that avoids possible point sources, which may be an influence at these sites.

Seasonal differences in OPE levels in air were not observed in 2014 or 2015. This is in line with other global regions monitored in the regular GAPS sampling campaign, in which seasonal differences were not observed in 2014.¹⁰ The

concentrations of OPEs in the GRULAC region in 2014 were significantly higher than concentrations seen in North America and Europe in 2014 in the core GAPS network ($p < 0.03$), as discussed in more detail in Rauert et al.¹⁰ Further monitoring is needed in the GRULAC region to determine if this is a trend or influenced by the high variability in concentrations at the sampling sites.

Polybrominated Diphenyl Ethers. The atmospheric concentrations of the PBDEs were at least an order of magnitude lower than the OPEs (Figure 2). The Σ_{14} PBDE concentrations were not significantly different ($p > 0.05$) between the two sampling years and ranged 0.40–18 pg/m³ in 2014 and 0.86–20 pg/m³ in 2015. The agricultural site of Sonora (Mexico) was the exception with highly elevated concentrations in both years (ranging 24 to 125 pg/m³), possibly associated with inputs from buildings near the sampling site.¹⁰ This will be evaluated in future sampling campaigns at this site.

Figure S2 presents box-and-whisker plots of the concentrations of the sum of the PBDE congeners that dominate in the PentaBDE, OctaBDE, and DecaBDE formulations, and the main congeners in the PentaBDE formulation (BDE28, 47, 99, 100) dominated the profiles at all 9 sites. The mean profile of the congeners in the PentaBDE formulation at each site is mapped in Figure S3. BDE209 was detected at three sites in this study, Sonora in 70% of samples, Celestún (Mexico) in 1 out of 8 samples, and Concepción (Chile) in 33% of samples. BDE209 has a higher detection limit than the other congeners, which contributes to the lower detection frequency of this congener, although the detected concentrations were lower than seen in North America in the core GAPS samples from 2014.¹⁰

PBDEs have been reported previously by Pegoraro et al.³⁶ from atmospheric samples collected in 2014 at Mar del Plata (Argentina) and off the coast of Argentina. Levels of PBDEs measured in air during this cruise were lower than in the GRULAC study and enriched in BDE209 (70% contribution). The lower levels of PBDEs in the off shore samples reflect their greater distance from point sources of PBDEs, whereas the higher proportion of BDE209 indicates a different source type. PBDEs have also been reported from PUF-PAS deployed in three Chilean cities in 2008–2009, including Concepción. Total PBDE concentrations at Concepción were similar between this Chilean study³⁷ (0.5–20 pg/m³) and this GRULAC study. BDE-209 was the dominant congener at the urban and industrial sites, suggesting there may be different sources between the sampling locations in the respective studies.

Seasonal differences of the PBDEs were not observed at these sites, which is in line with observations at other global regions sampled in the GAPS Network.¹⁰ As discussed in Rauert et al.¹⁰ concentrations of PBDEs in the GRULAC region in 2014 were significantly lower than concentrations from the core GAPS sites in North America in 2014. This is not surprising considering the higher historical use of PBDEs in North America,³⁸ as compared to developing regions.

Novel Flame Retardants. The novel FRs were detected less frequently and in generally lower concentrations than the PBDEs in both 2014 and 2015 (Figure 2). The profiles varied between sites and years, although 2-ethylhexyl-2,3,4,5-tetrabromobenzoate (EH-TBB), hexabromobenzene (HBB), and dechlorane 602 (Dec 602) generally dominated the profiles. Figure S4 presents box-and-whisker plots of the concentrations of EH-TBB, HBB, and Dec 602 in 2014 and 2015, also

highlighting that the detected concentrations were in the same range as the MDLs. Total novel FR (Σ_{15} novel FR) concentrations (excluding HBCD) ranged <0.1–13 pg/m³ in 2014 and <0.1–30 pg/m³ in 2015 except for an anomalously high concentration of EH-TBB and bis(2-ethyl-1-hexyl) tetrabromophthalate (BEH-TEBP) at Concepción (Chile) in Q4 of 2015 (2880 and 2890 pg/m³, respectively). This sample may have been contaminated during processing and is removed from further discussion. The Σ_{15} novel FR concentrations in 2015 were significantly higher than in 2014 ($p = 0.01$), primarily due to the higher detection frequencies of both EH-TBB and HBB in 2015 (86–93% in 2015 and 14–24% in 2014). Figure S5 maps the average concentrations of EH-TBB, HBB, and Dec 602 at the GRULAC sites, clearly showing the elevated concentrations of EH-TBB in 2015. Due to the low detection frequencies of the novel FRs, and the uncertainty due to detected concentrations being close to the detection limits, more monitoring is required to determine if an increasing trend of EH-TBB is being observed in this region.

Concepción was the only site where HBCD was detected, and only the γ -HBCD diastereomer was detected. Concentrations at this site in 2014 were the highest of all the 44 global sites in the GAPS network monitored in 2014.¹⁰ Concentrations in 2015, meanwhile, were an order of magnitude lower than in 2014. Figure S6 graphs the concentrations of γ -HBCD detected in 2014 and 2015, the diastereomer profile expected in the HBCD technical formulation and the profile expected after thermal rearrangement of the diastereomers.³⁹ There is no obvious explanation for why only the γ -HBCD diastereomer was detected and further monitoring will be conducted at this site.

Seasonal differences of the novel FRs were not observed at these sites in either 2014 or 2015. This is again in line with other global regions monitored in the regular GAPS sampling campaign, where seasonal differences were not observed in 2014.¹⁰ The low detection frequencies of these compounds, however, make comparisons difficult, and this is a challenge that should be addressed in future passive air sampling campaigns, as discussed further in Rauert et al.¹⁰

Flame-Retardant Summary. As the GRULAC region is developing, it is expected to become a major e-waste and waste electrical and electronic equipment (WEEE) producer in the next 10 years.^{38,40} Brazil in particular has a rapidly expanding market and is undergoing a “boom” in e-waste generation.⁴¹ The importation of e-waste is prohibited in Brazil under Brazil’s National Solid Waste Policy;⁴² however, Mexico is expected to receive significant quantities of e-waste, exported from developed countries.⁴⁰ Electronic equipment can contain high percentage levels of FRs, and the dismantling of e-waste (whether produced domestically or imported) is a major contributor of FRs to the surrounding environment as well as to the workers and the surrounding communities.⁴⁰ Limited controls over the disposal of e-waste in developing regions, where the majority of disposal areas are open area dumping sites, also contributes to environmental levels. Therefore, the GRULAC region is a key region for future monitoring studies for the assessment of a change in environmental FR levels with development.

SIP-PAS. SIP-PAS were deployed for one sampling quarter in 2015, at 7 sites. The SIP-PAS were only deployed at one urban site (Rio Gallegos, Argentina) and all of the background and agricultural sites. Concentrations of n-PFAS, perfluor-




Figure 4. Map of detected concentrations (pg/m^3) of (a) FOSA and FOSEs and (b) PFASs in the GRULAC region from SIP-PAS deployed in 2015. Note the different scales on the two maps. The one urban location is marked with an asterisk. NS indicates that a sample was not deployed at that site.

ocarboxylic acids (PFCAs), perfluorosulfonic acids (PFSA), and VMS from each sample are listed in Tables S23–S26.

Neutral Poly- and Perfluoroalkyl Substances. The fluorotelomer alcohols (FTOHs) dominated over the perfluorooctane sulfonamides and sulfonamido ethanols (FOSA and FOSEs) at the sites where they were detected (Figure 2). The 8:2 FTOH concentrations were 2–7 times higher than the 10:2 FTOH, while the 6:2 FTOH was below the MDL in all samples. The FOSAs were detected at all seven sites, and the FOSEs were only detected at three sites with concentrations close to method detection limits. Figure S7 presents box-and-whisker plots of individual n-PFAS.

Figure 4 maps the profiles of concentrations of the FOSA/FOSEs over the GRULAC region, and Figure S8 maps concentrations of the FTOHs. EtFOSA was detected at the highest concentrations at São Jose dos Ausentes (Brazil) and Manizales (Colombia) where concentrations were at least an order of magnitude higher than the other GRULAC sites (Figure 4). The concentrations at these two sites were also an order of magnitude higher than concentrations detected in North America and Europe in the core GAPS sampling campaign in 2015²⁸ (Table S23). EtFOSA is still produced and in use in Brazil as the active ingredient in the pesticide sulfluramid²⁵ and is exported from Brazil to other GRULAC countries for use, including Argentina, Colombia, and Costa Rica.²⁵ In Colombia, there are no specific regulations on the production or use of sulfluramid, and it is acknowledged in the Colombian National Implementation Plan of the Stockholm Convention that its use and registered sales have continued to at least 2015.⁴³ As such, the elevated atmospheric concentrations of EtFOSA detected in Brazil and Colombia are in line with the production and use of this pesticide.

Global n-PFAS levels (ng/PAS) have been previously reported from XAD-PAS deployed at up to 46 GAPS sites, including 8 GRULAC sites, in 2006–2011⁴⁴ (Table S23). The rural site of Indaiatuba (Brazil) and the background site of Saint Peter (Brazil) saw elevated concentrations of EtFOSA, which were also an order of magnitude higher than at the other

GRULAC and global sites sampled in 2006–2011. A calibration study was also conducted at the suburban site of San Antonio de Belen (Costa Rica). Elevated levels of EtFOSA (similar to the levels at Brazil) were again seen, which is in line with known export of sulfluramid to Costa Rica.²⁵ In this 2015 GRULAC study, concentrations of EtFOSA were not elevated at the background site of Tapanti (Costa Rica), indicating different use patterns of EtFOSA in suburban and background (national park) areas of Costa Rica.

Few studies have reported n-PFAS from the southern hemisphere, namely over the Atlantic and Southern Oceans, and are summarized in Table S23. Janke et al.,⁴⁵ Dreyer et al.,⁴⁶ and Wang et al.⁴⁷ reported concentrations of n-PFAS in 2005, 2007–2008, and 2010–2011, respectively, that were in line with concentrations seen in the GRULAC region. All three ship-based studies reported a strong concentration gradient with significantly lower concentrations in the southern hemisphere than the northern hemisphere. This concentration gradient was not observed at the sites sampled in the GAPS Network. As mentioned previously, the n-PFAS have been reported from North America and Europe in 2015 from the regular GAPS sampling campaign.²⁸ Concentrations of the FTOHs and the FOSA and FOSEs at the GRULAC sites (with the elevated EtFOSA concentrations at São Jose dos Ausentes and Manizales removed from the comparison) were in line with concentrations at the background and polar sites in North America and Europe in 2015²⁸ (Table S23). The concentrations between the GRULAC region, North America and Europe were not significantly different ($p > 0.05$), although there is a small data set for the European ($n = 4$) and the GRULAC sites ($n = 7$), reducing the statistical reliability. The similar concentrations of the n-PFAS detected in the developing and developed regions (where use and production is higher) demonstrates the global reach of these chemicals of concern and the long-range atmospheric transport properties of these volatile precursors of the PFAAs.^{48,49}

Perfluoroalkyl Acids. The PFASs were detected at every site with PFOS dominating, followed by perfluorobutanesulfonate

(PFBS) and PFHxS. PFDS was not detected in the GRULAC region. Figure S7 presents box-and-whisker plots of the individual PFASs. The PFCAs had elevated concentrations in the field blanks, as described further in Rauert et al.,²⁸ with target PFCAs only detected above MDLs at Tapanti (Costa Rica) (Table S24). Hence, the PFCAs are not included in the figures due to the low detection of these analytes.

PFSA concentrations were similar at all sites, including the urban site of Rio Gallegos (Argentina), except for Tapanti (Costa Rica), where lower concentrations were observed (Table S25). Figure 4 maps the profiles of atmospheric concentrations of the PFASs. The concentrations of the PFASs were 10 times higher than the precursor FOSAs/FOSEs, which is in line with observations at other global regions sampled in the GAPS Network.²⁸

PFOS is listed in the Stockholm Convention for reduction of production and use, but acceptable purposes are also listed, including the production of sulfluramid.^{22,24} Although elevated EtFOSA concentrations were detected in Brazil and Colombia, atmospheric concentrations of PFOS were similar in all countries in the GRULAC region (Figure 4). The numerous sources of PFOS to the atmosphere in this region, including the transformation of precursors, long-range transport, and localized emissions, all contribute to the similar regional background atmospheric concentrations observed.

PFAAs have been previously reported from a few atmospheric studies in the southern hemisphere, namely ship-based studies, summarized in Table S24–S25. Janke et al.⁴⁵ reported concentrations of PFOS in 2005 that were up to 2 orders of magnitude lower than the concentrations seen in the GRULAC region in 2015. Only the particulate phase was analyzed, and so concentrations may be underestimated. Dreyer et al.⁴⁶ also reported lower concentrations of the PFAS in 2007–2008 than in the GRULAC region in 2015. There are different sources to atmospheric samples from ship (marine)-based locations and land-based sites, with an increased number of potential local sources at land-based locations. The differences between these studies demonstrate that more monitoring is required from the southern hemisphere to determine the environmental fate and regional trends of the PFAS.

Similar concentrations of the PFASs were seen between the North American and European sites in the GAPS Network and the GRULAC sites. PFASs and PFCAs have been reported from North America and Europe in 2015 from the core GAPS sampling campaign²⁸ (Table S24–25). Concentrations of the PFASs were not statistically different between the GRULAC region, and the background and polar sites in North America or Europe ($p > 0.05$). This again demonstrates the global reach of the PFAAs and long-range transport properties.^{49–51}

Volatile Methyl Siloxanes. The SIP-PAS from São Jose (Brazil) had unacceptably low surrogate recoveries (<5%), and this sample has been removed from the data set. The following text compares the remaining six SIP-PAS that were deployed around this region.

Dodecamethylpentasiloxane (L5) was the only linear VMS (IVMS) detected, and concentrations were close to the detection limit. The cyclic VMS (cVMS), meanwhile, were detected at every site and had the highest concentrations of all of the analyte classes monitored (Figure 2). Figure S9 presents box-and-whisker plots of the individual cVMS. The urban site of Rio Gallegos (Argentina) had higher concentrations of the cVMS than the background and agricultural sites by at least a

factor of 2. The elevated concentrations at this urban site are in line with other studies that have reported increasing cVMS atmospheric concentrations with increasing population density.^{28,52} Figure 5 maps the profiles of atmospheric concentrations of the cVMS in the GRULAC region.




Figure 5. Map of detected concentrations (ng/m^3) of the cVMS in the GRULAC region from SIP-PAS deployed in 2015. The one urban location is marked with an asterisk. NS indicates that a sample was not deployed at that site.

This is the first report of VMS atmospheric concentrations from the GRULAC region as well as from the southern hemisphere (to the author's knowledge). The VMS have been reported previously from North America and Europe from the core GAPS sampling campaign in 2015.²⁸ Concentrations of the cVMS in the GRULAC region were not statistically different ($p > 0.05$) to the concentrations in North America or Europe, summarized in Table S26. The profiles were also similar between the three regions with octamethylcyclotetrasiloxane (D4) and decamethylcyclopentasiloxane (D5) dominating. It is estimated that Europe, North America, and China account for about one-quarter each of the world consumption market of the VMS.²⁶ The similar atmospheric concentrations in the developed regions (with higher production and use of the VMS) and the developing GRULAC region, highlights that long-range atmospheric transport is a significant contributor to atmospheric concentrations in developing regions and also shows the global coverage of these chemicals as shown in modeling studies.^{32,53,54}

To further investigate long-range transport as a source to the atmosphere at these sites, the ratio of D5 to D4 was investigated as an indicator of atmospheric inputs of cVMS to a location.^{28,52} A higher ratio (>2) indicates that local sources, such as emissions from wastewater treatment plants,²⁷ are the main contributors to atmospheric concentrations and a lower ratio (<2) indicates long-range atmospheric transport is the

588 main input.²⁸ At the GRULAC sites, the highest ratios were
589 seen at the urban site of Rio Gallegos and the agricultural site of
590 Sonora (Mexico) of 4.4 and 8.6, respectively, suggesting that
591 local inputs are a primary contributor to the air mass at these
592 areas. The background site of Manizales (Colombia) had a ratio
593 of 2.2 indicating a mix of sources to this location, and the other
594 sites all had ratios of <2, indicating that long-range atmospheric
595 transport is the main contributor to atmospheric concen-
596 trations. All ratios are listed in Table S26.

597 **Implications.** The time series of newer POPs (and emerging
598 chemicals of concern) from long-term monitoring programs in
599 North America and Europe are still too short to distinguish
600 trend changes from annual fluctuations.⁶ The time-series data
601 from developing regions is even shorter or nonexistent. This
602 study highlights the importance and need for monitoring data
603 from developing countries, providing regional-scale monitoring
604 data of the OPEs, PBDEs, novel FR, and n-PFAS and the first
605 atmospheric data of the PFAAs and VMS from the GRULAC
606 region. Further monitoring is required to investigate the
607 tendencies observed in this study, such as the significantly
608 higher concentrations of OPEs in the GRULAC region, as
609 compared to North America and Europe, and the elevated
610 concentrations of EtFOSA in Brazil and Colombia.

611 As discussed in Rauert et al.,¹⁰ there are a number of
612 challenges that passive air sampling monitoring networks may
613 face. In this GRULAC study, we have observed anomalous
614 results at select sites in which certain compounds were elevated
615 during one sampling year but not the next or over the entire
616 study. This highlights the need to conduct longer term
617 sampling and to exercise caution when deploying samplers to
618 avoid possible point sources such as indoor air from ventilation
619 systems. Placement of samplers should take advantage of the
620 electricity-free nature of PAS, with the samplers deployed well
621 away from buildings, electrical equipment and other potential
622 sources of target analytes. Furthermore, to obtain representative
623 results for urban areas, which tend to be more heterogeneous in
624 terms of sources compared to background sites, it is advisable
625 to deploy at multiple sites. Another uncertainty occurs with the
626 use of an average sampling rate (4 m³/day) instead of site-
627 specific sampling rates. The use of models of global sampling
628 rates, currently being developed, may improve the accuracy of
629 this measurement in future studies.⁵⁵

630 Another challenge, highlighted further in this GRULAC
631 study, is that for some compound classes (e.g., the PBDEs and
632 novel FRs), detection in air at background sites is difficult, as
633 shown in Figure 2. The concentrations at these sites are in the
634 same range as the MDLs, resulting in a lower confidence in the
635 data and trends produced. Higher MDLs can be partly
636 attributed to contamination issues and should be explored
637 through improved laboratory practices, although this can be
638 difficult and costly. Another option is to focus on representative
639 urban sites, with higher air concentrations, to help overcome
640 limitations related to MDLs and to develop meaningful regional
641 and temporal trends.

642 Lastly, the study highlights the versatility of the PUF disk and
643 SIP disk samplers for conducting region-scale sampling in a
644 simple and cost-effective way. These samplers are capable of
645 sampling multiple target chemical classes that span an
646 enormous range of volatilities. The GAPS-type sampler allows
647 for both gas-phase and particle-phase chemicals to be sampled
648 at similar rates,⁵⁶ which allows for the comparison of data
649 against results obtained using conventional high-volume
650 samplers.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the
ACS Publications website at DOI: 10.1021/acs.est.8b00995.

Tables showing site and sampling information, target
analyte information, and concentrations of target analytes
in each sample. Figures showing box and whisker plots of
target analytes and maps of target analytes over the
GRULAC region. (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Phone: +1-416-739-4837; e-mail: Tom.Harner@Canada.ca.

ORCID

Cassandra Rauert: 0000-0002-2543-9023

Tom Harner: 0000-0001-9026-3645

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial funding and support was provided by the United
Nations Environment Programme (UNEP) and the Chemicals
Management Plan (CMP). These data have been generated in
the frame of the Global Monitoring Plan for POPs under the
Stockholm Convention, with support from the Secretariat of
the Stockholm Convention and UN Environment. The
implementation of the project as part of the Global Monitoring
Plan for POPs has been made possible thanks to the generous
contribution to the Stockholm Convention Voluntary Trust
Fund by the European Commission through its Global Public
Goods and Challenges (GPGC) programme. Thanks go to the
team at Rio Blanco nature reserve, Aguas de Manizales in
Manizales, for the collaboration during the sampling period.
G.F. is a research fellow of the Brazilian Research Council
(CNPq PQ 312341/2013-0) and CAPES (88881.120814/
2016-01). K.P. is funded by the Fondecyt project no. 1161673.
Deep thanks go to the Chacaltaya GAW station team helping
on the deployment and recovery of the samples, especially
Fernando Velarde and Marcelo Peñaloza. Thanks go to
Clemens Ruepert from the Laboratory of Pesticide Residue
Analysis of IRET for his collaboration during field work.

■ REFERENCES

- (1) United Nations Environment Programme. Stockholm Convention-Global Monitoring Plan. <http://chm.pops.int/Implementation/GlobalMonitoringPlan/Overview/tabid/83/Default.aspx>.
- (2) Rauert, C.; Harner, T.; Schuster, J. K.; Quinto, K.; Fillmann, G.; Castillo, L. E.; Fentanes, O.; Villa Ibarra, M.; Miglioranza, K.; Moreno Rivadeneira, I.; Pozo, K.; Padilla Puerta, A.; Aristizabal Zuluaga, B. H. Towards a regional passive air sampling network and strategy for new POPs in the GRULAC region: Perspectives from the GAPS Network and first results for organophosphorus flame retardants. *Sci. Total Environ.* **2016**, *573*, 1294–1302.
- (3) Meire, R. O.; Lee, S. C.; Yao, Y.; Targino, A. C.; Torres, J. P. M.; Harner, T. Seasonal and altitudinal variations of legacy and current-use pesticides in the Brazilian tropical and subtropical mountains. *Atmos. Environ.* **2012**, *59*, 108–116.
- (4) Tominaga, M. Y.; Silva, C. R.; Melo, J. P.; Niwa, N. A.; Plascak, D.; Souza, C. A. M.; Sato, M. I. Z. PCDD, PCDF, dl-PCB and organochlorine pesticides monitoring in São Paulo City using passive air sampler as part of the Global Monitoring Plan. *Sci. Total Environ.* **2016**, *571*, 323–331.

- (5) Leslie, H. A.; van Bavel, B.; Abad, E.; de Boer, J. Towards comparable POPs data worldwide with global monitoring data and analytical capacity building in Africa, Central and Latin America, and the South Pacific. *TrAC, Trends Anal. Chem.* **2013**, *46*, 85–97.
- (6) Wöhrnschimmel, H.; Scheringer, M.; Bogdal, C.; Hung, H.; Salamova, A.; Venier, M.; Katsoyiannis, A.; Hites, R. A.; Hungerbühler, K.; Fiedler, H. Ten years after entry into force of the Stockholm Convention: What do air monitoring data tell about its effectiveness? *Environ. Pollut.* **2016**, *217*, 149–158.
- (7) Estellano, V. H.; Pozo, K.; Harner, T.; Franken, M.; Zaballa, M. Altitudinal and seasonal variations of persistent organic pollutants in the Bolivian Andes mountains. *Environ. Sci. Technol.* **2008**, *42*, 2528–2534.
- (8) Francisco, A. P.; Nardocci, A. C.; Tominaga, M. Y.; da Silva, C. R.; de Assunção, J. V. Spatial and seasonal trends of polychlorinated dioxins, furans and dioxin-like polychlorinated biphenyls in air using passive and active samplers and inhalation risk assessment. *Atmos. Pollut. Res.* **2017**, *8*, 979–987.
- (9) Pozo, K.; Harner, T.; Wania, F.; Muir, D. C. G.; Jones, K. C.; Barrie, L. A. Toward a global network for persistent organic pollutants in air: Results from the GAPS study. *Environ. Sci. Technol.* **2006**, *40*, 4867–4873.
- (10) Raurert, C.; Schuster, J. K.; Eng, A.; Harner, T. Global atmospheric concentrations of brominated, chlorinated flame retardants and organophosphate esters. *Environ. Sci. Technol.* **2018**, *52*, 2777–2789.
- (11) Schuster, J. K.; Harner, T.; Fillmann, G.; Ahrens, L.; Altamirano, J. C.; Aristizábal, B.; Bastos, W.; Castillo, L. E.; Cortés, J.; Fentanes, O.; Gusev, A.; Hernandez, M.; Ibarra, M. V.; Lana, N. B.; Lee, S. C.; Martínez, A. P.; Miglioranza, K. S. B.; Padilla Puerta, A.; Segovia, F.; Siu, M.; Tominaga, M. Y. Assessing polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in air across Latin American countries using polyurethane foam disk passive air samplers. *Environ. Sci. Technol.* **2015**, *49* (6), 3680–3686.
- (12) United Nations Environment Programme. The new POPs under the Stockholm Convention. <http://chm.pops.int/TheConvention/ThePOPs/TheNewPOPs/tabid/2511/Default.aspx>.
- (13) Covaci, A.; Harrad, S.; Abdallah, M. A.-E.; Ali, N.; Law, R. J.; Herzke, D.; de Wit, C. A. Novel brominated flame retardants: A review of their analysis, environmental fate and behaviour. *Environ. Int.* **2011**, *37*, 532–556.
- (14) Van der Veen, I.; de Boer, J. Phosphorus flame retardants: Properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* **2012**, *88*, 1119–1153.
- (15) Wei, G.-L.; Li, D.-Q.; Zhuo, M.-N.; Liao, Y.-S.; Xie, Z.-Y.; Guo, T.-L.; Li, J.-J.; Zhang, S.-Y.; Liang, Z.-Q. Organophosphorus flame retardants and plasticizers: Sources, occurrence, toxicity and human exposure. *Environ. Pollut.* **2015**, *196*, 29–46.
- (16) United Nations Environment Programme. SC-6/13: Listing of hexabromocyclododecane; <http://chm.pops.int/TheConvention/ThePOPs/TheNewPOPs/tabid/2511/Default.aspx>.
- (17) Government of Canada. Canada Consumer Product Safety Act (S.C. 2010, c.21). <http://laws-lois.justice.gc.ca/eng/acts/C-1.68/page-10.html>.
- (18) Salamova, A.; Hermanson, M. H.; Hites, R. A. Organophosphate and halogenated flame retardants in atmospheric particles from a European Arctic site. *Environ. Sci. Technol.* **2014**, *48*, 6133–6140.
- (19) Hung, H.; Katsoyiannis, A. A.; Brorström-Lundén, E.; Olafsdóttir, K.; Aas, W.; Breivik, K.; Bohlin-Nizzetto, P.; Sigurdsson, A.; Hakola, H.; Bossi, R.; Skov, H.; Sverko, E.; Barresi, E.; Fellin, P.; Wilson, S. Temporal trends of persistent organic pollutants (POPs) in Arctic air: 20 years of monitoring under the Arctic Monitoring and Assessment Programme (AMAP). *Environ. Pollut.* **2016**, *217*, 52–61.
- (20) Sühling, R.; Diamond, M. L.; Scheringer, M.; Wong, F.; Púcko, M.; Stern, G.; Burt, A.; Hung, H.; Fellin, P.; Li, H.; Jantunen, L. M. Organophosphate esters in Canadian Arctic air: Occurrence, levels and trends. *Environ. Sci. Technol.* **2016**, *50*, 7409–7415.
- (21) Buck, R. C.; Franklin, J.; Berger, U.; Conder, J. M.; Cousins, I. T.; de Voogt, P.; Jensen, A. A.; Kannan, K.; Mabury, S. A.; van Leeuwen, S. P. J. Perfluoroalkyl and polyfluoroalkyl substances in the environment: Terminology, classification, and origins. *Integr. Environ. Assess. Manage.* **2011**, *7*, 513–541.
- (22) United Nations Environment Programme. SC-4/17: Listing of perfluorooctane sulfonic acid, its salts and perfluorooctane sulfonyl fluoride. <http://chm.pops.int/TheConvention/ThePOPs/TheNewPOPs/tabid/2511/Default.aspx>.
- (23) United Nations Environment Programme. Chemicals proposed for listing under the Convention. <http://chm.pops.int/TheConvention/ThePOPs/ChemicalsProposedforListing/tabid/2510/Default.aspx>.
- (24) United Nations Environment Programme. Acceptable Purposes: PFOS, its salts and PFOSF. <http://chm.pops.int/Implementation/Exemptionsandacceptablepurposes/RegistersofAcceptablePurposes/AcceptablePurposesPFOSandPFOSF/tabid/794/Default.aspx>.
- (25) Löfstedt Gilljam, J.; Leonel, J.; Cousins, I. T.; Benskin, J. P. Is ongoing sulfuramid use in South America a significant source of perfluorooctanesulfonate (PFOS)? Production inventories, environmental fate and local occurrence. *Environ. Sci. Technol.* **2016**, *50*, 653–659.
- (26) Wang, D.-G.; Norwood, W.; Alaei, M.; Byer, J. D.; Brimble, S. Review of recent advances in research on the toxicity, detection, occurrence and fate of cyclic volatile methyl siloxanes in the environment. *Chemosphere* **2013**, *93*, 711–725.
- (27) Shoeib, M.; Schuster, J.; Raurert, C.; Su, K.; Smyth, S.-A.; Harner, T. Emission of poly and perfluoroalkyl substances, UV-filters and siloxanes to air from wastewater treatment plants. *Environ. Pollut.* **2016**, *218*, 595–604.
- (28) Raurert, C.; Shoeib, M.; Schuster, J. K.; Eng, A.; Harner, T. Atmospheric concentrations and trends of poly- and perfluoroalkyl substances (PFAS) and volatile methyl siloxanes (VMS) over 7 years of sampling in the GAPS Network. *Environ. Pollut.* **2018**, *238*, 94–102.
- (29) Government of Canada. Siloxane D4 (Cyclotetrasiloxane, octamethyl-). <https://www.canada.ca/en/health-canada/services/chemical-substances/challenge/batch-2/cyclotetrasiloxane-octamethyl.html>.
- (30) Shoeib, M.; Harner, T.; Vlahos, P. Perfluorinated chemicals in the Arctic atmosphere. *Environ. Sci. Technol.* **2006**, *40*, 7577–7583.
- (31) Genualdi, S.; Lee, S. C.; Shoeib, M.; Gawor, A.; Ahrens, L.; Harner, T. Global Pilot Study of Legacy and Emerging Persistent Organic Pollutant using Sorbent-Impregnated Polyurethane Foam Disk Passive Air Samplers. *Environ. Sci. Technol.* **2010**, *44*, 5534–5539.
- (32) Genualdi, S.; Harner, T.; Cheng, Y.; MacLeod, M.; Hansen, K. M.; van Egmond, R.; Shoeib, M.; Lee, S. C. Global Distribution of Linear and Cyclic Volatile Methyl Siloxanes in Air. *Environ. Sci. Technol.* **2011**, *45*, 3349–3354.
- (33) Stockholm Convention. Guidance on the Global Monitoring Plan for Persistent Organic Pollutants, January 2013; <http://chm.pops.int/Implementation/GlobalMonitoringPlan/Overview/tabid/83/Default.aspx>.
- (34) Shoeib, M.; Harner, T.; Lee, S. C.; Lane, D.; Zhu, J. Sorbent-impregnated polyurethane foam disk for passive air sampling of volatile fluorinated chemicals. *Anal. Chem.* **2008**, *80*, 675–682.
- (35) Harner, T. 2017_v1_5_Template for calculating PUF and SIP disk sample air volumes_September_15.
- (36) Pegoraro, C. A.; Harner, T.; Su, K.; Chiappero, M. S. Assessing levels of POPs in air over the South Atlantic ocean off the coast of South America. *Sci. Total Environ.* **2016**, *571*, 172–177.
- (37) Pozo, K.; Oyola, G.; Estellano, V. H.; Harner, T.; Rudolph, A.; Prybilova, P.; Kukucka, P.; Audi, O.; Klánová, J.; Metzdröff, A.; Focardi, S. Persistent organic pollutants (POPs) in the atmosphere of three Chilean cities using passive air samplers. *Sci. Total Environ.* **2017**, *586*, 107–114.
- (38) Hale, R. C.; Alaei, M.; Manchester-Neesvig, J. B.; Stapelton, H. M.; Ikonomou, M. G. Polybrominated diphenyl ether flame retardants in the North American environment. *Environ. Int.* **2003**, *29*, 771–779.
- (39) Covaci, A.; Gerecke, A. C.; Law, R. J.; Voorspoels, S.; Kohler, M.; Heeb, N. V.; Leslie, H.; Allchin, C. R.; de Boer, J. 47

- 848 Hexabromocyclododecanes (HBCDs) in the Environment and
849 Humans: A Review. *Environ. Sci. Technol.* **2006**, *40*, 3679–3688.
- 850 (40) Robinson, B. H. E-waste: an assessment of global production
851 and environmental impacts. *Sci. Total Environ.* **2009**, *408*, 183–191.
- 852 (41) Araújo, M. G.; Magrini, A.; Mahler, C. F.; Bilitewski, B. A model
853 for estimation of potential generation of waste electrical and electronic
854 equipment in Brazil. *Waste Manage.* **2012**, *32*, 335–342.
- 855 (42) Política nacional de resíduos sólidos. Lei n. 12.305, de 2 de
856 agosto de 2010. 3. ed., Brasília: Câmara dos Deputados, Edições
857 Câmara (Série legislação No 230); http://bd.camara.leg.br/bd/bitstream/handle/bdcamara/14826/politica_residuos_solidos_3ed.pdf?sequence=15.
- 858
859
- 860 (43) National Implementation Plan. *Colombia Stockholm convention*
861 *on persistent organic pollutants - POPs*; Ministry of Environment and
862 Sustainable Development: Bogota, Colombia, 2017.
- 863 (44) Gawor, A.; Shunthirasingham, C.; Hayward, S. J.; Lei, Y. D.;
864 Gouin, T.; Mmereki, B. T.; Masamba, W.; Ruepert, C.; Castillo, L. E.;
865 Shoeib, M.; Lee, S. C.; Harner, T.; Wania, F. Neutral polyfluoroalkyl
866 substances in the global Atmosphere. *Environmental Science Processes &*
867 *Impacts* **2014**, *16*, 404–413.
- 868 (45) Jahnke, A.; Berger, U.; Ebinghaus, R.; Temme, C. Latitudinal
869 gradient of airborne polyfluorinated alkyl substances in the marine
870 atmosphere between Germany and South Africa (53° N–33° S).
871 *Environ. Sci. Technol.* **2007**, *41*, 3055–3061.
- 872 (46) Dreyer, A.; Weinberg, I.; Temme, C.; Ebinghaus, R.
873 Polyfluorinated compounds in the atmosphere of the Atlantic and
874 Southern Oceans: Evidence for a global distribution. *Environ. Sci.*
875 *Technol.* **2009**, *43*, 6507–6514.
- 876 (47) Wang, Z.; Xie, Z.; Mi, W.; Möller, A.; Wolschke, H.; Ebinghaus,
877 R. Neutral poly/per-fluoroalkyl substances in air from the Atlantic to
878 the Southern Ocean and in Antarctic Snow. *Environ. Sci. Technol.* **2015**,
879 *49*, 7770–7775.
- 880 (48) Yarwood, G.; Kembal-Cook, S.; Keinath, M.; Waterland, R. L.;
881 Korzeniowski, S. H.; Buck, R. C.; Russell, M. H.; Washburn, S. T.
882 High-resolution atmospheric modeling of fluorotelomer alcohols and
883 perfluorocarboxylic acids in the North America troposphere. *Environ.*
884 *Sci. Technol.* **2007**, *41*, 5756–5762.
- 885 (49) Wang, Z.; Boucher, J. M.; Scheringer, M.; Cousins, I. T.;
886 Hungerbühler, K. Toward a comprehensive global emission inventory
887 of C₄–C₁₀ perfluoroalkanesulfonic acids (PFSA) and related
888 precursors: Focus on the life cycle of C₈-based products and ongoing
889 industrial transition. *Environ. Sci. Technol.* **2017**, *51*, 4482–4493.
- 890 (50) Wallington, T. J.; Hurley, M. D.; Xia, J.; Wuebbles, D. J.;
891 Sillman, S.; Ito, A.; Penner, J. E.; Ellis, D. A.; Martin, J.; Mabury, A.;
892 Nielsen, O. J.; Sulbaek Andersen, M. P. Formation of C₇F₁₅COOH
893 (PFOA) and other perfluorocarboxylic acids during the atmospheric
894 oxidation of 8:2 fluorotelomer alcohol. *Environ. Sci. Technol.* **2006**, *40*,
895 924–930.
- 896 (51) Armitage, J. M.; MacLeod, M.; Cousins, I. T. Comparative
897 assessment of the global fate and transport pathways of long-chain
898 perfluorocarboxylic acids (PFCAs) and perfluorocarboxylates (PFCs)
899 emitted from direct sources. *Environ. Sci. Technol.* **2009**, *43*, 5830–
900 5836.
- 901 (52) Yucuis, R. A.; Stanier, C. O.; Hornbuckle, K. C. Cyclic siloxanes
902 in air, including identification of high levels in Chicago and distinct
903 diurnal variation. *Chemosphere* **2013**, *92*, 905–910.
- 904 (53) Krogseth, I. S.; Kierkegaard, A.; McLachlan, M. S.; Breivik, K.;
905 Hansen, K. M.; Schlabach, M. Occurrence and seasonality of cyclic
906 volatile methyl siloxanes in Arctic air. *Environ. Sci. Technol.* **2013**, *47*,
907 502–509.
- 908 (54) Janecek, N. J.; Hansen, K. M.; Stanier, C. O. Comprehensive
909 atmospheric modeling of reactive cyclic siloxanes and their oxidation
910 products. *Atmos. Chem. Phys.* **2017**, *17*, 8357–8370.
- 911 (55) Herkert, N. J.; Spak, S. N.; Smith, A.; Schuster, J. K.; Harner, T.;
912 Martinez, A.; Hornbuckle, K. C. Calibration and evaluation of PUF-
913 PAS sampling rates across the global atmospheric passive sampling
914 (GAPS) network. *Environmental Science Processes & Impacts* **2018**, *20*,
915 210–219.
- (56) Markovic, M. Z.; Prokop, A.; Staebler, R. M.; Liggio, J.; Harner, T. Evaluation of the particle infiltration efficiency of three passive
916 samplers and the PS-1 active air sampler. *Atmos. Environ.* **2015**, *112*, 918
919 289–293.