

## Research Paper

**Cite this article:** Galíndez G, Ceccato D, Bubillo R, Lindow-López L, Malagrina G, Ortega-Baes P, Baskin CC (2018). Three levels of simple morphophysiological dormancy in seeds of *Ilex* (Aquifoliaceae) species from Argentina. *Seed Science Research* 1–9. <https://doi.org/10.1017/S0960258518000132>

Received: 30 November 2017  
Accepted: 2 April 2018

**Keywords:**

cold stratification; embryo length; *Ilex* species; morphophysiological dormancy; move-along experiment; seed length ratio; underdeveloped embryo; warm stratification

**Author for correspondence:**

Carol C. Baskin, Email: [ccbask0@uky.edu](mailto:ccbask0@uky.edu)

# Three levels of simple morphophysiological dormancy in seeds of *Ilex* (Aquifoliaceae) species from Argentina

Guadalupe Galíndez<sup>1</sup>, Diana Ceccato<sup>2</sup>, Rosana Bubillo<sup>3</sup>, Lucía Lindow-López<sup>1</sup>, Gisela Malagrina<sup>4</sup>, Pablo Ortega-Baes<sup>1</sup> and Carol C. Baskin<sup>5</sup>

<sup>1</sup>Laboratorio de Investigaciones Botánicas (LABIBO), Facultad de Ciencias Naturales, Universidad Nacional de Salta-CONICET, Av. Bolivia 5150, Salta 4400, Argentina; <sup>2</sup>Agencia de Extensión Rural San Julián Estación Experimental Santa Cruz, Instituto Nacional de Tecnología Agropecuaria (EEA Santa Cruz-INTA), Av. San Martín 1280, San Julián 9310, Santa Cruz, Argentina; <sup>3</sup>Estación Experimental Cerro Azul, Instituto Nacional de Tecnología Agropecuaria (EEA Cerro Azul-INTA), Ruta Nacional 14 Km 836, Cerro Azul 3313, Misiones, Argentina; <sup>4</sup>Banco Base de Germoplasma, Instituto de Recursos Biológicos, CIRN-INTA, De los Reseros y N. Repetto s/n, Hurlingham 1686, Buenos Aires, Argentina and <sup>5</sup>Department of Biology, University of Kentucky, Lexington, KY, USA, and Plant and Soil Sciences, University of Kentucky, Lexington, KY, USA

**Abstract**

As a contribution to understanding the world biogeography of seed dormancy in the cosmopolitan genus *Ilex*, we studied seeds of *I. argentina*, *I. brasiliensis*, *I. brevicuspis*, *I. dumosa*, *I. paraguariensis* and *I. theezans* from the subtropical region of Argentina. We hypothesized that seeds of these species have non-deep simple morphophysiological dormancy (MPD). Effects of temperature, cold stratification and gibberellic acid (GA<sub>3</sub>) on seed germination and embryo growth were tested. Regardless of incubation temperature, little or no germination occurred for any species until  $\geq 6$  weeks. There was an up to 3-fold increase in embryo length to seed length (E:S) ratio before seeds germinated, and embryos grew only during warm-stratifying conditions. Seeds of *I. brasiliensis*, *I. brevicuspis* and *I. theezans* had non-deep simple MPD and germinated to  $\geq 80\%$  after 12, 24 and 16 weeks, respectively. Cold stratification increased germination of *I. brasiliensis* and *I. brevicuspis*, and GA<sub>3</sub> increased the rate but not final germination percentage of *I. brasiliensis* and *I. theezans*. Fresh seeds of *I. dumosa* required 40 weeks of warm stratification to germinate to 53%, while those after-ripened for 2 months germinated to 81% after 30 weeks; this species has intermediate simple MPD. Seeds of *I. argentina* and *I. paraguariensis* germinated to 15 and 21%, respectively, after 40 weeks of warm stratification and did not after-ripen or respond to GA<sub>3</sub>; these seeds have deep simple MPD. This is the first report of intermediate and deep simple MPD that is broken by warm stratification, thereby increasing our knowledge of seed dormancy in *Ilex* and in subtropical regions.

**Introduction**

As a contribution to understanding the world biogeography of seed dormancy, genera such as *Ilex* with a cosmopolitan distribution merit study to determine if the kind of dormancy in individual species is correlated with geography. The *Ilex* genus (Aquifoliaceae) consists of more than 500 species of dioecious shrubs and trees, some of which are deciduous and others evergreen (Cuénoud *et al.*, 2000). The present distribution of *Ilex* includes Asia, Europe, North and South America, a few Pacific islands, northeastern Australia, sub-Saharan Africa and Madagascar (Loizeau *et al.*, 2005). According to Cuénoud *et al.* (2000), *Ilex* already had a cosmopolitan distribution long before the end of the Cretaceous period.

The embryo in seeds of *Ilex* species is differentiated but small (underdeveloped) (Martin, 1946; Hu, 1975; Ng, 1991; Young and Young, 1992; Chien *et al.*, 2011), and it grows inside the seed prior to radicle emergence (e.g. Hu, 1975; Chien *et al.*, 2011). Thus, seeds have morphological dormancy. Furthermore, the embryo in *Ilex* seeds has low growth potential, i.e. physiological dormancy (PD), and warm and/or cold stratification treatments are required to overcome the physiological inhibition of germination (Barton and Thornton, 1947; Nikolaeva *et al.*, 1985; Tsang and Corlett, 2005; Tezuka *et al.*, 2013).

The presence of both morphological and physiological dormancy in the same seed is called morphophysiological dormancy (MPD), and it has been reported in at least 90 families, including Aquifoliaceae (Baskin and Baskin, 2014). In seeds with MPD, embryo growth may occur at the same time that PD is being broken (Baskin and Baskin, 1994) or after part or all of the PD is broken (Baskin and Baskin, 1990). Depending on the species, PD is broken by warm and/or cold stratification, and embryo growth occurs during warm or cold stratification (Nikolaeva, 1977; Baskin and Baskin, 2014). Thus, depending on the temperature

requirements to break PD and promote embryo growth and response of seeds to gibberellic acid (GA<sub>3</sub>), nine levels of MPD have been described: non-deep simple, intermediate simple, deep simple, non-deep simple epicotyl, deep simple epicotyl, deep simple double, non-deep complex, intermediate complex and deep complex (see Baskin and Baskin, 2014). In the simple levels of MPD, embryos grow at temperatures suitable for warm stratification ( $\geq 15^{\circ}\text{C}$ ), and in the complex levels of MPD embryos grow at temperatures suitable for cold stratification (*ca* 0 to  $10^{\circ}\text{C}$ ).

Seeds of *Ilex aquifolium*, *I. glabra*, *I. montana*, *I. opaca*, *I. verticillata* and *I. vomitoria* from temperate areas of the Northern Hemisphere have deep simple MPD (Nikolaeva *et al.*, 1985) and require both warm and cold stratification for dormancy break and germination. On the other hand, seeds of *I. maximo-wicziana* from the tropical and subtropical parts of Taiwan have non-deep simple MPD (Chien *et al.*, 2011) and require only warm stratification for embryo growth and germination. Seeds of *I. asprella*, *I. graciliflora*, *I. hanceana*, *I. pubescens* and *I. viridis* collected in the region of Hong Kong and incubated in a greenhouse there began to germinate after 8–9 weeks (Tsang and Corlett, 2005) and are presumed to have non-deep simple MPD. Also, seeds of *I. glabra* from the southeastern USA germinated to 24, 56 and 70% after 49, 69 and 102 days incubation at  $15^{\circ}\text{C}$ , respectively, suggesting that the seeds had non-deep simple MPD (Hughes, 1964).

In South America, *Ilex* is represented by *ca* 250 species (Giberti, 1995), and seven occur in Argentina: *I. affinis*, *I. argentina*, *I. brasiliensis*, *I. brevicuspis*, *I. dumosa* var. *guaranina*, *I. paraguariensis* var. *paraguariensis* and *I. theezans* (Giberti, 2008), all of which are distributed in the subtropical region of the country (Keller and Giberti, 2011). Seed dormancy in South American species of *Ilex* is attributed to the presence of underdeveloped embryos, and seed germination is delayed for 3 to 9 months with germination percentages often being very low (Fontana *et al.*, 1990; Cuquel *et al.*, 1994; Dolce *et al.*, 2010, 2011). Thus, MPD is present and difficult to break. Several studies have been conducted on seed germination of these species, particularly *I. paraguariensis* (known locally as yerba mate, from which a hot beverage with the same name is made from the leaves), but the majority of them have been concerned with *in vitro* cultivation of embryos with the main objective of improving vegetative propagation of this species (Sansberro *et al.*, 1998, 2001; Dolce *et al.*, 2010, 2011, 2015; Wendling *et al.*, 2013). Only a little attention has been given to the other species of *Ilex* (Dolce *et al.*, 2015). No studies on *Ilex* species from Argentina have been concerned with identifying the level of MPD in the seeds and determining the environmental conditions required to break it. Development of effective dormancy-breaking protocols for plant propagation from seeds of these species is of high commercial and medicinal value in that the non-cultivated species serve as sources of resistance to disease and pests in interspecific hybridization with *I. paraguariensis* (Sansberro *et al.*, 2001) and therefore are an important genetic resource for this crop (Dolce *et al.*, 2015).

The main objective of our study was to determine the level of MPD in seeds of *Ilex argentina*, *I. brasiliensis*, *I. brevicuspis*, *I. dumosa* var. *guaranina*, *I. paraguariensis* var. *paraguariensis* and *I. theezans*. As (1) the six species are distributed in the subtropical zone of Argentina, and (2) seeds of other species of *Ilex* that have been investigated have underdeveloped embryos, we hypothesized that seeds of the Argentinian species have non-deep simple MPD. To test this hypothesis, we determined the effects of

(1) different alternating temperature regimes, (2) a sequence of simulated natural habitat temperatures, and (3) cold stratification on embryo growth and seed germination. We also tested the effects of GA<sub>3</sub> pre-treatments on dormancy break and seed germination.

## Materials and methods

### Study species

*Ilex argentina* Lillo, *I. brasiliensis* (Spreng.) Loes., *I. brevicuspis* Reissek, *I. dumosa* var. *guaranina* (Reissek) Loes., *I. paraguariensis* var. *paraguariensis* (A. St.-Hil.) and *I. theezans* Mart. ex Reissek. are shrubs or trees up to 25 m in height. Flowering typically occurs from October to March and fruiting from January to April (summer–autumn), depending on the species. The fruit is a drupe, and each fruit contains an average of four to five pyrenes, each of which consists of a seed surrounded by a woody endocarp (hereafter seeds). The fruits are red or black and are mainly consumed by birds. All the species are native to Argentina, Brazil, Paraguay and Uruguay (Giberti, 2008). In Argentina, they are distributed in the Paranaense and Yungas (only *I. argentina*) eco-regions, where the climate is mainly subtropical (Cabrera, 1971).

In Argentina, Paraguay, southern Brazil and Uruguay, *I. paraguariensis* is economically the most important perennial crop, and its leaves are used to prepare a traditional beverage called ‘mate’, which is appreciated for its flavour, stimulating properties (contains caffeine and theobromine) and content of vitamins, folic acid and many polyphenolic compounds (Filip *et al.*, 2001). Mate also is used as a phytotherapeutic drug against obesity, colon cancer and other illnesses (Filip *et al.*, 2001; Schinella *et al.*, 2005; de Mejía *et al.*, 2010). The other Argentinian *Ilex* species also have important phytochemical properties similar to those in *I. paraguariensis* and are used as substitutes or adulterants of yerba mate.

### Fruit collection and seed and embryo traits

Mature fruits of each species were collected from at least 10 individuals growing at the Estación Experimental Agropecuaria Cerro Azul– Instituto Nacional de Tecnología Agropecuaria in the locality of Cerro Azul (EEA Cerro Azul, INTA), Misiones, Argentina ( $27^{\circ}39' \text{ S}$ ;  $55^{\circ}26' \text{ W}$ ) in April 2011 for *I. brasiliensis*, *I. paraguariensis* var. *paraguariensis* (hereafter *I. paraguariensis*) and *I. theezans* and in January–February 2012 for *I. argentina*, *I. brevicuspis* and *I. dumosa* var. *guaranina* (hereafter *I. dumosa*). Seeds were manually separated from the pulp, and those that floated on water were discarded (<20%); only those that sank were used in the germination studies. Seeds were dried and stored at  $5^{\circ}\text{C}$  before studies were initiated ( $\leq 15$  days).

Seed mass was estimated by weighing four replicates of 100 seeds each, and seed moisture content (MC) was assessed gravimetrically by drying three replicates of 25 seeds each at  $103^{\circ}\text{C}$  for 17 h. Seed viability was assessed in five replicates of 20 seeds per species using the tetrazolium chloride (TZ) staining technique (ISTA, 2008). Embryo (E) and seed (S) lengths were determined for 20 fresh seeds of each species. Seeds were allowed to imbibe water for 24 h, and then they were cut in half longitudinally using a razor blade under a dissecting optical microscope (Leica ES2 Stereomicroscope, Meyer;  $10\times$  magnification). Seed and embryo images were obtained using a digital camera coupled

to the microscope (Sony Cyber-shot DSC HX100 V 16M), and embryo and seed lengths were measured using the image analysis software ImageJ 1.x (National Institutes of Health, Bethesda, MD, USA). Seed length was measured excluding the thickness of the endocarp, and the embryo (E): seed (S) ratio was calculated for each seed.

### Effect of temperature on germination and embryo growth

To determine the effect of temperature on seed germination, four replicates of 50 fresh seeds each of each species were sown in 90-mm diameter plastic Petri dishes on sterilized sand moistened with distilled water and incubated at 12 h/12 h alternating temperature regimes of 15/5, 20/10, 25/15 and 30/20°C and at a constant temperature of 25°C. At the alternating temperatures, a daily photoperiod of 12 h (30  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , 400–700 nm of cool white fluorescent light; hereafter light) was given during the high temperature period. Additionally, a move-along experiment was performed to evaluate the effects of starting the dormancy-breaking protocol with cold vs warm stratification. Four replicates of 50 seeds each for each species were sown in 90-mm diameter plastic Petri dishes on sterilized sand moistened with distilled water and incubated in light in two temperature sequences: (1) 15/5°C for 12 weeks → 25/15°C for 8 weeks → 30/20°C for 12 weeks → 20/10°C for 8 weeks, and (2) 30/20°C for 12 weeks → 20/10°C for 8 weeks → 15/5°C for 12 weeks → 25/15°C for 8 weeks. These temperatures approximate those that could occur at Cerro Azul in winter (15/5°C), spring (25/15°C), summer (30/20°C) and autumn (20/10°C) (Fig. 1). The control seeds were incubated continuously at 15/5, 20/10, 25/15 and 30/20°C. Germination was defined as radicle emergence  $\geq 1$  mm, and the number of germinated seeds was monitored weekly for 40 weeks. At the end of the germination tests, when no additional germination had occurred for 2 weeks, a cut test was carried out to determine the viability of remaining seeds (soft or firm), and the final germination percentage was calculated on the basis of the total number of firm (viable) seeds.

To determine the effect of temperature on embryo growth, extra dishes of 20 seeds each were incubated at the temperatures described above, and at bi-weekly intervals the E:S ratio was determined. The critical E:S ratio to which embryos must grow before

the radicle emerges (i.e. when the endocarp had split but no radicle protrusion had occurred) was also determined.

### Effect of cold stratification on germination and embryo growth

To evaluate the effect of cold stratification on germination, four replicates of 50 seeds for each species were sown in 90-mm diameter plastic Petri dishes on sterilized sand moistened with distilled water and stored at 5°C in darkness for 0 (control), 4, 8 and 12 weeks. After each treatment, seeds were incubated for 20 weeks in light at 25/15°C and germination was monitored weekly. Results are expressed as germination percentages (%).

The E:S ratio was determined for 20 seeds immediately after each cold stratification treatment and at bi-weekly intervals during the incubation period at 25/15°C for each treatment.

### Effect of gibberellic acid on germination

The effect of GA<sub>3</sub> on germination was evaluated in seeds soaked in 0 (water control), 26, 260 and 2600  $\mu\text{M}$  solutions of GA<sub>3</sub> (95% purity, Sigma, USA) for 24 h at room temperature. These are the concentrations of GA<sub>3</sub> used by Chien *et al.* (2011) for *I. maximo-wicziana*. After soaking in GA<sub>3</sub>, seeds were sown in 90-mm diameter plastic Petri dishes on sterilized sand moistened with distilled water and incubated in light at 25/15°C for 30 weeks. Germination was monitored weekly, and germination percentages were calculated for each treatment.

### Statistical analysis

To test for significant effects of incubation temperature, cold stratification and concentration of GA<sub>3</sub> on seed germination at the end of the experiment, we used a generalized linear model in R (R Core Team, 2016) with a binomial distribution and logit link function. The number of germinated seeds divided by the total number of viable seeds was used as the dependent variable; incubation temperatures, duration of cold stratification and concentration of GA<sub>3</sub> were treated as fixed factors. Significance was tested with a Type II Likelihood Ratio Test (car package; Fox and Weisberg, 2011) and differences among treatments were tested with the lsmeans function (Lenth, 2016). The effects of incubation temperatures and cold stratification on E:S ratio were analysed by a one-way analysis of variance (ANOVA). When significant differences ( $P < 0.05$ ) were detected, a *post-hoc* DGC (named for the authors Di Rienzo, Guzman and Casanoves) multiple comparison test was used to determine differences among treatment means (Di Rienzo *et al.*, 2002, 2014).

## Results

### Seed and embryo traits

Fresh seed mass of *I. brasiliensis*, *I. brevicuspis*, *I. dumosa* and *I. theezans* varied between 0.41 and 0.49 g, while that of *I. argentina* and *I. paraguayensis* was 0.61 and 0.64 g, respectively (Table 1). Seed MC varied from 7.1% in *I. brevicuspis* to 14.1% in *I. argentina*. Both TZ and cut tests indicated high seed viability for *I. brasiliensis*, *I. brevicuspis*, *I. dumosa* and *I. theezans* ( $\geq 85\%$ ), whereas 67 and 40% of the *I. paraguayensis* and *I. argentina* seeds, respectively, were viable. All species had small, heart-shaped (rudimentary) embryos that occupied a small cavity in the endosperm at the micropyle end of the seed, and initial embryo length




Fig. 1. Monthly mean (■), maximum (●) and minimum (▼) daily temperatures and accumulated rainfall (grey bars) for the period 2010 to 2012 at Cerro Azul, Misiones, Argentina.

**Table 1.** Measurements (mean  $\pm$  1SE) of various seed traits of the six *Ilex* species and embryo length of fresh seeds (initial) and embryo length before radicle emergence (critical)

Species	Seed mass (g)	Seed moisture content (%)	Seed viability (%)	Seed length (mm)	Embryo length (mm)	
					Initial	Critical
<i>Ilex argentina</i>	0.61 $\pm$ 0.01	14.1 $\pm$ 0.1	67 $\pm$ 3	2.01 $\pm$ 0.21	0.23 $\pm$ 0.02	1.08 $\pm$ 0.08
<i>Ilex brasiliensis</i>	0.42 $\pm$ 0.006	10.0 $\pm$ 0.9	95 $\pm$ 2	2.64 $\pm$ 0.06	0.24 $\pm$ 0.02	1.52 $\pm$ 0.09
<i>Ilex brevicuspis</i>	0.49 $\pm$ 0.004	7.1 $\pm$ 0.01	92 $\pm$ 5	3.01 $\pm$ 0.04	0.47 $\pm$ 0.03	2.46 $\pm$ 0.23
<i>Ilex dumosa</i>	0.44 $\pm$ 0.01	13.3 $\pm$ 0.1	85 $\pm$ 3	1.98 $\pm$ 0.11	0.44 $\pm$ 0.04	1.65 $\pm$ 0.05
<i>Ilex paraguariensis</i>	0.64 $\pm$ 0.03	10.6 $\pm$ 0.7	40 $\pm$ 5	2.28 $\pm$ 0.20	0.26 $\pm$ 0.02	1.00 $\pm$ 0.03
<i>Ilex theezans</i>	0.41 $\pm$ 0.02	10.1 $\pm$ 0.2	90 $\pm$ 2	2.23 $\pm$ 0.04	0.24 $\pm$ 0.01	2.05 $\pm$ 0.08

varied from 0.23 mm for *I. argentina* to 0.47 mm for *I. brevicuspis*. The critical embryo length for germination ranged from 1.00 mm for *I. paraguariensis* to 2.46 mm for *I. brevicuspis*. Thus, embryo length increased about 3.75- to 8.54-fold prior to germination, depending on the species.

#### Effect of temperature on germination and embryo growth

Incubation temperature had a significant effect on the final seed germination percentage for the study species ( $P < 0.05$ ; Fig. 2). For *I. argentina*, final germination percentages were low in all treatments, with seeds incubated continuously at 25/15°C showing the highest germination percentage (17%); at 15/5°C germination was null (Fig. 2a). Seeds of *I. brasiliensis* and *I. brevicuspis* incubated continuously at 20/10, 25/15 and 30/20°C germinated to 96–98% (Fig. 2b, c), while those of *I. dumosa* and *I. paraguariensis* germinated to a maximum of only 53 and 25%, respectively, at 25/15°C (Fig. 2d, e). Germination of *I. theezans* seeds was significantly higher at 20/10 and 25/15°C ( $\geq 93\%$ ) than at the other temperature regimes (Fig. 2f). Germination percentages of *I. brasiliensis* and *I. theezans* seeds increased when they were moved from high  $\rightarrow$  low temperatures. Germination percentages of *I. argentina*, *I. brasiliensis*, *I. brevicuspis* and *I. theezans* seeds increased when they were moved from low  $\rightarrow$  high temperatures; however, they did not differ significantly from those at continuous (control) temperature regimes (Fig. 2b, c, f).

E:S ratio in fresh seeds of *I. argentina* and *I. paraguariensis* was 0.11  $\pm$  0.01 and 0.10  $\pm$  0.01, respectively, and the critical E:S ratio for germination was 0.41  $\pm$  0.01 and 0.45  $\pm$  0.02, respectively, which was reached after 24 weeks at 25/15°C (Fig. 2g, k). For *I. brasiliensis*, the initial E:S ratio was 0.09  $\pm$  0.01, and the critical E:S ratio was 0.60  $\pm$  0.02, which was reached just before seeds were moved from 30/20 to 20/10°C (Fig. 2h). On the other hand, the E:S ratio for fresh seeds of *I. brevicuspis* was 0.16  $\pm$  0.01, and the critical E:S ratio was 0.78  $\pm$  0.05, which was reached 4 weeks after seeds were moved from 25/15 to 30/20°C, in the low  $\rightarrow$  high temperature move-along sequence (Fig. 2i). However, at the end of weeks 8 and 12 embryos of *I. brasiliensis* and *I. brevicuspis*, respectively, were longer at 25/15 and 30/20°C than in the other treatments, and 16–20% of the seeds had germinated (Fig. 2h, i). For *I. dumosa*, initial and critical E:S ratios were 0.22  $\pm$  0.02 and 0.71  $\pm$  0.01, respectively, whereas for *I. theezans* they were 0.11  $\pm$  0.01 and 0.78  $\pm$  0.05, respectively. The critical embryo size for *I. dumosa* and *I. theezans* was reached after 20 and 12 weeks of incubation at 25/15°C, respectively (Fig. 2j, l).

#### Effect of cold stratification on seed germination and embryo growth

In all treatments, germination of *I. argentina* and *I. paraguariensis* seeds was  $\leq 2\%$  (Fig. 3a, e), while seeds of *I. brasiliensis* and *I. brevicuspis* stratified for 4, 8 and 12 weeks germinated to a higher percentage than the control (Fig. 3b, c). In contrast, germination of *I. dumosa* seeds was higher after 0 and 4 weeks than after 8 and 12 weeks of cold stratification (Fig. 3d). Germination of *I. theezans* seeds was higher after 4 and 8 weeks of cold stratification ( $>90\%$ ; Fig. 3f) than after 0 and 12 weeks of cold stratification.

After 24 weeks of incubation at 25/15°C, the maximum E:S ratio prior to radicle emergence for *I. argentina* and *I. paraguariensis* was in the control (0.37  $\pm$  0.09 and 0.28  $\pm$  0.01, respectively; Fig. 3g, k). For *I. brasiliensis*, *I. brevicuspis*, *I. dumosa* and *I. theezans* the maximum E:S ratio was 0.36  $\pm$  0.04, 0.37  $\pm$  0.02, 0.37  $\pm$  0.02 and 0.39  $\pm$  0.02, respectively (Fig. 3h, i, j, l), for seeds cold stratified for 12 weeks. Thus, the E:S ratio increased about 1.68- to 3.54-fold prior to germination, depending on the species.


#### Effect of GA<sub>3</sub> on seed germination

Control and treated seeds did not differ in final germination percentage (Fig. 4). Germination was  $>80\%$  for *I. brasiliensis*, *I. brevicuspis*, *I. dumosa* and *I. theezans*, whereas it was  $<35\%$  for *I. argentina* and  $<10\%$  for *I. paraguariensis*. Germination of *I. theezans* seeds treated with 2600  $\mu\text{M}$  GA<sub>3</sub> was significantly (barely) higher than that of seeds in the other treatments (Fig. 4f).


#### Discussion

At the time of seed dispersal, embryos of the six *Ilex* species were small, heart-shaped, under-developed and occupied a small cavity in the endosperm. Regardless of incubation temperature, little or no germination occurred for any species until  $\geq 6$  weeks, indicating that seeds were dormant. Before germination, seeds of all species exhibited a 1.68- to 3.54-fold increase in the E:S ratio, and this growth occurred only during incubation at warm-stratifying temperatures (Fig. 2). Thus, seeds of the six *Ilex* species had simple MPD, but there were large differences among the species with regard to the speed of dormancy break during warm stratification.


Seeds of *I. brasiliensis*, *I. brevicuspis* and *I. theezans* germinated to  $\geq 80\%$  after 12, 24 and 16 weeks, respectively, with 50% of the seeds germinating after 10–12, 14–16 and 12–14 weeks,



**Fig. 2.** Cumulative germination percentage and E:S ratio of *I. argentina* (a,g), *I. brasiliensis* (b,h), *I. brevicuspis* (c,i), *I. dumosa* (d,j), *I. paraguariensis* (e,k) and *I. theezans* (f,l) incubated for 40 weeks at continuous 30/20°C (●), 25/15°C (○), 20/10°C (▼), 15/5°C (△), 25°C (◆), and at move-along 30/20°C (■) and 15/5°C (□). Different lower case letters ( $\alpha=0.05$ ) at the end of the experiment indicate significant differences between means ( $\pm 95\%$  binomial confidence intervals), and each value of E:S ratio is the mean ( $\pm 1SE$ ). Arrows indicate times when seeds were moved to the next temperature in the sequence.



**Fig. 3.** Effect of cold stratification at 5°C for 0 (●), 4 (○), 8 (▼) and 12 (△) weeks on seed germination percentage and E:S ratio of *I. argentina* (a,g), *I. brasiliensis* (b,h), *I. brevispis* (c,i), *I. dumosa* (d,j), *I. paraguariensis* (e,k) and *I. theezans* (f,l) incubated for 20 weeks at 25/15°C. Different lower case letters indicate significant differences between means ( $\alpha=0.05$ ) at the end of the experiment. Each value of germination is the mean ( $\pm 95\%$  binomial confidence intervals), and each value of E:S ratio is the mean ( $\pm 1SE$ ).



**Fig. 4.** Germination percentage of *I. argentina* (a), *I. brasiliensis* (b), *I. brevicuspis* (c), *I. dumosa* (d), *I. paraguariensis* (e) and *I. theezans* (f) seeds soaked in water (0, ●) and in 26 (○), 260 (▼) and 2600 (△)  $\mu\text{M}$  solutions of  $\text{GA}_3$  for 24 h at room temperature and then incubated for 30 weeks at 25/15°C. Different lower case letters indicate significant differences between means ( $\alpha = 0.05$ ) at the end of the experiment. Each value of germination is the mean ( $\pm 95\%$  binomial confidence intervals).

respectively. Furthermore, the E:S ratio of *I. brasiliensis*, *I. brevicuspis* and *I. theezans* had doubled after 4, 10 and 2 weeks, respectively. The responses of seeds of these three species to incubation temperatures were similar to those of *I. maximowicziana* from northern (subtropical) Taiwan in that 50% of the *I. maximowicziana* seeds (final germination was 95–100%) had germinated at 20/10, 25/15 and 30/20°C after 10–13 weeks (Chien *et al.*, 2011). Furthermore, like embryos in seeds of *I. maximowicziana* those in seeds of *I. brasiliensis*, *I. brevicuspis* and *I. theezans* did not grow during cold stratification but did so when seeds were subsequently incubated at 25/15°C.  $\text{GA}_3$  did not significantly increase final germination percentages of seeds of these three species, except for 2600  $\mu\text{M}$   $\text{GA}_3$ . However, similar to the responses of *I. maximowicziana*,  $\text{GA}_3$  treatment increased germination of *I. brasiliensis* and *I. theezans* seeds during the first 12 weeks of incubation at 25/15°C. Cold stratification increased final germination percentages of seeds of *I. brasiliensis* and *I. brevicuspis* but not those of *I. theezans*. As (1) warm stratification was the only condition required for embryo growth and germination, and (2) seeds germinated to high percentages in 12–24 weeks, depending

on the species, seeds of *I. brasiliensis*, *I. brevicuspis* and *I. theezans* have non-deep simple MPD, as hypothesized.

However, contrary to our hypothesis, seeds of *I. dumosa*, *I. argentina* and *I. paraguariensis* do not have non-deep simple MPD. Fresh seeds of *I. dumosa* reached a maximum germination of only 53% after 40 weeks of incubation at 25/15°C. However, seeds in the control for the cold stratification experiment of *I. dumosa* reached 64% germination after 20 weeks, and those in the control for the  $\text{GA}_3$  experiment reached 81% germination after 30 weeks. As the tests for the fresh, cold-stratified and  $\text{GA}_3$ -treated seeds were started in March, April and May, respectively, it is clear that after-ripening occurred during seed storage. It should be noted that *I. dumosa* seeds tested in March, April and May required 38, 12 and 18 weeks of warm stratification, respectively, to reach 50% germination. In seeds with intermediate PD, a period of after-ripening (or warm stratification) decreases the length of the treatment required to break dormancy (Nikolaeva, 1969). In the temperate zone, cold stratification for up to 20 or more weeks is required to break intermediate PD, but a period of dry after-ripening (or warm stratification) can greatly decrease

the length of this treatment required to break dormancy (e.g. Baskin *et al.*, 1988). In the case of *I. dumosa*, after-ripening decreased the length of the warm stratification period required to break dormancy. Thus, we conclude that seeds of *I. dumosa* have intermediate simple MPD. In seeds with intermediate PD, GA<sub>3</sub> may, or may not, be effective in breaking dormancy (Baskin and Baskin, 2014); it was not effective for *I. dumosa* seeds.

Several pieces of evidence indicate that seeds of *I. argentina* and *I. paraguariensis* have deep simple MPD. Firstly, embryo growth was slow. In both *I. argentina* and *I. paraguariensis*, 12 weeks of warm stratification were required for the E:S ratio to double, after which another 12-week period was required for it to double again (Fig. 2). Secondly, after 40 weeks of incubation, only 15 and 21% of the *I. argentina* and *I. paraguariensis* seeds had germinated (Fig. 2). For *I. paraguariensis* seeds collected in five different years in Argentina, the first seeds to germinate (at room temperature) did so after 120 days (17.2 weeks). Furthermore, some new seedlings appeared on day 360 (Fontana *et al.*, 1990). Thirdly, scarified seeds of *I. paraguariensis* germinated between days 120 and 360 (Fontana *et al.*, 1990). Fourthly, embryos in seeds of *I. argentina* and *I. paraguariensis* did not grow during cold stratification but grew slowly after seeds were transferred to 25/15°C (Fig. 3). Finally, treatment with GA<sub>3</sub> did not promote germination (Fig. 4).

In temperate regions, deep PD is broken by cold stratification for up to 20 or more weeks, and GA<sub>3</sub> does not promote germination (Nikolaeva, 1969; Baskin and Baskin, 2014). However, deep PD has been found in seeds of *Leptecophylla tameiameiae* (Ericaceae) from the montane zone in Hawaii (Baskin *et al.*, 2005). In *L. tameiameiae*, dormancy was broken by warm stratification, but 56 weeks were required for 50% of the seeds to germinate; some seeds had not germinated after 162 weeks when the study ended. Thus, we conclude that seeds of *I. argentina* and *I. paraguariensis* have deep simple MPD that is broken by warm stratification, and the germination percentages for these two species are low in our study because seeds were incubated for only 40 weeks. It should be noted that *I. argentina* belongs to the same clade as the temperate zone *I. opaca* (Gottlieb *et al.*, 2005), whose seeds have deep simple MPD and require a long period of both warm and cold stratification for embryo growth and germination (Ives, 1923; Barton and Thornton, 1947). Furthermore, the first seeds of *I. chapaensis*, *I. ficoidea* and *I. kwangtungensis* from Hong Kong germinated (in a greenhouse) only after 67, 46 and 49 weeks, respectively (Tsang and Corlett, 2005), suggesting that they also had deep simple MPD that was broken by long periods of warm stratification.

Breaking of non-deep simple MPD in temperate regions requires about half the time as the breaking of deep simple MPD. In seeds with non-deep simple MPD, the PD component can be broken during summer with embryo growth and germination occurring in autumn (e.g. Baskin and Baskin, 1990), or PD can be broken during winter with embryo growth and germination occurring in spring (e.g. Walck *et al.*, 1999). On the other hand, seeds with deep simple MPD require exposure to both summer and winter for dormancy break to occur and thus germinate in spring. For example, in seeds of *Jeffersonia diphylla* with deep simple MPD, part of the PD is broken in summer, embryo growth occurs in autumn, the second part of PD is broken in winter and seeds germinate in spring (Baskin and Baskin, 1989). What our study has revealed about MPD in *Ilex* species in the subtropical regions of Argentina is that, as expected, seeds of all species required warm stratification for embryo growth and germination

and thus have simple MPD. However, unexpectedly, seeds of some species require extensive periods of warm stratification before they germinate and thus have intermediate or deep simple MPD.

In relatively stable warm climates, it is not surprising that seeds require long periods of warm stratification for dormancy break, e.g. *L. tameiameiae* with deep PD in Hawaii, because this is a mechanism that can spread germination of a seed cohort over time. However, heretofore embryo growth and germination have not been studied in seeds of subtropical species that require more than about half a year of warm stratification to germinate. Our study contributes to a better understanding of the world biogeography of seed dormancy in *Ilex* in that we now know that some *Ilex* species in temperate as well as subtropical regions have deep simple MPD. We previously knew that seeds of some *Ilex* species in temperate regions had deep simple MPD broken by warm plus cold stratification and that about one year was required for germination. Now, we know that in subtropical regions there are *Ilex* species such as *I. argentina*, *I. paraguariensis* and *I. chapaensis* whose seeds require a year or more to germinate, but this dormancy is broken by extended periods of warm stratification, instead of warm plus cold stratification. The results of our study suggest that more attention now needs to be given to *Ilex* in the temperate zone to determine if there are species whose seeds have non-deep simple and intermediate simple MPD, in which the PD part of MPD is broken by cold stratification.

**Acknowledgments.** We especially acknowledge Ing. Agr. Sergio D. Prat Kricun and Ing. Agr. Luis D. Belingheri for the creation and maintenance of field collection of *Ilex* species from South America now available at the EEA Cerro Azul-INTA, Misiones. We also thank Professor Silvia Sühling and Professor Fernando Biganzoli for statistical assistance and Professor Jerry M. Baskin for a careful and critical reading of the manuscript. This work was supported by The National Institute of Agricultural Technology Project (INTA-AERG-PE 231221).

## References

- Barton LV and Thornton NC (1947) Germination and sex population studies of *Ilex opaca* Ait. *Contribution from the Boyce Thompson Institute* **14**, 405–410.
- Baskin CC and Baskin JM (1994). Deep complex morphophysiological dormancy in seeds of the mesic woodland herb *Delphinium tricorne* (Ranunculaceae). *International Journal of Plant Sciences* **155**, 738–743.
- Baskin CC and Baskin JM (2014) *Seeds: Ecology, Biogeography, and Evolution of Dormancy and Germination* (2nd edn). San Diego: Elsevier/Academic Press.
- Baskin CC *et al.* (2005) Germination of drupelets in multi-seeded drupes of the shrub *Leptecophylla tameiameiae* (Ericaceae) from Hawaii: a case for deep physiological dormancy broken by high temperatures. *Seed Science Research* **15**, 349–356.
- Baskin JM and Baskin CC (1989) Seed germination ecophysiology of *Jeffersonia diphylla*, a perennial herb of mesic deciduous forests. *American Journal of Botany* **76**, 1073–1080.
- Baskin JM and Baskin CC (1990) Germination ecophysiology of seeds of the winter annual *Chaerophyllum tainturieri*: a new type of morphophysiological dormancy. *Journal of Ecology* **78**, 993–1004.
- Baskin JM, Baskin CC and McCann MT (1988) A contribution to the germination ecology of *Floerkea proserpinacoides* (Limnaceae). *Botanical Gazette* **149**, 427–431.
- Cabrera AL (1971) Fitogeografía de la república Argentina. *Boletín de la Sociedad Argentina de Botánica* **14**, 1–42.
- Chien CT *et al.* (2011) Nondeep simple morphophysiological dormancy in seeds of *Ilex maximowicziana* from northern (subtropical) and southern (tropical) Taiwan. *Ecological Research* **26**, 163–171.



- Cuénoud P et al.** (2000). Molecular phylogeny and biogeography of the genus *Ilex* L. (Aquifoliaceae). *Annals of Botany* **85**, 111–122.
- Cuquel FL, Carvalho MLM and Chama HMCP** (1994) Avaliação de métodos de estratificação para a quebra de dormência de sementes de erva-mate. *Scientia Agricola* **51**, 415–421.
- de Mejía EG et al.** (2010). Yerba mate tea (*Ilex paraguariensis*): phenolics, antioxidant capacity and *in vitro* inhibition of colon cancer cell proliferation. *Journal of Functional Foods* **2**, 23–34.
- Di Rienzo JA et al.** (2014) InfoStat versión. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. Available at: <http://www.infostat.com.ar>
- Di Rienzo JA, Guzmán AW and Casanoves F** (2002) A multiple-comparisons method based on the distribution of the root node distance of a binary tree. *Journal of Agricultural, Biological and Environmental Statistics* **7**, 129–142.
- Dolce NR, Mroginski LA and Rey HY** (2010) Endosperm and endocarp effects on the *Ilex paraguariensis* A. St.-Hil. (Aquifoliaceae) seed germination. *Seed Science and Technology* **38**, 441–448.
- Dolce NR, Mroginski LA and Rey HY** (2011) Enhanced seed germination of *Ilex dumosa* R. (Aquifoliaceae) through *in vitro* culture of cut pyrenes. *HortScience* **46**, 278–281.
- Dolce NR et al.** (2015) Sowing pyrenes under aseptic conditions enhances seed germination of *Ilex brasiliensis*, *I. pseudoboxus* and *I. theezans* (Aquifoliaceae). *Seed Science and Technology* **43**, 273–277.
- Filip R et al.** (2001) Phenolic compounds in seven South American *Ilex* species. *Fitoterapia* **72**, 774–778.
- Fontana HP, Prat Krikum SD and Belingheri LD** (1990). Estudios sobre la germinación y conservación de semillas de yerba mate (*Ilex paraguariensis* St. Hil.). Informe Técnico 52 EEA INTA Cerro Azul, Misiones, Argentina.
- Fox J and Weisberg S** (2011) *An R companion to applied regression*. Thousand Oaks, California, USA: Sage.
- Giberti GC** (1995) Florística, sistemática y potencialidades con relación a un banco de germoplasma para la yerba mate. In Winge H, Ferreira A, Mariath J, Tarasconi L (eds), *Erva-Mate: Biología e Cultura no Cono Sul*. University Federal do Rio Grande do Sul, Brasil, pp. 303–312
- Giberti GC** (2008) Aquifoliaceae, pp. 1143–1146 in Zuloaga FO, Morrone O and Belgrano MJ (eds), *Catálogo de las Plantas vasculares del Cono Sur* (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay), vol. 2, Dicotyledoneae: Acanthaceae – Fabaceae (Abarema – Schizolobium). *Monographs of Systematic Botany Missouri Botanical Garden* **107**, 1143–1146.
- Gottlieb AM, Giberti GC and Poggio L** (2005) Molecular analyses of the genus *Ilex* (Aquifoliaceae) in southern South America, evidence from AFLP and ITS sequence data. *American Journal of Botany* **92**, 352–369.
- Hu CY** (1975) *In vitro* culture of rudimentary embryos of eleven *Ilex* species. *Journal of the American Society for Horticultural Science* **100**, 221–225.
- Hughes RH** (1964) Some observations on germination of gallberry seed. U.S. Forest Service Research Note SE-17, 1–3.
- ISTA** (2008) International Rules for Seed Testing. International Seed Testing Association, Bassersdorf, Switzerland.
- Ives SA** (1923) Maturation and germination of seeds of *Ilex opaca*. *Botanical Gazette* **76**, 60–77.
- Keller HA and Giberti GC** (2011) Primer registro para la flora Argentina de *Ilex affinis* (Aquifoliaceae), sustituto de la 'yerba mate'. *Boletín de la Sociedad Argentina de Botánica* **46**, 187–194.
- Lenth RV** (2016) Least-squares means: the R Package lsmmeans. *Journal of Statistical Software* **69**, 1–33.
- Loizeau P-A et al.** (2005) Towards an understanding of the distribution of *Ilex* L. (Aquifoliaceae) on a world-wide scale. *Biologiske Skrifter* **55**, 501–520.
- Martin AC** (1946). The comparative internal morphology of seeds. *American Midland Naturalist* **36**, 513–660.
- Nikolaeva MG** (1969). *Physiology of deep dormancy in seeds*. Leningrad: Izdatel'stvo Nauka. (Translated from Russian by Z Shapiro, NSF, Washington, DC.).
- Nikolaeva MG** (1977) Factors controlling the seed dormancy pattern. In Khan AA (ed), *The Physiology and Biochemistry of Seed Dormancy and Germination*. Amsterdam, North-Holland, pp. 51–74
- Nikolaeva MG, Rasumova MV and Gladkova VN** (1985) *Reference Book on Dormant Seed Germination*, ed. Danilova MF Leningrad: Nauka Publishers.
- Ng FSP** (1991) *Manual of Forest Fruits, Seeds and Seedlings*, vol. 1. Kuala Lumpur: Forest Research Institute of Malaysia.
- R Core Team** (2016) R: A language and environment for statistical computing. Vienna, Austria, R Foundation for Statistical Computing. Available at: <https://www.r-project.org/> (accessed September 2016).
- Sansberro PA et al.** (1998) *In vitro* culture of rudimentary embryos of *Ilex paraguariensis*: responses to exogenous cytokinins. *Journal of Plant Growth Regulation* **17**, 101–105.
- Sansberro PA, Rey HY and Mroginski LA** (2001) *In vitro* culture of zygotic embryos of *Ilex* species. *HortScience* **36**, 351–352.
- Schinella G, Fantinelli JC and Mosca SM** (2005) Cardioprotective effect of *Ilex paraguariensis* extract: evidence for a nitric oxide-dependent mechanism. *Clinical Nutrition* **24**, 360–366.
- Tezuka T et al.** (2013) Factors affecting seed germination of *Ilex latifolia* and *I. rotunda*. *HortScience* **48**, 352–356.
- Tsang AC and Corlett RT** (2005) Reproductive biology of the *Ilex* species (Aquifoliaceae) in Hong Kong, China. *Botany* **83**, 1645–1654.
- Walck JL, Baskin CC and Baskin JM** (1999) Seeds of *Thalictrum mirabile* (Ranunculaceae) require cold stratification for loss of nondeep simple morphophysiological dormancy. *Canadian Journal of Botany* **77**, 1769–1776.
- Wendling I et al.** (2013) Vegetative propagation of adult *Ilex paraguariensis* trees through epicormic shoots. *Acta Scientiarum Agronomy* **35**, 117–125.
- Young JA and Young CG** (1992). *Seeds of Woody Plants in North America*, revised and enlarged edition. Portland: Dioscorides Press.