

PAPER • OPEN ACCESS

Spin-1/2 Landau levels in the symmetric gauge from the zero energy modes

To cite this article: Lucas Sourrouille 2018 *J. Phys. Commun.* **2** 045030

View the [article online](#) for updates and enhancements.



PAPER

Spin-1/2 Landau levels in the symmetric gauge from the zero energy modes

OPEN ACCESS

RECEIVED

4 October 2017

REVISED

4 April 2018

ACCEPTED FOR PUBLICATION

13 April 2018

PUBLISHED

25 April 2018

Lucas Sourrouille^{1,2} ¹ IMDEA Nanociencia, Calle de Faraday, 9, Cantoblanco, 28049, Madrid, Spain² IFISUR, Departamento de Física (UNS-CONICET) Avenida Alem 1253, Bahía Blanca, 8000, Buenos Aires, ArgentinaE-mail: sourrou@df.uba.ar**Keywords:** Landau Levels, gauge fields, zero modes, quantum mechanics

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**Abstract**

Starting from the zero modes of the Dirac–Weyl equation for Landau levels in the symmetric gauge, we propose a novel mechanism to construct the eigenvalues and its eigenfunctions. We show that the problem may be addressed without numerical calculation and only solving the Dirac–Weyl equation for the zero modes. Specifically, the eigenstates associated to the negative magnetic field configurations may be constructed from the zero mode with positive chirality. In addition, we obtain that the eigenstates associated to the positive magnetic field configurations may be constructed from the zero mode with negative chirality. Finally, we show that our mechanism may be used to obtain the eigenvalues and eigenfunctions of the Hamiltonian corresponding to bilayer graphene system.

1. Introduction

The problem of a single electron confined to two dimensions and exposed to a magnetic field was explored by Darwin [1], Fock [2] and Landau [3]. They show that the electron kinetic energy is quantized, being the discrete kinetic energy levels ‘the Landau levels’.

In particular, the Landau levels become relevant in the Quantum Hall problem [4–14]. Indeed, the integral Quantum Hall effect is a direct consequence of the Landau level formation. In addition the explanation for the fractional quantum Hall effect, arises because the lowest Landau level splits into Landau-like energy levels [15–18].

In addition, deals with Landau levels in planar geometry and in the symmetric gauge, acquire special significance, since the physics of the FQHE would not have revealed itself in a gauge other than the symmetric gauge of the planar geometry [15, 16].

On the other hand, the experimental realization of monolayer graphene films [19–21] has allowed explore the physics of two-dimensional (2D) Dirac–Weyl fermions. This allows relativistic physics to be explored in a solid state system and physical phenomena such as the Klein–Gordon paradox, the anomalous Landau–Hall effect or nanoelectric materials [21–23] may be addressed. Also, the study of Dirac–Weyl electrons in magnetic fields has received much attention in the recent time in order to find a way for confining the charges [24–33].

In this note, we care to study the Landau levels for the Dirac–Weyl equation in the symmetric gauge. This problem was, previously, studied numerically in the Landau gauge (see for review see [34]). In particular, in reference [35] exact analytical solutions for the bound states of the Dirac–Weyl electron in magnetic fields with various q -parameters under an electrostatic potential were obtained. In addition, the Landau levels in the symmetric gauge were analyzed numerically for the nonrelativistic case [36, 37].

Here, we show that the problem may be addressed without numerical calculation, and only solving the Dirac–Weyl equation for the zero modes. Specifically, we will develop a formalism to construct the eigenvalues and its eigenfunctions of the Dirac–Weyl equation for the Landau problem in the symmetric gauge. This formalism is similar to the well known ladder operators mechanism, which is used to obtain the Landau levels for the Schrödinger Hamiltonian [8, 38, 39] and allows us to generate all the eigenstates of the Hamiltonian from any one energy eigenstate. The novelty, here, is to develop a mechanism of ladder operators for Dirac–Weyl

equation, which allows us obtain all Landau levels and their respective eigenfunctions. As we will see our formalism is not only applicable to Dirac-Weyl Hamiltonian, which govern the dynamics of monolayer graphene, but also to bilayer and multilayer graphene Hamiltonians. Specifically, we will show that, the eigenstates associated to the negative magnetic field configurations may be constructed from the zero mode with positive chirality. On the other hand, we obtain that the eigenstates associated to the positive magnetic field configurations may be constructed from the zero mode with negative chirality. In addition, we discuss how to generalize our mechanism to more complex problems such as the study of bilayer graphene system [40–46].

2. The framework and the Aharonov-Casher theorem

Let us start by considering a $(2 + 1)$ -dimensional Dirac-Weyl model whose Hamiltonian is described by

$$H = \sigma^i p_i = (\sigma^1 p_1 + \sigma^2 p_2), \quad (1)$$

Here, the σ^i ($i = 1, 2$) are 2×2 Pauli matrices, i.e.

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad (2)$$

and $p_i = -i\partial_i$ is the two-dimensional momentum operator. The massless Dirac-Weyl equation in $(2 + 1)$ dimensions is

$$\sigma^i p_i \Phi(x, y, t) = i\partial_t \Phi(x, y, t) \quad (3)$$

Here, $\Phi(x, y, t)$ is the two-component spinor

$$\Phi = (\phi_a, \phi_b)^T \quad (4)$$

where ϕ_a and ϕ_b represent the envelope functions associated with the probability amplitudes. Since, we are interested in stationary states, it is natural to propose a solution of the form

$$\Phi(x, y, t) = e^{-iEt} \Psi(x, y), \quad (5)$$

then, the time-independent Dirac-Weyl equation is

$$\sigma^i p_i \Psi(x, y) = E \Psi(x, y) \quad (6)$$

In the presences of a perpendicular magnetic field to the (x, y) -plane, we replace the momentum operator p_i by the covariant derivative, defined as $D_i = -i\partial_i + A_i$ ($i = 1, 2$), where A_i are components of the vector potential,

$$B = \partial_x A_y - \partial_y A_x \quad (7)$$

Thus, the equation (6) becomes,

$$\sigma^i D_i \Psi(x, y) = E \Psi(x, y) \quad (8)$$

We can develop this equation to get,

$$\begin{pmatrix} 0 & D_1 - iD_2 \\ D_1 + iD_2 & 0 \end{pmatrix} \begin{pmatrix} \psi_a \\ \psi_b \end{pmatrix} = E \begin{pmatrix} \psi_a \\ \psi_b \end{pmatrix} \quad (9)$$

where ψ_a and ψ_b are the components of the spinor Ψ (i.e. $\Psi = (\psi_a, \psi_b)^T$). From this equation we can write the two coupled equations for the components ψ_a and ψ_b

$$D_1 \psi_b - iD_2 \psi_b = E \psi_a \quad (10)$$

$$D_1 \psi_a + iD_2 \psi_a = E \psi_b \quad (11)$$

Here, we are interested to find the the eigenvalues and eigenstates corresponding to equation (8). In order to find these one needs to specify a gauge for the vector potential. Here, we will use a symmetric gauge,

$$\mathbf{A} = \frac{1}{2}(-By, Bx, 0) \quad (12)$$

Then, equations (10) and (11) becomes,

$$\left[-i\partial_x - \partial_y - z \frac{B}{2} \right] \psi_b = E \psi_a \quad (13)$$

$$\left[-i\partial_x + \partial_y - z^\dagger \frac{B}{2} \right] \psi_a = E \psi_b \quad (14)$$

where $z = ix + y$ and $z^\dagger = -ix + y$. The simplest solution of the equations (13) and (14) are the zero energy modes, that is the solutions for zero energy. This solution may be constructed explicitly following, the work done by Aharonov and Casher [47]. For this purpose we assume that the vector potential is divergenceless, which is

clearly satisfied by (12). Then, one can introduce a scalar potential $\lambda(x, y)$ such that,

$$A_x = -\partial_y \lambda, \quad A_y = \partial_x \lambda \quad (15)$$

and due to the equation (7),

$$B = \partial_x^2 \lambda + \partial_y^2 \lambda \quad (16)$$

Then, it is not difficult to find the solutions of the equations (13) and (14) for the energy zero case. Indeed, substituting in equation (13)

$$\psi_b = f_b e^{-\lambda} \quad (17)$$

and setting $E = 0$, we obtain,

$$[-i\partial_x - \partial_y]f_b = 0 \quad (18)$$

In similar way, if we propose

$$\psi_a = f_a e^{\lambda} \quad (19)$$

equation is reduced to

$$[-i\partial_x + \partial_y]f_a = 0 \quad (20)$$

Thus, f_a and f_b are analytic and complex conjugated analytic entire functions of $z = ix + y$, respectively.

The equation (16) has the following solution

$$\lambda(\mathbf{r}) = \int d\mathbf{r}' G(\mathbf{r}, \mathbf{r}') B(\mathbf{r}') \quad (21)$$

where

$$G(\mathbf{r}, \mathbf{r}') = \frac{1}{2\pi} \ln \left(\frac{|\mathbf{r} - \mathbf{r}'|}{r_0} \right) \quad (22)$$

is the Green function of the Laplace operator in two dimensions and r_0 is an arbitrary constant. According to [47] the magnetic flux Φ is localized in a restricted region so that for $r \rightarrow \infty$

$$\lambda(\mathbf{r}) = \frac{\Phi}{2\pi} \ln \left(\frac{r}{r_0} \right) \quad (23)$$

and

$$\psi_{a,b} = f_{a,b} \left(\frac{r}{r_0} \right)^{\frac{\gamma\Phi}{2\pi}} \quad (24)$$

where $\gamma = 1$ and -1 for ψ_a and ψ_b respectively. Since the entire function $f(z)$ cannot go to zero in all directions at infinity, $\psi_{a,b}$ can be normalizable only assuming that $\gamma\Phi < 0$, that is, zero-energy solutions can exist only for one spin direction, depending on the sign of the total magnetic flux.

Now, consider the case $\Phi > 0$, then in view of (24) we have $\psi_a = 0$ and

$$\psi_b = f_b e^{-\lambda} \simeq f_b \left(\frac{r}{r_0} \right)^{\frac{-\Phi}{2\pi}} \quad (25)$$

The function f_b is dictated by (18) and it is not difficult to check that the solutions are polynomials of the form

$$f_b = \sum_{i=0}^j a_i z^i \quad (26)$$

However, one can easily see from equation (25) that the solution is integrable with the square only assuming that $j \leq N$ (we count j from $j = 0$), where N is the integer part of $\frac{\Phi}{2\pi}$. For the case $\Phi < 0$ we have

$$\psi_a = f_a e^{\lambda} \simeq f_a \left(\frac{r}{r_0} \right)^{\frac{\Phi}{2\pi}}, \quad \psi_b = 0 \quad (27)$$

where

$$f_a = \sum_{i=0}^j \tilde{a}_i (z^\dagger)^i \quad (28)$$

and

$$j \leq N \quad (29)$$

Thus, the number of the independent states with zero energy for one spin projection is equal to $N + 1$, and there are no such solutions for another spin projection.

3. The first level of energy and its eigenstates for negative magnetic field

Let us now concentrate on the construction of eigenstates corresponding to eigenvalues different from zero. To proceed we start by considering the simplest case,

$$\Psi_{0,0} = \begin{pmatrix} \tilde{a}_0 e^\lambda \\ 0 \end{pmatrix} \quad (30)$$

Here, the first subindex denote the number of independent state with zero energy and the second index denote level of energy. In other words the spinor (30) is the first independent zero energy mode. According to what we have seen (30) is a solution of the set

$$\begin{aligned} \left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \psi_b &= 0 \\ \left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right] \psi_a &= 0 \end{aligned} \quad (31)$$

Then, we can take the operator $\left[-i\partial_x - \partial_y - z\frac{B}{2} \right]$ and apply it to ψ_a ,

$$\left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \psi_a = \tilde{a}_0 [-i\partial_x \lambda - \partial_y \lambda] e^\lambda - z\frac{B}{2} \tilde{a}_0 e^\lambda \quad (32)$$

Thus, in view of (12), (15) and the definition of z , it not difficult to arrive to following result

$$\left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \psi_a = -zB\tilde{a}_0 e^\lambda = -zB\psi_a \quad (33)$$

To continue we can take the state $-zB\tilde{a}_0 e^\lambda$ and apply the operator $\left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right]$. Then, we have,

$$\left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right] (-zB\tilde{a}_0 e^\lambda) = -2B\tilde{a}_0 e^\lambda = -2B\psi_a \quad (34)$$

From, (33), (34) we conclude that

$$\left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right] \left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \psi_a = -2B\psi_a \quad (35)$$

This is crucial point since indicates that ψ_a is an eigenstate of the operator $\left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right]$ with eigenvalue $-2B$. Thus, we can rename $-2B$ as E^2 so that

$$E = \pm \sqrt{-2B} \quad (36)$$

Then we can think the equations (33) and (34) as

$$\left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \psi_a = E \psi_{a,0} \quad (37)$$

$$\left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right] E \psi_{a,0} = E^2 \psi_a = -2B\psi_a \quad (38)$$

where $\psi_{a,0}$ may be obtained dividing $(-zB\tilde{a}_0 e^\lambda)$ by E ,

$$\psi_{a,0} = \frac{(-zB\tilde{a}_0 e^\lambda)}{\pm \sqrt{-2B}} = \pm \sqrt{\frac{-B}{2}} z \psi_a \quad (39)$$

Therefore, dividing the equation (38) by E , the equations (37) and (38) become

$$\left[-i\partial_x - \partial_y - z\frac{B}{2} \right] \psi_a = E \psi_{a,0} \quad (40)$$

$$\left[-i\partial_x + \partial_y - z^\dagger\frac{B}{2} \right] \psi_{a,0} = E \psi_a \quad (41)$$

Comparing these last two equations with (13) and (14), we see that

$$\Psi_{0,1} = \begin{pmatrix} \psi_{a,0} \\ \psi_a \end{pmatrix} \quad (42)$$

is an eigenstate of the Dirac-Weyl Hamiltonian with eigenvalue $\pm\sqrt{-2B}$. Here, it is important to point out that for the eigenvalue to make sense the magnetic field must be negative. This implies that $\Phi < 0$ which agrees with the result (27).

We can try to go even further and study the case in which ψ_a is the first order polynomial in z^\dagger , that is $\psi_a = (\tilde{a}_0 + \tilde{a}_1 z^\dagger)e^\lambda$, then we have the following zero mode,

$$\Psi_{1,0} = \begin{pmatrix} (\tilde{a}_0 + \tilde{a}_1 z^\dagger)e^\lambda \\ 0 \end{pmatrix} \quad (43)$$

Again, we can take the operator $\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]$ and apply it to ψ_a ,

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_a = -zB[(\tilde{a}_0 + \tilde{a}_1 z^\dagger)e^\lambda] - 2\tilde{a}_1 e^\lambda = -zB\psi_a - 2\tilde{a}_1 e^\lambda \quad (44)$$

Here we see that our result incorporates the term $-zB[(\tilde{a}_1 z^\dagger)e^\lambda] - 2\tilde{a}_1 e^\lambda$ to the result obtained in (33), of course, this term is due to the action of $\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]$ on $\tilde{a}_1 z^\dagger e^\lambda$. Also, it is interesting to note that, writing in terms of ψ_a , the result of (33) is $-zB\psi_a$ whereas in (44) we obtain $-zB\psi_a - 2\tilde{a}_1 e^\lambda$, that is, the result in (44) is not only reduced to $-zB\psi_a$, but also now appears the term $-2\tilde{a}_1 e^\lambda$.

Now, we can try to apply the operator $\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right]$ to $-zB\psi_a - 2\tilde{a}_1 e^\lambda$. This operation leads us to the following result,

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right](-zB\psi_a - 2\tilde{a}_1 e^\lambda) = -2B(\tilde{a}_0 + \tilde{a}_1 z^\dagger)e^\lambda = -2B\psi_a \quad (45)$$

Again the result obtained in (34) and (35) is repeated. Therefore, $-2B$ is an eigenvalue of $\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right]$ with eigenstate $\psi_a = (\tilde{a}_0 + \tilde{a}_1 z^\dagger)e^\lambda$. Thus, the equations (44) and (45) may be rewritten as

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_a = E \psi_{a,1} \quad (46)$$

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right](E \psi_{a,1}) = E^2 \psi_a = -2B\psi_a, \quad (47)$$

with,

$$E = \pm\sqrt{-2B}, \quad (48)$$

and where we have renamed $-zB\psi_a - 2\tilde{a}_1 e^\lambda$ as $E \psi_{a,1}$, which implies,

$$\psi_{a,1} = \frac{-zB\psi_a - 2\tilde{a}_1 e^\lambda}{\pm\sqrt{-2B}} = \pm\frac{1}{\sqrt{2}}\left[\sqrt{-B}z\psi_a - \frac{2\tilde{a}_1 e^\lambda}{\sqrt{-B}}\right] \quad (49)$$

So, the equations (46) and (47) reads as,

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_a = E \psi_{a,1} \quad (50)$$

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right]\psi_{a,1} = E \psi_a \quad (51)$$

Thus, we have constructed a new eigenstate

$$\Psi_{1,1} = \begin{pmatrix} \psi_{a,1} \\ \psi_a \end{pmatrix} \quad (52)$$

of the Dirac-Weyl Hamiltonian with eigenvalue $\pm\sqrt{-2B}$.

In order to obtain a general formula for the eigenstates of the Dirac-Weyl Hamiltonian, let us explore the case in which ψ_a is the second order polynomial in z^\dagger , that is $\psi_a = (\tilde{a}_0 + \tilde{a}_1 z^\dagger + \tilde{a}_2 (z^\dagger)^2)e^\lambda$. Then, after a bit of algebra, we have

$$\begin{aligned} \left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_a &= -zB[(\tilde{a}_0 + \tilde{a}_1 z^\dagger + \tilde{a}_2 (z^\dagger)^2)e^\lambda] + (-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda \\ &= -zB\psi_a + (-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda \end{aligned} \quad (53)$$

Notice that the result is composed by the term $-zB\psi_a$ plus a polynomial of first order in z^\dagger . If we compare this result with the results of formulas (33) and (44), we will notice that the result to apply the operator $\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]$ to the state ψ_a is composed by the term $-zB\psi_a$ plus a polynomial of a lower order in z^\dagger than ψ_a .

To proceed, we apply the operator $\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right]$ to the state $-zB\psi_a + (-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda$. Thus, we arrive to

$$\begin{aligned} & \left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right](-zB\psi_a + (-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda) \\ & = -2B(\tilde{a}_0 + \tilde{a}_1 z^\dagger + \tilde{a}_2 (z^\dagger)^2)e^\lambda = -2B\psi_a \end{aligned} \quad (54)$$

Thereby, we can construct the state $\psi_{a,2}$, dividing $-zB\psi_a + (-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda$ by $\pm\sqrt{-2B}$,

$$\psi_{a,2} = \frac{-zB\psi_a + (-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda}{\pm\sqrt{-2B}} = \pm \frac{1}{\sqrt{2}} \left[\sqrt{-B} z\psi_a + \frac{(-2\tilde{a}_1 - 4\tilde{a}_2 z^\dagger)e^\lambda}{\sqrt{-B}} \right] \quad (55)$$

and the eigenstate of the Dirac-Weyl Hamiltonian as

$$\Psi_{2,1} = \begin{pmatrix} \psi_{a,2} \\ \psi_a \end{pmatrix} \quad (56)$$

It is not difficult to generalize this idea to the case

$$\psi_a = \sum_{i=0}^j \tilde{a}_i (z^\dagger)^i e^\lambda \quad (57)$$

with

$$j \leq N \quad (58)$$

The result is

$$\psi_{a,j} = \pm \frac{1}{\sqrt{2}} \left[\sqrt{-B} z\psi_a - \frac{1}{\sqrt{-B}} \sum_{i=0}^j 2i \tilde{a}_i (z^\dagger)^{i-1} e^\lambda \right] \quad (59)$$

Therefore, we can create another new eigenstate of the Dirac-Weyl Hamiltonian,

$$\Psi_{j,1} = \begin{pmatrix} \psi_{a,j} \\ \psi_a \end{pmatrix} \quad (60)$$

with eigenvalue $\pm\sqrt{-2B}$. When $j = N$ we have

$$\Psi_{N,1} = \begin{pmatrix} \psi_{a,N} \\ \psi_a \end{pmatrix} \quad (61)$$

Here, it is important to remark that the state (61) is $(N + 1)$ -th eigenstate that we create and due to the Aharonov-Casher theorem we cannot create another independent eigenstate. So, for a level of energy $E = \sqrt{-2B}$ we can only have $(N + 1)$ -th independent eigenstates and in similar way for the level $E = -\sqrt{-2B}$ we have $(N + 1)$ -th independent eigenstates associated to it. Another interesting aspect shows that the eigenstates $\Psi_{j,1}$ depend only on ψ_a i.e. an eigenstate of positive chirality. Hence, it is not surprising that for the eigenvalue to make sense the magnetic field must be negative. This agrees with the result (27) and involves a extension of the Aharonov-Casher result.

4. The first level of energy and its eigenstates for positive magnetic field

Let us consider how to construct eigenstates of the Dirac-Weyl Hamiltonian starting now by a simplest spinor of negative chirality, i.e.

$$\Psi_{0,0} = \begin{pmatrix} 0 \\ a_0 e^{-\lambda} \end{pmatrix} \quad (62)$$

as the spinor of (30) it is a zero mode of the Dirac-Weyl Hamiltonian. The idea, now, is to apply the operator $\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right]$ to ψ_b . This can be done easily and leads us to,

$$\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right]\psi_b = -a_0 B z^\dagger e^{-\lambda} = -z^\dagger B\psi_b \quad (63)$$

Following the same idea we take $-z^\dagger B\psi_b$ and apply the operator $\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right]$,

$$\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right](-z^\dagger B\psi_b) = 2a_0 B e^{-\lambda} = 2B\psi_b \quad (64)$$

Thus, if we compare with (34) and (35) we see that the Schrödinger equation

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right]\psi_b = 2B\psi_b \quad (65)$$

has now an eigenvalue with the same absolute value but different sign, i.e. $2B$. Therefore, we have that the energy, now reads

$$E = \pm\sqrt{2B}, \quad (66)$$

so that the equations (63) and (81), can be arranged to give,

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right]\psi_b = E\psi_{b,0} \quad (67)$$

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]E\psi_{b,0} = E^2\psi_b \quad (68)$$

where

$$\psi_{b,0} = \frac{-z^\dagger B\psi_b}{\pm\sqrt{2B}} = \mp\sqrt{\frac{B}{2}}z^\dagger\psi_b \quad (69)$$

Hence, dividing (68) by E , we have, the set following set of equations

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2}\right]\psi_b = E\psi_{b,1} \quad (70)$$

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_{b,0} = E\psi_b \quad (71)$$

We can compare this equations with (13) and (14). Then, it is clear that

$$\Psi_{0,1} = \begin{pmatrix} \psi_b \\ \psi_{b,0} \end{pmatrix} \quad (72)$$

is an eigenstate of the Dirac-Weyl Hamiltonian with eigenvalue $\pm\sqrt{2B}$. In contrast with the previous section, here, the magnetic field must be positive, otherwise the eigenvalue becomes meaningless. This implies that $\Phi > 0$, and again, it agrees with the result of the Aharonov and Casher theorem.

As in the previous section the idea may be generalized. Indeed, if we take

$$\psi_b = \sum_{i=0}^j a_i (z^\dagger)^i e^{-\lambda} \quad (73)$$

and follow the same procedure of the previous section, we can create a j -th eigenstate, with eigenvalue $E = \pm\sqrt{2B}$, of the Dirac-Weyl Hamiltonian

$$\Psi_{j,1} = \begin{pmatrix} \psi_b \\ \psi_{b,j} \end{pmatrix} \quad (74)$$

where

$$\psi_{b,j} = \mp\frac{1}{\sqrt{2}}\left[\sqrt{B}z^\dagger\psi_b - \frac{1}{\sqrt{B}}\sum_{i=0}^j 2i a_i (z^\dagger)^{i-1} e^{-\lambda}\right] \quad (75)$$

Again, due to the Aharonov and Casher theorem, $j \leq N$, so that we can, only, create $N + 1$ independent eigenstates associated to the eigenvalue $E = \sqrt{2B}$ and another $N + 1$ associated to level $E = -\sqrt{2B}$

5. The generalization for higher energy levels

Let us concentrate in the generalization of the procedure studied in the previous section. We can start by considering the second level of energy. In order to proceed consider the equations (70) and (71) and repeat the procedure of the previous section, i.e. take the second component of the spinor

$$\Psi_{0,1} = \begin{pmatrix} \psi_b \\ \psi_{b,0} \end{pmatrix} \quad (76)$$

and apply the operator $\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right]$ to this state. Then, we have,

$$\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right] \psi_{b,0} = \mp \sqrt{\frac{B}{2}} \left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right] (z^\dagger \psi_b) \quad (77)$$

Since, $[-i\partial_x + \partial_y] z^\dagger = 0$ and

$$\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right] \psi_b = E_1 \psi_{b,0} \quad (78)$$

where $E_1 = \pm \sqrt{2B}$, it is not difficult to arrive to

$$\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right] \psi_{b,0} = -Bz^\dagger \psi_{b,0} \quad (79)$$

We can continue with the procedure and take the state $-Bz^\dagger \psi_{b,0}$ and apply the operator $\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right]$,

$$\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right] (-Bz^\dagger \psi_{b,0}) = 2B\psi_{b,0} - Bz^\dagger \left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right] \psi_{b,0} \quad (80)$$

Here, we can consider the formula (68). Then, the term of (80) can be rewritten as

$$\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right] (-Bz^\dagger \psi_{b,0}) = 2B\psi_{b,0} \mp B\sqrt{2B} z^\dagger \psi_b \quad (81)$$

The second term of (81) may be arranged as

$$\mp B\sqrt{2B} z^\dagger \psi_b = 2B\psi_{b,0}, \quad (82)$$

so that

$$\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right] (-Bz^\dagger \psi_{b,0}) = 4B\psi_{b,0} \quad (83)$$

If we rename $4B$ as E_2^2 , and create a new state

$$\psi_{b,1} = \mp \frac{\sqrt{B}}{2} z^\dagger \psi_{b,0}, \quad (84)$$

then, the equations (79) and (83) reads as

$$\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right] \psi_{b,0} = E_2 \psi_{b,1} \quad (85)$$

$$E_2 \left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right] \psi_{b,1} = E_2^2 \psi_{b,0} \quad (86)$$

Dividing the last equation by E_2 , it becomes clear that

$$\Psi_{0,2} = \begin{pmatrix} \psi_{b,0} \\ \psi_{b,1} \end{pmatrix} \quad (87)$$

is an eigenstate of the Dirac-Weyl Hamiltonian with eigenvalue $\pm 2\sqrt{B}$. In general, we can get the spinor (74).

Since, it is a solution of the Dirac-Weyl equation, it satisfy

$$\left[[-i\partial_x + \partial_y] - z^\dagger \frac{B}{2}\right] \psi_b = E_1 \psi_{b,j} \quad (88)$$

$$\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right] \psi_{b,j} = E_1 \psi_b \quad (89)$$

We can repeat the previous steps to obtain

$$\begin{aligned} \left[\bar{\partial} - z^\dagger \frac{B}{2}\right] \psi_{b,j} &= \mp \sqrt{\frac{B}{2}} \left[\bar{\partial} - z^\dagger \frac{B}{2}\right] (z^\dagger \psi_b) \\ &\pm \frac{1}{\sqrt{2B}} \sum_{i=0}^j 2i a_i \left[\bar{\partial} - z^\dagger \frac{B}{2}\right] ((z^\dagger)^{i-1} e^{-\lambda}) \end{aligned} \quad (90)$$

Here ψ_b is dictated by the formula (73) and $\bar{\partial} = [-i\partial_x + \partial_y]$. The formula (90) may be developed easily if we consider the following equalities,

$$\begin{aligned}
\bar{\partial}z^\dagger &= 0 \\
\left[\bar{\partial} - z^\dagger \frac{B}{2}\right]\psi_b &= E_1 \psi_{b,j} \\
\left[\bar{\partial} - z^\dagger \frac{B}{2}\right]e^{-\lambda} &= -Bz^\dagger e^{-\lambda}
\end{aligned} \tag{91}$$

then, it is not difficult to show that,

$$\left[\bar{\partial} - z^\dagger \frac{B}{2}\right]\psi_{b,j} = -Bz^\dagger \psi_{b,j} \mp \sqrt{2B} \sum_{i=0}^j i a_i (z^\dagger)^i e^{-\lambda} \tag{92}$$

Now, the application of the operator $\left[\partial - z \frac{B}{2}\right]$, where we have named $\partial = [-i\partial_x - \partial_y]$, on the state $-Bz^\dagger \psi_{b,j} \mp \sqrt{2B} \sum_{i=0}^j i a_i (z^\dagger)^i e^{-\lambda}$ leads us to

$$\begin{aligned}
\left[\partial - z \frac{B}{2}\right] \left(-Bz^\dagger \psi_{b,j} \mp \sqrt{2B} \sum_{i=0}^j i a_i (z^\dagger)^i e^{-\lambda}\right) &= -B \left[\partial - z \frac{B}{2}\right] (z^\dagger \psi_{b,j}) \\
&\mp \sqrt{2B} \sum_{i=0}^j i a_i \left[\partial - z \frac{B}{2}\right] ((z^\dagger)^i e^{-\lambda})
\end{aligned} \tag{93}$$

To proceed we use the following identities

$$\begin{aligned}
\partial z^\dagger &= -2 \\
\left[\partial - z \frac{B}{2}\right]\psi_{b,j} &= E_1 \psi_b \\
\left[\partial - z \frac{B}{2}\right]e^{-\lambda} &= 0,
\end{aligned} \tag{94}$$

then equation (93) may be developed to give

$$\left[\partial - z \frac{B}{2}\right] \left(-Bz^\dagger \psi_{b,j} \mp \sqrt{2B} \sum_{i=0}^j i a_i (z^\dagger)^i e^{-\lambda}\right) = 4B\psi_{b,j} \tag{95}$$

The equations (92) and (97) are arranged, so we have

$$\left[\bar{\partial} - z^\dagger \frac{B}{2}\right]\psi_{b,j} = E_2 \psi_{b,j+1} \tag{96}$$

$$\left[\partial - z \frac{B}{2}\right](E_2 \psi_{b,j+1}) = 4B\psi_{b,j} \tag{97}$$

where, as in (85) and (86), $E_2 = \pm 2\sqrt{B}$ and the state $\psi_{b,j+1}$ is defined as

$$\psi_{b,j+1} = \frac{1}{\pm 2\sqrt{B}} \left(-Bz^\dagger \psi_{b,j} \mp \sqrt{2B} \sum_{i=0}^j i a_i (z^\dagger)^i e^{-\lambda}\right) \tag{98}$$

Thus, we have that the state,

$$\Psi_{j,2} = \begin{pmatrix} \psi_{b,j} \\ \psi_{b,j+1} \end{pmatrix} \tag{99}$$

is an eigenstate of the Dirac-Weyl Hamiltonian with eigenvalue $\pm 2\sqrt{B}$. In this way, we can obtain all independent state of the second Landau level of the energy $\pm 2\sqrt{B}$. Again, it is interested to note that there are $N + 1$ independent states with energy $E_2 = 2\sqrt{B}$ and another $N + 1$ associated to the energy $E_2 = -2\sqrt{B}$. We can repeat the procedure to create the eigenstates associated to the third levels of energy and so on. In addition, it is not difficult to imagine the same procedure for the case of eigenvalues and eigenstates associated to negative magnetic field configuration. In this case, we would take the first component of the spinor (60) and would apply the operator $\left[[-i\partial_x - \partial_y] - z \frac{B}{2}\right]$, in order to obtain a state $\psi_{a,j+1}$ and a new spinor

$$\Psi_{j,2} = \begin{pmatrix} \psi_{a,j+1} \\ \psi_{a,j} \end{pmatrix} \tag{100}$$

which is the $j + 1$ -th eigenstate of the Dirac-Weyl Hamiltonian associated to the second level of energy $E_2 = \pm 2\sqrt{-B}$. Of course, as for the states associated positive magnetic field configuration, there are $N + 1$ independent states with energy $E_2 = 2\sqrt{-B}$ and another $N + 1$ associated to the energy $E_2 = -2\sqrt{-B}$.

6. Application to bilayer graphene

To conclude, we discuss the application of our mechanism to study the case of bilayer Hamiltonian. The bilayer graphene [40–46] in the simplest approximation can be considered as a zero-gap semiconductor with parabolic touching of the electron and hole bands described by the single-particle Hamiltonian. By exfoliation of graphene one can obtain several layers of carbon atoms. Its electronic structure can be understood in the framework of a tight-binding model.

Let us then consider the Hamiltonian for the bilayer graphene,

$$\begin{pmatrix} 0 & (D_1 - iD_2)^2 \\ (D_1 + iD_2)^2 & 0 \end{pmatrix} \begin{pmatrix} \psi_a \\ \psi_b \end{pmatrix} = E \begin{pmatrix} \psi_a \\ \psi_b \end{pmatrix} \quad (101)$$

This description is accurate at the energy scale larger than a few meV, otherwise a more complicated picture including trigonal warping takes place; we will restrict ourselves only by the case of not too small doping when the approximate Hamiltonian (123) works. Two components of the wave function originated from the crystallographic structure of graphite sheets with two carbon atoms in the sheet per elementary cell. There are two touching points per Brillouin zone: K and K'. For smooth enough external potential, no Umklapp processes between these points are allowed and thus they can be considered independently.

Thus, equation (123) may be rewritten as

$$(D_1 - iD_2)^2 \psi_b = E \psi_a \quad (102)$$

$$(D_1 + iD_2)^2 \psi_a = E \psi_b \quad (103)$$

Following the same procedure for the single-layer Hamiltonian we should find the eigenfunctions of the Hamiltonian (123) for zero energy in the symmetric gauge, that is,

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2} \right]^2 \psi_b = 0 \quad (104)$$

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2} \right]^2 \psi_a = 0 \quad (105)$$

The solution of these equations can be found easily from the development done in section 3. Indeed, we can check that

$$\psi_b = z^\dagger \sum_{i=0}^j a_i z^i e^{-\lambda} \quad (106)$$

and

$$\psi_a = z \sum_{i=0}^j \tilde{a}_i (z^\dagger)^i e^{\lambda} \quad (107)$$

satisfy (104) and (105). In order to construct the eigenstates different from the zero modes we can start by considering the simplest solution in a negative field background. According to what we see in the previous sections, this solution should be a spinor

$$\Psi_{0,0} = \begin{pmatrix} z\tilde{a}_0 e^{\lambda} \\ 0 \end{pmatrix} \quad (108)$$

which is the simplest zero mode of the Hamiltonian (123). Again, we can take the operator $\left[[-i\partial_x - \partial_y] - z\frac{B}{2} \right]$ and apply it to ψ_a . Thus, using the result of section 3 we obtain

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2} \right] \psi_a = -z^2 B \tilde{a}_0 e^{\lambda} \quad (109)$$

Applying the operator $\left[[-i\partial_x - \partial_y] - z\frac{B}{2} \right]$ one more time, lead us to

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2} \right]^2 \psi_a = z^3 B^2 \tilde{a}_0 e^{\lambda} \quad (110)$$

Following the same reasoning carried out in the section 3, we should apply the operator $\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2} \right]^2$ to the polynomial $z^3 B^2 \tilde{a}_0 e^{\lambda}$. Again, using the results of the section section 3 and after a bit of algebra we arrive to

$$\left[[-i\partial_x + \partial_y] - z^\dagger\frac{B}{2} \right] (z^3 B^2 \tilde{a}_0 e^{\lambda}) = 6B^2 z^2 \tilde{a}_0 e^{\lambda} \quad (111)$$

The application of the operator $\left[[-i\partial_x + \partial_y] - z^{\dagger\frac{B}{2}}\right]^2$ for the second time lead us to following result

$$\left[[-i\partial_x + \partial_y] - z^{\dagger\frac{B}{2}}\right](6B^2z^2\tilde{a}_0e^\lambda) = 24B^2z\tilde{a}_0e^\lambda = 24B^2\psi_a \quad (112)$$

Renamed, $24B^2$ as E^2 we have,

$$E = \pm\sqrt{24}|B| \quad (113)$$

Then, the equations (110) and (112) may be rewritten as follow

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]^2\psi_a = E\psi_{a,0} \quad (114)$$

$$\left[[-i\partial_x + \partial_y] - z^{\dagger\frac{B}{2}}\right]^2\psi_{a,0} = E\psi_a \quad (115)$$

where $\psi_{a,0}$ is obtained by dividing $(z^3B^2\tilde{a}_0e^\lambda)$ by E ,

$$\psi_{a,0} = \frac{(z^3B^2\tilde{a}_0e^\lambda)}{\pm\sqrt{24}B} = \pm\frac{z^3B\tilde{a}_0e^\lambda}{\sqrt{24}} \quad (116)$$

This show that the spinor

$$\Psi_{0,1} = \begin{pmatrix} \psi_{a,0} \\ \psi_a \end{pmatrix} \quad (117)$$

is an eigenstate of the Hamiltonian (123) with eigenvalue $\pm\sqrt{24}B$. In general, we can start by considering a zero mode of the form

$$\Psi_{0,1} = \begin{pmatrix} z\sum_{i=0}^j \tilde{a}_i(z^\dagger)^i e^\lambda \\ 0 \end{pmatrix} \quad (118)$$

Then, we apply the operator $\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]$ to $z\sum_{i=0}^j \tilde{a}_i(z^\dagger)^i e^\lambda$. Thus, in view of the equation (59) and $[-i\partial_x - \partial_y]z = 0$ it not difficult to see that

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]z\sum_{i=0}^j \tilde{a}_i(z^\dagger)^i e^\lambda = \pm z\sqrt{-2B}\psi_{a,j} \quad (119)$$

where $\psi_{a,j}$ is dictated by the formula (59). The application of the operator $\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]$ to $\pm z\sqrt{-2B}\psi_{a,j}$ leads to

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]^2z\sum_{i=0}^j \tilde{a}_i(z^\dagger)^i e^\lambda = \pm z\sqrt{-2B}\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_{a,j} \quad (120)$$

where

$$\left[[-i\partial_x - \partial_y] - z\frac{B}{2}\right]\psi_{a,j} = \pm\frac{1}{\sqrt{2}}\left[|B|\sqrt{2}z\psi_{a,j} - \frac{1}{\sqrt{-B}}z^{-1}2i(\pm\sqrt{-2B})\psi_{a,j}\right] \quad (121)$$

The same mechanism can be repeated starting from the zero mode

$$\Psi_{j,0} = \begin{pmatrix} 0 \\ \sum_{i=0}^j a_i z^i e^{-\lambda} \end{pmatrix} \quad (122)$$

In addition, following the same steps of the section 5, we can obtain the higher levels of energy and its eigenstates.

The generalization of this formalism to a multilayer graphene, can be easily understood, from the form of the multilayer Hamiltonian equation [42–46]

$$\begin{pmatrix} 0 & (D_1 - iD_2)^J \\ (D_1 + iD_2)^J & 0 \end{pmatrix} \begin{pmatrix} \psi_a \\ \psi_b \end{pmatrix} = E \begin{pmatrix} \psi_a \\ \psi_b \end{pmatrix} \quad (123)$$

where J is a positive integer. Then we can consider the solutions of the bilayer graphene and repeat the same formalism to obtain the solutions for a 3-layer graphene system and so on for an arbitrary number of layers.

7. Conclusion

In summary, we have developed a formalism to construct the eigenvalues and its eigenfunctions of the Dirac-Weyl equation for the Landau problem in the symmetric gauge. We have shown that for negative magnetic field configurations the eigenstates may be constructed from the zero mode with positive chirality whereas for positive magnetic field the eigenstates are constructed from the zero mode with negative chirality. This is a consequence of the Aharonov-Casher theorem, which establishes that for a negative magnetic flux the zero-energy solutions can exist only for positive spin direction, whereas if the magnetic flux is positive the zero-energy solutions can exist only for negative spin direction.

In addition, we showed that our mechanism may be generalized to study the Hamiltonians of multilayer graphene systems, in such a way to obtain the eigenfunctions and eigenvalues for arbitrary numbers of graphene layers. This is important due to the role that Landau levels play in graphene. The bilayer Hamiltonian is different both from nonrelativistic (Schrodinger) and from relativistic (Dirac) cases. The eigenstates of this Hamiltonian have very special chiral properties [42], resulting in a special Landau quantization and special scattering. In this sense, our method becomes a useful tool in order to obtain the Landau levels for bilayer graphene. Although, in this note we do not study in detail the bilayer graphene, it would be interesting to deal with more emphasis the study of Landau levels in bilayer and multilayer graphene. This becomes of particular importance due to the unconventional quantum Hall effect in bilayer graphene [40, 42], where Landau quantization of the fermions results in plateaus in Hall conductivity at standard integer positions, being missing the last (zero-level) plateau.

Acknowledgments

This work is supported by CONICET.

ORCID iDs

Lucas Sourrouille  <https://orcid.org/0000-0002-8399-6346>

References

- [1] Darwin C G 1930 *Proc. Cambridge Philos. Soc.* **27** 86
- [2] Fock V 1928 *Z. Phys.* **47** 446
- [3] Landau L D 1930 Paramagnetism of Metals *Z. Phys.* **64** 623–37
- [4] Girvin S M 2000 The Quantum Hall Effect: Novel Excitations and Broken Symmetries *Topological Aspects of Low Dimensional Systems* (Berlin: Springer-Verlag)
- [5] Goerbig M O arXiv:0909.1998
- [6] Prange and Girvin 1990 *The Quantum Hall Effect* (Springer-Verlag New York Inc.)
- [7] Jain K 2007 *Composite Fermions* (Cambridge: Cambridge University Press)
- [8] Tong D Lectures on the Quantum Hall Effect arXiv:1606.06687
- [9] Iordansky S V 1982 On the conductivity of two-dimensional electrons in a strong magnetic field *Solid State Commun.* **43** 1
- [10] Kazarinov R F and Luryi S 1982 Quantum percolation and quantization of Hall resistance in two-dimensional electron gas *Phys. Rev. B* **25** 7626
- [11] Prange R E and Joynt R 1982 Conduction in a strong field in two dimensions: the quantum Hall effect *Phys. Rev. B* **25** 2943
- [12] Trugman S A 1983 Localization, percolation, and the quantum Hall effect *Phys. Rev. B* **27** 7539
- [13] Laughlin R B 1981 Quantized Hall conductivity in two dimensions *Phys. Rev. B* **23** 5632
- [14] Halperin B I 1982 Quantized Hall conductance, current-carrying edge states, and the existence of extended states in a two-dimensional disordered potential *Phys. Rev. B* **25** 2185
- [15] Laughlin R B 1983 Anomalous quantum Hall effect: an incompressible quantum fluid with fractionally charged excitations *Phys. Rev. Lett.* **50** 1395
- [16] Laughlin R B 2005 *A Different Universe: Reinventing Physics from the Bottom Down* (New York: Basic Books)
- [17] Jain K 2000 The composite fermion: a quantum particle and its quantum fluids *Phys. Today* **53** 39
- [18] Jain J K 2003 The role of analogy in unraveling the fractional-quantum-Hall-effect mystery *Physica E* **20** 79
- [19] Novoselov K S *et al* 2004 *Science* **306** 666
Novoselov K S *et al* 2005 *Nature (London)* **438** 197
- [20] Zhang Y, Tan Y W, Stormer H and Kim P 2005 *Nature (London)* **438** 201
- [21] Berger C *et al* 2006 *Science* **312** 1191
- [22] Castro Neto A H, Guinea F, Peres N M, Novoselov K S and Geim A K 2009 *Rev. Mod. Phys.* **81** 109
- [23] Gusynin V P and Sharapov S G 2005 *Phys. Rev. Lett.* **95** 146801
- [24] Peeters F M and Matulis A 1993 *Phys. Rev. B* **48** 15166
- [25] Pereira J M, Mlinar V, Peeters F M and Vasilopoulos P 2006 *Phys. Rev. B* **74** 045424
- [26] Lukose V, Shankar R and Baskaran G 2007 *Phys. Rev. Lett.* **98** 116802
- [27] De Martino A, Dell'Anna L and Egger R 2007 *Phys. Rev. Lett.* **98** 066802
- [28] Ramezani Masir M, Vasilopoulos P, Matulis A and Peeters F M 2008 *Phys. Rev. B* **77** 235443
- [29] Ghosh T K, DeMartino A, Hausler W, DellAnna L and Egger R 2008 *Phys. Rev. B* **77** 081404(R)
- [30] Oroszlany L, Rakytá P, Kormanyos A, Lambert C J and Cserti J 2008 *Phys. Rev. B* **77** 081403

- [31] Ghosh T K 2009 *J. Phys. Condens. Matter* **21** 045505
- [32] Giavaras G, Maksym P A and Roy M 2009 *J. Phys. Condens. Matter* **21** 102201
- [33] Kuru S, Negro J and Nieto L M 2009 *J. Phys. Condens. Matter* **21** 455305
- [34] Katsnelson M 2012 *Graphene: Carbon in two Dimensions* (Cambridge: Cambridge University Press)
- [35] Eshghi M and Mehraban H 2017 *Comptes Rendus Physique* **18** 47
- [36] de Brito P E and Nazareno H N 2007 *Eur. J. Phys.* **28** 9
- [37] de Brito P E and Nazareno H N 2001 *Phys. Rev. B* **64** 045112
- [38] Feldman A and Kahn A H 1969 *Phys. Rev. B* **1** 4584
- [39] Jain J K 2007 *Composite Fermions* (Cambridge: Cambridge University Press)
- [40] McCann E and Falco V I 2006 *Phys. Rev. Lett.* **96** 086805
- [41] Guinea F, Castro A H and Peres N M R 2006 *Phys. Rev. B* **73** 245426
- [42] Novoselov K S, McCann E, Morozov S V, Falco V I, Katsnelson M I, Zeitler U, Jiang D, Schedin F and Geim A K 2006 *Nat. Phys.* **2** 177–80
- [43] Koshino M and Ando T 2007 *Phys. Rev. B* **76** 085425
- [44] McCann E, Abergel D S L and Falco V I 2007 *Solid State Commun.* **143** 110–5
- [45] Min H and MacDonald A H 2008 *Phys. Rev. B* **77** 155416
- [46] McCann E and Koshino M 2013 *Rep. Prog. Phys.* **76** 056503
- [47] Aharonov Y and Casher A 1979 *Phys. Rev. A* **19** 2461