



Published in final edited form as:

Biochem Biophys Res Commun. 2015 November 6; 467(1): 1–6. doi:10.1016/j.bbrc.2015.09.144.

Intracellular Ca²⁺ oscillations generated via the extracellular Ca²⁺-sensing receptor (CaSR) in response to extracellular Ca²⁺ or L-phenylalanine: impact of the highly conservative mutation Ser170Thr

Steven H. Young^a, Osvaldo Rey^b, and Enrique Rozengurt^{a,c}

^aDivision of Digestive Diseases, Department of Medicine; CURE: Digestive Diseases Research Center, David Geffen School of Medicine and Molecular Biology Institute, University of California at Los Angeles, Los Angeles, CA 90095, USA

^cVeterans Affairs West Los Angeles Healthcare System, Los Angeles, CA, USA

^bInstitute of Immunology, Genetics and Metabolism, Consejo Nacional de Investigaciones Cientificas y Tecnicas, Universidad de Buenos Aires, Buenos Aires, Argentina

Abstract

The extracellular Ca²⁺-sensing receptor (CaSR) is an allosteric protein that responds to changes in the extracellular concentration of Ca²⁺ ([Ca²⁺]_e) and aromatic amino acids with the production of different patterns of oscillations in intracellular Ca²⁺ concentration ([Ca²⁺]_i). An increase in [Ca²⁺]_e stimulates sinusoidal oscillations in [Ca²⁺]_i whereas aromatic amino acid-induced CaR activation in the presence of a threshold [Ca²⁺]_e promotes transient oscillations in [Ca²⁺]_i. Here, we examined spontaneous and ligand-evoked [Ca²⁺]_i oscillations in single HEK-293 cells transfected with the wild type CaSR or with a mutant CaSR in which Ser170 was converted to Thr (CaSR_{S170T}). Our analysis demonstrates that cells expressing CaSR_{S170T} display [Ca²⁺]_i oscillations in the presence of low concentrations of extracellular Ca²⁺ and respond to L-Phe with robust transient [Ca²⁺]_i oscillations. Our results indicate that the S170T mutation induces a marked increase in CaSR sensitivity to [Ca²⁺]_e and imply that the allosteric regulation of the CaSR by aromatic amino acids is not only mediated by an heterotropic positive effect on Ca²⁺ binding cooperativity but, as biased agonists, aromatic amino acids stabilize a CaSR conformation that couples to a different signaling pathway leading to transient [Ca²⁺]_i oscillations.

Keywords

Ca²⁺ oscillations; amino acid signaling; CaSR mutant S170T; allosteric regulation

*To whom correspondence should be addressed at: 900 Veteran Avenue, Warren Hall Room 11-124, Department of Medicine, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA 90095-1786, USA, Tel: (310) 794-6610; erozengurt@mednet.ucla.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

The extracellular Ca^{2+} -sensing receptor (CaSR), a member of the C family of heptahelical G protein-coupled receptors (GPCRs), is an allosteric protein that plays a major physiological role in correcting small changes in extracellular concentration of Ca^{2+} ($[\text{Ca}^{2+}]_e$) by inhibiting parathyroid hormone secretion and renal Ca^{2+} absorption [1]. Interestingly, subsequent studies demonstrated that the CaSR recognizes other ligands and is expressed in many other tissues and organs, including the gastrointestinal tract, brain, pituitary, thyroid, skin, breast, pancreas, lung, bone, and heart [2], suggesting that this receptor plays additional, yet less well defined, physiological roles in the regulation of normal and abnormal cell function [3,4,5,6]. Indeed, recent evidence indicates that the CaSR is implicated in the negative control of colon cell proliferation [7] and cancer [8], nutrient sensing [4], epithelial transport [9], inflammation [10], bone turnover [11] and stem cell differentiation [6]. Thus, the signaling mechanisms triggered via CaSR activation are attracting intense attention.

A number of studies of CaSR activation in individual living cells have shown that intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) oscillates upon stimulation of CaSR by an elevation in $[\text{Ca}^{2+}]_e$ within a physiological range [12,13,14,15,16,17,18]. Oscillatory changes in $[\text{Ca}^{2+}]_i$ in response to receptor stimulation is a fundamental mechanism of cell signaling implicated in the regulation of Ca^{2+} - and calmodulin-dependent protein kinase II [19], conventional protein kinase C (PKC) isoforms [15,20], mitochondrial function [21,22], and nuclear transcriptional activity leading to differential gene expression [23,24]. We proposed that $[\text{Ca}^{2+}]_i$ oscillations induced by activation of the CaSR in response to an increase in extracellular Ca^{2+} results from negative feedback involving PKC-mediated phosphorylation of the CaSR at the inhibitory residue Thr⁸⁸⁸ [14,25]. In addition to its role as sensor of $[\text{Ca}^{2+}]_e$, the CaSR is also stimulated by aromatic amino acids [26] which, like $[\text{Ca}^{2+}]_e$, induce striking and lasting CaSR-mediated $[\text{Ca}^{2+}]_i$ oscillations [13,15,16]. However, the patterns of $[\text{Ca}^{2+}]_i$ oscillations induced by these agonists are different. Aromatic amino acid stimulation of the CaSR in the presence of a threshold $[\text{Ca}^{2+}]_e$ induces repetitive, low frequency $[\text{Ca}^{2+}]_i$ spikes that return to the base-line level, a pattern known as transient oscillations [13,15,16]. The amplitude, frequency, and duration of $[\text{Ca}^{2+}]_i$ oscillations are increasingly recognized to encode important information for a variety of biological processes, including metabolism and gene expression.

The most striking structural feature of the family C GPCR is its large extracellular domain, which consists of ~600 amino acid residues arranged in two predicted lobes referred as the Venus Flytrap Domain (VFTD), the site of numerous disease-causing mutations in this receptor [27,28]. The VFTD contains multiple binding sites for extracellular Ca^{2+} [29] and the allosteric binding site for amino acids. The residues Ser-147, Ser-170, Asp-190, Tyr-218 and Glu-297 of the VFTD comprise the first Ca^{2+} binding site which is considered the most important and operational under physiological conditions [27,30]. A number of studies identified Ser-170 as playing a critical role in Ca^{2+} binding and signal generation since mutation of Ser-170 to Ala completely abolished the response of the mutant CaSR to even very high $[\text{Ca}^{2+}]_e$ [31] without affecting its membrane expression [30,31]. In contrast, a conservative mutation of Ser-170 to Thr (S170T) appeared to have very little effect on CaSR

responsiveness to an increase in $[Ca^{2+}]_e$ but impaired L-phenylalanine (L-Phe) sensing [29,31] though the ability of this CaSR mutant to mediate transient oscillations in single cells was not examined. The detailed study of the impact of mutations in the N-terminal VFTD is of importance to advance understanding of the molecular mechanisms of CaSR regulation.

Here, we examined spontaneous and ligand-evoked $[Ca^{2+}]_i$ oscillations in single HEK-293 cells transfected with the wild type CaSR or with CaSR in which Ser170 was mutated to Thr (CaSR_{S170T}). Our analysis demonstrates that cells expressing CaSR_{S170T} display $[Ca^{2+}]_i$ oscillations in the presence of low concentrations of extracellular Ca^{2+} and respond to L-Phe with robust transient $[Ca^{2+}]_i$ oscillations. We propose that the S170T mutation increases CaSR sensitivity to $[Ca^{2+}]_e$ and separates the positive heterotropic effect that enhances the apparent affinity of the CaSR for extracellular Ca^{2+} from the stabilization of a receptor conformation that promotes transient oscillations in the presence of a threshold $[Ca^{2+}]_e$.

2. Materials and Methods

2.1 Materials

Fura 2-AM, Dulbecco's Modified Eagle's Medium (DMEM), and Hanks Balanced Salt Solution (HBSS) were obtained from Invitrogen (Carlsbad, CA).

2.2 Cell culture and transfection

Human Embryonic Kidney (HEK-293) cells were maintained in DMEM supplemented with 10% fetal bovine serum (FBS) in a humidified incubator under 10% CO₂ and 90% air at 37°C, as described previously [13,25,32]. For experimental purposes, cells were plated onto 18-mm diameter glass coverslips inside 35-mm plastic dishes, where they could be dually transiently transfected with a plasmid encoding the human CaSR or a mutant CaSR receptor (CaSR_{S170T}). Site-directed mutagenesis to convert Ser170 of the CaSR into Thr was performed as previously described [33]. Identification of cells transiently transfected with pCR3.1-CaSRwt (0.5 µg/dish) or pCR3.1-CaSR_{S170T} (0.5 µg/dish) was achieved by co-transfection with pDsRed-Express (BD Biosciences) (0.5 µg/dish), a vector that encodes a red fluorescent protein. After 16 h, the cultures were loaded with the Ca^{2+} indicator Fura-2 as described below.

2.3 Measurement of $[Ca^{2+}]_i$

$[Ca^{2+}]_i$ was measured in single cells loaded with the calcium indicator fura-2 as previously described [13]. Briefly, cells were incubated in saline solution containing 138 mM NaCl, 4 mM NaHCO₃, 0.3 mM Na₂HPO₄, 5 mM KCl, 0.3 mM KH₂PO₄, 1.5 mM CaCl₂, 0.5 mM MgCl₂, 0.4 mM MgSO₄, 5.6 mM D-glucose, 20 mM HEPES, pH: 7.4 which was supplemented with 5 µM fura-2 AM for 45–60 min at 37°C before $[Ca^{2+}]_i$ imaging. The cells were then washed and placed in an experimental chamber that was perfused with saline solution at 1.5 ml/min at 37°C. The chamber in turn was placed on the stage of an inverted microscope connected to a digital imaging system. Ratios of images (340 nm excitation/ 380 nm excitation, emission filter 520 nm) were obtained at 1.5 sec intervals. A region of interest covering 15 µm × 15 µm was defined over each cell, and the average ratio intensity

over the region was converted to $[Ca^{2+}]_i$ using a standard curve constructed with a series of calibrated buffered calcium solutions (Calcium Calibration Buffer Kit #2, Invitrogen Corp).

2.4 Western blots and indirect immunofluorescence

HEK-293 cells were transfected with vectors encoding wild type human CaSR or CaSR_{S170T} at 0.5 μ g/ dish and analyzed by Western blot 16 h post-transfection using a murine monoclonal antibody against the CaSR (Affinity BioReagents). The images were captured using a luminescent image analyzer LAS-4000 mini (Fujifilm Life Sciences). For indirect immunofluorescence, the transfected cells were fixed 16 h post-transfection in 4% paraformaldehyde in phosphate-buffered saline (PBS) for 7 min at 24°C to only reveal surface expression of the receptor. The fixed cells were first incubated during 2 h at 24°C in blocking buffer (PBS-1% gelatin) and then 4 h with an anti-CaSR murine monoclonal antibody (Affinity BioReagents) diluted in blocking buffer. The cells were then washed with PBS at 25°C and incubated during 2 h at 24°C with Alexa 488-conjugated rabbit-anti mouse (Invitrogen) diluted in blocking buffer. After extensive washes at 24°C with PBS, the samples mounted with a gelvatol-glycerol solution containing 2.5% 1,4-diazobicyclo-[2.2.2]octane. The samples were examined with a epifluorescence microscope (Zeiss Axioskop) as previously described [33]. The selected cells displayed in the appropriate figures were representative of 80% of the population of positive cells.

4. Results

In order to examine the impact of mutations in the 1st Ca²⁺ binding site of the VF_{TD}, we analyzed the pattern of $[Ca^{2+}]_i$ response in individual HEK-293 cells transiently co-transfected with a plasmid encoding the human wild type CaSR or with a CaSR containing a highly conservative substitution in Ser-170 to Thr (CaSR_{S170T}) and a plasmid encoding a red fluorescent protein (pDsRed-Express) to facilitate the identification of the transfected cells. After 16 h, the cells were loaded with the fluorescent Ca²⁺ indicator fura 2-AM and incubated in the presence of medium containing 1.5 mM $[Ca^{2+}]_e$. Single cell imaging of fura-2 loaded HEK-293 cells expressing the CaSR or CaSR_{S170T} revealed that a rise in the $[Ca^{2+}]_e$ from 1.5 mM to 5 mM produced a rapid elevation in $[Ca^{2+}]_i$ followed by a sustained plateau in cells transfected with either the wild type CaSR (Fig. 1 A) or mutant CaSR_{S170T} (Fig. 1 B, C). The maximal response of CaSR_{S170T} to the increase in $[Ca^{2+}]_e$ was similar to that of the wild type receptor (Fig. 1 A, B). Most transfected cells with CaSR or CaSR_{S170T} responded to the increase in $[Ca^{2+}]_e$ (Fig. 1 D). Accordingly, equivalent levels of CaSR and CaSR_{S170T} were expressed on the cell membrane of HEK-293 cells, as judged by indirect immunofluorescence (Fig. 1 F, G) and total cellular expression of CaSR and CaSR_{S170T} was also equal, as shown by immunoblotting (Fig. 1 H).

Although the results in Fig. 1 indicated that the responses of wild type and mutant CaSR to an increase in the $[Ca^{2+}]_e$ were virtually identical, analysis of $[Ca^{2+}]_i$ prior to stimulation revealed an important difference between CaSR and CaSR_{S170T} in the dynamics of basal $[Ca^{2+}]_i$. In agreement with our previous results, most cells transfected with wild type CaSR and incubated in the presence of medium containing 1.5 mM $[Ca^{2+}]_e$ exhibited a stable

$[Ca^{2+}]_i$ with only a small proportion of cells (~10%) exhibiting spontaneous oscillatory activity.

In striking contrast, a substantial proportion of cells transfected with CaSR_{S170T} display spontaneous oscillatory fluctuations in $[Ca^{2+}]_i$. The difference in spontaneous $[Ca^{2+}]_i$ oscillations is illustrated in Fig. 2 A, showing a pseudo-colored images of clusters of cells transfected with wild type CaSR or CaSR_{S170T} incubated for various times in medium containing 1.5 mM $[Ca^{2+}]_e$. Cells labeled O show a spontaneous increase in $[Ca^{2+}]_i$ which declines, and then increases again (oscillatory response) while other cells exhibited a stable $[Ca^{2+}]_i$. Examples of tracings corresponding to HEK-293 cells transfected with CaSR_{S170T} displaying spontaneous $[Ca^{2+}]_i$ oscillations are illustrated in Fig. 2 B. The examples also illustrate that the oscillatory frequency is markedly higher in cells expressing CaSR_{S170T}. The difference in basal $[Ca^{2+}]_i$ oscillations between CaSR and CaSR_{S170T} was substantiated by analyzing 714 individual cells. We found a marked increase in the proportion of cells that display spontaneous $[Ca^{2+}]_i$ oscillations when transfected with CaSR_{S170T} (Fig. 2 C, bars). These results demonstrate, for the first time, that a sub-population of HEK-293 cells expressing CaSR_{S170T} (~27%) displays spontaneous $[Ca^{2+}]_i$ oscillations while the majority of cells transfected with wild type CaSR exhibited a stable $[Ca^{2+}]_i$. In agreement with our previous results [13], we did not detect any $[Ca^{2+}]_i$ oscillations in non-transfected HEK-293 cells or in cells transfected with vector.

The results shown in Fig. 2 prompted us to hypothesize that HEK-293 cells transfected with CaSR_{S170T} have increased sensitivity to extracellular Ca^{2+} . To examine this possibility, we monitored basal $[Ca^{2+}]_i$ oscillations in HEK-293 cells transfected with wild type CaSR or CaSR_{S170T} incubated in medium containing different $[Ca^{2+}]_e$. Cells incubated in medium containing 0.8 mM Ca^{2+} show striking differences, as shown in Fig. 3. We found that ~20% of HEK-293 cells expressing CaSR_{S170T} exhibit spontaneous $[Ca^{2+}]_i$ oscillations with a frequency of 3 spikes/min. In sharp contrast, only ~3% of cells transfected with wild type CaSR exhibited oscillatory activity of lower frequency (2 spikes/min) whereas 97% of transfected cells showed stable $[Ca^{2+}]_i$. A further decrease in $[Ca^{2+}]_e$ to 0.4 mM completely eliminated any spontaneous $[Ca^{2+}]_i$ oscillations in HEK 293 cells transfected with wild type CaSR while 10% of HEK-293 cells expressing CaSR_{S170T} exhibited spontaneous $[Ca^{2+}]_i$ oscillations with a frequency of 2.7 spikes/min (Fig. 3). A small subpopulation of HEK 293 cells expressing CaSR_{S170T} show spontaneous $[Ca^{2+}]_i$ oscillations at a $[Ca^{2+}]_e$ as low as 0.2 mM. The threshold $[Ca^{2+}]_e$ required for spontaneous $[Ca^{2+}]_i$ oscillations in individual HEK-293 cells transfected with either CaSR or CaSR_{S170T} was 0.6mM and 0.2 mM, respectively, as extrapolated from the data in Fig. 3. These results support the hypothesis that a highly conserved mutation of the CaSR at Ser170 increases the sensitivity of the CaSR for extracellular Ca^{2+} .

In addition to its role as sensor of $[Ca^{2+}]_e$, the CaSR is also stimulated by aromatic amino acids [26] that induce striking and lasting transient $[Ca^{2+}]_i$ oscillations characterized by low frequency $[Ca^{2+}]_i$ spikes that return to the base-line level [13]. These oscillations are mediated by a phospholipase C-independent pathway that involves Rho, filamin and TRPC1 [13,15,16,18]. In contrast, CaSR activation in response to modest increase in $[Ca^{2+}]_e$ (e.g. from 1.5 to 3 mM) produces high frequency sinusoidal oscillations upon a raised plateau

level of $[Ca^{2+}]_i$ [13,15,16]. Consequently, we next determined whether CaSR_{S170T} mediates different oscillatory patterns, i.e. sinusoidal and transient oscillations in response to different agonists in cells initially incubated in low $[Ca^{2+}]_e$. To examine this possibility, HEK-293 cells transfected with CaSR_{S170T} were incubated in medium containing 0.8 mM $[Ca^{2+}]_e$ to diminish spontaneous $[Ca^{2+}]_i$ oscillations and then stimulated by an increase of $[Ca^{2+}]_e$ to 3 mM or by addition of 5 mM L-Phe. As shown in Fig. 4, HEK-293 cells expressing CaSR_{S170T} displayed sinusoidal oscillations in response to an increase in $[Ca^{2+}]_e$ to 3 mM and striking transient oscillations in response to 5 mM L-Phe in the presence of 0.8 mM $[Ca^{2+}]_e$ (Fig. 4). The responses were comparable to those showed by HEK-293 cells expressing wild type CaSR but incubated initially in medium containing 1.5 mM $[Ca^{2+}]_e$ and thus substantiating that the Ser170 to Thr mutation decreases the $[Ca^{2+}]_e$ required for detecting transient oscillations in response to aromatic amino acids.

5. Discussion

Oscillatory changes in $[Ca^{2+}]_i$ in response to receptor stimulation is a fundamental mechanism of cell signaling implicated in the regulation of signal transduction [15,19,20], metabolism [21,22], and differential gene expression [23,24]. Previous studies identified Ser-170 in the N-terminal region of the VFTD as a critical residue in Ca^{2+} binding and signal generation since mutation of Ser-170 to Ala completely abolished the response to even very high concentrations of extracellular Ca^{2+} [31] without affecting its membrane expression [30,31]. In the current study we analyzed $[Ca^{2+}]_i$ in single HEK-293 cells expressing wild type CaSR or a receptor with a highly conservative mutation of Ser-170 to Thr. Our results demonstrated that cells expressing CaSR_{S170T} exhibit spontaneous oscillatory activity when incubated in medium containing 1.5 mM $[Ca^{2+}]_e$. Analysis of spontaneous $[Ca^{2+}]_i$ oscillations as a function of $[Ca^{2+}]_e$ revealed that HEK-293 cells expressing CaSR_{S170T} display $[Ca^{2+}]_i$ oscillations of higher frequency at markedly lower $[Ca^{2+}]_e$ as compared with HEK-293 cells expressing wild type CaSR. Furthermore, HEK-293 cells expressing CaSR_{S170T} display sinusoidal $[Ca^{2+}]_i$ oscillations in response to an increase in $[Ca^{2+}]_e$ or transient oscillations when challenged by L-Phe, provided that cells are initially incubated in medium containing a low $[Ca^{2+}]_e$. We conclude that a small increase in the size of the side chain of Ser170 (Ser-to-Thr substitution in the VTFD) is sufficient to induce a marked increase in sensitivity to $[Ca^{2+}]_e$ of the CaSR.

Previous studies concluded that mutation of Ser-170 to Thr impairs amino acid sensing [30,31], as judged by experiments in which the EC_{50} value for $[Ca^{2+}]_e$ remains unchanged in the presence of L-Phe. In contrast, we demonstrated that L-Phe induced transient oscillations in cells expressing CaSR_{S170T} incubated in medium containing a low $[Ca^{2+}]_e$. We propose that the S170T mutation separates two different allosteric effects of aromatic amino acids on the CaSR. Specifically, (1) the stabilization of a conformation that promotes transient oscillations in the presence of a threshold level of $[Ca^{2+}]_e$ via a multi-protein complex that includes Rho, filamin-A, and TRPC1 from (2) the positive heterotropic effect that causes an increase in the affinity of the CaSR for extracellular Ca^{2+} . We obtained similar results using a different CaSR mutant in which Glu127 was replaced by Ala, an activating mutation. As with CaSR_{S170T}, cells expressing the CaSR_{E127A} mutation displayed $[Ca^{2+}]_i$ transient oscillations in response to L-Phe. We conclude that the allosteric regulation

of the CaSR by aromatic amino acids is not only mediated by an heterotropic positive effect on Ca²⁺ binding cooperativity but, as a biased agonist, aromatic amino acids stabilize a CaSR conformation that couples to a different signaling pathway leading to transient [Ca²⁺]_i oscillations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The work was supported by the Veterans Affairs Merit grant (I01BX001473) and by National Institutes of Health Grants R01DK100405, P30-DK41301 and P01CA163200 (all to ER). Cell Imaging Services were provided by the Imaging and Stem Cell Biology Core, CURE: Digestive Diseases Research Center supported by P30DK41301. E.R. holds the Ronald S. Hirschberg Chair of Pancreatic Cancer Research. O.R. is the recipient of PICT 2012-0875 and PICT 2013-0891, FONCyT-MINCYT, Argentina.

References

1. Brown EM, MacLeod RJ. Extracellular calcium sensing and extracellular calcium signaling. *Physiol Rev.* 2001; 81:239–297. [PubMed: 11152759]
2. Alfadda TI, Saleh AMA, Houillier P, Geibel JP. Calcium-sensing receptor 20 years later. *Am J Physiol Cell Physiol.* 2014; 307:C221–C231. [PubMed: 24871857]
3. Hofer AM, Brown EM. Extracellular calcium sensing and signalling. *Nat Rev Mol Cell Biol.* 2003; 4:530–538. [PubMed: 12838336]
4. Geibel JP, Hebert SC. The functions and roles of the extracellular Ca²⁺-sensing receptor along the gastrointestinal tract. *Annu Rev Physiol.* 2009; 71:205–217. [PubMed: 19575679]
5. Saidak Z, Mentaverri R, Brown EM. The Role of the Calcium-Sensing Receptor in the Development and Progression of Cancer. *Endocr Rev.* 2009; 30:178–195. [PubMed: 19237714]
6. Riccardi D, Kemp PJ. The calcium-sensing receptor beyond extracellular calcium homeostasis: conception, development, adult physiology, and disease. *Annu Rev Physiol.* 2012; 74:271–297. [PubMed: 22017175]
7. Rey O, Chang W, Bikle D, Rozengurt N, Young SH, Rozengurt E. Negative Cross-talk between Calcium-sensing Receptor and beta-Catenin Signaling Systems in Colonic Epithelium. *J Biol Chem.* 2012; 287:1158–1167. [PubMed: 22094462]
8. Aggarwal A, Prinz-Wohlgenannt M, Tennakoon S, Höbaus J, Boudot C, Mentaverri R, Brown EM, Baumgartner-Parzer S, Kállay E. The calcium-sensing receptor: A promising target for prevention of colorectal cancer. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research.* 2015; 1853:2158–2167. [PubMed: 25701758]
9. Tang L, Peng M, Liu L, Chang W, Binder HJ, Cheng SX. Calcium-sensing receptor stimulates Cl[−] and SCFA-dependent but inhibits cAMP-dependent HCO₃[−] secretion in colon. *Am J Physiol Gastrointest Liver Physiol.* 2015; 308:G874–G883. [PubMed: 25792563]
10. Lee GS, Subramanian N, Kim AI, Aksentijevich I, Goldbach-Mansky R, Sacks DB, Germain RN, Kastner DL, Chae JJ. The calcium-sensing receptor regulates the NLRP3 inflammasome through Ca²⁺ and cAMP. *Nature.* 2012; 492:123–127. [PubMed: 23143333]
11. Goltzman D, Hendy GN. The calcium-sensing receptor in bone—mechanistic and therapeutic insights. *Nat Rev Endocrinol.* 2015; 11:298–307. [PubMed: 25752283]
12. Breitwieser GE, Gama L. Calcium-sensing receptor activation induces intracellular calcium oscillations. *Am J Physiol Cell Physiol.* 2001; 280:C1412–C1421. [PubMed: 11350736]
13. Young SH, Rozengurt E. Amino acids and Ca²⁺ stimulate different patterns of Ca²⁺ oscillations through the Ca²⁺-sensing receptor. *Amer J Physiol-Cell Physiol.* 2002; 282:C1414–C1422. [PubMed: 11997256]

14. Young SH, Wu SV, Rozengurt E. Ca²⁺-stimulated Ca²⁺ Oscillations Produced by the Ca²⁺-sensing Receptor Require Negative Feedback by Protein Kinase C. *J. Biol. Chem.* 2002; 277:46871–46876. [PubMed: 12356761]
15. Rey O, Young SH, Yuan J, Slice L, Rozengurt E. Amino acid-stimulated Ca²⁺ oscillations produced by the Ca²⁺-sensing receptor are mediated by a phospholipase C/inositol 1,4,5-trisphosphate-independent pathway that requires G12, Rho, filamin-A, and the actin cytoskeleton. *J Biol Chem.* 2005; 280:22875–22882. [PubMed: 15837785]
16. Rey O, Young SH, Papazyan R, Shapiro MS, Rozengurt E. Requirement of the TRPC1 Cation Channel in the Generation of Transient Ca²⁺ Oscillations by the Calcium-sensing Receptor. *J. Biol. Chem.* 2006; 281:38730–38737. [PubMed: 17046820]
17. Davies SL, Ozawa A, McCormick WD, Dvorak MM, Ward DT. Protein Kinase C-mediated Phosphorylation of the Calcium-sensing Receptor Is Stimulated by Receptor Activation and Attenuated by Calyculin-sensitive Phosphatase Activity. *J Biol Chem.* 2007; 282:15048–15056. [PubMed: 17376781]
18. Rey O, Young SH, Jacamo R, Moyer MP, Rozengurt E. Extracellular calcium sensing receptor stimulation in human colonic epithelial cells induces intracellular calcium oscillations and proliferation inhibition. *J Cell Physiol.* 2010; 225:73–83. [PubMed: 20648625]
19. Soderling TR, Chang B, Brickey D. Cellular Signaling through Multifunctional Ca²⁺/Calmodulin-dependent Protein Kinase II. *J. Biol. Chem.* 2001; 276:3719–3722. [PubMed: 11096120]
20. Oancea E, Meyer T. Protein kinase C as a molecular machine for decoding calcium and diacylglycerol signals. *Cell.* 1998; 95:307–318. [PubMed: 9814702]
21. Hajnoczky G, Robb-Gaspers LD, Seitz MB, Thomas AP. Decoding of Cytosolic Calcium Oscillations in the Mitochondria. *Cell.* 1995; 82:415–424. [PubMed: 7634331]
22. Drago I, Pizzo P, Pozzan T. After half a century mitochondrial calcium in- and efflux machineries reveal themselves. *EMBO J.* 2011; 30:4119–4125. [PubMed: 21934651]
23. Li W, Llopis J, Whitney M, Zlokarnik G, Tsien RY. Cell-permeant caged InsP₃ ester shows that Ca²⁺ spike frequency can optimize gene expression. *Nature.* 1998; 392:936–941. [PubMed: 9582076]
24. Hu Q, Deshpande S, Irani K, Ziegelstein RC. (Ca²⁺)_i oscillation frequency regulates agonist-stimulated NF-κB transcriptional activity. *J Biol Chem.* 1999; 274:33995–33998. [PubMed: 10567364]
25. Young SH, Rey O, Sinnott-Smith J, Rozengurt E. Intracellular Ca²⁺ oscillations generated via the Ca²⁺-sensing receptor are mediated by negative feedback by PKCα at Thr888. *Am J Physiol Cell Physiol.* 2014; 306:C298–C306. [PubMed: 24336654]
26. Conigrave AD, Quinn SJ, Brown EM. L-amino acid sensing by the extracellular Ca²⁺-sensing receptor. *Proc Natl Acad Sci USA.* 2000; 97:4814–4819. [PubMed: 10781086]
27. Hannan FM, Nesbit MA, Zhang C, Cranston T, Curley AJ, Harding B, Fratter C, Rust N, Christie PT, Turner JJO, Lemos MC, Bowl MR, Bouillon R, Brain C, Bridges N, Burren C, Connell JM, Jung H, Marks E, McCredie D, Mughal Z, Rodda C, Tollefsen S, Brown EM, Yang JJ, Thakker RV. Identification of 70 calcium-sensing receptor mutations in hyper- and hypo-calcaemic patients: evidence for clustering of extracellular domain mutations at calcium-binding sites. *Hum Mol Genet.* 2012; 21:2768–2778. [PubMed: 22422767]
28. Hu J, Spiegel AM. Naturally occurring mutations of the extracellular Ca²⁺-sensing receptor: implications for its structure and function. *Trends Endocrinol Metabol.* 2003; 14:282–288.
29. Huang Y, Zhou Y, Castiblanco A, Yang W, Brown EM, Yang JJ. Multiple Ca²⁺-Binding Sites in the Extracellular Domain of the Ca²⁺-Sensing Receptor Corresponding to Cooperative Ca²⁺ Response†. *Biochemistry.* 2009; 48:388–398. [PubMed: 19102677]
30. Zhang C, Huang Y, Jiang Y, Mulpuri N, Wei L, Hamelberg D, Brown EM, Yang JJ. Identification of an l-Phenylalanine Binding Site Enhancing the Cooperative Responses of the Calcium-sensing Receptor to Calcium. *J Biol Chem.* 2014; 289:5296–5309. [PubMed: 24394414]
31. Mun H-C, Culverston EL, Franks AH, Collyer CA, Clifton-Bligh RJ, Conigrave AD. A Double Mutation in the Extracellular Ca²⁺-sensing Receptor's Venus Flytrap Domain That Selectively Disables l-Amino Acid Sensing. *J Biol Chem.* 2005; 280:29067–29072. [PubMed: 15888439]

32. Needham LK, Rozengurt E. G α 12 and G α 13 stimulate Rho-dependent tyrosine phosphorylation of focal adhesion kinase, paxillin and p130 Crk-associated substrate. *J Biol Chem.* 1998; 273:14626–14632. [PubMed: 9603980]
33. Rey O, Young SH, Cantrell D, Rozengurt E. Rapid Protein Kinase D Translocation in Response to G Protein-coupled Receptor Activation. Dependence on Protein Kinase C. *J. Biol. Chem.* 2001; 276:32616–32626. [PubMed: 11410587]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

HIGHLIGHTS

- $[Ca^{2+}]_i$ oscillations were analyzed in HEK-293 cells expressing CaSR and CaSR_{S170T}
- CaSR_{S170T} mediated basal $[Ca^{2+}]_i$ oscillations even in medium with low $[Ca^{2+}]_e$
- CaSR_{S170T} produced transient $[Ca^{2+}]_i$ oscillations in response to L-phenylalanine.
- CaSR_{S170T} separates enhanced sensitivity to $[Ca^{2+}]_e$ from transient oscillations





Figure 1. Changes in $[Ca^{2+}]_i$ in HEK-293 cells transfected with CaSR (WT) **A**, or CaSR_{S170T} (S170T) **B,C**, in response to an increase in $[Ca^{2+}]_e$ from 1.5 mM to 5 mM. The addition of Ca^{2+} is indicated by upward arrows whereas the return of $[Ca^{2+}]_e$ to 1.5 mM is indicated by the downward arrows. **D**, Bar chart: Percentage of HEK-293 cells transfected with CaSR (WT) or CaSR_{S170T} which responded to an increase in $[Ca^{2+}]_e$ from 1.5 mM to 5 mM. **E, F, G**. Fixed cells were labeled with an anti-CaSR murine monoclonal antibody to detect surface expressed receptor. **E**. Control, no transfection. **F**. Transfected with CaSR(WT). **G**. Transfected with CaR_{S170T}. Scale Bar 30 μ M. **H**. Western Blot analysis of control, no transfection (-), CaSR (WT), and CaSR_{S170T} (S170T) cell lysates.

**Figure 2.**

Spontaneous $[Ca^{2+}]_i$ oscillations of HEK-293 cells transfected with wild type CaSR (WT) or CaSR_{S170T} (S170T) and incubated in medium containing 1.5 mM $[Ca^{2+}]_e$. **A.** Pseudo-colored images of clusters of cells showing changes in $[Ca^{2+}]_i$ with time. Shown is one cycle of rest to peak response. Spontaneously oscillating cells (O) are marked with an arrow. **B.** Typical traces of $[Ca^{2+}]_i$ as a function of time (s) for a single HEK-293 cell transfected with the wild type CaSR or a single HEK-293 cells transfected with CaSR_{S170T}. **C.** Percentage of spontaneously oscillating HEK-293 cells from a total of 544 transfected with the wild type CaSR and from 170 cells transfected with CaSR_{S170T}.




Figure 3.

Percentage of spontaneously oscillating HEK-293 cells transfected with wild type CaSR (WT) or CaSR_{S170T} (S170T), (left panels) and their oscillating frequencies (right panels) as a function of varying $[Ca^{2+}]_e$. The data is a summary derived from a total of 1,289 HEK-293 cells transfected with wild type CaSR and 1,386 HEK-293 cells transfected with CaSR_{S170T}. Numbers show number of cells measured at each $[Ca^{2+}]_e$.




Figure 4.

Oscillatory responses in $[Ca^{2+}]_i$ in HEK-293 expressing $CaSR_{S170T}$ (S170T) cells when challenged with increases in $[Ca^{2+}]_e$ or L-phenylalanine (L-Phe). The cells were initially incubated in medium containing $[Ca^{2+}]_e$ at 0.8 mM. After recording basal $[Ca^{2+}]_i$ the cells were challenged by either increasing the $[Ca^{2+}]_e$ to 3mM or 5 mM L-Phe (upward arrows). Downward arrow marks return to medium containing $[Ca^{2+}]_e$ at 1.5 mM.