

Original Article

Activation of human smooth muscle BK channels by hydrochlorothiazide requires cell integrity and the presence of BK β_1 subunit

Pedro MARTÍN^{1,*}, Melisa MONCADA¹, Guruprasad KUNTAMALLAPPANAVAR², Alex M DOPICO², Verónica MILESI¹

¹IIFP, UNLP, CONICET – Instituto de estudios Inmunológicos y Fisiopatológicos, Universidad Nacional de La Plata, Consejo Nacional de Investigaciones Científicas y Tecnológicas, La Plata, Argentina. 47 y 115 S/N, La Plata (1900), Buenos Aires, Argentina; ²Department of Pharmacology, College of Medicine, The University of Tennessee Health Science Center, 71 South Manassas St, Memphis, TN, 38103, USA

Abstract

Thiazide-like diuretics are the most commonly used drugs to treat arterial hypertension, with their efficacy being linked to their chronic vasodilatory effect. Previous studies suggest that activation of the large conductance voltage- and Ca^{2+} -dependent K^+ (BK) channel (Slo 1, MaxiK channel) is responsible for the thiazide-induced vasodilatory effect. But the direct electrophysiological evidence supporting this claim is lacking. BK channels can be associated with one small accessory β -subunit (β_1 – β_4) that confers specific biophysical and pharmacological characteristics to the current phenotype. The β_1 -subunit is primarily expressed in smooth muscle cells (SMCs). In this study we investigated the effect of hydrochlorothiazide (HCTZ) on BK channel activity in native SMCs from human umbilical artery (HUASMCs) and HEK293T cells expressing the BK channel (with and without the β_1 -subunit). Bath application of HCTZ (10 $\mu\text{mol/L}$) significantly augmented the BK current in HUASMCs when recorded using the whole-cell configurations, but it did not affect the unitary conductance and open probability of the BK channel in HUASMCs evaluated in the inside-out configuration, suggesting an indirect mechanism requiring cell integrity. In HEK293T cells expressing BK channels, HCTZ-augmented BK channel activity was only observed when the β_1 -subunit was co-expressed, being concentration-dependent with an EC_{50} of 28.4 $\mu\text{mol/L}$, whereas membrane potential did not influence the concentration relationship. Moreover, HCTZ did not affect the BK channel current in HEK293T cells evaluated in the inside-out configuration, but significantly increases the open probability in the cell-attached configuration. Our data demonstrate that a β_1 -subunit-dependent mechanism that requires SMC integrity leads to HCTZ-induced BK channel activation.

Keywords: thiazide; hydrochlorothiazide; BK channel; Slo1; β_1 -subunit; vascular smooth muscle cells; human umbilical artery; patch-clamp electrophysiology

Acta Pharmacologica Sinica advance online publication, 30 Nov 2017; doi: 10.1038/aps.2017.133

Introduction

Thiazide diuretics remain as one of the drugs most commonly used to treat systemic arterial hypertension. In particular, hydrochlorothiazide (HCTZ) is one of the first-line antihypertensive therapy drugs^[1]. HCTZ and related agents exert antihypertensive effects by primarily acting on the kidney, where these drugs block the Na^+/Cl^- cotransporter (NCC) and thus produce acute diuresis^[2]. However, long-term antihypertensive therapy with thiazides has been linked to the agents' ability to lower vascular resistance rather than to their diuretic

action^[2,3]. Remarkably, despite 58 years of demonstrated anti-hypertensive effectiveness, the precise mechanism(s) by which HCTZ and related thiazides evoke vasodilation and therefore reduce vascular resistance are not fully characterized.

Thiazide-induced vasorelaxation has been well documented through *in vivo* and *ex vivo* experiments in humans^[4-6] and other species^[5,7-9]. Although studies have established that thiazide-induced vasodilation contributes to the clinical benefit of these agents in chronically treated hypertensive patients, several studies have shown that such thiazide action is independent of NCC blockade^[3,10,11]. Different mechanisms have been proposed to mediate thiazide-induced artery dilation by targeting smooth muscle cells, including activation of large conductance, voltage- and $[\text{Ca}^{2+}]_i$ -gated potassium (BK) chan-

*To whom correspondence should be addressed.

E-mail pedromartin@biol.unlp.edu.ar

Received 2017-04-24 Accepted 2017-08-28

nels^[4, 5, 7, 8, 12], inhibition of voltage-operated calcium channels (VOCCs)^[5, 7, 8, 13] and/or inhibition of the RhoA/Rho kinase pathway^[9]. The latter would induce a Ca²⁺ desensitization of the smooth muscle contractile machinery. Making the overall scenario more complex, it has been argued that the relative contribution of each of the proposed mechanisms to thiazide-induced vasodilation depends on the species from which vascular tissue was obtained for *ex vivo* assays^[5].

The importance of BK channels in the regulation of vascular smooth muscle cell (VSMC) contractility, peripheral resistance and blood pressure has been extensively established^[14, 15]. These channels are activated by membrane depolarization and/or an increase in intracellular Ca²⁺ concentration. Since both events are associated with VSMC contraction and BK channel activation, the evoked outward K⁺ current acts as a negative feedback mechanism on VSMC contraction, thereby favoring VSMC relaxation^[15, 16]. Moreover, several studies have suggested that BK channel activity is altered in hypertension (see ^[17] and ^[18] reviews for more detail). Thus, the activation of this ion channel has emerged as a novel molecular target for treating diseases where increased tone and/or contractility of smooth muscle play a relevant pathophysiological role, such as hypertension^[19].

In most mammalian tissues, native BK channels are homotetramers of pore-forming α -subunits (encoded by the *KCNMA1* gene, also named *Slo1*) that can be associated with one of four auxiliary β -subunits (encoded by *KCNMB1-4* genes)^[20-25]. Unlike α -subunits, β -subunits do not form functional channels but modify several gating processes^[26-28]. The differential expression of auxiliary subunits in different cell types explains the multiplicity of functions and regulatory mechanisms of BK channels. In VSMCs, the β_1 -subunit is a main partner of the BK channel^[25, 29]. Several studies have shown that endogenous and exogenous compounds can modulate BK channels through β -subunits^[30-33]. Additionally, mutations in the β_1 -subunit that confer a gain in BK activity are associated with the reduced prevalence of hypertension in humans^[34], and the opposite effect occurs in the case of a loss of function mutation^[35].

HCTZ did not activate BK channels in skeletal muscle and BK channel α -subunits expressed in HEK cells, two preparations in which the functional expression of BK β_1 is negligible^[36, 37]. Moreover, the hypothesis that the activation of vascular smooth muscle BK channels, which contain β_1 -subunits, is involved in HCTZ-induced vasodilation is supported by *ex vivo* experiments in which this effect is abolished in the presence of BK channel inhibitors^[5, 8, 12]. The lack of electrophysiological studies on VSMCs makes it difficult to establish whether HCTZ-induced BK activation reflects a direct drug interaction with channel subunits or indirect drug interactions, i.e., requiring additional cell signals. This question requires electrophysiological studies on VSMCs using different patch-clamp configurations under controlled conditions of voltage and [Ca²⁺]_i.

In the present study, using patch-clamp electrophysiology, we determined the effects of HCTZ on (i) BK channels

from native human umbilical artery smooth muscle cells (HUASMCs) in "whole cell" (WCR) and cell-free, "inside-out" (IO) configurations and (ii) recombinant BK (slo1) channels with or without β_1 -subunits co-expressed in HEK293T cells. These results demonstrated that HCTZ effectively induces BK channel activation, and this effect requires both cell integrity and the presence of β_1 -subunits.

Materials and methods

Smooth muscle cell isolation for patch-clamp experiments

Umbilical cords were obtained from normal term pregnancies after vaginal and cesarean deliveries. The umbilical cords were placed in a transport solution with the following composition (in mmol/L): 130 NaCl, 4.7 KCl, 1.17 KH₂PO₄, 1.16 MgSO₄, 24 NaCO₃H, and 2.5 CaCl₂, pH 7.4 at 4 °C. The umbilical cords were immediately transferred to the laboratory, stored at 4 °C and used within the next 24 h. All procedures were performed in accordance with the Declaration of Helsinki (1975).

Human umbilical arteries (HUAs) were dissected from Wharton's jelly just prior to cell isolation. HUA smooth muscle cells (HUASMCs) were isolated as previously described^[38], with further modifications to diminish the enzyme content in the dissociation medium (DM)^[32]. Briefly, a segment of HUA was cleaned of any residual connective tissue (Wharton's jelly), cut into small strips and placed for 15 min in DM containing the following (in mmol/L): 140 NaCl, 5 KH₂PO₄, 5 MgCl₂, 6 glucose, and 5 HEPES; the pH was adjusted to 7.4 with NaOH. Vessel strips were subsequently placed in DM containing 3 mg/mL collagenase type I for 15–30 min, with gentle agitation at 35 °C. After the incubation period, the strips were washed with DM, and single HUASMCs were obtained by gentle aspiration of the tissue through a Pasteur pipette to render cell dispersion. The supernatant containing the isolated cells was stored at room temperature (~22 °C) until further use. Immediately prior to electrophysiological recordings, HUASMCs were allowed to settle onto the coverglass bottom of a 3-mL experimental chamber. Only well-relaxed, spindle-shaped smooth muscle cells were used for patch-clamp electrophysiology. Data were collected within 4–6 h after cell isolation.

Clones and transient transfection

HEK293T cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and subsequently split when reaching 70%–80% confluence. The pcDNA3 plasmids harboring the human Slo1 α -subunit (A#U11058), the auxiliary β_1 -subunit (A#AF035046) and enhanced green fluorescent protein (eGFP) cDNAs were transfected into HEK cells using FuGene 6 (Promega, Madison, WI, USA) and transiently expressed. Transfections were performed in HEK cells grown in 30-mm Petri dishes using cDNAs encoding the α -subunits alone or both α - and β_1 -subunits. The α - and β_1 -subunits were mixed at 1:3 molar ratios to ensure that all α -subunits were saturated with β_1 -subunits (total amount of DNA: 2.5 μ g distributed in 0.5, 0.7 and 1.3 μ g of plasmids containing eGFP, α -subunits and

β_1 -subunits, respectively). When α -subunits were transfected alone, β_1 cDNA was replaced with empty plasmid. After transfection, the cells were cultured for 18 h. The electrophysiological measurements were performed 18–24 h after transfection. BK α and β_1 clones were kind gifts of Dr Carlos GONZÁLEZ (CINV, Chile). The eGFP clone and vector were kindly provided by Dr J RAINGO (IMBICE, Argentina).

Electrophysiology

Isolated cells were observed using a mechanically stabilized inverted epifluorescence microscope (Arcano XYL 403 YAT PLAN, China) equipped with a 40 \times objective lens. Test solutions were applied through a multibarreled pipette positioned close to the target cell (approximated 1 mm), which enabled bathing the cell or the pipette tip (in the inside-out configuration) with the test solution for only 5 s. The perfusion rate was approximately 1 mL/min. After each experiment on a single cell, the experimental chamber was replaced with another containing a new sample of cells. All experiments were performed at room temperature ($\sim 22^\circ\text{C}$). For all experimental data included in the present study, we prolonged the exposure time to HCTZ to at least 15 min to ensure drug permeation, reflecting its poor lipophilicity. The concentration-response curve to HCTZ (10^{-5} – 3×10^{-4} mol/L) in HEK293T cells was constructed by applying a single HCTZ concentration to each cell since it took at least 15 min for HCTZ to increase channel activity.

The standard tight-seal IO and WCR configurations of the patch-clamp technique^[39] were used to record single-channel and macroscopic currents, respectively. Glass pipettes were pulled from WPI PG52165-4 glass on a two-stage vertical micropipette puller (PP-83, Narishige Scientific Instrument Laboratories, Tokyo, Japan). Pipette resistance ranged from 1.7 to 3.5 M Ω when filled with electrode solution. All currents were filtered with a 4-pole low-pass Bessel filter at 2 kHz (Axopatch 200A amplifier, Axon Instruments, Foster City, CA, USA) and digitized (Digidata 1440, Molecular Devices, Sunnyvale, CA, USA) at 20 kHz. Recordings were stored on a computer hard disk for subsequent analysis.

Single-channel recordings

Single-channel currents were recorded in the inside-out configuration (IO) in HUASMCs and the cell-attached configuration (CA) in HEK293T cells co-transfected with the α - and β_1 -subunits of the BK channel. Voltage-clamp recordings (up to 30–60 s) were obtained at different membrane potentials to measure the single channel current amplitude and steady-state activity, which results from the product of the single channel open probability (Po) and the number of functional channels in the membrane patch (N). Identification of BK channel presence in the patch was based on its unitary conductance value, voltage-dependent activity and activation by an increase in intracellular Ca^{2+} concentration, as previously described^[40]. The control pipette solution used for single-channel recordings contained the following components (in mmol/L): 140 KCl, 0.5 MgCl₂, 10 HEPES, 6 glucose, and 1 CaCl₂; the pH was adjusted

to 7.4 with KOH. The bath solution (BS) contained the following components (in mmol/L): 140 KCl, 0.5 MgCl₂, 0.4 CaCl₂, 10 HEPES, 6 glucose, and 1 EGTA; the pH was adjusted to 7.4 with KOH. The free Ca^{2+} concentration was calculated as 50 nmol/L using Maxchelator software from Stanford University (<http://maxchelator.stanford.edu>).

HCTZ's action on BK channels was examined by directly adding the adequate drug quantity to the BS to reach the desired final concentration. Single-channel currents were analyzed using Clampfit software (Molecular Devices, version 10.3). Steady-state activity (NPo) was derived from the equation below, where N is the number of single channels present in each patch.

$$NPo = \frac{\sum_{j=1}^n j t_j}{T}$$

where T is the recording duration and t_j is the time spent with $j=1,2,3,\dots,n$ channels open. Stationary conditions of single channel recordings were controlled by plotting the NPo values calculated for intervals of 30 s of recording as a function of time (i.e., a stability plot).

Whole-cell recordings in HUASMCs

Whole-cell current stability was monitored by applying successive 500 ms voltage steps, from a holding potential of -50 mV to a test potential of +40 mV. Thus, cells in which the current amplitude did not remain constant in time were discarded. After the current was stabilized, the same voltage-clamp step protocol was applied under control (solvent) conditions or in the presence of HCTZ. A voltage-clamp step protocol, including 500 ms voltage steps spaced 10 mV between -70 and +70 mV from a holding potential of -50 mV, was applied under control conditions and in the presence of HCTZ for further current amplitude-voltage relationship (I - V) analysis. The pipette solution contained the following components (in mmol/L): 130 KCl, 10 HEPES, 0.1 EGTA, 0.1 CaCl₂, 1 MgCl₂, and 5 ATP-Na₂; the pH was adjusted to 7.4 with KOH. The BS contained the following components (in mmol/L): 130 NaCl, 5.4 KCl, 1.2 MgCl₂, 2.5 CaCl₂, 5 HEPES, 6 glucose, and 5 4-aminopyridine (4-AP); the pH was adjusted to 7.4 with HCl. The compound 4-AP was added to block Kv channels, which have been detected in HUASMCs^[41]. Cell membrane capacitance was calculated from the capacity current obtained from the recording of a single 6-ms voltage step from a holding potential of -50 mV to -60 mV. The mean capacitance and series resistance of the recorded HUASMCs were 24.8 ± 2.0 pF and 3.1 ± 0.3 M Ω , respectively.

Inside-Out macroscopic current recordings in heterologously expressed BK channels

Macroscopic currents of recombinant BK channels expressed in HEK293T cells were recorded using the IO configuration. Current stability was monitored by applying successive 150 ms voltage steps (from a holding potential of 0 mV to a test potential of +90 mV). Cells in which the current amplitude did not remain constant in time were discarded. The same

voltage-clamp step protocol was applied under control conditions and in the presence of HCTZ. After current stabilization, a voltage-clamp step protocol, including a family of 150 ms voltage steps spaced 20 mV between -90 and 190 mV from a holding potential of 0 mV followed by a 50 ms voltage step to -150 mV, was applied under control conditions and in the presence of 10, 30 and 100 $\mu\text{mol/L}$ HCTZ for *I-V* relationship analysis. Since channel expression (estimated by current amplitude) was extremely variable and the current saturated the amplifier at the more positive potentials examined in most of the evaluated cells, the current values were normalized to the control current obtained at +130 mV. The pipette and bath solutions were identical to those used in the IO experiments in HUASMCs.

Whole-cell current recordings in heterologously expressed BK_{Ca} channels

The effects of HCTZ on whole-cell currents mediated by BK channel-forming slo1 proteins (α -subunit) expressed in HEK293T cells with and without the β_1 -subunit were examined. Current stability was monitored by applying successive 100 ms voltage steps (from a holding potential of -50 mV to a test potential of +60 mV), discarding cells in which the current amplitude did not remain constant in time. The same voltage-clamp step protocol was applied under control conditions or in the presence of HCTZ. After current stabilization, a voltage-clamp step protocol, including a family of 100 ms voltage steps spaced 10 mV between -60 and +90 mV from a holding potential of -50 mV, was applied under control conditions and in the presence of 10, 30, 100 and 300 $\mu\text{mol/L}$ HCTZ to conduct *I-V* plots. Since channel expression was extremely variable and the current saturated the amplifier at the more positive potentials examined in most of the evaluated cells, current values were normalized to the control values obtained at +90 mV. The pipette solution contained the following components (in mmol/L): 130 KCl, 10 HEPES, 1 EGTA, 0.4 CaCl₂, 1 MgCl₂, and 5 ATP-Na₂; the pH was adjusted to 7.3 with KOH. The BS contained the following components (in mmol/L): 130 NaCl, 5.4 KCl, 1 EGTA, 1.2 MgCl₂, 0.0094 CaCl₂, 5 HEPES, and 6 glucose; the pH was adjusted to 7.4 with NaOH. Both solutions contained a nominal free Ca²⁺ concentration of 50 nmol/L, calculated using Maxchelator. The mean series resistance of the recorded HEK293T cells was 2.7±0.2 M Ω .

Drugs and reagents used

HCTZ, 4-AP, EGTA, Na₂ATP and collagenase were purchased from Sigma Chemical Co (St Louis, USA). All other reagents, including DMEM medium and FBS, were of analytical grade and purchased from local suppliers. HCTZ was dissolved in dimethyl sulfoxide (DMSO). Fresh aliquots of stock solutions of HCTZ in DMSO were added to the bath solution on the day of the experiment. Appropriate amounts of DMSO were added to all control solutions without HCTZ.

Statistics

The results are expressed as the means±standard error of the

mean. Paired Student's *t* tests were used to compare two groups. ANOVA was used to compare more than two groups. In all cases, a *P* value less than 0.05 was considered as statistically significant. The concentration-response curve data for BK channel activation by HCTZ were fitted using GraphPad Prism (version 5.03) to a Hill relationship with a variable slope. The equation used for this fit is:

$$\text{Current increase} = \text{Bottom} + \frac{\text{Top} - \text{Bottom}}{1 + 10^{((\log \text{EC}_{50} - x) \times \text{Hill slope})}}$$

where EC₅₀ is the concentration of agonist that gives a response halfway between bottom and top, Hill slope describes the steepness of the curves, and top and bottom are plateaus of current increase. In the present study, we used the parameter $pD_{2,2}$, which is the $-\log \text{EC}_{50}$. The fitting parameters at different voltages were compared using the extra sum of squares principle (*F* method available in GraphPad Prism).

Results

Hydrochlorothiazide activates whole-cell BK current in human umbilical artery smooth muscle cells (HUASMCs)

Based on the *ex vivo* results from Calder *et al*^[8, 12] and Pickkers *et al*^[4, 7], BK channel activation was postulated as the mechanism underlying vasodilation mediated by thiazide-like diuretics. However, the effect of HCTZ on the BK currents from native vascular smooth muscle cells has not actually been evaluated. Furthermore, when this diuretic was tested on skeletal muscle cells and HEK cells expressing the α -subunit of this channel^[36, 37], it failed to activate BK channels in both native and heterologous systems in the absence of any β -subunit. Thus, we evaluated the effect of HCTZ in freshly dissociated HUASMCs as a model of human VSMCs in which the BK channel is associated with the β_1 -subunit^[32] and involved in the regulation of the basal tone^[42]. We obtained the isolated voltage-activated macroscopic BK currents from HUASMCs in the standard patch-clamp WCR configuration as previously described^[32, 42] and examined the effect of HCTZ on these cells (Figure 1). The data demonstrated that after 7 min of bath (extracellular) application of 10 $\mu\text{mol/L}$ HCTZ, the amplitude of the outward whole-cell current was significantly augmented at membrane potential values between 20 and 70 mV (Figure 1; *P*<0.05). The effect of HCTZ was fully reversed upon drug washout with solvent-containing bath solution, demonstrating that 10 $\mu\text{mol/L}$ HCTZ is able to activate BK channels in human VSMCs.

HCTZ failed to modulate BK channel activity in the absence of cell integrity

To determine whether the HCTZ-induced increased ionic BK current in HUASMCs was due to direct drug action on the BK channel proteins or required cell integrity and intact signaling, we next examined the effect of HCTZ on BK currents at a single channel resolution using the IO configuration of the patch-clamp technique. Thus, we recorded steady-state channel activity (NPo; see methods) at a constant voltage (+40 mV) in the presence or absence of 10 $\mu\text{mol/L}$ HCTZ. Interestingly, in




Figure 1. Hydrochlorothiazide (HCTZ) increases the large-conductance voltage- and Ca^{2+} -activated K^+ (BK) channel component of whole-cell (WC) K^+ currents in human umbilical artery smooth muscle cells (HUASMCs). (A) Superimposed representative recordings of WC currents of HUASMCs (recorded in 5 mmol/L 4-aminopyridine) obtained by applying 10 mV voltage steps from -70 mV to +70 mV from a holding potential of -50 mV in control conditions, after 7 min of 10 $\mu\text{mol/L}$ HCTZ perfusion and after washout. (B) Mean current density vs voltage ($I/\text{Cm-V}$) curves, corresponding to the control conditions and after 7 min of 10 $\mu\text{mol/L}$ HCTZ perfusion obtained in the same conditions as in a) ($n=6$ cells from 3 umbilical cord donors). The symbol * indicates statistically significant difference from control (paired t test, $P<0.05$).

this cell-free condition, HCTZ failed to modulate channel function: neither unitary current amplitude (Figure 2B, $P>0.05$) nor NPo was modified by HCTZ concentrations that increased the macroscopic BK current (Figure 2C, $P>0.05$). Collectively, the results shown in Figures 1 and 2 indicate that HCTZ requires cell integrity to increase BK currents.

HCTZ-mediated activation of recombinant BK channels in a mammalian heterologous system requires cell integrity and the presence of BK regulatory β_1 -subunits

The results shown above demonstrate that HCTZ is able to activate BK channels in HUASMCs. In contrast, this drug

failed to activate BK channels in skeletal muscle and HEK cells expressing only the α -subunit^[36, 37]. Thus, since endogenous β_1 -subunits are highly expressed in vascular smooth muscle cells but not in skeletal muscle cells or HEK cells, we hypothesized that the β_1 -subunit was necessary for HCTZ-induced BK channel activation in HUASMCs^[21, 25, 29]. To test this hypothesis, we expressed the channel α -subunit in HEK293T cells with or without the β_1 -subunit and examined the effect of HCTZ on these cells using the WCR configuration. Consistent with our hypothesis, HCTZ failed to modulate BK channel activity in the absence of auxiliary subunits (Figure 3, $P>0.05$). In contrast, HCTZ significantly increased the ionic current mediated




Figure 2. HCTZ failed to activate BK channel in inside-out (IO) configuration in HUASMCs. (A) Representative recording of BK channel current obtained at single channel resolution in an IO configuration at 40 mV in absence or presence of 10 $\mu\text{mol/L}$ HCTZ. The closed level is indicated as “c”. (B and C) Mean current amplitude (B) and steady state activity (NPo) values (C) of BK channels recorded as shown in A) in control conditions and with 10 $\mu\text{mol/L}$ HCTZ ($n=7$ patches from 3 umbilical cord donors). No statistically significant differences were observed (paired t test, $P>0.05$).




Figure 3. HCTZ failed to modulate BK channel activity in the absence of regulatory β -subunits in HEK293T cells. (A) Superimposed representative WCR currents of BK channels expressed in HEK293T cells without the β_1 -subunit, before (left) and after 15 min of 100 $\mu\text{mol/L}$ HCTZ perfusion (right). The currents were elicited in response to 10 mV voltage steps from -60 mV to +90 mV from a holding potential of -50 mV. (B) I - V curves corresponding to the control conditions and 15 min of 100 $\mu\text{mol/L}$ HCTZ perfusion. The BK channel currents were normalized to the ones elicited by the +90 mV voltage step under control conditions. (C) Fractional increases in current size at +40 mV induced by 100 and 300 $\mu\text{mol/L}$ HCTZ. No statistically significant differences were observed (ANOVA, Ranks test, $P>0.05$).

by $\alpha\beta_1$ heteromeric channels after 10–15 min of application (Figure 4A, B; $P<0.05$).

Next, we studied the HCTZ-induced BK activation in the heterologous system in more detail. Thus, the effect of the diuretic on the whole-cell BK current resulting from $\alpha\beta_1$ expression was examined at 10, 30, 100 and 300 $\mu\text{mol/L}$. HCTZ induced a left-shift in the I - V curves (Figure 4A, B; $P<0.05$). Consistently, higher increases in fractional activity were observed at less depolarized voltages, at which the basal open probability is lower (Figure 4 C). Additionally, we plotted the concentration-response curve at all membrane potentials where the HCTZ-induced current increase was statistically significant (from +20 to +90 mV) and fitted these data to a Hill equation. The current-enhancing effect of HCTZ presents an EC_{50} of 28.4 $\mu\text{mol/L}$ (95% CI: 10.4 - 77.7 $\mu\text{mol/L}$) measured at +40 mV, rendering a $\text{pD}_2=4.546\pm 0.211$ (n : 5–8, Figure 4 D). Moreover, both pD_2 and the Hill slope of HCTZ activation were voltage independent (Figures 4E and 4F, respectively, $P>0.05$). These results showed that HCTZ increased the BK current when the channel emulates the basic composition of this channel in HUASMCs^[32].

The results, showed above on native HUASMCs demonstrated that cellular integrity is essential for HCTZ-induced BK channel activation (Figure 2). To validate these findings, we determined whether cellular integrity is required for the

HCTZ-induced activation of recombinant BK channels containing the β_1 -subunit in a mammalian expression system. First, we obtained macroscopic currents (in the IO configuration) evoked from β_1 -containing BK channels in the presence or absence of HCTZ in HEK cells. HCTZ had no effect on BK current at any examined concentration (10, 30 and 100 $\mu\text{mol/L}$) (Figure 5 C, $P>0.05$). Figure 5 B shows the I - V relationships obtained for each condition. The negative results from the IO configuration could indicate that this diuretic requires cell integrity to activate the BK channel. However, considering the poor lipophilicity of HCTZ (log P : -0.07), these results could also indicate that the region responsible for BK channel activation is located on the extracellular side of the channel. To clarify this issue, we examined the effect of HCTZ on recombinant BK channels containing the β_1 -subunit in HEK293T cells at single channel resolution using the cell-attached (CA) configuration. A concentration of 100 $\mu\text{mol/L}$ HCTZ increased the steady state BK channel open probability at a 40 mV membrane potential. Figure 5D shows a typical stability plot of channel activation, where the open probability (expressed as NPo) was calculated every 30 s and plotted as a function of the recording time. HCTZ slowly increased the NPo of the BK channel, reaching maximal steady state activation after approximately 15 min of HCTZ perfusion. Typical recordings of BK channel activity before and after diuretic perfusion are




Figure 4. HCTZ activates recombinant BK channels expressed with regulatory β_1 -subunits in HEK293T cells in a concentration-dependent manner. Superimposed representative WCR currents of BK channels expressed in HEK293T cells with the β_1 -subunit, before (left) and after 15 min of 10 (A) and 100 $\mu\text{mol/L}$ (B) HCTZ perfusion (middle). The currents were elicited in response to 10 mV voltage steps from -60 mV to +90 mV from a holding potential of -50 mV. Mean current vs voltage (I - V) curves corresponding to the control conditions and 15 min of 10 and 100 $\mu\text{mol/L}$ HCTZ perfusion (right). The BK channel currents were normalized to the ones elicited by the +90 mV voltage step under control conditions. The symbol * indicates statistically significant difference from control (paired t test, $P < 0.05$). (C) Fractional increases in current size at different voltages ($n=6-8$) induced by 10 and 100 $\mu\text{mol/L}$ HCTZ. (D) Concentration-dependence of HCTZ-induced BK channel activation. Currents were measured at +40 mV and a single concentration of HCTZ shown was tested in each cell. Data points were fitted by a Hill function (solid line). Half-maximal effective concentration (EC_{50}): 28.4 $\mu\text{mol/L}$ (95%CI: 10.4–77.7 $\mu\text{mol/L}$), Hill slope: 1.6 (95% CI: 0.586 to 2.781). (E, F) Voltage-dependence of the EC_{50} , expressed as pD_2 (\pm SEM), and Hill Slope (\pm SEM) of BK channel activation. Each value was obtained as in D. No statistically significant differences were observed in both parameters (F method, $P > 0.05$).




Figure 5. HCTZ requires cell integrity to activate recombinant BK channels expressed with the β_1 -subunit in HEK293T cells. (A) Superimposed representative cell-free IO currents of BK channels expressed in HEK293T cells with the β_1 -subunit, before (left) and after 15 min of 100 $\mu\text{mol/L}$ HCTZ perfusion (right). The currents were elicited in response to 20 mV voltage steps from -90 mV to +130 mV from a holding potential of 0 mV. (B) I - V curves corresponding to the control conditions and 15 min of 100 $\mu\text{mol/L}$ HCTZ perfusion recorded as A. The BK channel currents were normalized to the ones elicited by the +130 mV voltage step under control conditions. (C) Fractional increases in current size at +90 mV induced by 10, 30 and 100 $\mu\text{mol/L}$ HCTZ. No statistically significant differences were observed (ANOVA, Ranks test, $P > 0.05$). (D) Temporal course of 100 $\mu\text{mol/L}$ HCTZ effect on BK channels expressed in HEK293T cells with the β_1 -subunit recorded in the Cell-Attached configuration at 40 mV. Single-channel activity was expressed as NPo, which values were calculated every 30 s and plotted as a function of recording time. The inset corresponds to representative traces of BK channel activity in control condition and during 100 $\mu\text{mol/L}$ HCTZ perfusion. (E) Mean changes in activity values (NPo) of BK channels recorded as described in Figure 5D in control conditions and with 100 $\mu\text{mol/L}$ HCTZ. The symbol * indicates statistically significant difference from control (paired t test, $P < 0.05$).

included in Figure 5D. HCTZ induced a significant 4.72 ± 1.24 -fold increase in the NPo values (Figure 5E, $P < 0.05$, $n = 3$). This change in channel activity recorded in the CA configuration can only occur through indirect mechanisms or the direct activity of membrane permeable drugs that can access the ion channel present in the patch only after crossing the plasma membrane from the bath solution to the cytoplasmic milieu. The results in the IO configuration enabled the rejection of the latter mechanism, confirming the cell integrity requirement for BK channel activation by HCTZ.

Discussion

The present study represents the first use of patch-clamp electrophysiology to evaluate the effect of HCTZ, a thiazide diuretic, on BK currents in a VSMC. Evaluating the effect of HCTZ on whole-cell BK currents in both native and heterologous systems demonstrated that 1) HCTZ activates the BK channel present in human VSMCs, confirming this channel as a probable target for the diuretic's vasodilator effect; 2) The presence of the β_1 accessory subunit is necessary for HCTZ to activate BK channels; and 3) HCTZ-induced BK channel activation requires cellular integrity.

The reduction in peripheral vascular resistance is a common property of thiazide-like diuretics that is not related to the NCC blockage^[2]. However, evidence suggests that different thiazides do not share the same mechanisms for the induction of vasodilation^[5, 8]. In particular, Calder *et al* previously proposed the HCTZ-induced BK channel activation^[5, 8], as a probable mechanism to explain the dilation of guinea pig mesenteric and human subcutaneous arteries through the *in vitro* application of HCTZ. Indeed, in the guinea pig model, the relaxation effect of this diuretic was independent of the presence or absence of the endothelium^[8]. Additionally, HCTZ induced Rb^{86} efflux^[12] and vessel relaxation, both actions being largely blocked by the selective BK channel blocker iberiotoxin^[8]. Subsequently, Pickkers *et al*^[7], also studying the mesenteric guinea pig artery, demonstrated that HCTZ reduced the increase in intracellular Ca^{2+} concentration evoked by norepinephrine without affecting the Ca^{2+} release from intracellular Ca^{2+} stores. These authors speculated that HCTZ activates BK channels, and this activation, acting as a hyperpolarizing mechanism, reduces the Ca^{2+} influx through the VOCC. Collectively, these previous studies have suggested that BK channel activation contributes to HCTZ-induced vasodilation. However, whether HCTZ increases the smooth muscle BK potassium current through direct or indirect interactions with this channel protein has remained unknown. Addressing this dichotomy acquired particular relevance in light of studies showing that BK channel activity could be enhanced by vasodilator substances acting directly on the channel protein^[30, 43] and via intracellular signaling and shuttling of BK proteins between intracellular organelles and the plasmalemma^[16]. In the present study, using HUASMCs, we propose that HCTZ activates the BK channel not through a direct interaction between the diuretic and the channel protein but rather through an indirect mechanism that requires cell integrity.

This idea is sustained by the fact that BK current enhancement was observed only in whole-cell (Figures 1 and 4) and cell-attached configurations (Figure 5D), while in the cell-free inside-out configuration (Figures 2 and 5A), neither the unitary channel conductance nor its open probability were modified by HCTZ. The similar outcomes between studies using vascular smooth muscle and HEK cells strongly suggests that the indirect mediator(s) of HCTZ-thiazide interaction on BK channels is conserved between two different cell types.

Several mechanisms can be proposed to explain how HCTZ indirectly alters channel activity. There is a growing list of interacting proteins and mediators associated with the C-terminus of the BK α -subunit. Moreover, these molecules depend on each particular cellular environment, α -subunit splicing variants, and co-expression with auxiliary subunits^[44]. This list can include different intracellular ions (such as Mg^{2+} , Ca^{2+} and H^+); protein kinases (such as PKA, PKC, PKG, PyK2 and FAK); intracellular mediators (such as GMPc, PIP_2 and arachidonic acid); receptors that regulate channel activity independent of downstream pathways (such as β_2 -adrenergic and thromboxane A_2 receptor); and structural cell proteins (*eg*, caveolin and actin)^[44]. It has been hypothesized that HCTZ-induced increases in intracellular pH in VSMCs (via carbonic anhydrase inhibition) is responsible for BK channel activation^[45]. This hypothesis contrasts with the low antihypertensive efficacy of specific carbonic anhydrase inhibitors^[1], such as acetazolamide. The present data also contrast with the "pH hypothesis": we demonstrated that HCTZ action is conserved under conditions in which changes in pH_i in the electrode solution (*ie*, WCR configuration) were precluded by strong buffering. Thus, it is highly likely that the mediators of HCTZ BK channel activation in HUASMCs are membrane bound (*eg*, PKC translocation and BK subunit shuttling).

The experiments on BK channel expressed in HEK293T cells further clarify the mechanism implicated in channel activation by HCTZ. In the whole cell configuration, HCTZ produces a concentration-dependent BK current enhancement with an EC_{50} value of $28 \mu\text{mol/L}$ (Figure 4D). Interestingly, this concentration was similar to that measured by Calder *et al*^[8] in *ex vivo* experiments and the concentration used in previous studies^[5-7, 12, 45]. Additionally, the EC_{50} and Hill slope values were both voltage independent (Figures 4E and 4F), indicating that putative intracellular mediators involved in the indirect effect of HCTZ, are insensitive to the membrane voltage field and likely membrane associated (similar to observations in HUASMCs, and the effect is lost in IO patches).

The physiological and pharmacological modulation of the BK channel, including by HCTZ, is variable among different cell types^[5, 19]. The differential expression of the auxiliary β -subunits can explain, in part, this variability, as these subunits have been reported to interact with the α -subunit, promoting or decreasing the effects of endogenous or exogenous substances on channel activity^[19, 30, 31, 43]. Moreover, the expression of these four subtypes is different among tissues^[21, 46, 47] and even among different arteries from the same species^[48]. Thus, β -subunits would be key structures to obtain a tissue-

selective pharmacological modulation of the BK channel^[19]. In the case of HCTZ, we demonstrated that the presence of the β_1 -subunit, which is primarily expressed in smooth muscle cells, is a requirement for BK channel activation (Figures 3 and 4). This result also explains why Tricarico *et al* did not observe BK channel current enhancement by HCTZ in skeletal muscle cells^[37], where the channel is expressed without any β -subunit, and in HEK cells transfected only with the α -subunit^[36]. Collectively, the present results, which include both the complexity of the native cells and the simplicity of the heterologous system, demonstrated that the activation of the BK channels by HCTZ is effective in human vascular cells, likely involving an intracellular factor, and shows selectivity for the α - β_1 -complex without affecting the channel formed only by four α -subunits.

Notably, the HCTZ effect depends on the species employed for the experimental procedures^[5]. Particularly, in human vasculature, the activation of a K^+ channel by thiazide-like diuretics was supported by previous *in vivo* and *ex vivo* experiments. HCTZ relaxes human subcutaneous and internal mammary arteries, and this effect is blocked by charybdotoxin, which inhibits BK and intermediate Ca^{2+} -activated channel^[5, 6]. Moreover, tetraethylammonium (a nonselective K^+ channels blocker) avoids the *in vivo* vasodilator effect of HCTZ in the human forearm artery^[4]. Here, we demonstrated BK channel activation induced by HCTZ in human umbilical artery smooth muscle cells. Taken together, these results suggest that HCTZ-induced BK activation is a relevant process in human vasculature and support the idea that this activation is likely involved in the vasorelaxant effect of this commonly used antihypertensive drug.

In conclusion, we propose that BK channel activation by HCTZ could contribute to its vasodilator effects, an action resulting from HCTZ acting on the SMC itself, while requiring cell integrity but not direct binding to the BK channel proteins. Thus, the present experimental approach represents significant progress in determining the molecular target or signaling pathway responsible for HCTZ-induced SMC relaxation. Moreover, we speculate that BK channel activation in vascular SMCs may be, at least partially, responsible for the efficacy of hypertension treatment with thiazide diuretics. Finally, we provide additional evidence for the growing acceptance of using the BK channel as a target for hypertension treatment, focusing on the β_1 -subunit as the key to obtaining selectivity in VSMCs.

Acknowledgments

The authors gratefully acknowledge the excellent technical assistance of Mr Ángel Flores CASTRO and Ms Julia de SANTIS. The authors would also like to thank Dr Miriam MACHUCA and the staff of the Hospital Gutierrez from La Plata for the collection of umbilical cords. This work was financially supported through a grant from the Agencia Nacional de Promoción Científica y Tecnológica (ANPCYT), Argentina (PICT 2014-0603). The authors are grateful to Dr Alejandro REBOLLEDO and Nicolás ENRIQUE for critical feedback on the manuscript.

Author contribution

Pedro MARTÍN designed the research, performed the research, analyzed the data and drafted the manuscript; Melisa MONCADA performed the research; Guruprasad KUNTA-MALLAPPANAVAR drafted the manuscript; Alex M DOPICO revised the manuscript for important intellectual content; and Verónica MILESI designed the research and drafted the manuscript.

Abbreviations

HCTZ: hydrochlorothiazide; NCC: Na^+/Cl^- cotransporter; BK channel: large conductance voltage- and Ca^{2+} -dependent potassium channel; VOCCs: voltage-operated calcium channels; VSMCs: vascular smooth muscle cells; HUAs: human umbilical arteries; HUASMCs: human umbilical artery smooth muscle cells; DM: dissociation medium; DMEM: Dulbecco's modified Eagle's medium; FBS: fetal bovine serum; eGFP: enhanced green fluorescent protein; IO: inside-out configuration; Po: open probability; BS: bath solution; WCR: whole-cell recording; 4-AP: 4-aminopyridine; *I-V*: current-voltage relationship; DMSO: dimethyl sulfoxide; K_v : voltage-dependent K^+ channels

References

- 1 James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, *et al*. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014; 311: 507–20.
- 2 Duarte JD, Cooper-DeHoff RM. Mechanisms for blood pressure lowering and metabolic effects of thiazide and thiazide-like diuretics. *Expert Rev Cardiovasc Ther*. 2010; 8: 793–802.
- 3 van Brummelen P, Man IVA, Schalekamp MA. Hemodynamic changes during long-term thiazide treatment of essential hypertension in responders and nonresponders. *Clin Pharmacol Ther* 1980; 27: 328–36.
- 4 Pickkers P, Hughes AD, Russel FG, Thien T, Smits P. Thiazide-induced vasodilation in humans is mediated by potassium channel activation. *Hypertension* 1998; 32: 1071–6.
- 5 Calder JA, Schachter M, Sever PS. Vasorelaxant actions of thiazides and related drugs. *Br J Pharmacol* 1992; 105: 307P.
- 6 Afsar S, Hemsinli D, Ozyazgan S, Akkan AG, Arslan C. The effects of potassium channels in human internal mammary artery. *Pharmacology* 2016; 97: 72–7.
- 7 Pickkers P, Hughes AD. Relaxation and decrease in $[Ca^{2+}]_i$ by hydrochlorothiazide in guinea-pig isolated mesenteric arteries. *Br J Pharmacol* 1995; 114: 703–7.
- 8 Calder JA, Schachter M, Sever PS. Ion channel involvement in the acute vascular effects of thiazide diuretics and related compounds. *J Pharmacol Exp Ther* 1993; 265: 1175–80.
- 9 Zhu Z, Zhu S, Liu D, Cao T, Wang L, Tepel M. Thiazide-like diuretics attenuate agonist-induced vasoconstriction by calcium desensitization linked to Rho kinase. *Hypertension* 2005; 45: 233–9.
- 10 Lake C, Ziegler M, Coleman M, IJ K. Hydrochlorothiazide-induced sympathetic hyperactivity in hypertensive patients. *Clin Pharmacol Ther* 1979; 26: 428–32.
- 11 Wilson I, Freis E. Relationship between plasma and extracellular fluid volume depletion and the antihypertensive effect of chlorothiazide. *Circulation* 1959; 20: 1028–36.
- 12 Calder JA, Schachter M, Sever PS. Potassium channel opening prop-

- erties of thiazide diuretics in isolated guinea pig resistance arteries. *J Cardiovasc Pharmacol* 1994; 24: 158–64.
- 13 Mironneau J, Savineau JP, Mironneau C. Compared effects of indapamide, hydrochlorothiazide and chlorthalidone on electrical and mechanical activities in vascular smooth muscle. *Eur J Pharmacol* 1981; 75: 109–13.
- 14 Nelson M, Brayden J. Regulation of arterial tone by calcium-dependent K^+ channels and ATP-sensitive K^+ channels. *Cardiovasc Drugs Ther* 1993; 7 Suppl 3: 605–10.
- 15 Ledoux J, Werner M, Brayden J, Nelson M. Calcium-activated potassium channels and the regulation of vascular tone. *Physiology (Bethesda)* 2006; 21: 69–78.
- 16 Jaggar J, Porter V, Lederer W, Nelson M. Calcium sparks in smooth muscle. *Am J Physiol Cell Physiol* 2000; 278: C235–56.
- 17 Cox RH. Changes in the expression and function of arterial potassium channels during hypertension. *Vascul Pharmacol* 2002; 38: 13–23.
- 18 Eichhorn B, Dobrev D. Vascular large conductance calcium-activated potassium channels: functional role and therapeutic potential. *Naunyn-Schmiedeberg's Arch Pharmacol* 2007; 376: 145–55.
- 19 Torres YP, Granados ST, Latorre R. Pharmacological consequences of the coexpression of BK channel alpha and auxiliary beta subunits. *Front Physiol* 2014; 5: 383.
- 20 Xia XM, Ding JP, Lingle CJ. Molecular basis for the inactivation of Ca^{2+} - and voltage-dependent BK channels in adrenal chromaffin cells and rat insulinoma tumor cells. *J Neurosci* 1999; 19: 5255–64.
- 21 Brenner R, Jegla TJ, Wickenden A, Liu Y, Aldrich RW. Cloning and functional characterization of novel large conductance calcium-activated potassium channel beta subunits, hKCNMB3 and hKCNMB4. *J Biol Chem* 2000; 275: 6453–61.
- 22 Behrens R, Nolting A, Reimann F, Schwarz M, Waldschutz R, Pongs O. hKCNMB3 and hKCNMB4, cloning and characterization of two members of the large-conductance calcium-activated potassium channel beta subunit family. *FEBS Lett* 2000; 474: 99–106.
- 23 Uebele VN, Lagrutta A, Wade T, Figueroa DJ, Liu Y, McKenna E, et al. Cloning and functional expression of two families of beta-subunits of the large conductance calcium-activated K^+ channel. *J Biol Chem* 2000; 275: 23211–8.
- 24 Wallner M, Meera P, Toro L. Molecular basis of fast inactivation in voltage and Ca^{2+} -activated K^+ channels: a transmembrane beta-subunit homolog. *Proc Natl Acad Sci U S A* 1999; 96: 4137–42.
- 25 Knaus HG, Folander K, Garcia-Calvo M, Garcia ML, Kaczorowski GJ, Smith M, et al. Primary sequence and immunological characterization of beta-subunit of high conductance Ca^{2+} -activated K^+ channel from smooth muscle. *J Biol Chem* 1994; 269: 17274–8.
- 26 Gonzalez C, Baez-Nieto D, Valencia I, Oyarzun I, Rojas P, Naranjo D, et al. K^+ channels: function-structural overview. *Compr Physiol* 2012; 2: 2087–149.
- 27 Sweet TB, Cox DH. Measuring the influence of the BK_{ca} {beta}1 subunit on Ca^{2+} binding to the BKCa channel. *J Gen Physiol* 2009; 133: 139–50.
- 28 Latorre R, Brauchi S. Large conductance Ca^{2+} -activated K^+ (BK) channel: activation by Ca^{2+} and voltage. *Biol Res* 2006; 39: 385–401.
- 29 Knaus HG, Garcia-Calvo M, Kaczorowski GJ, Garcia ML. Subunit composition of the high conductance calcium-activated potassium channel from smooth muscle, a representative of the mSlo and slowpoke family of potassium channels. *J Biol Chem* 1994; 269: 3921–4.
- 30 Bukiya AN, Liu J, Toro L, Dopico AM. Beta1 (KCNMB1) subunits mediate lithocholate activation of large-conductance Ca^{2+} -activated K^+ channels and dilation in small, resistance-size arteries. *Mol Pharmacol* 2007; 72: 359–69.
- 31 Bukiya AN, Vaithianathan T, Toro L, Dopico AM. Channel beta2-4 subunits fail to substitute for beta1 in sensitizing BK channels to lithocholate. *Biochem Biophys Res Commun* 2009; 390: 995–1000.
- 32 Martín P, Moncada M, Enrique N, Asuaje A, Valdez Capuccino JM, Gonzalez C, et al. Arachidonic acid activation of BKCa (Slo1) channels associated to the β 1-subunit in human vascular smooth muscle cells. *Pflugers Arch* 2013; 466: 1779–92.
- 33 Hoshi T, Tian Y, Xu R, Heinemann SH, Hou S. Mechanism of the modulation of BK potassium channel complexes with different auxiliary subunit compositions by the omega-3 fatty acid DHA. *Proc Natl Acad Sci U S A* 2013; 110: 4822–7.
- 34 Fernandez-Fernandez JM, Tomas M, Vazquez E, Orio P, Latorre R, Senti M, et al. Gain-of-function mutation in the KCNMB1 potassium channel subunit is associated with low prevalence of diastolic hypertension. *J Clin Invest* 2004; 113: 1032–9.
- 35 Yang Y, Li PY, Cheng J, Mao L, Wen J, Tan XQ, et al. Function of BKCa channels is reduced in human vascular smooth muscle cells from Han Chinese patients with hypertension. *Hypertension* 2013; 61: 519–25.
- 36 Tricarico D, Mele A, Calzolaro S, Cannone G, Camerino GM, Dinardo MM, et al. Emerging role of calcium-activated potassium channel in the regulation of cell viability following potassium ions challenge in HEK293 cells and pharmacological modulation. *PLoS One* 2013; 8: e69551.
- 37 Tricarico D, Mele A, Conte Camerino D. Carbonic anhydrase inhibitors ameliorate the symptoms of hypokalaemic periodic paralysis in rats by opening the muscular Ca^{2+} -activated- K^+ channels. *Neuromuscul Disord* 2006; 16: 39–45.
- 38 Klockner U. Intracellular calcium ions activate a low-conductance chloride channel in smooth-muscle cells isolated from human mesenteric artery. *Pflugers Arch* 1993; 424: 231–7.
- 39 Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch* 1981; 391: 85–100.
- 40 Martín P, Enrique N, Roldán Palomo AR, Rebolledo A, Milesi V. Bupivacaine inhibits large conductance, voltage- and Ca^{2+} -activated K^+ channels in human umbilical artery smooth muscle cells. *Channels (Austin)* 2012; 6: 174–80.
- 41 Martín P, Rebolledo A, Palomo AR, Moncada M, Piccinini L, Milesi V. Diversity of potassium channels in human umbilical artery smooth muscle cells: a review of their roles in human umbilical artery contraction. *Reprod Sci* 2014; 21: 432–41.
- 42 Milesi V, Raingo J, Rebolledo A, Grassi de Gende AO. Potassium channels in human umbilical artery cells. *J Soc Gynecol Investig* 2003; 10: 339–46.
- 43 Bukiya A, McMillan J, Fedinec AL PS, Miller DD, Leffler CW, Parrill AL, et al. Cerebrovascular dilation via selective targeting of the cholane steroid-recognition site in the BK channel β 1-subunit by a novel nonsteroidal agent. *Mol Pharmacol* 2013; 83: 1030–44.
- 44 Lu R, Alioua A, Kumar Y, Eghbali M, Stefani E, Toro L. MaxiK channel partners: physiological impact. *J Physiol* 2006; 570: 65–72.
- 45 Pickkers P, Garcha RS, Schachter M, Smits P, Hughes AD. Inhibition of carbonic anhydrase accounts for the direct vascular effects of hydrochlorothiazide. *Hypertension* 1999; 33: 1043–8.
- 46 Brenner R, Perez GJ, Bonev AD, Eckman DM, Kosek JC, Wiler SW, et al. Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature* 2000; 407: 870–6.
- 47 Behrens R, Nolting A, Reimann F, Schwarz M, Waldschütz R, Pongs O. hKCNMB3 and hKCNMB4, cloning and characterization of two members of the large-conductance calcium-activated potassium channel beta subunit family. *FEBS Lett* 2000; 474: 99–106.
- 48 Kuntamallappanavar G, Bisen S, Bukiya AN, Dopico AM. Differential distribution and functional impact of BK channel beta1 subunits across mesenteric, coronary, and different cerebral arteries of the rat. *Pflugers Arch* 2016; 469: 263–77.