

1 **First report of *Leptosphaeria biglobosa* 'brassicae' as causal agent of Phoma leaf spot in**
2 ***Brassica napus* (Canola) in Argentina.**

3 **Franco Rubén Rossi^{a#}, Fernando Matías Romero^a, Andrés Gárriz^a and Oscar Adolfo Ruíz^{a,b}.**

4 ^a- Instituto de Investigaciones Biotecnológicas-Instituto Tecnológico Chascomús, Universidad
5 Nacional de General San Martín-Consejo Nacional de Investigaciones Científicas y Técnicas (IIB-
6 INTECH/UNSAM-CONICET), Chascomús, Argentina. ^b- Instituto de Fisiología y Recursos
7 Genéticos Vegetales, Instituto Nacional de Tecnología Agropecuaria (IFRGV-INTA), Córdoba,
8 Argentina. [#] Corresponding author. E-mail: francorossi@intech.gov.ar

9

10 Canola (*Brassica napus* L.) is the second largest oilseed crop in the world providing 13% of the
11 world's oil supply. This crop has been grown in Argentina since the 1930s, and the area
12 devoted to its cultivation varies every year, reaching a maximum of 95000 Ha in the 2012/2013
13 growing season.

14 Phoma stem canker is considered the most important and devastating disease in *Brassica*
15 *napus* and other *Brassicaceae* species [1]. The causal agent is a complex of two closely related
16 fungal species, *Leptosphaeria maculans* and *L. biglobosa*. In Argentina, the presence of *L.*
17 *maculans* in canola plants was reported for the first time in 2004 [2], but the existence of *L.*
18 *biglobosa* has not been recorded so far.

19 During the 2015/2016 season, we collected several leaf samples with typical Phoma leaf spot
20 symptoms obtained from canola crops from the north and northeastern regions of the *Buenos*
21 *Aires* province. The necrotic lesions were circular to irregularly oval, 8 to 15 mm in diameter,
22 pale brown in the center, grayish green at the margin and characterized with the presence of
23 pycnidia. Several leaf pieces with lesions were rinsed twice with deionized sterile water and
24 placed in a humid chamber during 2-3 days to induce the pycnidia to exude cirri of conidia.
25 After this period, one cirrus per sample was transferred onto PDA plates supplemented with
26 antibiotics (15 mg/L STR, 15 mg/L GEN and 12 mg/L TET) using an inoculation needle under
27 stereoscopic microscope. Thus, several isolates were obtained, some of them showing rapid
28 mycelial growth rate and pigment production on PDA medium, as showed by the isolate
29 Tapidor of *L. biglobosa* used as control (kindly provided by Professor Bruce Fitt, University of
30 Hertfordshire-UK). Microscopic analysis showed hyaline conidias, subcylindrical with an
31 average size of 4-5 x 1.5-2 µm [3]. In order to confirm the identity of these isolates, a PCR
32 assay using genomic DNA as template was performed to distinguish *L. maculans* from *L.*
33 *biglobosa* with the species-specific primers *LmacR*, *LmacF*, and *LbigF* in a three-primers
34 strategy [4]. These reactions gave a 444-bp amplicon as expected for *L. biglobosa* 'brassicae'.
35 In addition, these results were confirmed by sequencing the nuclear ribosomal internal
36 transcribed spacer (ITS) region, which showed a 99% of identity with the sequence of *L.*
37 *biglobosa* 'brassicae' deposited at the GenBank database (FO905468). *L. biglobosa* isolates
38 were then tested for pathogenicity on the canola cultivars Westar and Bioaureo 2286
39 (Nuseed). With this purpose, cotyledons of 10-day-old seedlings were pricked with a needle,
40 and each wound inoculated with 10 µl of a conidial suspension (10⁷ conidia/ml). Fourteen days
41 after inoculation, irregular and brown necrotic lesions were visible at the site of inoculation.
42 These cotyledons were detached and placed in a humid chamber to induce pycnidia formation.
43 After three days cirri of conidia were transferred to a plate with PDA supplemented with
44 antibiotics as mentioned above. The identity of these isolates of *L. biglobosa* were confirmed
45 by pigment production on PDA medium and by PCR assay using species-specific primers. To
46 our knowledge, this is the first report of *L. biglobosa* 'brassicae' as a pathogen of canola in
47 Argentina. This finding shows that not only *L. maculans* but also *L. biglobosa* are the causal
48 agents of Phoma leaf spot in Argentina's canola cropping areas. Therefore, precise molecular
49 phenotyping techniques are necessities to identify the causal agent of this disease.

Acknowledgments.

We are grateful to J. L. Burgos and J. P. Ezquiaga (IIB-INTECH) for technical assistance. This work was supported by grant PICT2015-2956 from the National Agency for the Promotion of Science and Technology (ANPCyT, Argentina). F.R.R., A.G. and O.A.R. are staff researchers of the National Research Council (CONICET, Argentina). F.M.R is a post-doctoral fellow of ANPCyT.

1. Fitt B, Brun H, Barbetti M, Rimmer S (2006) World-wide importance of phoma stem canker (*Leptosphaeria maculans* and *L. biglobosa*) on oilseed rape (*Brassica napus*). *Eur J Plant Pathol* 114:3–15. doi: 10.1007/s10658-005-2233-5
2. Gaetan SA (2005) First Outbreak of Blackleg Caused by *Phoma lingam* in Commercial Canola Fields in Argentina. *Plant Dis* 89:435. doi: 10.1094/PD-89-0435B
3. Boerema G, Gruyter J, Noordellos M, Hamers M (2004) PHOMA IDENTIFICATION MANUAL. Differentiation of Specific and Infra-specific Taxa in Culture.
4. Liu SY, Liu Z, Fitt BDL, Evans N, Foster SJ, Huang YJ, Latunde-Dada AO, Lucas JA (2006) Resistance to *Leptosphaeria maculans* (phoma stem canker) in *Brassica napus* (oilseed rape) induced by *L. biglobosa* and chemical defence activators in field and controlled environments. *Plant Pathol* 55:401–412. doi: 10.1111/j.1365-3059.2006.01354.x