

# The thickening of the thin disk in the third Galactic quadrant<sup>1</sup>

Giovanni Carraro<sup>1</sup>

European Southern Observatory, Alonso de Cordova 3107, Casilla 19001, Santiago 19, Chile

`gcarraro@eso.org`

Rubén A. Vázquez

Facultad de Ciencias Astronómicas y Geofísicas (UNLP), Instituto de Astrofísica de La Plata (CONICET, UNLP), Paseo del Bosque s/n, La Plata, Argentina

`rvazquez@fcaglp.unlp.edu.ar`

Edgardo Costa

Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile

`costa@das.uchile.cl`

Javier A. Ahumada

Observatorio Astronómico, Universidad Nacional de Córdoba, Laprida 854, 5000 Córdoba, Argentina

`javier@oac.uncor.edu`

and

Edgar E. Giorgi

Facultad de Ciencias Astronómicas y Geofísicas (UNLP), Instituto de Astrofísica de La Plata (CONICET, UNLP), Paseo del Bosque s/n, La Plata, Argentina

`egiorgi@fcaglp.unlp.edu.ar`

Received \_\_\_\_\_;    accepted \_\_\_\_\_

---

<sup>1</sup>On leave from Dipartimento di Astronomia, Università di Padova, Italy.

## ABSTRACT

In the third Galactic quadrant ( $180^\circ \leq l \leq 270^\circ$ ) of the Milky Way, the Galactic thin disk exhibits a significant warp —shown both by gas and young stars— bending down a few kpc below the formal Galactic plane ( $b = 0^\circ$ ). This warp shows its maximum at  $l \sim 240^\circ$ , in the direction of the Canis Major constellation. In a series of papers we have traced the detailed structure of this region using open star clusters, putting particular emphasis on the spiral structure of the outer disk. We noticed a conspicuous accumulation of young star clusters within 2-3 kpc from the Sun and close to  $b = 0^\circ$ , that we interpreted as the continuation of the Local (Orion) arm towards the outer disk. While most clusters (and young stars in their background) follow closely the warp of the disk, our decade-old survey of the spiral structure of this region led us to identify three clusters, Haffner 18(1 and 2) and Haffner 19, which remain very close to  $b = 0^\circ$  and lie at distances (4.5,  $\sim 8.0$ , and 6.4 kpc) where most of the material is already significantly warped.

Here we report on a search for clusters that share the same properties as Haffner 18 and 19, and investigate the possible reasons for such an unexpected occurrence. We present *UBVRI* photometry of 5 young clusters, namely NGC 2345, NGC 2374, Trumpler 9, Haffner 20, and Haffner 21, which also lie close to the formal Galactic plane. With the exception of Haffner 20, in the background of these clusters we detected young stars that appear close to  $b = 0^\circ$ , and are located at distances up to  $\sim 8$  kpc from the Sun, thus deviating significantly from the warp. These populations define a structure that distributes over almost the entire third Galactic quadrant. We discuss this structure in the context of a possible thin disk flaring, in full similarity with the Galactic thick disk.

*Subject headings:* Galaxy: disk — Galaxy: structure — open clusters and associations: general — open clusters and associations: individual (Haffner 20, Haffner 21, NGC 2345, NGC 2374, Trumpler 9)

## 1. Introduction

The relevance of the study of the open star clusters of the Milky Way is unique for many areas of modern astronomy such as stellar astrophysics, stellar and Galactic evolution, and Galactic structure, and is widely recognised as such.

With this motivation, during the last decade we have been securing homogeneous  $UBVRI_{kc}$  observations (same telescope, set-up, and strategy of data analysis) of young open clusters in the third Galactic quadrant (3GQ:  $180^\circ \leq l \leq 270^\circ$ ) of the Milky Way, in order to study its complicated structure. In this fashion, we have contributed significantly to the present understanding of the spiral structure in this Galactic region (Carraro et al. 2005, Moitinho et al. 2006, Vázquez et al. 2008, Carraro et al. 2010).

Recent summaries of the status of this investigation and of the questions that still remain unanswered are given in Carraro (2014) and Carraro et al. (2014). Briefly, and using Fig. 1 as reference, our extensive campaign showed that the Local (Orion) arm (LOA) is an inter-arm feature, a bridge most probably connecting the Carina-Sagittarius arm in the first Galactic quadrant with the outer (Norma-Cygnus) arm in the 3GQ. The LOA also apparently breaks the Perseus arm, which is not prominent at all in young stars or gas in this region of the Galaxy. The extension of the LOA into the 3GQ occurs towards  $l \sim 240^\circ$ , that is in the direction of the Canis Major over-density ( $b \sim -8^\circ$ , Martin et al. 2004; Momany et al 2006; Moitinho et al. 2006). This picture confirms independent findings based on HI studies in the same region (Levine et al. 2006). Because our sample extends up to 20 kpc from the Galactic center, we were also able to trace the progressive bending of the thin disk caused by the warp, and did not find any indication of a disk cutoff or

---

<sup>1</sup>Based on observations carried out at Cerro Tololo Inter-American Observatory, under programs CHILE-0008B-017 and CHILE-0010A-006

truncation at 14 kpc from the Galactic center, as previously postulated by Galactic models (Robin et al. 1992).

Two of the main open questions that arise from Fig. 1 are: (i) at what distance from the Sun the disk starts to bend down, and (ii) whether flaring (the thickening of the disk in the vertical direction), clearly visible in the old disk population (Momany et al. 2006, López-Corredoira & Molgó 2014), has some impact on the structure of the Galactic thin disk as well.

Our previous studies indicate that the warp becomes important beyond 12 kpc from the Galactic center ( $\sim 4$  kpc from the Sun). From the works of Vázquez et al. (2008, Fig. 6), and Vázquez et al. (2010, Fig. 15), it is evident that the disk keeps close to the formal ( $b = 0^\circ$ ) plane up to about 12 kpc, and that it is very thin ( $\pm 100$  pc). Beyond this distance, the thin disk starts bending, and at the same time its vertical extent gets more and more significant. Although the bulk of young star clusters follows the warp closely, there is a group of young star clusters, like Haffner 18(1/2) and Haffner 19 (at 4.5,  $\sim 8$ , and 6.4 kpc, see Fig. 1) that clearly and un-expectedly deviate from the main disk locus. Although these clusters are relatively nearby, we found young stellar populations in their background, at distances up to 9 kpc from the Sun, which also do not follow the warp. These latter might be indicating a flaring of the thin disk, or, more conservatively, they could be tracing local irregularities (e.g. disk corrugations) of the same kind as the Big Dent (Alfaro et al. 1991, Moitinho et al 2006).

In this paper we report on an extensive search for other presumably young clusters in the same Galactic sector, and close to the formal Galactic plane, like Haffner 18 and 19.. The aim of this study is twofold: we want (1) to improve the basic parameter of a set of poorly studied, but potentially interesting clusters, and (2) to use them to consolidate our

knowledge of the thin disk structure in the 3GQ. Specifically, we present results for five open clusters (see Table 1): NGC 2345 (C0706–130), NGC 2374 (C0721–131), Haffner 20 (C0754–302), Haffner 21 (C0759–270), and Trumpler 9 (C0753–258), based on new  $UBVRI_{kc}$  data.

Since all these five clusters have been very poorly studied, a significant part of this paper is dedicated to the description of the data and the derivation of their properties. Therefore, the outline of the paper is as follows. In Section 2 we present a brief literature overview of the 5 clusters; in Section 3 we present the observations, the data reduction procedure and the photometry; in Section 4 we make a comparison of our data-set with previous investigations; in Section 5 we describe the star count procedure and derivation of the cluster’s sizes; and in Section 6 we present the analysis of the clusters’ photometric diagrams. In Section 7 we discuss the properties and spatial distribution of early type field stars in the clusters back- and fore-ground. Finally, in Section 8 we summarise the main conclusions of our work.

## 2. Literature overview

A search of the literature shows that the data for the five clusters are truly scanty, and that no modern (CCD) studies are available.

### NGC 2345:

NGC 2345 is an open cluster projected towards the Monoceros constellation. Moffat (1974) carried out photoelectric photometry of 62 stars down to  $V = 14$ . He detected variable reddening ( $E_{(B-V)} = 0.5$  to  $\sim 1.10$ ), and placed the cluster at a distance of 1.75 kpc. By means of star counts, Moffat found that the cluster diameter is 10.5 arcmin, but indicated

that, using only faint stars, it could be as large as 12.3 arcmin. Interestingly for the purpose of the present study, Moffat also isolated a small number of blue stars which he classified as OB by means of objective prism observations, and that could be at distances as large as 10 kpc. Lastly, from the inspection of the photometric diagrams he inferred for the cluster an age of  $\sim 6 \times 10^6$  yr.

#### **NGC 2374:**

Like NGC 2345, this cluster is projected towards the Monoceros constellation. Photographic and photoelectric photometry for 39 stars was secured by Babu (1985), who estimated a uniform color excess  $E(B - V) = 0.175$ , an age of  $75 \times 10^6$  yr, and a distance of 1.2 kpc. We note that the age derived by Babu (1985) is significantly younger than previous estimates, such as those from Fenkart et al. (1972) and Lyngå (1980), who report ages of  $3.5 \times 10^8$  and  $2 \times 10^9$  yr, respectively.

#### **Trumpler 9:**

Trumpler 9 is a Puppis association object. Pişmiş (1970) found, from photoelectric and photographic data, that the earliest spectral type is A0, that its reddening in  $(B - V)$  is at most 0.2 mag, and that its spectroscopic distance is approximately 900 pc. Vogt & Moffat (1972) used photoelectric  $UBV-H\beta$  photometry and derived a larger reddening  $E(B - V) = 0.29$ , an apparent distance modulus of 11.98, and a distance of 1.62 kpc. Their estimated bluest spectral type is B1, very different from Pişmiş’s results.

#### **Haffner 21 and 20:**

These two clusters also belong to Puppis association. Based on photographic photometry, Fitzgerald & Moffat (1974) established B9 as the earliest spectral type present in Haffner 20 and Haffner 21. They also determined a distance and reddening of 2.4 kpc and 0.55 mag for Haffner 20, and of 3.3 kpc and 0.20 mag for Haffner 21, respectively.



### 3. Observations, data reduction and photometry

In this Section we briefly introduce the observation strategy, the collected data, and the technique employed to reduce them. Since this work is part of a long series of papers, full details about the observations, data reduction procedure, and the photometry can be found in previous papers from our group (see, e.g., Carraro et al. 2010a,b).

#### 3.1. Observations

Observations were carried out with the Y4KCAM camera attached to the Cerro Tololo Inter-American Observatory (CTIO, Chile) 1-m telescope, operated by the SMARTS consortium<sup>2</sup>, in February 2008 and December 2010. This camera is equipped with an STA  $4064 \times 4064$  CCD<sup>3</sup> with  $15\text{-}\mu\text{m}$  pixels, yielding a scale of  $0.289''/\text{pixel}$  and a field-of-view (FOV) of  $20' \times 20'$  at the Cassegrain focus of the telescope.

In Table 2 we present the log of our *UBVRI* observations. All observations were carried out in photometric, good-seeing conditions. Our *UBVRI* instrumental photometric system was defined by the use of a standard broad-band Kitt Peak *UBVRI<sub>kc</sub>* set of filters.<sup>4</sup> To determine the transformation from our instrumental system to the standard Johnson-Kron-Cousins system, and to correct for extinction, each night we observed Landolt’s area SA 98 (Landolt 1992) multiple times (3 or 4), and with different air-masses ranging from  $\sim 1.1$  to  $\sim 2.2$ . Field SA 98 includes over 40 well-observed standard stars, with a good magnitude and color coverage:  $9.5 \leq V \leq 15.8$ ,  $-0.2 \leq (B - V) \leq 2.2$ ,

---

<sup>2</sup><http://http://www.astro.yale.edu/smarts>

<sup>3</sup><http://www.astronomy.ohio-state.edu/Y4KCam/detector.html>

<sup>4</sup><http://www.astronomy.ohio-state.edu/Y4KCam/filters.html>

$$-0.3 \leq (U - B) \leq 2.1.$$

### 3.2. Reductions

Basic reduction of the CCD frames was done using the Yale/SMARTS y4k reduction script based on the IRAF<sup>5</sup> package CCDRED, which includes bias and sky-flats correction. Photometry was then performed using IRAF’s DAOPHOT and PHOTCAL packages. Instrumental magnitudes were extracted following the point spread function (PSF) method (Stetson 1987) using a quadratic, spatially variable master PSF (PENNY function). We typically use a minimum of 50 PSF stars spread over the detector. Finally, the PSF photometry was aperture corrected using aperture corrections determined making aperture photometry of bright, isolated stars in the field.

### 3.3. The photometry

To transform our instrumental magnitudes to the standard system we used equations of the form:

$$M = m + c_0 + c_1 \times X + c_2 \times (B - V), \quad (1)$$

where  $M$  and  $m$  are the standard and instrumental magnitudes,  $X$  is the airmass, and  $c_0$ ,  $c_1$ ,  $c_2$  are the zero point, first-order extinction, and color term coefficients, respectively. We note that the use of second-order extinction terms did not improve the fit. In Table 3 we list the coefficients of the transformation equations, and their corresponding errors, for

---

<sup>5</sup>IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

each night.

In this way we produced calibrated  $UBVRI_{kc}$  photometric catalogs for NGC 2345, NGC 2374, Haffner 20, Haffner 21, and Trumpler 9, containing 6326, 3266, 3740, 6725, and 6119 entries, respectively.

These optical catalogs were cross-correlated with 2MASS (Skrutskie et al. 2006) to convert pixel (i.e., detector coordinated) into equatorial RA and DEC for equinox J2000.0, thus providing 2MASS-based astrometry.

#### 4. Comparison with previous studies

As mentioned previously, no modern CCD studies exist for the five clusters we are presenting in this study, so we want to note that we are making a comparison of our CCD data just with photographic and photoelectric data. In general, photoelectric data are precise, but suffer from two major problems: they are not very deep, and in crowded fields they cannot resolve blends because of the relatively large fixed apertures adopted. On the other hand, photographic data are in general somewhat deeper, but are less precise, and quickly lose efficiency near the sensitivity limit because of poor quantum efficiency.

Figs. 2 and 3 show the results of the comparison; below we comment on a cluster-by-cluster basis. We note that all mean differences presented were computed in the sense our measurements minus others and errors on the mean are standard deviations.

##### **NGC 2345:**

Photometric differences with photoelectric measures for 62 stars in common with Moffat (1974) are shown in the upper panels of Fig. 2. Mean differences are:  $\Delta V = -0.086 \pm 0.07$ ,  $\Delta(B - V) = -0.015 \pm 0.03$ , and  $\Delta(U - B) = +0.04 \pm 0.08$ . While colors are in fine agreement, our  $V$  magnitudes are somewhat brighter, but still comparable within the errors.

Moffat explains that his observations were carried out using a small telescope. Therefore we believe this may lead to probable contamination in the star light due to the presence of faint neighbours in the diaphragm area and/or difficulties in removing the sky contribution onto the star light. These are the typical limitations associated with photo-electric photometry. In this respect, CCD imaging combined with PSF photometry allow more precise removal of the contamination sources and also a better isolation of the star light.

#### **NGC 2374:**

Photometric differences for 36 stars in common with Babu (1985) are shown in the lower panels of Fig. 2. If we take all the 36 stars into account (photoelectric and photographic data taken together), the mean differences are:  $\Delta V = -0.16 \pm 0.16$ ,  $\Delta(B - V) = +0.108 \pm 0.23$ , and  $\Delta(U - B) = +0.11 \pm 0.22$ . Excluding the 10 stars with photographic photometry (all fainter than  $V \sim 14.0$ ), the mean differences become:  $\Delta V = -0.15 \pm 0.09$ ,  $\Delta(B - V) = -0.01 \pm 0.05$ , and  $\Delta(U - B) = +0.03 \pm 0.11$ . In this last case the scatter around the means significantly decreases and the color agreement is fine. However, our  $V$  magnitudes still appear slightly brighter with large scatter. We could not find any detailed information either about the quality of the photoelectric observing runs by Babu (1985) [made at the Kevalur 102-cm telescope in India] or related to the errors involved in the measures, but we draw the attention to Babu who suggests photographic mean errors above 0.2 for  $B$  and  $V$  and even larger in  $U$ . This said, one can understand the zero point offset between Babu photoelectric magnitudes and ours.

#### **Trumpler 9:**

The photometric comparison is shown in the upper panels of Fig. 3. Empty triangles depict the comparison with the photographic data from Pişmiş (1970), while the comparison to her photoelectric data is depicted by filled circles. Filled squares show the comparison against the photoelectric photometry by Vogt & Moffat (1972). The comparison with photographic

photometry from Pişmiş (1970) yielded:  $\Delta V = +0.17 \pm 0.08$ ,  $\Delta(B - V) = -0.09 \pm 0.13$ , and  $\Delta(U - B) = -0.06 \pm 0.18$  for 13 stars in common. Both the offset in  $V$  and the scatter around the mean are strong. Something worse happens when comparing with photoelectric measures for 10 stars also from Pişmiş (1970). In this case we found the following differences:  $\Delta V = -0.50 \pm 0.64$ ,  $\Delta(B - V) = 0.01 \pm 0.08$ , and  $\Delta(U - B) = 0.01 \pm 0.12$ . The mean  $\Delta V$  is absolutely unrealistic. Such a curious mean difference leads us to conclude that some of her photoelectric  $V$  measures are simply wrong. There is also, in our opinion, a propagation effect producing the large  $V$  off-set with photographic and photoelectric photometry since they come from bad data calibration by Pişmiş (1970). In brief, she used this wrong photoelectric sequence to put her photographic data into the standard system. This scenario is supported by the comparison with Vogt & Moffat (1972) photometry. We found 20 stars in common with their photoelectric measures. As seen in Fig. 3, there is a significant discrepancy in  $V$  for six of them. Once these six problematic stars have been removed, the mean differences with Vogt & Moffat (1972) are:  $\Delta V = +0.05 \pm 0.06$ ,  $\Delta(B - V) = 0.00 \pm 0.04$ , and  $\Delta(U - B) = 0.03 \pm 0.12$ . The  $V$  offset is relatively small and well within the error. The six deviating stars can be either mis-identified, or variable, objects.

### **Haffner 20 and 21:**

The comparison against the photographic photometry of Fitzgerald and Moffat (1974) is shown in Fig. 3; in the middle panels for Haffner 21 (47 stars), and in the bottom panels for Haffner 20 (31 stars). Mean differences are:  $\Delta V = -0.09 \pm 0.08$ ,  $\Delta(B - V) = +0.05 \pm 0.09$ , and  $\Delta(U - B) = +0.03 \pm 0.15$  for Haffner 21. For Haffner 20:  $\Delta V = +0.06 \pm 0.09$ ,  $\Delta(B - V) = +0.06 \pm 0.11$ , and  $\Delta(U - B) = -0.13 \pm 0.14$ . In both clusters most of the scatter comes from the faint tail of the photometry, as one can readily see inspecting Fig. 3. However, and particularly in the case of Haffner 20, there is a distinctive group of largely deviating stars, some of them for more than 0.8 mag in  $V$ . These stars are distributed over

the whole magnitude range, which makes it impossible to separate, in the photographic plates, images of close stars and obtain their true photometric values. It must also be added that the calibration of photographic magnitudes by Fitzgerald and Moffat (1974) was done using (secondary) photoelectric standards from some nearby open cluster sequences. Fitzgerald and Moffat (1974) mention that the external errors of their calibration is of the order of 0.09 in  $V$ ,  $B$  and  $U$  filters. As for the internal standard errors of the mean they claim they are all above 0.05 in  $V$ ,  $B - V$  and  $U - B$ . This large uncertainty in the old photographic measures can easily explain the differences with our photometry.

## 5. Star counts and clusters's size

The linear size is a fundamental quantity for a comprehensive characterisation of a star cluster. It is necessary to estimate its evolutionary status through the analysis of its density profile and other related parameters (see Aarseth 1996; de La Fuente Marcos 1997; Kroupa et al. 2001) as the slope of the mass function. It is also routinely used as a criterion for separating cluster-dominated from field-dominated regions in photometric diagrams (Baume et al. 2004).

The usual way to estimate the radius of a star clusters is to assume a centrally peaked spherical stellar density distribution and determine the distance at which the density converges to the mean field background . This determination is done either by visually setting the limit in a plot or by fitting King (1962) or other profiles to the run of radial density. However, both approaches assume a spherically symmetric distribution and, in the case of the King profiles, it also assumes that the system is dynamically relaxed. This is not at all the case for Galactic open clusters.

Since we are mostly interested in determining the cluster radii to reduce field contamination in photometric diagrams and derive the best possible cluster parameters, the following approach has therefore been adopted: contour maps were computed in each field by choosing adequate kernel sizes and grid steps. Once a notorious stellar density is identified, a circle enclosing its full range is drawn by eye and the cluster limit is taken as that defined by the radius of this circle. We adopt the cluster center to be the geometric center of the cluster as opposed to the position of the density peak. We want to emphasise that unlike what usually happens with globular clusters, the center of an open cluster can rarely be unambiguously identified. Furthermore, assigning a unique center value to an open cluster with complicated non-circular morphology or poorly populated may often carry unclear implications of what precisely that center represents, in terms of the cluster’s structure.

Fig. 4 shows contour maps of the five clusters under analysis. The insert in each figure, the colour-coding represents the number of stars per arcmin<sup>2</sup>. With the exception of NGC 2374, we detect in all cases a clear over-density around the nominal cluster coordinates. As for NGC 2374, if the cluster exists, it might simply be larger than the area we covered with our photometry since no clear over-density is appreciable in Fig. 4. NGC 2345 and Haffner 21 emerge clearly as star clusters, but their regions suffer from variable extinction, exemplified by the conspicuous dust lanes, in both cases in the north-west area of the corresponding images. The most evident peak is that of Haffner 20, while Trumpler 9 shows several peaks, mostly in the northern region. The central peak in Trumpler 9, however, is evident enough to consider it as the star cluster since it coincides with the cluster location in earlier studies and we use it to set its limits. We recall the readers that our method is intended to just yield an approximation to the cluster size because of the loose and asymmetric nature of open star clusters, and the often wrong impression of an over density induce by variable visual absorption. This is particularly important for Trumpler 9, Haffner 21 and NGC 2345 where the variable extinction in the

stellar field introduces important fluctuations in the star density background which, in turn, alters the cluster extension.

We then used star counts to derive the radial density profile, and measure the cluster radial extent, improving over previous, mostly visual, estimates. The results are shown in Fig. 5. In all cases, a distinct over-density is detected, with radii of  $\sim 4.0, 2.5, 4.0, 2.0$ , and  $1.0$  arcmin for NGC 2345, NGC 2374, Trumpler 9, Haffner 21, and Haffner 20 (from top to bottom of Fig. 5) respectively. In the case of NGC 2374, the radius of  $1.4$  arc min refers to the central over-density as seen in Fig. 4 and commented on in Sect .6.

## 6. Analysis of the clusters' photometric diagrams

To derive the basic parameters of the star clusters under study, we have performed a detailed inspection of the two color (TCD) and color-magnitude (CMD) diagrams after having verified its reliability and defined the angular cluster sizes according to the results found in Section 5. The next step in the analysis consists in the identification of the cluster sequences in all the photometric diagrams, determine the color excesses by shifting the Schmidt-Kaler (1982) to fit the sequences in the TCD and estimate the distance by superposition of the ZAMS onto the blue envelope of the corresponding star sequence in the CMD. In each case, both the fit and its error have been estimated by eye on a trial and error basis. Once mean reddening and distance have been determined we fit isochrones (Marigo et al. 2008) computed with solar metallicity. Again, this fitting has been performed by eye but instead of error in the fitting we suggest a range of values. As for the value of the absorption law (the  $R$ -value) in each cluster we have adopted  $A_V = R \times E_{(B-V)}$  with  $R = 3.1$ . Our adoption of this  $R$ -value is supported in the long series of articles see for instance, Moitinho (2001), Carraro et al. (2007) or Vázquez et al. (2010). we have performed in this quadrant



of the Galaxy where we have demonstrated, using the  $B - V, V - I$  color-color diagrams, that the absorption law is normal with very few exceptions, as Haffer 18/19 discussed in Vázquez et al. (2010). Finally, we reiterate once again our methodology is a classical and well-known procedure (see Straizys 1991 for an exhaustive review). Error analysis has been conducted following Carraro et al. (2007) and Carraro (2011) propagation formulae, which take into account the effects of spectral mis-classification on distance and reddening. In the following we only comment on the results, on a cluster-by-cluster basis.

### NGC 2345:

The TCDs and CMDs shown in the upper panels of Fig. 6 were constructed using all stars within 3.75 arcmin from the cluster center. The TCD shows a clear and slightly blue sequence partially contaminated by field interlopers as seen in the corresponding CMDs. Indeed, a superficial view of the CMD can lead to the wrong conclusion that variable reddening is present amongst the stars in NGC 2345 in line with earlier findings from Moffat. However, a careful inspection of the position of these stars in the TCD and CMDs simultaneously is enough to conclude that most of the stars causing the spread are, indeed, foreground stars. The fit of the intrinsic color relation (Zero Age Main Sequence, ZAMS) from Schmidt-Kaler (1983) to the mean cluster sequence in the TCD (upper left panel) provides a reddening  $E_{(B-V)} = 0.59 \pm 0.04$ . Absorption is obviously highly variable across the cluster. This is evident in Fig. 4 (upper left panel) where the number of stars decreases strongly to the north-west. However, the effect of variable reddening is not so strong in the cluster area, in agreement with Fig. 6 and the low dispersion around the mean color excess. In the CMD (middle upper panel in Fig. 6) we show the same ZAMS fitted vertically to the blue envelope of the cluster sequence, which yields an apparent distance modulus  $(V - M_V) = 14.2 \pm 0.1$ . When removing the effects of visual absorption onto the apparent distance modulus we derive a distance of  $3.0 \pm 0.5$  kpc from the Sun for NGC 2345. A similar fitting to the  $V$  vs.  $(V - I)$  CMD (right upper panel) confirms previous findings.

In both CMDs, an estimate of the age is obtained by superposing the cluster sequence with models from the Padova group (Marigo et al. 2008). The super-imposed isochrones bracketing the cluster sequence are for  $\log(\text{age}) = 7.8\text{--}7.9$  (63–70 Myr).

From our analysis, NGC 2345 turns out to be significantly more distant than the 1.75 kpc given by Moffat (1974). Such a large difference in the cluster distance comes from the fact that Moffat’s data only reach  $V \sim 14$  mag, a magnitude level where confusion with field interlopers is particularly high (see Figs. 3 in Moffat paper). At the level of the magnitude cut-off of Moffat data he had no chance to see the run of the lower MS sequence as we instead do in Fig. 6. Therefore, our ZAMS fitting is much more robust because it encompasses an almost 6 mag range, down to  $V \simeq 18$ .

An interesting, final, comment is necessary for the sake of clarity and its relevance for the main aim of this work. A close inspection of the lower panels of Fig. 6, where supposedly only field stars are present, shows that several stars fall between the reference lines for stars of spectral types O4 and B5. Because these stars have unique reddening solution in the TCD, we applied the Q-method (Straizys 1991) to extract their intrinsic colors, color excesses, and distances, as we have done successfully many times in the past for several other fields (see e.g., Carraro et al. 2007; Vázquez et al. 2010; Carraro et al 2010). We will return to these stars again in Section 7.

#### **NGC 2374:**

In the previous Section we noted that this cluster is not obvious at all from the contour plot in Fig. 4 upper right panel. However, when looking at the upper panels in Fig. 7, it is quite evident the presence of a slightly evolved sequence in the CMD. The earliest spectral type is  $\sim$  B9. This sequence extends for more than five magnitudes in the CMDs and is affected

by a uniform reddening  $E_{(B-V)} = 0.07 \pm 0.03$ . Fitting a ZAMS yields an apparent distance modulus of  $(V - M_V) = 10.90 \pm 0.10$ , which, after correcting by visual absorption, places this cluster at a distance of  $1.33 \pm 0.25$  kpc, in agreement with the estimate of Babu (1985). In Fig 7 (upper panels) we have over-plotted isochrones from the Padova group (Marigo et al. 2008) on the  $V$  vs.  $(B - V)$  and  $V$  vs.  $(V - I)$  CMDs. They indicate that the cluster is 220–280 Myr ( $\log(\text{age}) = 8.35\text{--}8.45$ ). This age range indicates that the cluster is older than assumed by Babu (1985), younger than the estimates from Lynga (1980),  $2 \times 10^9$  yr, and comparable to the age obtained by Fenkart et al (1972),  $3.5 \times 10^8$  yr.

As we have already mentioned, NGC 2374 is a sparse open cluster and probably it covers a larger area than the one we have surveyed. We adopt as field stars those ones that do not belong to the cluster sequence we have defined in the TCD and CMD upper panels. Inspection of the TCD and CMDs for the stellar field in the direction of NGC 2374 (lower panels in Fig. 7) shows the presence of a population of stars (indicated by red symbols) which defines a Blue Plume (BP). In other words, the BP stars do not share the average locus occupied by field stars but they are blue-wards of the field sequence intersecting it at  $V \sim 17 - 18$  mag.

We remind the reader that previous studies from our group have shown that these stars, the BP stars, do not constitute a physical sequence of bound stars, but a superposition of stars along the line of sight (Carraro et al. 2005). We will return to these stars in the following section.

### **Haffner 20:**

This cluster stands out notoriously in the contour plot of Fig. 4. Visual inspection of the TCD and CMD (upper panels of Fig. 8) for stars within the adopted cluster radius ( $r < 2$  arcmin) confirms that Haffner 20 is a physical group. The Main Sequence (MS) of this cluster is quite clear in the TCD, and is affected by a uniform reddening of  $E(B - V$

) =  $0.65 \pm 0.03$ . Still, like in the case of NGC 2345, contamination by field interlopers is evident in the cluster area, although to a lesser degree. Fitzgerald and Moffat (1974) give a maximum  $E_{(B-V)}$  for this cluster of about 0.55. Regrettably, they do not show the corresponding TCD of this object (nor the one of Haffner 21) but we have redrawn the TCD (not shown for saving space) with their data finding a strong data dispersion that makes impossible to assign a trustable reddening to Haffner 20 (this is fully in line with our photometric comparison in Section 4). The apparent distance modulus ( $V - M_V$ ) that we infer from ZAMS fitting of the CMD is  $15.70 \pm 0.30$ , which, in combination with the reddening estimated above, places the cluster at a distance of  $5.5 \pm 1.0$  kpc from the Sun. This new value is twice the obtained by Fitzgerald and Moffat (1974) and is the result of the use of a deep photometric data-set that allows to follow the lower cluster MS and perform a more solid ZAMS fitting. The earliest spectral type, as estimated from the TCD, is about B5-B7, which suggests an age around 100 Myr. This is supported by the two isochrones from Marigo et al. (2008), for  $\log(\text{age})$  between 7.95 and 8.00, that we over-plotted on the CMDs. This range lowers the cluster age with respect to the 200 My suggested by Fitzgerald and Moffat (1974). The field toward Haffner 20 does not reveal any presence of a young population in the background. There might be various explanations for such an occurrence. One possibility is that the background population is extremely reddened, and we do not detect it because it is confused with the cluster and field MSs. Another possibility is that there is indeed no background populations. Several times in previous works (see e.g. Carraro et al. 2005) we show cases of presence and absence of background stellar populations.

#### **Haffner 21:**

A similar analysis for Haffner 20 leads to an estimate of the reddening of  $E(B - V) = 0.21 \pm 0.03$  (see Fig. 9, upper left panel). The MS for the probable members of Haffner 21 is well defined in both the TCD and the CMD, and indicates that Haffner 21 may be

slightly older than Haffner 20, because its earliest spectral type would be around B8–B9. From the CMDs (Fig 9, upper right panels) we infer an apparent distance modulus  $(V - M_V) = 13.35 \pm 0.35$ , which puts the cluster at a distance of  $3.5 \pm 0.5$  kpc from the Sun. In this case, our reddening and distance are in agreement with previous studies from Fitzgerald and Moffat (1974). The superposition of isochrones from Marigo et al. (2008) indicates that the age range for  $\log(\text{age})$  is between 8.05 and 8.1 or  $1.1$  to  $1.25 \times 10^8$  yr, significantly younger than the  $2 \times 10^8$  yr found by Fitzgerald and Moffat (1974).

In the control field (lower panels of Fig. 9, in red symbols) we detect a young population in the background of the cluster. It is not possible to position accurately this population, because it seems to be composed of stars located at very different distances, from close to the Sun to all the way up to about 10 kpc from it. We further analyze this population in Section 7.

### **Trumpler 9:**

The photometric diagrams for the stars within the cluster radius (see Fig. 10, upper panels) indicate that we are dealing with an extremely young object. The MS in all the diagrams is very tight, and the reddening solution provides a color excess  $E(B - V) = 0.20 \pm 0.02$ , with no spread. By fitting the CMDs with a ZAMS we derive an apparent distance modulus  $(V - M_V) = 12.90 \pm 0.30$ . This yields an heliocentric distance of  $2.9 \pm 0.5$  kpc, it larger than the values from Vogt & Moffat (1972) and Pişmiş (1970). This distance is consistent with Trumpler 9 being part of the Puppis OB1 association, which is also corroborated by our estimate of the cluster age ( $\sim 10$  Myr) from the isochrone fitting. A glance at the lower panels in Fig. 10 allows us to conclude that there is also an evident young stellar population in the background of Trumpler 9, as in the case of Haffner 21. Moreover, from the TCD one can conclude that young stars (red symbols) show a wide range of reddening, which

conservatively implies that they are located at very different distances, and hence they are tracing an elongated structure in this particular Galactic direction (see below).

### 6.1. Trumpler 9, Haffner 20 and Haffner 21 and their relation to the Puppis association

These three clusters are located in a region of the Galaxy historically identified as Puppis constellation, and that contains two prominent OB associations: Puppis OB1 and Puppis OB2. The approximate borders of the area are between  $l^\circ \sim 242$  and  $l^\circ \sim 246$  and from  $b^\circ \sim +2$  to  $b^\circ \sim -1$ . Although the nature and properties of these two OB associations is out of the scope of this work, still we would like to comment on the relationship of Trumpler 9, and Haffner 20 and 21 with them.

We first of all emphasise that no clear understanding of this region is available. Humphrey (1978) found various components of what is designated as Puppis OB1 association at 2.5 kpc from the Sun, while Havlen (1976) first reported a second, more distant, OB association, Puppis OB2 at 4.3 kpc. The existence of these two separate associations have been questioned by Kaltcheva and Hilditch (2000), who could not find evidences of them, but, instead, suggested the presence in the same Galaxy direction of two other star groups (at 1 kpc the first, and 3.2 the second), but significantly lower, at  $b^\circ \sim -4$ . These would surround the star cluster NGC 2439, located at 3.5-4.5 kpc from the Sun.

Since Trumpler 9 is located at 2.9 and it is  $10^6$  yrs old, we argue it is a probable distant member of Puppis OB1. On the other hand, Haffner 20, at 5.5 kpc and about  $10^8$  yrs old, would be placed in front of Puppis OB2 and be an old member of this association.

Something similar happens with Haffner 21. Its distance of 3.5 and its comparable age to Haffner 20 would make it a distant member of Puppis OB2.

Lastly, let us emphasise that the assessment of these three clusters membership to these associations (even if they really exist) requires an extensive spectroscopic and radial velocity study. In line with Kaltcheva and Hilditch (2000) study, the existence of two young stellar groups at  $b^\circ \sim -4$  and with a distance range of 1 to 3.2 kpc in addition to the more distant NGC 2439 open cluster is clearly telling us that star formation processes is still vigorous in this region of the Galaxy.

## 7. Early-type field stars in the fore- and back-ground of the clusters

Groups of evenly distributed early-type stars are frequently found in the background of Galactic open clusters (see, for instance, Carraro et al. 2005) in the third Galactic quadrant. This occurs because Galactic open clusters are mostly located at low Galactic latitudes in this Milky Way regions, and the line of sight to them often intersects parts of distant spiral arms. These stars manifest themselves in classical photometric diagrams in two ways. First, we can spot them in the *UBV* TCD in the position corresponding to stars for which a unique reddening solution is derived (Straizys 1991). The reddening solution is achieved moving the stars along the reddening lines back to the position of blue-types on the intrinsic (zero reddening) line (see Perren et al. 2012, for a vivid example and a detailed description of the method). Second, in the CMD they tend to define tight sequences, emerging from the left (the blue) side of the more prominent MS of the thin (or thick) disc stars. These features are well known as BPs (Carraro et al. 2005; Moitinho et al. 2006), and their exact location in the CMD depends on the line of sight. In this study, field blue stars have been detected in NGC 2345 (Fig 6, lower panels) and Trumpler 9 (Fig. 10, lower panels), while

BPs are evident in NGC 2374 (Fig. 7, lower panels) and Haffner 21 (Fig. 9, lower panels). In what follows we discuss the properties of these groups of stars individually.

### 7.1. NGC 2345

NGC 2345 and NGC 2374 are only 4 degrees apart on sky (see Table 1) FNGC 2345 is a young open cluster located at intersection of the Perseus arm with the LOA. The TCD of stars outside the limits of NGC 2345 (lower panels of Fig. 6) shows a large number of stars above the reference line for stars of spectral type B5, with different reddening values roughly along the color range  $-0.1 \leq (B - V) \leq 1.0$ . As we mentioned in Section 6, field blue stars in NGC 2345 were de-reddened using the Q-method to derive their intrinsic colors  $(B - V)_0$  and individual color excesses  $E(B - V)$ . We remind the reader that once these two quantities are obtained for a star, a spectral type and an absolute magnitude can be assigned using the relationships given in Schmidt-Kaler (1982). This procedure is a variation of the spectroscopic parallax method applied to get individual star distances, that has been proven to be quite solid (see, e.g., Carraro et al. 2007; Carraro et al. 2010).

In Fig. 11 we show the trend of reddening versus distance for the four fields in which blue stars have been detected. To minimise the effects of an erroneous spectral type assignment due to photometric errors, we have only taken into account stars with spectral types earlier than B4. In the case of NGC 2345, Fig. 11 shows the existence of large  $E(B - V)$  values, from 0.5 to 1.2, from the Sun to about 3.5 kpc, distance at which the reddening levels off at a mean value of  $E(B - V) = 1.2 \pm 0.2$ .

Most blue stars in the field of NGC 2345 are found south-ward the cluster. At a distance of 3.0 kpc, the reddening is larger than the reddening of the cluster itself



( $E(B - V) = 0.59$ ), as if the cluster were seen through a dust window. We note that, as shown by the contour plot presented in Fig. 4, that absorption significantly increases towards the north-west of the cluster.

Lastly, from Fig. 12 one can notice that these early-type stars are found at any distance up to 8 kpc, and that a few of them lie at distances larger than 10 kpc. All of them are located below the formal  $b = 0^\circ$  plane, roughly following the trend defined by the majority of young open clusters in the first 3 kpc from the Sun.

## 7.2. NGC 2374

In the CMD of NGC 2374 a clear BP emerges at  $V \sim 17$ . This BP corresponds to the group of stars with  $0.0 \leq (B - V) \leq 0.5$  in the TCD (lower panels of Fig. 7), where we have depicted in red about 50 BP candidate stars. The bulk of these BP stars has a reddening value of  $0.40 \pm 0.10$  (Fig. 7, left panel), and an apparent distance modulus of  $15.2 \pm 0.2$ , or a distance of  $6.1^{+0.7}_{-0.5}$  kpc (Fig. 7, middle and right panels).

This BP structure clearly deserves closer attention, so we applied the same method for determining individual distances and color excesses to the 50 BP candidate stars that we used for the blue stars in the field of NGC 2345. Inspection of Fig. 11 shows in this case that:

- BP stars begin to appear at  $d \sim 1.5$  kpc and extend for more than 8 kpc;
- color excess increases steadily from 0.25 to 0.5;
- the earliest photometric spectral type found among the BP stars in NGC 2374 corresponds to B5-type stars, while latest types are around A0.

At odds with NGC 2374 itself, early-type stars in its background are confined above the formal Galactic disk plane.

### 7.3. Haffner 21

The TCD of field stars in the direction of Haffner 21 (lower left panel of Fig. 9) reveals the presence of a strip of stars immediately below the O4 reddening line, from  $(B - V) = 0.4$  to 1.2. They are depicted by black symbols.

Haffner 21 exhibits a notorious BP (see Fig. 9, lower panel) extending for more than five magnitudes. We marked some of the stars in the BP with red symbols and fitted them with the Schmidt-Kaler ZAMS. This procedure yielded an apparent distance modulus of  $15.2 \pm 0.4$  that, for a reddening value of  $0.30 \pm 0.12$ , corresponds to a distance of  $7.1_{-1.2}^{+1.4}$  kpc. We also applied the Q-method to obtain the intrinsic properties of these stars which we assume are representative of the BP. Fig. 11 shows that these stars are distributed starting at 1.5 kpc from the Sun, and that the reddening steadily increases up to a value of about 0.25.

### 7.4. Trumpler 9

Trumpler 9 is at 1.3 degrees above the galactic plane and at 1.8 degrees away from Haffner 21; it is also above the galactic plane. These two clusters are located close to the northern edge of the Vela Gum SN remnant. As in the case of NGC 2345, in the field of Trumpler 9 there is an evident group of blue stars outside the cluster’s boundary (depicted by red symbols in Fig. 10, lower panels). In the TCD they define a strip between the reddening lines for O4- and A0-type stars, thus admitting a unique reddening solution. This situation is reflected in the CMDs (Fig. 10, mid and right lower panels), where BP stars

are colour-coded in red as well. From our analysis, the earliest photometric spectral type found among these stars is about B0.5; Fig. 11 shows that their reddening  $E(B - V)$  has a mean value of  $0.20 \pm 0.10$ , which stays constant up to 4 kpc. Beyond 4 kpc, a few stars show  $E(B - V)$  values near 0.9 and helio-distance up to 8 kpc. An outlier - having large reddening- is at 2 kpc from the Sun. Both these group are located north-east of Trumpler 9, the most absorbed region covered by our images.

## 8. Discussion and Conclusions

In this work we have presented new *UBVRI* CCD data for five poorly known Galactic star clusters: NGC 2345, NGC 2374, Haffner 20, Haffner 21, and Trumpler 9. Our data supersede previous photographic and photoelectric studies, both in depth and quality, which allowed us to constrain the clusters’s basic parameters more solidly. We report in Table 4 the results of this new analysis, in comparison with previous studies.

We stress, however, that the main goal of this work is to study the spatial distribution of early-type field stars that often happen to lie in the background of open clusters in the 3GQ of the Milky Way (Moitinho et al. 2006; Vázquez et al 2008). To summarise our conclusions along this vein, we refer the reader to Fig. 12 —which is an updated version of Fig. 1—, where we have added the new findings that have resulted from the analysis of the cluster sample presented in this paper.

Before interpreting our data, we would like to further stress that our sample is mostly made of *bona fide* OB stars. In fact, one way wonder that our sample can be significantly contaminated by old hot sub-dwarf stars, or that the OB stars we detect out of the plane are mainly OB run-away stars. We would like to comment on these two possibilities.

Hot sub-dwarf stars are often found in the Milky Way. They are associated with old stellar populations, and they are in most cases binaries. The statistics of these stars are poorly known. Early estimates (Downes 1986) count one hot sub-dwarf every ten million stars, and one B sub-dwarf every million stars, when considering one cubic parsec volume. Recent studies (Breedt & Gansicke 2011, and references therein) confirm these figures. These stars are very faint, and, when detected nearby to the Sun, can be confused with distant OB stars, in absence of any spectroscopic information. The typical magnitude of these stars is  $M_V \approx +5.0$ . This means that with our limiting magnitude of  $V \approx 17.5$  for the BPs, we can detect them at a distance modulus up to  $(m - M) \approx 12.0$ , assuming no reddening. This turns into a distance from the Sun of about 3.0 kpc, thus defining a volume of  $\sim 340,000$  cubic parsec, since our detectors covers  $20 \times 20$  arcmin<sup>2</sup> on the sky. This implies that we expect 0.2 O sub-dwarf per cubic parsec, and 0.7 B sub-dwarf per cubic parsec (according to Downes 1986) in each of our pointings. An additional point is that OB sub-dwarf stars do not define the typical tilted BP sequences we have found, which, on the other hand, resemble more the MS of nearby young open clusters.

OB run-away stars are as well often found in the Milky Way disk. They are typically associated with massive young open clusters (e.g. Westerlund 2, Carraro et al. 2013; Roman-Lopes et al. 2011), from which they escaped (Fujii & Portegies Zwart 2011). First of all we stress that the escaping direction is not always vertical with respect to the disk plane, but it is clearly random. In Westerlund 2 we believe there are two run-away OB stars out of about 50 OB stars in the cluster. No OB run-away stars have been reported in young open clusters in the anti-center direction, most probably because clusters are not massive enough to produce this phenomenon (Fujii & Portegies Zwart 2011), which indeed is mostly

seen in the nearby of massive star cluster located within the solar ring. Even assuming that the proportion of run-away stars in the outer disk is the same as in the inner, say about 20%, it is difficult to reproduce the high number of off-plane OB stars we are finding.

In Fig. 12, which covers the whole 3GQ, from  $180^\circ$  to  $270^\circ$  in Galactic longitude, we depict the positions of the young open clusters studied by our group, and of over-densities of young stars that we have discovered in their background or foreground. In this figure the Sun is located at  $(0, 0, 0)$ . Black stars are young open clusters from our previous studies (Moitinho et al. 2006, Vázquez et al. 2008, and Vázquez et al. 2010), the two black stars surrounded by a blue circle are Haffner 18 and Haffner 19, and the five red stars are the clusters analyzed in this work. Crosses mark the distribution of early-type stars in the directions of NGC 2345 (green), NGC 2374 (brown), Trumpler 9 (red), and Haffner 21 (black), respectively. Open circles are BPs detected in previous studies. The solid logarithmic curves are the approximate extrapolated locations of the Perseus and Outer arms, according to Vallee (2005) model.

The addition of the five clusters studied here has added a wealth of new information on the structure of the Galactic thin disk in the 3GQ, thanks to the fact that four of them (the exception being Haffner 20) lie in fields rich in early-type stars located at very different distances from the Sun.

NGC 2345 and the associated blue stars follow the well-established structure of the disk in the 3GQ: the disk keeps close the formal  $b = 0^\circ$  Galactic plane up to  $\sim 4$  kpc from the Sun, and then starts to bend down following the Galactic warp. However, field stars in the direction of NGC 2374, Trumpler 9 and Haffner 21 are found all the way up to the location of the outer arm, and, being these clusters at positive Galactic latitudes, they define distributions of young stars that, at odds with any previous expectations, lie above

or close to the formal  $b = 0^\circ$  Galactic plane. In other words, as illustrated in the lower panel in Fig. 12, they do not follow the Galactic warp. Beyond  $\sim 4$  kpc from the Sun, the Galactic thin disk, as defined by young stars and open clusters, thickens significantly, and this thickening keeps increasing because of the sudden appearance of the Galactic warp.

We interpret this situation as an indication that the Galactic thin disk is not only warped, but also flared, like the thick disk (Momany et al. 2006; López-Corredoira & Molgó 2014). In other words, the Galactic thin disk scale height seems to increase moving outwards from the Sun. This interpretation nicely explains the unusual, un-expected location of Haffner 18 and Haffner 19 (see Fig. 12): they are very young (NGC 2374, Trumpler 9, and Haffner 21) and distant, but lie far from the main -warped- thin disk location. The addition of background blue stars- associate with three more clusters (T- sharing the same position as Haffner 18 and 19 — lends further support to this scenario.

One may wonder whether other independent observational evidences of this thickening of the disk are available, in addition to the few stellar fields discussed here. We note that in their recent all-sky survey of HII regions in the Wide-Field Infrared Survey Explorer (WISE) database, Anderson et al. (2014) detected quite a large number of HII regions in the Galactic anti-center. Their Fig. 4 shows very clearly both the warp and the flare of the thin disk in the second and third Galactic quadrants ( $90^\circ \leq l \leq 270^\circ$ ), nicely confirming our findings.

To conclude, we have shown in our series of papers, including the present one, that early type stars and very young open clusters not only follow the warp in the 3GQ, but they also contribute to the flaring of the thin disk beyond  $\sim 4$  kpc from the Sun. This might imply that in the outer disk spiral arms are thicker than commonly believed. In fact a large fraction of the stars we have studied are younger than 100 Myr, hence it is conceivable to

think that they did not displace much from their birthplaces. This implies that these young stars formed out of vertically dispersed material, which is not concentrated into the plane, but can be found even at large distances from the mean plane.

This interpretation is supported by the evidence (Carraro et al 2005, Vázquez et al. 2008) that, besides early type stars, also CO clouds are found at any vertical distances in the 3GQ quadrant, and their spatial distribution closely young stars and star clusters. In a very recent study, Suad et al. (2014) have performed a search of HI super-shells in the second and third quadrant of the Galaxy showing that HI super-shells can be found up to 3 kpc above the galactic plane (50% of them are at less than 500 pc above the galactic plane). Early type stars, young clusters, CO clouds and HI super-shells show all the same smoothed distribution that thickens the outer Galactic disk.

GC thanks Yazan Momany for many fruitful discussions. RAV and JAA acknowledge ESO for granting a visitor-ship at ESO premises in Santiago, where part of this work was done. EC acknowledges support by the Fondo Nacional de Investigación Científica y Tecnológica (Project No. 1110100 Fondecyt), and the Chilean Centro de Excelencia en Astrofísica y Tecnologías Afines (PFB06). RAV and EEG acknowledge financial support from the PIP 1359 from CONICET. This research used the WEBDA database, maintained at the Institute for Astronomy of the University of Vienna. Lastly, we greatly appreciate the input of the referee.

## REFERENCES

- Aarseth, S. J., 1996, in *The Origins, Evolution, and Destinies of Binary Stars in Clusters*, ed. E. F. Milone, & J.-C. Mermilliod, ASP Conf. Ser., 90, 423
- Alfaro, E.J., Cabrera-Cano, J., Delgado, A.J., 1991, *ApJ*, 378, 106
- Anderson, L. D., Bania, T. M., Balser, D. S., Cunningham, V., Wenger, T. V., Johnstone, B. M., Armentrout, W. P., 2014, *ApJS*, 212, 1
- Babu, S., 1985, *Journal of Astrophys. Astr.*, 6, 61
- Baume, G., Moitinho, A., Giorgi, E. E., Carraro, G., & Vázquez, R. A. 2004, *A&A*, 417, 961
- Breedt, E., Gansicke, B.T., 2011, *ASPC*, 447, 203
- Carraro, G., Vázquez, R.A., Moitinho, A., Baume, G., 2005, *ApJ*, 630, L153
- Carraro, G., Moitinho, A., Zoccali, M., Vázquez, R.A., Baume, G., 2007, *AJ*, 133, 1058
- Carraro, G., Costa, E., & Ahumada, J. A. 2010a, *AJ*, 140, 954
- Carraro, G., 2011, *A&A*, 536, 101
- Carraro, G., Vázquez, R.A., Costa, E., Perren, G., Moitinho, A., 2010b, *ApJ*, 718, 683
- Carraro, G., Turner, D., Majaess, D., Baume, G., 2013, *A&A*, 555, 50
- Carraro, G., Perren, G., Vázquez, R.A., Moitinho, A., in "Structure and Dynamics of Disk Galaxies" , 2014, M. Seigar and P. Treuthardt, eds, *ASPC*, 480, 10
- Carraro, G., *Proceedings of the IAU Symposium No. 298, "Setting the scene for Gaia and LAMOST"*, eds. S. Feltzing, G. Zhao, N. A. Walton, and P. A. Whitelock, 2014, *IAUS* 298, 7



- de La Fuente Marcos, R. 1997, *A&A*, 322, 764
- Downes, R.A., 1986, *ApJS*, 61, 569
- Fenkart, R.P., Buser, R., Ritter, H., Schmitt, H., Steppe, H., Wagner, R., Weidemann, D.,  
1972, *A&AS*, 7, 48
- Fitzgerald, M. P., & Moffat, A. F. J. 1974, *PASP*, 86, 480
- Fujii, M.S., Portegies Zwart, S., 2011, *Science*, 334, 1380
- Havlen R. J., 1976, *A&A*, 47, 193
- Humphreys R. M., 1978, *ApJS*, 38, 309
- Kaltcheva, N.T., & Hilditch, R.W., 2000, *MNRAS* 312, 753
- King, I. 1962, *AJ*, 67, 471
- Kroupa, P., Aarseth, S., & Hurley, J. 2001, *MNRAS*, 321, 699
- Landolt, A. U. 1992, *AJ*, 104, 372
- Levine, E.S., Blitz, L., Heiles, C., 2006, *Science*, 312, 1773
- Lyngå, G. 1980, *A Computer Readable Catalogue of Open Cluster Data*, Stellar Data  
Centre, Observatoire de Strasbourg, France
- López-Corredoira, M., Molgó, J, 2014, *A&A*, 567, 106
- Marigo, P., Girardi, L., Bressan, A., Groenewegen, M.A.T., Silva, L., Granato, G.L., 2008,  
*A&A*, 482, 883
- Martin N., Ibata R.A., Bellazzini M., Irwin M.J., Lewis G.F., Dehnen W., 2004, *MNRAS*  
348, 12

- Moffat, A.F.M., 1974, A&AS 16, 33
- Moitinho, A., 2001, A&A 370, 436
- Moitinho, A., Vázquez, R.A., Carraro, G., Baume, G., Giorrgi, E.E., Lyra, W., 2006, MNRAS, 368, L77
- Momany, Y., Zaggia, S., Gilmore, G., Piotto, G., Carraro, G., Bedin, L.R., de Angeli, F., 2006, A&A, 451, 515
- Perren, G., Vázquez, R.A., Carraro, G., 2012, A&A, 548, 125
- Pişmiş, P. 1970, Bol. Obs. Tonantzintla y Tacubaya, 5, 293
- Robin, A., Creze, M., Mohan, V., 1992, ApJ, 400, L25
- Roman-Lopes, A., Barba, R.H., Morrell, N.I., 2011, MNRAS, 416, 501
- Schmidt-Kaler, Th., 1982 , Landolt-Börnstein, Numerical data and Functional Relationships in Science and Technology, New Series, Group VI, Vol. 2(b), K. Schaifers & H. H. Voigt, eds., Springer-Verlag, Berlín, p. 14
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163
- Stetson, P. B. 1987, PASP, 99, 191
- Straižys, V., 1991, in Multicolor Photometry, Astronomy and Astrophysics series 15, Pachart Publishing House
- Suad, L.A., Caiafa, C.F., Arnal, W.M., Cichowolski, S., 2014, A&A, in press
- Vallée, J. P. 2008, AJ, 135, 1301
- Vázquez, R.A., May, J., Carraro, G., Bronfman, L., Moitinho, A., Baume, G., 2008, ApJ, 672, 930

Vázquez, R.A., Moitinho, A., Carraro, G., Dias, W.S., 2010, A&A, 511, 38

Vogt N., Moffat, A.F.J., 1972, A&AS, 7, 133




Fig. 1.— The thin-disk structure in the third Galactic quadrant as traced by young open clusters and background populations. The large panel shows the plane of the disk, while the two smaller panels show the  $X$ - $Z$  and  $Y$ - $Z$  projections. Notice the particular location of the star clusters Haffner 18(1/2) and 19, above the formal ( $b = 0^\circ$ ) plane of the Galaxy. Symbols are as follows: stars are young open clusters, and empty circle background populations of early type stars.




Fig. 2.— Comparison of our CCD photometry with literature sources for NGC 2345 and NGC 2374. Differences are in the sense: our measurements minus published data. See text for details.




Fig. 3.— Comparison of our CCD photometry for Trumpler 9, Haffner 20, and Haffner 21. Differences are in the sense: our measurements minus published data. See text for symbols in Trumpler 9.




Fig. 4.— Iso-density contour maps for the five clusters studied.




Fig. 5.— Star counts as a function of radius. See text for details.






Fig. 6.— Color-color and colour-magnitude diagrams for NGC 2345 (upper panels), and for the surrounding field (lower panels). Dashed lines are isochrones from Marigo et al. (2008) shifted by color excess and distance modulus. The solid line in the two color diagram is a reddening-free ZAMS, while the dashed line is the same ZAMS shifted by the color excess. Two reference spectral types are indicated in the lower left panel, together with the reddening vector. See text for details.




Fig. 7.— Color-color and colour-magnitude diagrams for NGC 2374 (upper panels), and for the surrounding field (lower panels). Yellow dashed lines are isochrone from Marigo et al. (2008) shifted by color excess and distance modulus. The solid line in the two color diagram is a reddening-free ZAMS, while the dashed yellow line is the same ZAMS shifted by color excess to fit the cluster’s probable members (filled yellow symbols). In the lower panels, red colours are used to identify a background young population, and the ZAMS that better fits its sequence. See text for details.




Fig. 8.— Color-color and color-magnitude diagrams highlighting Haffner 20 members (upper panels), and the surrounding OB field stars (lower panels). Dashed lines are isochrone from Marigo et al. (2008) shifted by color excess and distance modulus. The solid line in the two color diagram is a reddening-free ZAMS, while the dashed line is the same ZAMS shifted by the color excess to fit the cluster’s probable members. Two reference spectral types are indicated in the lower left panel, together with the reddening vector. See text for details.




Fig. 9.— Color-color and color-magnitude diagrams for Haffner 21 (upper panels), and for the surrounding field (lower panels). Dashed lines are isochrone from Marigo et al. (2008) shifted by color excess and distance modulus. The solid line in the two color diagram is a reddening-free ZAMS, while the dashed line is the same ZAMS shifted by the color excess to fit the cluster’s probable members. In the lower panels, red colors are used to identify a background young population, and the ZAMS that better fits its sequence. See text for details.




Fig. 10.— Color-color and color-magnitude diagrams for Trumpler 9 (upper panels), and for the surrounding field (lower panels). Dashed lines are isochrone from Marigo et al. (2008) shifted by color excess and distance modulus. The solid line in the two color diagram is a reddening-free ZAMS, while the dashed line is the same ZAMS shifted by the color excess to fit the cluster’s probable members. In the lower panels, red symbols are used to identify a background young population, and the ZAMS that better fits its sequence. See text for details.




Fig. 11.— Trend of reddening as a function of distance from the Sun for the five directions studied in this work.




Fig. 12.— Updated version of Fig. 1, where we have added the results of the analysis from the cluster sample presented in this paper. Symbols are as in Fig. 1, except that Haffner 18 (1/2) and 19 are highlighted by cyan circles. Red stars indicated the five additional clusters studied in this work. Green, brown, red and black small crosses correspond to early type stars in the line of sight to NGC 2345, NGC 2374, Trumpler 9 and Haffner 21, respectively.

Table 1: Cluster coordinates for J2000.0

Cluster	RA	DEC	$l$	$b$
NGC 2345	07:08:18	-13:11:36	226.58	-2.31
NGC 2374	07:23:56	-13:15:48	228.41	1.02
Trumpler 9	07:55:40	-25:53:00	243.07	1.28
Haffner 21	08:01:09	-27:13:00	244.85	1.63
Haffner 20	07:56:15	-30:22:00	246.97	-0.93



Table 2. Log of observations

Date	Cluster	Filter	Exposure Time (sec)	Airmass	Seeing
2010 Dec 30	Haffner 20	<i>V</i>	2x10,60,900	1.00–1.01	0.9–1.2
		<i>B</i>	20, 3x150	1.04–1.06	0.9–1.2
		<i>U</i>	30, 300	1.06–1.07	0.9–1.2
		<i>I</i>	3x10,60,900	1.02–1.03	0.9–1.2
	Trumpler 9	<i>V</i>	2x10,60,900	1.00	0.9–1.2
		<i>B</i>	20,150,1500	1.12–1.19	0.9–1.2
		<i>U</i>	2x30,300,2000	1.00–1.03	0.9–1.2
		<i>I</i>	2x10,60,900	1.20–1.28	0.9–1.2
2010 Dec 31	Haffner 20	<i>U</i>	2x20,2000	1.25–1.27	0.8–1.2
		<i>B</i>	20, 1500	1.37–1.38	0.8–1.2
	Haffner 21	<i>U</i>	30,300,2000	1.12–1.14	0.8–1.2
		<i>B</i>	20,150,1500	1.02–1.04	0.8–1.2
		<i>V</i>	10,60,900	1.00	0.8–1.2
		<i>I</i>	2x10,60,900	1.00–1.01	0.8–1.2
2008 Feb 01	NGC 2345	<i>U</i>	5,20,100,200	1.09–1.10	1.0–1.4
		<i>B</i>	5,20,100,200	1.07–1.08	1.0–1.4
		<i>V</i>	5,10,60,120	1.08–1.09	1.0–1.4
		<i>R</i>	5,10,60,120	1.09–1.12	1.0–1.4
		<i>I</i>	5,10,100,200	1.06–1.07	1.0–1.4
	NGC 2374	<i>U</i>	5,20,100,200	1.11–1.12	1.0–1.4
		<i>B</i>	5,20,100,200	1.14–1.15	1.0–1.4

Table 2—Continued

Date	Cluster	Filter	Exposure Time (sec)	Airmass	Seeing
2008 Feb 03	NGC 2345	<i>V</i>	5,10,60,120	1.22–1.23	1.0–1.4
		<i>R</i>	5,10,60,120	1.25–1.27	1.0–1.4
		<i>I</i>	5,10,100,200	1.17–1.19	1.0–1.4
		<i>U</i>	60,600,1500	1.05–1.07	0.9–1.1
		<i>B</i>	30,600,1500	1.05–1.08	0.9–1.1
		<i>V</i>	30,600,1200	1.08–1.09	0.9–1.1
	NGC 2374	<i>R</i>	30,600,1200	1.09–1.12	0.9–1.1
		<i>I</i>	30,600,1200	1.07–1.08	0.9–1.1
		<i>U</i>	60,600,1500	1.06–1.07	0.9–1.1
		<i>B</i>	30,600,1500	1.11–1.13	0.9–1.1
		<i>V</i>	30,600,1200	1.31–1.38	0.9–1.1
		<i>R</i>	30,600,1200	1.50–1.60	0.9–1.1
		<i>I</i>	30,600,1200	1.19–1.24	0.9–1.1

Table 3. Coefficients of the transformation equations

Mag	$c_0$	$\sigma$	$c_1$	$\sigma$	$c_2$	$\sigma$
CTIO, 1-m telescope, 2008 February 01						
$U$	+1.35	0.01	−0.49	0.01	+0.015	0.016
$B$	+2.27	0.01	−0.25	0.01	−0.100	0.015
$V$	+2.36	0.01	−0.17	0.01	−0.010	0.005
$R$	+2.10	0.01	−0.10	0.01	0.048	0.005
$I$	+1.36	0.01	−0.07	0.01	−0.043	0.005
CTIO, 1-m telescope, 2008 February 03						
$U$	+1.35	0.01	−0.49	0.01	+0.014	0.020
$B$	+2.20	0.02	−0.25	0.01	−0.097	0.014
$V$	+2.32	0.01	−0.16	0.01	−0.012	0.006
$R$	+2.14	0.01	−0.09	0.01	0.050	0.010
$I$	+1.35	0.01	−0.08	0.01	−0.045	0.005
CTIO, 1-m telescope, 2010 December 30						
$U$	+0.89	0.02	−0.52	0.01	+0.019	0.008
$B$	+2.09	0.02	−0.27	0.01	−0.115	0.007

Table 3—Continued

Mag	$c_0$	$\sigma$	$c_1$	$\sigma$	$c_2$	$\sigma$
$V$	+2.30	0.01	−0.15	0.01	+0.016	0.005
$I$	+1.29	0.01	−0.06	0.01	−0.045	0.005
CTIO, 1-m telescope, 2010 December 31						
$U$	+0.90	0.01	−0.50	0.01	+0.032	0.006
$B$	+2.10	0.01	−0.25	0.01	−0.112	0.005
$V$	+2.26	0.01	−0.11	0.01	+0.030	0.005
$I$	+1.30	0.01	−0.07	0.01	−0.048	0.005

Table 4: New and earlier results for the present cluster sample

Cluster	Size	$E_{B-V}$	$R_{Sun}$	$D$	Age	Reference
	arcmin		kpc	pc	( $\times 10^6$ ) yr	
NGC 2345	5.25	0.48–1.16	1.75		60	Moffat (1974)
	3.75	0.59	3.00	6.6	63–70	This work
NGC 2374		0.175	1.20		75	Babu (1985)
					2000	Lyngå (1980)
					350	Fenkart et al. (1972)
		0.07	1.33		220– 280	This work
Trumpler 9	3.6	0.20	0.90			Pişmiş (1970)
	3.5	0.25	2.23			Vogt & Moffat (1972)
	2.5	0.20	2.90	4.2	10	This work
Haffner 20	1.1	0.55	2.40		200	FitzGerald & Moffat (1974)
	< 2.0	0.65	5.50	6.2	90– 100	This work
Haffner 21	1.1	0.20	3.30		200	FitzGerald & Moffat (1974)
	<2.0	0.21	3.50	4.0	110-120	This work