

A non Abelian effective model for ensembles of magnetic defects in 3D Yang–Mills theory

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2013 J. Phys. A: Math. Theor. 46 335401

(<http://iopscience.iop.org/1751-8121/46/33/335401>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 200.0.233.52

This content was downloaded on 25/04/2014 at 12:20

Please note that [terms and conditions apply](#).

A non Abelian effective model for ensembles of magnetic defects in 3D Yang–Mills theory

C D Fosco¹ and L E Oxman²

¹ Centro Atómico Bariloche and Instituto Balseiro, Comisión Nacional de Energía Atómica, R8402AGP Bariloche, Argentina

² Instituto de Física, Universidade Federal Fluminense, Campus da Praia Vermelha, Niterói, 24210-340, RJ, Brazil

E-mail: oxman@if.uff.br

Received 12 March 2013, in final form 5 July 2013

Published 29 July 2013

Online at stacks.iop.org/JPhysA/46/335401

Abstract

We construct a non Abelian model for $SU(2)$ Yang–Mills theory in an Euclidean three-dimensional spacetime and study its different phases. The model contains a center vortex sector coupled to a dual effective field encoding information about how the vortices are paired in the ensemble. The possible phases in the parameter space are interpreted in terms of the proliferation of either closed center vortices or closed chains, where the endpoints of open vortices are attached in pairs to monopole-like defects.

PACS numbers: 11.15.–q, 11.15.Tk

1. Introduction

One of the most promising physical explanations for quark confinement is based on the role played by magnetic objects when analyzing the vacuum structure of $SU(N)$ Yang–Mills theories. These scenarios have been analyzed in the lattice, relying only on monopoles [1–4], only on $Z(N)$ center vortices [5–9], or on chains. In the latter case, the ensemble includes configurations where N center vortices are attached to a monopole [10–12]. In the continuum, the difficulties encountered when trying to explore these ideas from first principles, by following controlled steps on the partition function of pure Yang–Mills theories, are daunting. Indeed, three related questions come to mind, the first being how quantum effects can generate a dimensionful scale to characterize these ensembles. Then, one should know how to define a path integral measure including those magnetic configurations. Finally, one should compute the corresponding ensemble integration. These stumbling blocks are precisely what make semiheuristic effective models interesting. Using as a phenomenological input the possibility that magnetic defects could play a relevant role, and based on general principles of symmetry, different models can be proposed to be contrasted with experiments or the lattice. In this direction, effective field models based on the monopole component, to

describe confinement in 4D, have been extensively discussed (see [13–18]). More recently, a non Abelian model containing Higgs fields in the adjoint describing confining strings and junctions has been presented [19]. While the string confines a quark–antiquark pair to form normal mesons, junctions confine quarks in a color nonsinglet state together with a valence gluon, to form an overall neutral object, identified with a hybrid meson. These states are allowed by QCD, and current experimental collaborations are aimed at identifying them [20–22].

In 4D, monopoles are stringlike so that ensembles only formed by them can be naturally related to effective field models. This comes about as an ensemble of stringlike objects can be thought of as a sum over different numbers of particle worldlines, which corresponds to a second quantized field theory [23–28]. On the other hand, the inclusion of center vortices, which in 4D are two-dimensional objects, poses great difficulties. They are naturally associated with random surface models [29] rather than with an effective field description. To gain insight into the possible mechanisms based on magnetic objects and explore their possible relation with effective models, it would be interesting to take into account the role played by all the different types of magnetic defects. Models defined in 3D are unique in this sense since here vortices are stringlike, thus opening the possibility of discussing correlated monopoles and center vortices by means of a field theory. In this work, our main interest will be constructing and analyzing an $SU(2)$ model whose possible phases correspond to the different ensembles.

The vortex model introduced by t' Hooft [30] is a low energy effective field theory that successfully describes some aspects of the confinement mechanism in 3D $SU(N)$ Yang–Mills theories. It is defined in terms of a dynamical variable which is a complex scalar field V equipped with a discrete $Z(N)$ symmetry, whose Lagrangian is

$$\mathcal{L} = \partial_\mu \bar{V} \partial_\mu V + \mu^2 \bar{V} V + \alpha (\bar{V} V)^2 + \beta (V^N + \bar{V}^N). \quad (1)$$

Its form is based on a study of the possible nontrivial vortex correlation functions in the original theory. In particular, the confining phase is described as one where the discrete $Z(N)$ symmetry is spontaneously broken ($Z(N)$ SSB), due to the presence of a vortex condensate. In [31], we derived this type of model by considering an ensemble of chains, where the vortex endpoints are attached in pairs to monopole-like defects, and following recent polymer techniques to compute the vortex end-to-end probability. The obtained model is in fact a generalization of equation (1) where ∂_μ is substituted by the covariant derivative D_μ , which depends on a dual vector field λ_μ describing the off-diagonal sector. The dynamics is completed with a Proca action term for λ_μ (see also [32, 33]). 3D effective field models can also be obtained in scenarios based just on the monopole (instanton) component. In this case, the assumption of Abelian dominance and the associated monopole ensemble is encoded in a sine-Gordon type model for a scalar dual field (see [34, 35] and references therein), as occurs in the case of compact QED(3), discussed by Polyakov in [36].

In spite of the fact that the initial theory is a non Abelian one, these effective models are *Abelian*, that is, additional information regarding this transition is already incorporated, while it would be desirable to see it appearing as a phase transition in a previous non Abelian model.

In this paper, we construct a non Abelian effective model which encompasses a description of interacting effective gluons and center vortices. Depending on the choice of parameters, the vortices can be found in different states, including a phase where they are closed, and a phase where their endpoints become paired to form closed chains. To that aim, we use a parametrization that treats the different color components in a symmetric way [37], and describes correlated monopoles and center vortices as defects of a local color frame \hat{n}_a , $a = 1, 2, 3$. This parametrization is based on the usual manner to introduce thin center vortices in Yang–Mills theories [29, 38], and corresponds to a symmetric form of the Cho–Faddeev–Niemi (CFN) decomposition [39–41], used to represent monopoles as defects

of the third component $\hat{n} = \hat{n}_3$. For a description of center vortices in the CFN framework, and related consequences in the continuum, see [32, 33].

Since center vortices can be joined in pairs to pointlike monopoles, the natural non Abelian field content of the model is given by a scalar field with one (magnetic) color index, generalizing the vortex field V in equation (1), and a scalar field with two color indices, generalizing the scalar dual field in scenarios only involving the monopole component³. The order parameters present in the effective model bear a relation to the nature of the phase transition one may describe. In this respect, an interesting point has been raised [35] about whether the confining/deconfining phase transition is of the Kosterlitz–Thouless or Ising model type. The former involves the monopole sector: at high temperatures, the instanton magnetic flux is distributed along the two spatial directions, thus leading to effective logarithmic interactions. Then, because of dimensional reduction, instantons and anti-instantons tend to be suppressed by forming pairs. On the other hand, the latter naturally involves the vortex degrees of freedom, as they are the objects where the discrete symmetry transformations act.

In the model we construct and study below, since it does contain order parameters for both the center vortices and the distribution of monopole-like defects they can concatenate to form chains, an interesting framework to discuss the competition between different phases shall emerge, originating a phase diagram with a rich structure. This paper is organized as follows: in section 2, we deal with the topological defects included in the model, in particular, their parametrization, and the functional and ensemble integration over them. In section 3, based on the results of the previous section, and after discussing the possible symmetries, we construct an action for the effective model in terms of the fields introduced therein. Finally, in section 4, we present a study of the phase structure of the model, based on some assumptions about the relative strength of its different terms.

2. Non Abelian defects in YM theories

We shall begin from the $SU(2)$ Yang–Mills action, S_{YM} , which may be written as follows:

$$S_{YM} = \frac{1}{4} \int d^3x \vec{F}_{\mu\nu} \cdot \vec{F}_{\mu\nu}, \quad (2)$$

where $\vec{F}_{\mu\nu}$ is the non Abelian field-strength tensor. We use an arrow on top of any object to denote the 3-component vector formed by its components on the $su(2)$ Lie algebra basis, whose elements are the (Hermitian) generators $(T^a)_{a=1}^3$. In the concrete case we are considering, they can be conveniently realized as $T^a = \tau^a/2$, where τ^a denotes a Pauli matrix; they satisfy $[T^a, T^b] = i\epsilon^{abc} T^c$, and $\text{tr}(T^a T^b) = \frac{1}{2}\delta^{ab}$.

Thus, with this notation, we may write down the defining equation for $\vec{F}_{\mu\nu}$, as follows:

$$\vec{F}_{\mu\nu} \cdot \vec{T} = \frac{i}{g} [D_\mu, D_\nu], \quad D_\mu = \partial_\mu - ig\vec{A}_\mu \cdot \vec{T}, \quad (3)$$

where D_μ has been used to denote the covariant derivative operator, when acting on fields in the fundamental representation.

It goes without saying that a ‘canonical color basis’ $(\hat{e}^a)_{a=1}^3$ (with color components $\hat{e}_b^a = \delta_{ab}$) can be introduced, so that $\vec{A}_\mu = \vec{A}_\mu^a \hat{e}^a$. This seemingly trivial remark is made in order to highlight the next step; namely, that one could have used a different basis. Indeed, in order to describe configurations with defects, in a symmetric way that admits its extension to finite size objects, we introduce a space-dependent color basis $(\hat{n}_a)_{a=1}^3$, related to the original one by: $S T^a S^{-1} = \hat{n}_a \cdot \vec{T}$ (with $S \in SU(2)$). Thus the new basis is connected to the canonical

³ Interestingly, isospin two-order parameters appear in models for liquid crystals [42].

one by an orthogonal space-dependent matrix $R(S)$: $\hat{n}_a = R(S)\hat{e}_a$, which belongs to the adjoint representation. In this representation, the corresponding infinitesimal generators shall be denoted by M^a , with $(M^a)^{bc} \equiv -i\epsilon^{abc}$. They satisfy $[M^a, M^b] = i\epsilon^{abc}M^c$, $\text{tr}(M^a M^b) = 2\delta^{ab}$. At this point, and equipped with the local basis, we consider the parametrization of the gauge field [37]:

$$\vec{A}_\mu = (\mathcal{A}_\mu^a - C_\mu^a)\hat{n}_a, \quad (4)$$

where the frame-dependent fields

$$C_\mu^a = -\frac{1}{2g}\epsilon^{abc}\hat{n}_b \cdot \partial_\mu \hat{n}_c, \quad (5)$$

satisfy the properties:

$$\hat{n}_b \cdot \partial_\mu \hat{n}_c = -g\epsilon^{abc}C_\mu^a, \quad C_\mu^a M^a = \frac{i}{g}R^{-1}\partial_\mu R. \quad (6)$$

This corresponds to a symmetric form of the (CFN) decomposition [39–41]. In terms of the parametrization (4) of the gauge field, we note that the field-strength tensor becomes:

$$\vec{F}_{\mu\nu} = G_{\mu\nu}^a \hat{n}_a, \quad G_{\mu\nu}^a = \mathcal{F}_{\mu\nu}^a(\mathcal{A}) - \mathcal{F}_{\mu\nu}^a(C), \quad (7)$$

with $\mathcal{F}_{\mu\nu}^a(\mathcal{A}) \equiv \partial_\mu \mathcal{A}_\nu^a - \partial_\nu \mathcal{A}_\mu^a + g\epsilon^{abc}\mathcal{A}_\mu^b \mathcal{A}_\nu^c$ (and an analogous expression for $\mathcal{F}_{\mu\nu}^a(C)$), while the Yang–Mills action is given by:

$$S_{\text{YM}} = \int d^3x \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a. \quad (8)$$

Regarding the color components of the frame-dependent tensor $\mathcal{F}_{\mu\nu}^a(C)$, they can also be obtained by commuting covariant derivatives in the adjoint representation:

$$\mathcal{F}_{\mu\nu}^a(C)M^a = \frac{i}{g}[D_\mu, D_\nu], \quad D_\mu \equiv \partial_\mu - igC_\mu^a M^a, \quad (9)$$

so that the second equality in (6) implies the alternative expression for $\mathcal{F}_{\mu\nu}^a(C)$:

$$\mathcal{F}_{\mu\nu}^a(C) = \frac{i}{2g}\text{tr}(M^a R^{-1}[\partial_\mu, \partial_\nu]R). \quad (10)$$

This equation highlights the meaning of $\mathcal{F}_{\mu\nu}^a(C)$, by showing that it can only be different from zero where R has defects; these, are characterized here by the noncommutativity of the mixed partial derivatives. These defects are zero measure objects; in other words, the partial derivatives will fail to commute on zero measure regions. Being this an effective theory, this should be interpreted as the assumption that the model describes physics at distances much larger than the size of the defects.

Of course, there are infinitely many different local frames, and corresponding fields \mathcal{A}_μ^a , that can be used to describe one and the same gauge field configuration, A_μ^a . One can use that large amount of freedom in order to split it into its ‘regular’ and ‘singular’ parts. Indeed, the \mathcal{A}_μ^a measure will represent topologically trivial fluctuations. The singular configurations, described by the frames, will have a measure representing an ensemble integration over defects.

In [37], one of us has shown that the configuration in (4) is tantamount to the usual way [29, 38] to introduce thin center vortices on top of a trivial field configuration $\mathcal{A}_\mu^a \hat{e}_a$, namely,

$$\vec{A}_\mu \cdot \vec{T} = S\vec{\mathcal{A}}_\mu \cdot \vec{T}S^{-1} + \frac{i}{g}S\partial_\mu S^{-1} - \vec{I}_\mu(S) \cdot \vec{T}. \quad (11)$$

Because of the presence of the last term, this is not just a gauge transformation of the topologically trivial gauge field. Indeed, the $\vec{I}_\mu(S)$ field corresponds to the so called ideal center vortex, and is localized on a hypersurface Σ . This is the region which, when traversed,

makes S change by a center element. $\vec{I}_\mu(S)$ is designed to cancel the contribution in the second term originated from the discontinuity of S^{-1} , only leaving the effect of the border of Σ where the thin center vortices are located. That is, we can write

$$\frac{i}{g} S \partial_\mu S^{-1} |_\Sigma = \vec{I}_\mu(S) \cdot \vec{T}, \quad (12)$$

where the subscript in the left-hand side amounts to just keeping in the calculation the term originated from the derivative of the discontinuity in S^{-1} . Considering two regular mappings U, \tilde{U} , the ideal vortex satisfies

$$\vec{I}_\mu(US\tilde{U}^{-1}) \cdot \vec{T} = U\vec{I}_\mu(S) \cdot \vec{T}U^{-1}, \quad (13)$$

obtained from $\partial_\mu(\tilde{U}^{-1}S^{-1}U^{-1})|_\Sigma = \tilde{U}^{-1}\partial_\mu S^{-1}|_\Sigma U^{-1}$, as the term localized on Σ is only generated when ∂_μ acts on S^{-1} . The gauge field $\vec{A}_\mu = \vec{A}_\mu(\vec{\mathcal{A}}, S)$ in (11) enjoys the following properties,

$$\vec{A}^U(\vec{\mathcal{A}}, S) = \vec{A}(\vec{\mathcal{A}}, US), \quad \vec{A}(\vec{\mathcal{A}}, S) = \vec{A}(\vec{\mathcal{A}}^{\tilde{U}}, S\tilde{U}^{-1}). \quad (14)$$

Then, in terms of the $\vec{\mathcal{A}}, S$ variables we have a double redundancy, the usual one associated with invariance of the Yang–Mills action under gauge transformations, $\vec{A}_\mu \cdot \vec{T} = U\vec{A}_\mu \cdot \vec{T}U^{-1} + \frac{i}{g}U\partial_\mu U^{-1}$, represented by $S \rightarrow US$, and other originated from the different ways to express the same vector field, combining the transformation $\vec{A}_\mu \cdot \vec{T} = \tilde{U}\vec{A}_\mu \cdot \vec{T}\tilde{U}^{-1} + \frac{i}{g}\tilde{U}\partial_\mu \tilde{U}^{-1}$, together with a right multiplication of S .

At this point, we would like to emphasize that a nonperturbative definition of the path integral in Yang–Mills theory is still lacking. This comes about as a gauge fixing procedure generally leads to Gribov copies [43] in that regime, so that it is difficult to define an appropriate object where each physical situation is counted only once. The restriction to the modular region has been usually implemented by means of the Zwanziger action [44]. In this framework, in the infrared regime, the path integral has been shown to be dominated by configurations near the Gribov horizon. On the other hand, as is well known, configurations containing magnetic objects proliferate at the horizon [45–47]. From this perspective, it is natural to fix the redundancy by introducing the identity $1 = \Delta_{\text{FP}}[\mathcal{A}] \int [d\tilde{U}] \delta[f(\mathcal{A}^{\tilde{U}})]$, in the perturbative sector where the Faddeev–Popov procedure is well defined. In addition, as the S sector parametrizes correlated monopoles and center vortices, it represents configurations at the horizon, relevant to describe the large distance physics. Giving a configuration S , gauge fixing amounts to choose a representative of the orbit US . Any condition imposed on $\vec{A}(\vec{\mathcal{A}}, S)$ will be invariant under the \tilde{U} -transformations in equation (14). This is also the case for conditions depending on $\vec{I}(S)$, as it is invariant under right multiplication (cf equation (13)).

In this paper, we shall not attempt to derive a precise construction of the integration measure. Rather, having the previous remarks, notation and conventions in mind, we argue that it is quite natural to propose the following path integral

$$Z_{\text{YM}} = \int [d\mathcal{A}][dS] \Delta_{\text{FP}}[\mathcal{A}] \delta[f(\mathcal{A})] e^{-S_{\text{YM}}[\mathcal{A}]}, \quad (15)$$

where dS represents the ensemble integration over monopoles and thin center vortices, that is supposed to include its own appropriate gauge fixing condition. Note that, in the trivial sector, where $S = S_r$ is regular, we have $\vec{A}(\vec{\mathcal{A}}, S_r) = \vec{A}^{S_r}$ and the associated contribution to (15) is the usual, perturbative one. Only in that sector $\vec{A}(\vec{\mathcal{A}}^{\tilde{U}}, S) = \vec{A}(\vec{\mathcal{A}}, S\tilde{U})$ may be identified with a gauge transformation.

As a final step, and as a guide to the construction of the effective model, we rewrite the partition function in the equivalent form:

$$Z_{\text{YM}} = \int [d\mathcal{A}][dS][d\lambda] \Delta_{\text{FP}}[\mathcal{A}] \delta[f(\mathcal{A})] e^{-\int d^3x [\frac{1}{2}\lambda_\mu^a \lambda_\mu^a + i\lambda_\mu^a (\mathcal{F}_\mu^a(\mathcal{A}) - \mathcal{F}_\mu^a(C))]}, \quad (16)$$

where $\mathcal{F}_\mu^a = \frac{1}{2}\epsilon_{\mu\nu\rho}\mathcal{F}_{\nu\rho}^a$, and we have introduced a color-valued auxiliary field λ_μ^a to deal with a first-order version of (8).

3. The effective theory

Let us now derive a non Abelian effective field theory for the sector of defects. The derivation will become possible by relying on the symmetries exhibited by the ensemble integration. This effective theory shall contain mass parameters, which we assume are originated from those present in a (phenomenological) ansatz for the action of the defects. In this regard, we note that up to now we have considered thin center vortices, parametrized as in (4). However, lattice simulations [5] point to the idea that they become thick objects, characterized by some finite radius of the order of 1 fm. Moreover, as discussed in [37], the stable objects in the continuum could in fact correspond to some deformation of the thin objects given in (4), where the ‘thin’ quantities C_μ^a are replaced by some smooth finite radius profiles \mathcal{C}_μ^a . If this is assumed to be the case, rather than equations (7) and (8), the Yang–Mills action would have the form

$$S_{\text{YM}} = \int d^3x \frac{1}{4} (\mathcal{F}_{\mu\nu}^a(\mathcal{A}) - \mathcal{F}_{\mu\nu}^a(\mathcal{C}))^2 + \mathcal{R}, \quad (17)$$

where \mathcal{R} vanishes for thin center vortices. Note that the first term can be linearized, as we did before, by introducing the fields λ_μ^a . Besides, at large distances, approximating \mathcal{C}_μ^a by C_μ^a , this term shall originate the terms appearing in the exponent of equation (16), when the center vortices were considered to be thin. On the other hand, the second term (\mathcal{R}), will be concentrated on the center vortices and at large distances will produce instead an additional action S_d for the defects. Therefore, in the general case, the ensemble integration must be written in the form

$$e^{-S_{v,m}[\lambda]} = \int [dS] e^{-S_d + i \int d^3x \lambda_\mu^a \mathcal{F}_\mu^a(C)}. \quad (18)$$

The second term in the exponent above has a local $SO(3)$ symmetry under right multiplication: $S \rightarrow S\tilde{U}$, changing the color basis from $\hat{n}_a \cdot \vec{T} = ST^a S^{-1}$ to $\hat{n}'_a \cdot \vec{T} = S\tilde{U}T^a\tilde{U}^{-1}S^{-1}$, that is, $\hat{n}'_a = R(S)R(\tilde{U})\hat{e}_a$. Note that, using (10), and that $R(\tilde{U})$ contains no defects, we have

$$\mathcal{F}_{\mu\nu}^a(C') = \frac{i}{2g} \text{tr}(R(\tilde{U})M^a R^{-1}(\tilde{U})R^{-1}(S)[\partial_\mu, \partial_\nu]R(S)). \quad (19)$$

In other words, a regular local rotation of λ_μ^a can be translated to a regular local transformation of S . Then, if S_d were nullified, that is, if we were dealing with strictly thin center vortices, $S_{v,m}[\lambda]$ would be invariant under local $SO(3)$ rotations, as the transformation $S \rightarrow S\tilde{U}$ could be absorbed by the integration measure dS . In this regard, we would like to underline that this measure is to be accompanied by an appropriate gauge fixing condition that is invariant under right multiplication (see the discussion at the end of the previous section). However, in $S_{v,m}[\lambda]$, that symmetry will be broken to a global one because of the thick character expected for center vortices. To have a simple picture about this statement we note that an action for thick center vortices will typically contain a Nambu–Goto term plus other terms describing the center vortex rigidity [29, 48]. These pieces can be generated, for instance, from a large distance approximation of the more symmetric term (in color space)

$$\int d^3x d^3y \mathcal{F}_{\mu\nu}^a(C)|_x G_M(x-y) \mathcal{F}_{\mu\nu}^a(C)|_y,$$

where G_M is a kernel localized on a scale $1/M$. Now, as we have seen in equation (19), the field strength $\mathcal{F}_{\mu\nu}^a(C)$ will rotate under local \tilde{U} transformations. Therefore, as for any finite M the integrand above depends on the field strength at different spacetime points, it will change under the local transformations, only leaving a symmetry under the global ones.

Based on purely geometrical/mathematical grounds, the possible kinds of defects can be straightforwardly classified as follows.

- (i) Closed center vortices.
- (ii) Monopoles and antimonopoles, joined by center vortices (each pointlike object is joined by a pair of center vortices).
- (iii) A particular limit of (ii): a coincident pair of center vortices, which should correspond to an unobservable Dirac string.

To proceed, let us consider a type (ii) configuration (correlated monopoles and center vortices). To that end we recall that, in [32, 31], we have considered a particular case of that situation, namely, when the center vortex color points along the (locally) diagonal direction \hat{n}_3 . In that case, the effective field describing these objects corresponds to a complex vortex field V . In particular, in [31], we have shown how the ensemble integration over open center vortices, whose endpoints are joined in pairs to form closed chains, leads to an Abelian $Z(2)$ effective theory that can be written in terms of V , thus making contact between the initial representation and the final effective field theory. With this aim, we applied recent polymer techniques [49, 50] to deal with the end-to-end probability associated with center vortices interacting with a general vector field λ_μ , and a scalar field needed to represent vortex–vortex interactions. However, it is far from straightforward to extend this type of derivation to the non Abelian context. Therefore, in our case, we will propose a model relying on the symmetries displayed by the initial representation, that strongly constraints the possible associated effective theories.

In our case, the candidate for a vortex field has to be a real 3-component field ϕ^a ($a = 1, 2, 3$), because of the global $SO(3)$ symmetry of the action $S[\lambda]$. We shall also introduce an isospin-2 field Q , where Q is a traceless symmetric 3×3 real matrix, encoding information about how the monopole-like defects that center vortices can concatenate are distributed. We may then consider in the effective theory, an invariant term V_I that couples the monopole and vortex sectors:

$$V_I = \zeta \phi^T Q \phi, \quad \zeta \equiv \text{constant} \tag{20}$$

which is invariant under the local $SO(3)$ transformations:

$$\phi(x) \rightarrow R(x)\phi(x), \quad Q(x) \rightarrow R(x)Q(x)R^T(x). \tag{21}$$

There are also invariant terms involving just either the vortex or the monopole field. Regarding the former, we may include a ‘potential’ term V_ϕ , with the general structure:

$$V_\phi = \frac{\mu^2}{2} \phi^T \phi + \frac{\lambda}{4} (\phi^T \phi)^2, \tag{22}$$

where μ and λ are arbitrary constants. On the other hand, for the case of the monopole field, we recall that an order parameter Q , with a similar structure, is well known in the context of liquid crystals. Thus, we expect the relevant terms in the effective theory to be of the same kind, namely, we may include a potential V_Q [42]:

$$V_Q = \frac{A}{2} \delta + \frac{B}{3} \Delta + \frac{C}{4} \delta^2 + \frac{D}{5} \delta \Delta + \frac{E}{6} \Delta^2 + F \delta^3, \tag{23}$$

where A, \dots, F are constants, and we have introduced two independent $SO(3)$ invariants⁴ that can be built in terms of Q :

$$\delta = \text{Tr } Q^2, \quad \Delta = \text{Tr } Q^3. \tag{24}$$

⁴ Being a traceless real symmetric matrix, the invariant content of Q can be generated by two real invariants. For example, two of its eigenvalues.

Thus, the three terms V_ϕ , V_Q and V_I have the local symmetry (21). This local symmetry will be broken to its global counterpart by the kinetic terms; however, these terms shall be constructed in such a way that they are compatible with a local discrete gauge symmetry. This symmetry must be present, at least in a phase where the vacuum is symmetric (no spontaneous symmetry breaking).

In this regard, the field-strength tensor $\mathcal{F}_\mu^a(C)$ can be written as

$$\begin{aligned}\mathcal{F}_\mu^a(C) &= \frac{1}{2}\epsilon_{\mu\nu\rho}\mathcal{F}_{\nu\rho}^a(C) \\ &= \epsilon_{\mu\nu\rho}\partial_\nu C_\rho^a + \frac{g}{2}\epsilon_{\mu\nu\rho}\epsilon^{abd}C_\nu^b C_\rho^d,\end{aligned}\quad (25)$$

where

$$\begin{aligned}C_\mu^1 &= -\frac{1}{g}\hat{n}_2 \cdot \partial_\nu \hat{n}_3 \\ C_\mu^2 &= -\frac{1}{g}\hat{n}_3 \cdot \partial_\nu \hat{n}_1 \\ C_\mu^3 &= -\frac{1}{g}\hat{n}_1 \cdot \partial_\nu \hat{n}_2.\end{aligned}\quad (26)$$

We will show that $\mathcal{F}_\mu^a(C)$ can be rewritten as

$$\mathcal{F}_\mu^a(C) = \tilde{h}_\mu^a - h_\mu^a, \quad (27)$$

$$\tilde{h}_\mu^a = \epsilon_{\mu\nu\rho}\partial_\nu C_\rho^a, \quad h_\mu^a = -\frac{1}{2g}\epsilon_{\mu\nu\rho}\hat{n}_a \cdot (\partial_\nu \hat{n}_a \times \partial_\rho \hat{n}_a), \quad (28)$$

where, in the second tensor, no summation over a is understood.

Let us take, for example, the third component of the field-strength tensor

$$\mathcal{F}_\mu^3 = \tilde{h}_\mu^3 - h_\mu^3, \quad (29)$$

$$\tilde{h}_\mu^3 = \epsilon_{\mu\nu\rho}\partial_\nu C_\rho^3, \quad h_\mu^3 = -g\epsilon_{\mu\nu\rho}C_\nu^1 C_\rho^2. \quad (30)$$

In order to show that

$$h_\mu^3 = -\frac{1}{2g}\epsilon_{\mu\nu\rho}\hat{n}^3 \cdot (\partial_\nu \hat{n}^3 \times \partial_\rho \hat{n}^3), \quad (31)$$

we can simply note that

$$\begin{aligned}\partial_\nu \hat{n}_3 &= (\hat{n}_1 \cdot \partial_\nu \hat{n}_3)\hat{n}_1 + (\hat{n}_2 \cdot \partial_\nu \hat{n}_3)\hat{n}_2 + (\hat{n}_3 \cdot \partial_\nu \hat{n}_3)\hat{n}_3 \\ &= (\hat{n}_1 \cdot \partial_\nu \hat{n}_3)\hat{n}_1 + (\hat{n}_2 \cdot \partial_\nu \hat{n}_3)\hat{n}_2 \\ &= g(C_\nu^2 \hat{n}_1 - C_\nu^1 \hat{n}_2).\end{aligned}\quad (32)$$

Then, replacing in the second member of (31), and using $\hat{n}_1 \times \hat{n}_2 = \hat{n}_3$, etc., it is straightforward to make contact with (30).

The important point is that (27) and (28) imply that for a fixed monopole background correlated with center vortices, the integral of each component over a closed surface $\partial\vartheta$ (given as the border of a three-volume ϑ)

$$\oint_{\partial\vartheta} dS_\mu \mathcal{F}_\mu^a(C) = \oint_{\partial\vartheta} dS_\mu (\tilde{h}_\mu^a - h_\mu^a) = \frac{1}{2g} \oint_{\partial\vartheta} dS_\mu \epsilon_{\mu\nu\rho} \hat{n}_a \cdot (\partial_\nu \hat{n}_a \times \partial_\rho \hat{n}_a), \quad (33)$$

gives the Π_2 topological charge for the mapping $\partial\vartheta \rightarrow \hat{n}_a$. More precisely,

$$\oint_{\partial\vartheta} dS_\mu \mathcal{F}_\mu^a(C) = \frac{4\pi}{g}(n_+(\vartheta) - n_-(\vartheta)), \quad (34)$$

where $n_+(\vartheta)$ ($n_-(\vartheta)$) is the number of monopole (antimonopole) defects inside ϑ , for the component \hat{n}_a .

It may appear that the previous expression sets a preferred direction in the color space. This impression can be dispelled by considering the effect that a space independent change of color basis has on the expression (34). To that end, we consider a new basis $(\hat{n}'_a)_{a=1}^3$, related to the original one by a (constant) matrix R' , namely: $\hat{n}'_a = R' \hat{n}_a$, $a = 1, 2, 3$. In components, and using an obvious notation, the last relation means:

$$(\hat{n}'_a)_b = (R')_{bc} (\hat{n}_a)_c. \quad (35)$$

Thus, we see that:

$$\begin{aligned} \oint_{\partial\theta} dS_\mu \epsilon_{\mu\nu\rho} \hat{n}'_a \cdot (\partial_\nu \hat{n}'_a \times \partial_\rho \hat{n}'_a) \\ = \oint_{\partial\theta} dS_\mu \epsilon_{\mu\nu\rho} \epsilon_{b_1 b_2 b_3} (\hat{n}'_a)_{b_1} \partial_\nu (\hat{n}'_a)_{b_2} \partial_\rho (\hat{n}'_a)_{b_3} \\ = \oint_{\partial\theta} dS_\mu \epsilon_{\mu\nu\rho} \epsilon_{b_1 b_2 b_3} (R')_{b_1 c_1} (R')_{b_2 c_2} (R')_{b_3 c_3} (\hat{n}_a)_{c_1} \partial_\nu (\hat{n}_a)_{c_2} \partial_\rho (\hat{n}_a)_{c_3} \\ = (\det R') \oint_{\partial\theta} dS_\mu \epsilon_{\mu\nu\rho} \hat{n}_a \cdot (\partial_\nu \hat{n}_a \times \partial_\rho \hat{n}_a), \end{aligned} \quad (36)$$

where we have used the property:

$$\epsilon_{b_1 b_2 b_3} (R')_{b_1 c_1} (R')_{b_2 c_2} (R')_{b_3 c_3} = (\det R') \epsilon_{c_1 c_2 c_3}. \quad (37)$$

A similar relation holds if one changes the original canonical basis by a constant rotation matrix. What this proves is that it is possible to generate a monopole charge along any color direction as long as one needs how to do that for, say, the third one.

Thus, coming back to the discussion on the possible form of the kinetic terms, they must be—at least in the symmetric phase—compatible with the local discrete gauge symmetry:

$$\lambda_\mu^a \rightarrow \lambda_\mu^a + \partial_\mu \omega^a, \quad (38)$$

where ω^a is a *discontinuous* function taking values $\pm \frac{\xi}{2}$ inside a three-volume ϑ^a , and 0 outside. Note also that in a phase only containing closed center vortices, a larger symmetry,

$$\lambda_\mu^a \rightarrow \lambda_\mu^a + \partial_\mu \varphi^a, \quad (39)$$

for any smooth φ^a , is expected: in this case, the absence of monopoles would imply $\partial_\mu \mathcal{F}_\mu^a(C) = 0$.

Then, the kinetic terms must have the global $SO(3)$ symmetry plus the local Abelian one. The simplest choice, which, in the effective theory approach spirit we shall consistently adopt, is to minimally couple ϕ and Q to λ_μ^a (note that these couplings do not have the local $\vec{\lambda}(x) \rightarrow R(x)\vec{\lambda}(x)$ symmetry). Thus, the structure of the kinetic term K is as follows:

$$K = K_\phi + K_Q, \quad (40)$$

where:

$$\begin{aligned} K_\phi &= \frac{1}{2} (\nabla_\mu \phi)^a (\nabla_\mu \phi)^a \\ K_Q &= \frac{1}{2} (\nabla_\mu Q)^{ab} (\nabla_\mu Q)^{ab}, \end{aligned} \quad (41)$$

where ∇_μ denotes the covariant derivative operator (consistent with the symmetries mentioned above), which shall adopt a different expression when acting on each one of the fields. Explicitly:

$$\begin{aligned} (\nabla_\mu \phi)^a &= \partial_\mu \phi^a - ig_\phi \lambda_\mu^b \epsilon^{abc} \phi^c \\ (\nabla_\mu Q)^{ab} &= \partial_\mu Q^{ab} - ig_Q \lambda_\mu^c \epsilon^{acd} Q^{db} + ig_Q Q^{ad} \epsilon^{dcb} \lambda_\mu^c, \end{aligned} \quad (42)$$

where g_ϕ and g_Q are constants.

The global $SO(3)$ symmetry is evident, while by imposing $g_\phi = g_Q$, the effective action will display a non Abelian gauge symmetry, and the different phases for the ensemble of monopoles and center vortices will arise as different possible vacua when the system undergoes SSB. Note also that as the field ϕ represents center vortices that in the case of Abelian configurations possess a magnetic charge $2\pi/g$, the natural choice is $g_\phi = 2\pi/g$, which also matches the correct dimensions in equation (42) as $[\lambda] = 3/2$, $[g] = 1/2$.

Then, joining the different pieces and taking into account equation (16), the following model, encoding a general ensemble of magnetic defects, can be proposed

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{v,m} + \frac{1}{2}\lambda_\mu^a \lambda_\mu^a + i\lambda_\mu^a \mathcal{F}_\mu^a(\mathcal{A}), \quad (43)$$

$$\mathcal{L}_{v,m} = K_Q + K_\phi + V_\phi + V_Q + V_I. \quad (44)$$

We would like to underline that according to the discussion at the end of section 2, and beginning of section 3, the symmetry displayed by the second and third terms in equation (43), namely a transformation $\vec{A}^{\vec{U}}$, accompanied by the local $SO(3)$ rotation of λ_μ^a , is not the gauge symmetry that operates on \vec{A}_μ . Therefore, the noninvariance of $\mathcal{L}_{v,m}$ under local $SO(3)$ rotations of λ_μ^a is not an explicit breaking of the gauge symmetry in our effective model. Only in the trivial sector $\vec{A}^{\vec{U}}$ may be associated with a gauge transformation, in other words, our model refers to the interaction of effective fields, parametrizing a general ensemble, with effective gluons represented by \vec{A}_μ .

4. Phase structure

In order to analyze the possible phases of the model, it is necessary to study all the possible scenarios regarding both the ϕ and Q dependent potentials, V_ϕ and V_Q , as well as the interaction V_I . This will yield information about the possible translation invariant configurations that will determine the properties of each phase. Non translation invariant configurations, on the other hand, are important to understand the mechanism driving the phase transitions between them. Of course, that will require the inclusion of the derivative terms into the game.

Thus we consider the minima of

$$V_T = V_\phi + V_Q + V_I. \quad (45)$$

This analysis is greatly simplified if we note that Q can always be diagonalized by a similarity transformation $Q = R^T D R$, with

$$D = \begin{pmatrix} -\frac{q}{2} - \frac{\eta}{2} & 0 & 0 \\ 0 & -\frac{q}{2} + \frac{\eta}{2} & 0 \\ 0 & 0 & q \end{pmatrix}. \quad (46)$$

Defining $R\phi = \psi$, the potential V_T adopts the form

$$V_T = \frac{A}{2}\delta + \frac{B}{3}\Delta + \frac{C}{4}\delta^2 + \frac{D}{5}\delta\Delta + \frac{E}{6}\Delta^2 + \frac{\mu^2}{2}\psi^T\psi + \frac{\lambda}{4}(\psi^T\psi)^2 + \zeta\psi^T D\psi, \quad (47)$$

$$\delta = (3q^2 + \eta^2)/2$$

$$\Delta = 3q(q^2 - \eta^2)/4$$

$$\psi^T D\psi = -\frac{q}{2}(\psi_1^2 + \psi_2^2) + \frac{\eta}{2}(\psi_2^2 - \psi_1^2) + q\psi_3^2. \quad (48)$$

Here, the term δ^3 that was present in equation (23) has been discarded, as it does not modify the qualitative structure of the minima [42].




Figure 1. A-B phase diagram for the monopole sector, when $CE > 6D^2/25$, $D < 0$. The isotropic, N^+ or N^- , and N_b monopole phases trigger corresponding vortex models displaying non Abelian, $U(1)$ and $Z(2)$ symmetry, respectively. In particular, the N_b phase may lead to a confining vortex state, where $Z(2)$ is spontaneously broken. Solid lines represent first-order transitions while dashed lines are second order.

Now, to find the minima of the potential, we will suppose that the chain of spontaneous breaking of the symmetries is dominated by the monopole sector. Concretely, this approximation amounts to finding the minima of V_Q , and using the configurations q_0, η_0 that Q adopts in those minima as a fixed background where we look for the vortex field configuration that minimizes the remaining potential. Then, the whole space of minima is generated by means of R -rotations of the former.

The minima of V_Q are determined by:

$$\partial_q V_Q|_{q_0, \eta_0} = 0, \quad \partial_\eta V_Q|_{q_0, \eta_0} = 0, \tag{49}$$

plus the usual conditions on the second derivatives. We will follow the discussion in [42] where the different kinds of minima are obtained by varying A and B .

When $CE > 6D^2/25$, the potential, considered as a function of δ, Δ , contains a positive definite quadratic form. In addition, changing the independent variables q and η , the region $\delta^3 \geq 6\Delta^2$ is mapped, and the strict inequality occurs for $\eta \neq 0$. Then, the points obtained by simply minimizing with respect to δ, Δ taken as independent variables lead to possible minima as long as they correspond to $\eta_0 \neq 0$. Otherwise, the potential must be minimized with the constraint $\delta^3 = 6\Delta^2$, in which case two different situations are obtained, $q_0 = 0, \eta_0 = 0$ or $q_0 \neq 0, \eta_0 = 0$.

Then, the ψ -field minima follow from the study of the ‘effective potential’ $\mathcal{V}(\psi)$, defined by:

$$\mathcal{V}(\psi) = V_\phi + V_I(q_0, \eta_0; \psi), \tag{50}$$




Figure 2. A-B phase diagram for the monopole sector, when $CE < 6D^2/25$, $D > 0$. In this case, the N_b phase and the associated $Z(2)$ SSB vortex state are lacking. When $D < 0$, the diagram is similar, with the N^+N^- line having opposite tilt.

and of which explicit form is:

$$\mathcal{V}(\psi) = \frac{1}{2}[\mu^2 - \zeta(q_0 + \eta_0)]\psi_1^2 + \frac{1}{2}[\mu^2 - \zeta(q_0 - \eta_0)]\psi_2^2 + \frac{1}{2}(\mu^2 + 2\zeta q_0)\psi_3^2 + \frac{\lambda}{4}(\psi_1^2 + \psi_2^2 + \psi_3^2)^2. \tag{51}$$

For $D < 0$, the monopole phase diagram is that of figure 1 [42].

When $CE < 6D^2/25$, the minimization of the potential requires again the constraint $\delta^3 = 6\Delta^2$, thus leading to $q_0 = 0, \eta_0 = 0$ or $q_0 \neq 0, \eta_0 = 0$ minima. In this case, the monopole phase diagram is depicted in figure 2, so the analysis will be similar to the former case, with the important difference that the $\eta_0 \neq 0$ phase is lacking. Then, from now on we will consider $CE > 6D^2/25$. It is clear what kind of vacua may emerge, depending on the relative values of the parameters. We first note that stability requires $\lambda \geq 0$. If the parameters A, B are initially in region I , we have $Q = 0$ ($q_0 = \eta_0 = 0$), and the effective vortex potential results, $\mathcal{V}^I(\psi) = \frac{\mu^2}{2}\psi^2 + \frac{\lambda}{4}(\psi^2)^2$. Then, if $\mu^2 \geq 0$, the minimization gives $\psi = 0$. With this vacuum, $S_{v,m}$ displays a non Abelian gauge symmetry, much larger than the Abelian symmetries in equations (38) and (39), typically obtained when monopoles and center vortices are present. Therefore, this phase represents a situation where monopoles and center vortices do not proliferate (deconfining phase). Still in the $Q = 0$ phase, but with $\mu^2 < 0$, the system undergoes SSB leaving an Abelian symmetry. If the mass scale generated for the off-diagonal fields is large, they will be suppressed and then the effective theory will essentially be invariant under Abelian gauge transformations of the form

$$\vec{\lambda}_\mu \cdot \hat{\phi}_0 \rightarrow \vec{\lambda}_\mu \cdot \hat{\phi}_0 + \partial_\mu \varphi.$$

Thus, recalling equation (39), this phase describes an ensemble of closed center vortices.

Now, in order to continue the analysis, it is convenient to define a complex field $V = \frac{1}{\sqrt{2}}(\psi_1 + i\psi_2)$, and rewrite equation (51) in the form (we consider $\zeta < 0$)

$$\mathcal{V}(\phi) = (\mu^2 + |\zeta|q_0)\bar{V}V + \frac{1}{2}|\zeta|\eta_0(V^2 + \bar{V}^2) + \frac{1}{2}(\mu^2 - 2|\zeta|q_0)\psi_3^2 + \lambda(\bar{V}V + \frac{1}{2}\psi_3^2)^2. \quad (52)$$

When A is lowered from positive to negative values, after a first-order transition, we will enter the uniaxial nematic phase N^+ ($q_0 > 0$) or the N^- ($q_0 < 0$), depending on whether $B < 0$ or $B > 0$. These phases are characterized by $\eta_0 = 0$. In what follows, to simplify the analysis, we will suppose $\mu^2 > 0$. Then, if we enter the N^- phase, after a discontinuous transition, the effective potential $\mathcal{V}(\psi)$ will be minimized by $\psi_3 = 0$. In the monopole sector, the vacuum will be invariant under rotations around the third axis, while in the V -sector this symmetry will undergo a $U(1)$ SSB or not, depending on the sign of $(\mu^2 + |\zeta|q_0)$. In addition, the N^- phase will induce a mass of order q_0^2 for the charged dual vector fields λ_μ^1 and λ_μ^2 , originated from the covariant derivative of Q in equation (42). If we assume this mass to be large when compared with the other mass scales in the problem, these dual vector fields will become suppressed.

If we further diminish A , after a second-order phase transition, we will eventually reach the biaxial phase N_b , where $\eta_0 \neq 0$. As this transition is continuous, and we are approaching from the N^- phase, we will begin with ψ_3 and η_0 small. In the N_b phase, the $U(1)$ symmetry of the effective action in the former N^- phase will be broken to a discrete one under π -rotations along the third axis. Again, in the monopole sector the vacuum is invariant, while in the vortex sector it will display SSB of the discrete π -rotations depending on the sign of $(\mu^2 + |\zeta|q_0)$. When this quantity is negative, at the minima, the V field can take a pair of values $V_0, -V_0$ connected by a $Z(2)$ symmetry. That is, the obtained effective potential coincides with the confining phase of the vortex model introduced by t' Hooft, relying on the possible nontrivial vortex correlators in the initial theory. In this phase, the spontaneous $Z(2)$ symmetry breaking leads to domain walls attached to Wilson loops, thus providing an area law. Still in the $(\mu^2 + |\zeta|q_0) < 0$ case, in the intermediate N^- phase, the vacuum no longer displays the Abelian symmetry present in the initial phase, where center vortices are only closed objects, nor the discrete symmetry of the last phase, typical of open center vortices whose endpoints are joined in pairs to monopole-like objects that proliferate. From this perspective, we speculate that the N^- phase might be associated with one where monopoles and antimonopoles are still bound in pairs.

5. Conclusions

We have constructed a novel non Abelian effective model for $SU(2)$ Yang–Mills theory in an Euclidean three-dimensional spacetime that allows for the description of a phase diagram with a rich structure. The construction is based on a special parametrization of the gauge field configurations \vec{A}_μ in terms of a vector field $\vec{\mathcal{A}}_\mu$, representing a topologically trivial sector of smooth fluctuations, and a local color frame \hat{n}_a containing defects, the nontrivial sector describing monopoles and thin center vortices. The frame can be written as a local $SO(3)$ rotation R of the canonical basis \hat{e}_a , which can be also expressed in the form $R = R(S)$, where S is in the fundamental representation.

This parametrization is used to write the Yang–Mills action, which defines the weight assigned to each configuration. On the other hand, as in any non-perturbative definition of the functional integration measure in a non Abelian gauge theory, one is faced with the usual stumbling blocks, related to the Gribov problem. We do not attempt to tackle this problem; rather, since we use the functional integral just as a guide for the subsequent derivation of the effective model, we use instead a definition of the measure which: (a) reduces to the proper one for topologically trivial configurations and (b) is consistent with (although not uniquely determined by) the properties of the gauge field parametrization used.

The next step in the construction of the effective model proceeds with the introduction of an auxiliary field $\vec{\lambda}_\mu$ that linearizes the Yang–Mills action, and the incorporation of a

phenomenological weight S_d that senses the geometry of the defects. It is at this point where the real reduction to an effective theory is implemented. Indeed, the symmetries are identified here, for a given classification of defects, what allows us to construct an effective model.

If S_d were nullified (thin objects), the partition function for the sector of defects should be invariant under local $SO(3)$ rotations of $\vec{\lambda}$, as they could be absorbed by a frame redefinition, transforming S under right multiplication by an appropriate regular $SU(2)$ matrix \tilde{U}^{-1} . In the Yang–Mills partition function, the symmetry should also be accompanied by the transformation $\vec{A}_\mu \rightarrow \tilde{A}_\mu$. However, this symmetry is the one associated with the many different ways a given gauge field \vec{A}_μ containing thin defects can be decomposed, so that it is expected to be broken as soon as center vortices become thick. Alternatively, this could be seen as the noninvariance of the effective phenomenological action S_d under local frame rotations, only leaving a global $SO(3)$.

An interesting point is that in order to guide the construction of the effective model for the ensemble integration, not only the global $SO(3)$ symmetry is important but also a new symmetry comes into play. At least in the symmetric phase, due to the topological structure of monopoles, the model should be invariant under a local discrete gauge symmetry. This led us to propose a non Abelian model describing the interaction of the natural order parameters for monopoles and center vortices with the effective gluon field \vec{A}_μ . As center vortices can be attached in pairs to the non Abelian monopoles, the corresponding order parameters are given by fields ϕ and Q , carrying isospin one and two, respectively. The effective character of the gluons is due to the fact that gauge transformations of the Yang–Mills fields \vec{A}_μ act as a left multiplication of the S sector, leaving \vec{A}_μ invariant.

The effective model we introduced exhibits a rich phase diagram. For instance, the monopole sector of the effective potential depends on two invariants, $\delta = \text{Tr } Q^2$, $\Delta = \text{Tr } Q^3$. If this sector is supposed to dominate the transitions, the phase diagram inherits, by construction, some of the properties found in liquid crystals. In this case, if the quadratic form in the quantities δ and Δ is positive definite, and the coefficient of the linear Δ -term is positive, an interesting chain of phase transitions is obtained.

Initially, when the coefficient of the linear δ -term (A) is varied from positive to negative values, a first-order transition from the isotropic deconfining phase to a uniaxial monopole condensate takes place. In this process, in the vortex sector, the ‘third’ component becomes suppressed, while the other two components can be arranged as an Abelian complex vortex field V displaying $U(1)$ SSB. In this example, the vortex mass scales have been supposed to be negligible when compared with those generated in the monopole sector. The further reduction of A produces a second-order phase transition, and the monopole condensate becomes biaxial. Here, center vortices are left in a global $Z(2)$ SSB phase, thus making contact with the ’t Hooft vortex model, and arriving at the confining phase expected in 3D Yang–Mills theories.

Acknowledgments

The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil) and PROPPi-UFF are acknowledged for the financial support. CDF acknowledges financial support from CONICET and UNCuyo.

References

- [1] Chernodub M N and Polikarpov M I 1997 *Lectures given at the Workshop ‘Confinement, Duality and Non-Perturbative Aspects of QCD’ (Cambridge, England)* arXiv:[hep-th/9710205](https://arxiv.org/abs/hep-th/9710205)
- [2] Chernodub M N, Gubarev F V, Polikarpov M I and Veselov A I 1998 *Prog. Theor. Phys. Suppl.* **131** 309

- [3] Giacomo A Di, Lucini B, Montesi L and Paffuti G 2000 *Phys. Rev. D* **61** 034503
- [4] Kronfeld A S, Laursen M L, Schierholz G and Wiese U J 1987 *Phys. Lett. B* **198** 516
- [5] Del Debbio L, Faber M, Greensite J and Olejnik S 1997 *Phys. Rev. D* **55** 2298
- [6] Del Debbio L, Faber M, Giedt J, Greensite J and Olejnik S 1998 *Phys. Rev. D* **58** 094501
- [7] de Forcrand P and D'Elia M 1999 *Phys. Rev. Lett.* **82** 4582
- [8] Greensite J 2003 *Prog. Part. Nucl. Phys.* **51** 1
- [9] Engelhardt M, Quandt M and Reinhardt H 2004 *Nucl. Phys. B* **685** 227
- [10] Ambjørn J, Giedt J and Greensite J 2000 *Nucl. Phys. Proc. Suppl.* **83** 467
- [11] de Forcrand Ph and Pepe M 2001 *Nucl. Phys. B* **598** 557–77
- [12] Gubarev F V, Kovalenko A V, Polikarpov M I, Syritsyn S N and Zakharov V I 2003 *Phys. Lett. B* **574** 136
- [13] Bali G S 2001 *Phys. Rep.* **343** 1
- [14] Baker M, Ball J S and Zachariassen F 1995 *Phys. Rev. D* **51** 1968
- [15] Baker M, Ball J S, Brambilla N, Prospero G M and Zachariassen F 1996 *Phys. Rev. D* **54** 2829
Baker M, Ball J S, Brambilla N, Prospero G M and Zachariassen F 1997 *Phys. Rev. D* **56** 2475 (erratum)
- [16] Maedan S and Suzuki T 1989 *Progr. Theor. Phys.* **81** 229
- [17] Shiba H, Kamisawa S, Matsubara Y and Suzuki T 1993 *Nucl. Phys. B* **389** 563
- [18] Chernodub M N and Komarov D A 1998 *JETP Lett.* **68** 117
- [19] Oxman L E 2013 *J. High Energy Phys.* [JHEP03\(2013\)038](#)
- [20] Ketzer B 2012 *PoS(QNP2012)* 025
- [21] Dudek J J, Edwards R G, Peardon M J, Richards D G and Thomas C E 2009 *Phys. Rev. Lett.* **103** 262001
- [22] Dudek J J and Edwards R G 2012 [arXiv:1201.2349](#)
- [23] Ambjørn J 1994 *Quantization of Geometry, in Les Houches* ed F David, P Ginsparg and J Zinn-Justin, [arXiv:hep-th/9411179](#)
- [24] Bardakci K and Samuel S 1978 *Phys. Rev. D* **18** 2849
- [25] Kiometzis M, Kleinert H and Schakel A M J 1995 *Fortschr. Phys.* **43** 697
- [26] Halpern M B, Jevicki A and Senjanovic P 1977 *Phys. Rev. D* **16** 2476
- [27] Halpern M B and Siegel W 1977 *Phys. Rev. D* **16** 2486
- [28] Kleinert H 2006 *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics, and Financial Markets* (Singapore: World Scientific)
- [29] Engelhardt M and Reinhardt H 2000 *Nucl. Phys. B* **567** 249
- [30] 't Hooft G 1978 *Nucl. Phys. B* **138** 1
- [31] de Lemos A L L, Oxman L E and Teixeira B F I 2012 *Phys. Rev. D* **85** 125014
- [32] Oxman L E 2008 *J. High Energy Phys.* [JHEP12\(2008\)089](#)
- [33] Oxman L E 2010 *Phys. Rev. D* **82** 105020
- [34] Antonov D 2000 *Surveys High Energy Phys.* **14** 265
- [35] Kogan I I and Kovner A 2002 Monopoles, vortices and strings: confinement and deconfinement in 2+1 dimensions at weak coupling [arXiv:hep-th/0205026](#)
- [36] Polyakov A M 1975 *Phys. Lett. B* **59** 82
Polyakov A M 1977 *Nucl. Phys. B* **120** 429
- [37] Oxman L E 2011 *J. High Energy Phys.* [JHEP07\(2011\)078](#)
- [38] Reinhardt H 2002 *Nucl. Phys. B* **628** 133
- [39] Cho Y M 1980 *Phys. Rev. D* **21** 1080
Cho Y M 1981 *Phys. Rev. Lett.* **46** 302
Cho Y M 1981 *Phys. Rev. D* **23** 2415
- [40] Faddeev L and Niemi A J 1999 *Phys. Rev. Lett.* **82** 1624
- [41] Shabanov S V 1999 *Phys. Lett. B* **458** 322
- [42] de Gennes P G and Prost J 1993 *The Physics of Liquid Crystals* (Oxford: Clarendon)
- [43] Gribov N 1978 *Nucl. Phys. B* **139** 1
- [44] Zwanziger D 1989 *Nucl. Phys. B* **323** 513
Zwanziger D 1993 *Nucl. Phys. B* **399** 477
- [45] Bruckmann F, Heinzl T, Wipf A and Tok T 2000 *Nucl. Phys. B* **584** 589
- [46] Greensite J, Olejnik S and Zwanziger D 2005 *J. High Energy Phys.* [JHEP05\(2005\)070](#)
- [47] Maas A 2007 *Nucl. Phys. A* **790** 566
- [48] Buividovich P V, Polikarpov M I and Zakharov V I 2007 *PoS(LATTICE 2007)* 324
- [49] Morse D C and Fredrickson G H 1994 *Phys. Rev. Lett.* **73** 24
- [50] Fredrickson G 2006 *The Equilibrium Theory of Inhomogeneous Polymers* (Oxford: Clarendon)