

## Current recommendations and novel strategies for sustainable management of soybean sudden death syndrome

Maria Cecilia Rodriguez<sup>a\*</sup>, Francisco Sautua<sup>b</sup>, Mercedes Scandiani<sup>c</sup>, Marcelo Carmona<sup>b#</sup>, Sebastián Asurmendi<sup>a</sup>

<sup>a</sup> Instituto de Agrobiotecnología y Biología Molecular (IABIMO), Instituto Nacional de Tecnología Agropecuaria (INTA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina.

<sup>b</sup> Fitopatología, Facultad de Agronomía, Universidad de Buenos Aires, Buenos Aires, Argentina.

<sup>c</sup> Centro de Referencia de Micología (CEREMIC), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina.

\*corresponding autor: MC Rodriguez (e-mail: mcecirodri@gmail.com)

#co-corresponding autor: MA Carmona (e-mail: carmonam@agro.uba.ar)

### ABSTRACT

The increase in food production requires reduction of the damage caused by plant pathogens, minimizing the environmental impact of management practices. Soil-borne pathogens are among the most relevant pathogens that affect soybean crop yield. Soybean sudden death syndrome (SDS), caused by several distinct species of *Fusarium*, produces significant yield losses in the leading soybean producing countries in North and South America. Current management strategies for SDS are scarce since there are no highly resistant cultivars, and only a few fungicide seed treatments are available. Because of this, innovative approaches for SDS management need to be developed. Here, we summarize recently explored strategies based on plant nutrition, biological control, priming of plant defenses, host-induced gene silencing, and the development of new SDS-resistance cultivars using precision breeding techniques. Finally, sustainable management of SDS should also consider cultural control practices with minimal environmental impact.

**Keywords:** Sudden death syndrome, *Fusarium virguliforme*, *Fusarium tucumaniae*, disease control, integrated disease management, *Glycine max*.

### INTRODUCTION

Soybean sudden death syndrome (SDS) is one of the most yield-limiting soybean diseases in North and South America. Multiple variables influence SDS development, such as edaphic properties of each field plot, the interaction with other soil-borne pathogens like nematodes, the agronomic practices carried out by each producer (rotations, tillage system, fertilization), and the associations with certain field-specific environmental factors of each particular year. These variables interact in many ways, explaining why numerous field experiments have obtained contradictory results in the last 50 years.

The disease is caused by several species of the soil-borne fungus *Fusarium*: *F. virguliforme* (*F.v.*), *F. tucumaniae* (*F.t.*), *F. brasiliense* (*F.b.*), *F. crassistipitatum* (*F.cr.*), and a novel undescribed *Fusarium*

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ps.6458](https://doi.org/10.1002/ps.6458)

spp. (Table 1) □.<sup>1-5</sup> *F. cuneirostrum* was reported to cause SDS in earlier literature, but it was recharacterized and removed from the list in recent classification<sup>6</sup>. These species are classified into clade 2 of the *Fusarium solani* species complex (FSSC).<sup>7</sup> In North America, *F.v.* is widely prevalent, but *F.b.* and *F.cu.* have also been detected in some regions of the USA.<sup>8</sup> In South America, *F.t.* and *F.b.* are the most prevalent species in Argentina and Brazil, respectively, but other *Fusarium* species have also been isolated.<sup>1-4</sup> □ Members of the FSSC are identified based on phenotypic analyses of macro- and microscopic characters and phylogenetic analyses of multilocus DNA sequence data.<sup>1,3-</sup> □<sup>4</sup> Additionally, the comparison of whole-genome sequences has allowed a better characterization of the FSSC, establishing differences in their life cycle,<sup>9,10</sup> □ and in their sensitivity to chemical compounds.<sup>11</sup> Finally, differences in the reproduction between members of FSSC in clade 2 have been described based on the structural organization of the mating-type locus, indicating that some species could be sexually propagated. However, only *F.t.* has been shown to exhibit this type of propagation to date.<sup>10,12-13</sup>

*F. tucumaniae* and *F. virguliforme* overwinter primarily as chlamydospores both in infested crop residues and free in the soil. These resistance structures can survive in the soil for several years. Sexual recombination of *F. tucumaniae* was confirmed to have occurred. In contrast, *F. virguliforme* never produced perithecia<sup>12</sup>. Therefore, *F. tucumaniae* life cycle in S. America includes a sexual reproductive mode while the *F. virguliforme* population in the U.S. may be exclusively asexual. Other sources of primary inoculum are residues from other crops (such as corn) and alternative hosts. Germinating soybeans plus warming temperatures lead to the germination of the chlamydospores allowing for the infection cycle to begin (Figure1). After planting, the roots of the newly emerging seedlings are particularly susceptible to *Fusarium* and root infection in soybean can occur within days of planting. Primary infection is favored by high soil moisture. Initially, symptoms generated by SDS-causing pathogens of the FSSC include root discoloration, necrosis, and dieback. Later, the pathogen colonizes the vascular tissue in the basal area of the plant, a few centimeters above the soil line. Once established in the xylem, the pathogen produces toxins which are systemically transferred throughout the plant leading to the foliar symptoms. Thus, foliar symptoms are not typically seen until later in the season around the flowering stage. Various phytotoxins were isolated from soybean xylem sap.<sup>14-</sup> <sup>16</sup> Among them, a 13.5 kDa protein named Fvtox1 causes interveinal chlorosis and necrosis in leaves of susceptible soybean varieties.<sup>14,17</sup> Interestingly, foliar SDS-like symptoms development requires light, which suggests that FvTox1 alters the photosynthetic process.<sup>14,□18</sup> Besides, toxin-associated symptoms are affected by other environmental conditions such as temperature and media substrate<sup>19-</sup> <sup>21</sup> and, consequently, environmental changes over the years can explain the appearance of various degrees of foliar symptoms in the field.<sup>22-23</sup> In addition to this, some reports have shown inconsistencies between SDS foliar symptom severity and yield reduction,<sup>23-24</sup> □ making the screening for SDS-resistant germplasm and evaluation of other management tools more complex. Recently, Kandel *et al.* (2020) found a mean yield decline of 0.51% for each unit increase in a foliar disease index that combines foliar disease incidence and disease severity, providing an estimated yield reduction based on foliar symptoms.<sup>25</sup>

Yield losses caused by severe SDS infection are the result of premature defoliation of soybean plants and decreased seed size.<sup>23,26</sup> Flower and pod abortion can also be observed when severe SDS occurs during flowering and pod-filling.<sup>23,27</sup> □ However, yield losses due to SDS are variable and dependent on the interaction of soybean genotype, *Fusarium* pathotypes present in the field plots, weather and

edaphic conditions such as relative abundance of multiple microbial taxa in the soil, macro- and micronutrients availability, soil pH, macroporosity and cultural practices, among the main factors.<sup>28-</sup>

<sup>30</sup> In terms of its economic impact, SDS is considered to be one of the top ten soybean diseases of North America<sup>31</sup> with average yield losses of 0.82 million metric tons per year between 1996 to 2009<sup>32</sup> and 321,5 million USD per year between 1996 to 2016<sup>31</sup>. Additionally, SDS generates significant yield losses in Argentina and Brazil, two other leading soybean-producing countries<sup>33-34</sup> (Table 1).

Particularly in the USA, the synergism between *F.v.* and *Heterodera glycines*, the soybean cyst nematode (SCN) has been reported to cause a decrease in soybean yield when both pathogens are present in a given field plot.<sup>35-36</sup> According to a three-year research study, Westphal *et al.* (2014) suggested that *F.v.* depended on infections by *H. glycines* to cause highly severe damage.<sup>37</sup> Similarly, Roth *et al.* (2019) reported that the additional presence of SCN can increase the risk of severe SDS epidemics.<sup>38</sup> In Argentina, the presence of *H. glycines* and other nematode species such as *Meloidogyne incognita*, *M. javanica*, and *Helicotylenchus* spp. has been reported,<sup>39</sup> but their association with SDS is not common (Scandiani M, pers. comm.). In Brazil, *H. glycines* has been reported in at least 10 states,<sup>40-41</sup> but there are no reports of an association between these two pathogens.

Due to the increase in human population and dietary changes, food demand is expected to grow by 60 percent by the year 2050.<sup>42</sup> In parallel, soil depletion, disease risk associated with monoculture, and increasing costs due to fungicide resistance, negatively impact food production and affordability.<sup>43-44</sup> Therefore, improvement in the efficiency of management practices is needed to reduce the environmental impact of agriculture and to increase food production. In the last decades, new practices, such as reduced tillage strategies that use less fuel and preserve organic carbon closer to the surface of the soil, have been implemented in soybean production. However, no-till or reduced tillage can cause an increase in disease pressure, especially of those diseases caused by soil-borne fungal pathogens.<sup>45</sup> Also, the demand for sustainable agriculture calls for the development of alternative strategies for the control of fungal diseases. In particular, the management of SDS is limited due to the lack of highly resistant cultivars and the availability of only a few fungicides.<sup>25,46</sup> Therefore, the development of new tools for the sustainable management of SDS is required. Here, we discuss different strategies currently being tested for SDS management, together with some recent approaches developed for other fungal pathogens based on the use of gene-editing techniques and RNA silencing mechanisms. Finally, we evaluate the efficacy of these tools for the management of SDS (Table 2).

## CHEMICAL CONTROL

Infection of soybean root at early seedling stages allows the invasion of xylem and phloem tissues by SDS-causing *Fusarium* species and the translocation of the toxins that cause the aboveground symptoms.<sup>47</sup> Thus, delaying the pathogen infection by seed-applied fungicides can be an effective way to reduce the appearance of foliar symptoms. Recently, fluopyram (ILEVO™, a succinate dehydrogenase inhibitor developed by BASF, Ludwigshafen, Germany) was registered for the treatment of soybean seed to manage SDS. This fungicide reduces the initial establishment of the disease by inhibiting the early stages of the infection process and has proved to be effective against *F.v.*, *F.b.*, and *F.t.*<sup>11,48-49</sup> In particular, Sjarpe *et al.*<sup>48</sup> observed that fluopyram seed or in-furrow

treatment reduced SDS and increased yield relative to the control. Moreover, these authors evaluated the effect of fluopyram and other seed treatment and foliar products that have been registered for management of SDS and found that fluopyram provided the highest level of control of root and foliar symptoms of SDS among all the treatments.<sup>49</sup> Although phytotoxicity has been observed in cotyledons when fluopyram was applied as a seed treatment, likely causing a small reduction in plant population, fluopyram seed treatment has an overall positive impact on soybean yield under SDS disease pressure<sup>50</sup>. A second active ingredient, pydiflumetofen (Saltro®, another succinate dehydrogenase inhibitor developed by Syngenta), was also recently introduced to the market. A preliminary report has indicated promising results.<sup>51</sup> A third active ingredient, cyclobutrifluram (TYMIRIUM™), presumed to be an inhibitor of the mitochondrial electron transport chain complex II based on its similarity in chemical structure to fluopyram,<sup>52</sup> has shown to be effective for the control of SDS,<sup>53</sup> but it is not yet commercially available.<sup>54</sup>

Since the only two active ingredients currently available against SDS to date have the same single-site mode of action, the risk of fungicide resistance is high.<sup>55</sup> Based on this, their use should be carefully evaluated to avoid the appearance of resistance. In this sense, sustainable management should consider and quantify the disease risk to bypass the unnecessary use of fungicide seed-treatment.<sup>56</sup> Additionally, fluopyram and pydiflumetofen fungicide sensitivities should be monitored to evaluate and detect changes in the sensitivity to these compounds.<sup>57</sup>

## CULTURAL CONTROL

Cold temperatures and wet conditions during the early reproductive stages of soybean increase SDS foliar symptoms.<sup>58-59</sup> Agricultural practices that reduce the exposure to wet and cool weather, such as late planting, have been suggested to lower disease severity.<sup>60-61</sup> However, late-planted soybean produces less yield because of the shorter daylight hours during flowering and seed filling and, hence, it is not recommended as an effective management practice.<sup>61-62</sup>

Another factor influencing SDS development is the degree of soil compaction affecting the root system of soybean plants. Compacted soils tend to promote SDS development<sup>63-65</sup>. Therefore, any measure that allows for more aeration, porosity, and less soil compaction will be beneficial. This will reduce soil moisture and will avoid low soil temperatures, making the disease less intense.<sup>65</sup> In this regard, conventional tillage may be effective to lower the incidence and severity of SDS foliar symptoms.<sup>65-66</sup> However, other authors reported little or no effect of tillage on the intensity of SDS.<sup>67</sup> No-till systems usually depend on herbicide applications, particularly glyphosate, the most widely-used herbicide worldwide since 2001.<sup>68</sup> The effect of glyphosate on SDS development is still unclear. A report indicated that glyphosate can increase the incidence and severity of SDS foliar symptoms,<sup>69</sup> while others claimed that it has no effect.<sup>70-72</sup> Finally, it seems that when assessing the impact of tillage, crop rotation should also be taken into consideration.<sup>73-74</sup>

*Fusarium* species have several survival strategies such as resting conidia or chlamydospores in soil, debris infestation, and infecting a wide range of host plant species.<sup>23,75</sup> *Fusarium* chlamydospores can survive in the soil for many years.<sup>76</sup> Crop rotation has been proposed as an environment-friendly approach for soil-borne pathogen management.<sup>77</sup> Experiments testing short-term rotations have obtained mixed results, without a clear trend. For example, Pérez-Brandán *et al.*<sup>78</sup> found that SDS incidence was significantly higher under soybean monoculture than when soybean was grown in rotation with maize. Other studies reported that the current corn-soybean rotation is insufficient to

reduce the risk of damage by SDS,<sup>79,80</sup> probably because *F.v.* can remain asymptomatic in maize tissues.<sup>81</sup> These authors suggested that soil suppressiveness is an important component that can impact the SDS-pathogens independently of short-term crop rotations. The impact of long-term crop rotation practices on SDS severity has also been evaluated.<sup>82-83</sup> In particular, SDS incidence and severity were evaluated in a 6-year study including a 3 year-cropping system that incorporated corn-soybean-oat + red clover and a 4-year cropping system that included corn-soybean-oat + alfalfa-alfalfa rotation compared with a 2-year cropping system consisting of corn-soybean rotation.<sup>82</sup> Interestingly, the diversification of the cropping system reduced SDS incidence and severity and increased soybean yield in 5 of 6 years compared with the 2-year system. In this study, the crop rotations were confounded with fertility regime, and may have been affected by the crop rotation as well as the fertility regime used. Thus, the factors involved in the reduction of SDS severity need further characterization. Long-term rotations may help the development of suppressiveness and improvement of soil health which is a key factor to reducing the onset of SDS.<sup>30,78,84</sup> The succession of crops also influences the outcome of the rotation, since not all non-soybean crops are effective in reducing the soil population densities of *Fusarium* species causing SDS. For example, when fescue was incorporated in the rotation, *F.v.* density in soil was higher than when wheat and sorghum were included.<sup>83</sup> In connection with this, other cultural practices that can impact SDS development are growing cover crops and green manures. Cover crops are increasingly being used as a soil conservation practice when planted to improve soil structure, suppress the growth of weeds, and protect the soil from erosion caused by wind and water. However, while several species of legumes used as cover crops, such as *Trifolium* spp, *Medicago sativa*, and *Pisum sativum*, are considered hosts of *F.v.*, other legumes, grasses, cereals, and Brassicaceae cover crops, such as *Vicia villosa*, *Camelina sativa*, *Brassica juncea*, *Pennisetum glaucum*, *Secale cereale*, *Lolium multiflorum*, *Triticale hexaploide*, and *Triticum aestivum*, are nonhosts or poor hosts.<sup>85</sup> Therefore, knowledge of the susceptibility of cover crops to SDS-causing *Fusarium* species can help farmers choose which species should be planted in field plots with a history of SDS epidemics. Similarly, recent research reported promising results when green manure amendments were tested for SDS suppression in experimental plots in the greenhouse.<sup>86</sup> Amendments of oat and rye reduced root rot severity of soybean by 85% and 67%, respectively. However, these experiments involved a short period, until the soybean plants reached growth stage V3 (approximately 40 days after plating), and thus long-term field studies, involving rotations with different crops, are necessary to evaluate the real impact of this practice.

Along with these agronomic measures, precision farming practices, such as the quantification of inoculum level and the determination of the disease risk, are being implemented in some countries.<sup>38,56, 87-91</sup> For example, Roth *et al.*<sup>38</sup> showed that the quantification of *F.v.* abundance in soil at-planting provides valuable information that can be used to develop SDS risk prediction models. This tool may allow farmers to minimize the risk of yield loss, reducing treatment costs and the environmental impact associated with chemical treatments by applying these treatments only where they are required.<sup>56</sup> Additionally, detecting SDS-diseased patches using remote sensing methods is a promising tool that would allow to identify infected fields and adapt the management strategy for the following growing seasons in specific field-patches.<sup>87,89-91</sup>

## PLANT NUTRITION



Soil-borne pathogens cause root rot symptoms that affect nutrient uptake. Additionally, fungal toxins and enzymes with hydrolytic and catalytic activities can alter nutrient translocation and utilization<sup>92</sup>. Consequently, nutrition can influence crop diseases and can help to reduce infection symptoms in plants.<sup>93</sup> For example, high calcium (Ca) concentration in plant tissues is correlated with resistance to root diseases caused by *Fusarium solani* (F.s.).<sup>94</sup> Besides, potassium fertilization has been linked to a significant reduction in the incidence of many infectious diseases.<sup>93</sup> Nevertheless, while the addition of KCl to the soil notably reduced the SDS severity, different potassium salts had different effects on the growth of *F.v.*<sup>28</sup>. Therefore, the role of nutrients such as K in disease increase or suppression must be examined in conjunction with other mineral elements.

Additionally, sulfur (S) and micronutrients (MNs) such as Mn, copper (Cu), boron (B), zinc (Zn), and molybdenum (Mo) have been suggested as capable of priming systemic defense responses.<sup>93,95</sup> In particular, it was observed that treatment of soybean plants with nanomaterials (NMs) of CuO, B, MoO<sub>3</sub> or ZnO significantly decreases the impact of SDS, causing a reduction of 17-25% in root rot severity.<sup>96</sup> The direct effect of NPs over the pathogen was tested *in vitro*, showing no effect. Therefore, the authors suggest that the NPs may act through the increase of plant defenses. Another recent investigation confirmed that foliar applications of Cu-based NMs at the seedling stage significantly reduce SDS symptoms.<sup>97</sup> In addition, the authors demonstrated that Cu treatment alleviates the increased expression of antioxidant enzymes together with the changes in the fatty acid profile, and induces the expression of a set of defense genes. These investigations show that the use of Cu, Zn, B and Bo-based NMs have the potential to be used for management of SDS. Although these responses have been known for many years, it is not yet clear how specifically these ions can trigger defense responses in plants. Even though many studies discuss the impact of macro and micronutrients on host-pathogen relationships, very little research examines the role of nutrition in the control of SDS.

## BIOLOGICAL CONTROL

Biological pesticides employ a wide range of beneficial microorganisms (biological control agents, BCAs) that control or suppress populations of plant pathogens, reducing plant disease incidence.<sup>98</sup> In recent years, biological pesticides have gained increased interest due to their eco-friendly properties compared to the use of chemical pesticides. Biopesticides are less prone to generate tolerance in the target pathogens, are less harmful to other beneficial microorganisms, and show an extended persistence due to their natural reproduction capacity.<sup>99</sup> Because of these properties, various investigations have evaluated the use of BCAs and their mode of action for the control of SDS. For example, *Trichoderma harzianum* and two arbuscular mycorrhizal fungus (AMF), *Rhizophagus irregularis* and *R. intraradices*, reduced SDS severity caused by *F.v.*<sup>100-102</sup>. On the other hand, plant growth-promoting bacteria (PGPB) also proved to be effective in controlling SDS-causing *Fusarium* species. Pin viso et al.<sup>103</sup> reported a significant *F.t.* and *F.v.* mycelial growth inhibition caused by *Bacillus subtilis*, *Pseudomonas fluorescens* and *Chryseobacterium vietnamense* *in vitro*. In the case of *Lysobacter enzymogenes* strain C3, inoculated plants show increased biomass compared to non-inoculated plants.<sup>104</sup> Regarding the mode of action, both *T. harzianum* and *Lysobacter enzymogenes* strain C3 directly suppress *F.v.* mycelial growth.<sup>100,104</sup> Additionally, *T. harzianum* induces the activation of defense responses in soybean plants, increasing the expression of defense-related genes associated with the salicylic acid (SA) and jasmonic acid/ethylene (JA/ET) pathways.

The up-regulation of SA and JA defense-related genes was also detected in the soybean/AMF/*F.v.* interaction,<sup>105</sup> highlighting the importance of these defense pathways in the control of SDS. For the *R. intraradices*/soybean interaction, it was proposed that AMF-mediated root growth may alleviate SDS symptoms.<sup>101</sup> Altogether, these studies show that biological control may be an effective tool for the management of SDS, although additional studies should be performed to evaluate the efficacy of these treatments in field conditions.

## PRIMING OF PLANT DEFENSES

Priming is an adaptation mechanism that increases plant defense responses after exposure to abiotic or biotic stimuli. Following the stimulus perception, induced defense mechanisms are activated.<sup>106-107</sup> In general, after the pathogen perception by plant cells systemic acquired resistance (SAR), a long-lasting broad-spectrum systemic defense response, is induced.<sup>108</sup> Generally, priming is triggered by molecules produced by pathogens (PAMPs, from pathogen-associated molecular patterns) that are perceived by plants through appropriate protein receptors. Typical fungal PAMPs include chitosan, a natural biopolymer composed of D-glucosamine and N-acetyl-D-glucosamine that is found in fungal cell walls<sup>109</sup> and generates an immune response in plants called chitosan triggered immunity (CTI).<sup>110</sup> It has been shown that chitosan delayed SDS symptom expression when applied preventively on soybean leaves. Interestingly, chitosan treatment increased the level of chitinase activity in fungal infected leaves, suggesting that it can induce defense responses through CTI against SDS-causing *Fusarium* species in soybean plants. Additionally, chitosan effectively inhibited the *in vitro* growth of *Fusarium*.<sup>111</sup> Priming can also be induced by treatment with defense hormones or some natural secondary metabolites. The effect of salicylic acid (SA) treatment in the induced resistance against *F.s.* has been characterized. Pre-treatment of soybean seedlings with 200  $\mu$ M of SA reduced the disease symptoms caused by *F.s.* and the relative levels of fungal biomass in soybean seedlings after 3 days post-inoculation. Moreover, SA treatment significantly increased the activity of antioxidant enzymes and reduced the levels of malondialdehyde and hydrogen peroxide. Additionally, the level of SA-responsive genes in fungal infected seedlings was increased compared to non-treated infected seedlings. These results indicate that SA plays a role in the defense against *F.s.*, alleviating SDS symptoms.<sup>112</sup> Similarly, the efficacy of ethephon, an ethylene inducer, was evaluated by Abdelsamad *et al.*<sup>113</sup>. This compound was able to lower SDS foliar symptoms compared to control treatments in both susceptible and SDS-resistant cultivars. Moreover, ethephon treatment induced the ethylene biosynthesis genes, ethylene synthase and ethylene oxidase, and defense response genes like pathogenesis-related proteins, basic peroxidase (*IPER*), chalcone synthase, and defense-associated transcription factors. Particularly, *IPER* was highly upregulated 24 hours after ethephon application compared to the control treatment. Although the precise role of this enzyme was not characterized, additional studies may shed light on its role during *F.v.* infection. Finally, levels of pipecolic acid (Pip), a non-proteinaceous product of lysine catabolism that can induce SAR,<sup>114</sup> were increased in the xylem sap of leaves of *F.v.* infected plants.<sup>115</sup> Thus, this compound may be implicated in the defense response against SDS. Further studies evaluating the L-Pip exogenous effect are needed to clarify the role of this compound in the defense response against *F.v.* Altogether, these results indicate that treatments with natural compounds such as plant hormones and PAMPs could be a promising sustainable approach for partial control of SDS.

Other defense inducers like phosphites ( $\text{PO}_3^{3-}$ ) may be effective for the priming of plant defenses against SDS. Phosphites induce transcriptional changes that primes plants to defend themselves against several fungal pathogens, as was demonstrated in the case of *Phakopsora pachyrhizi* in soybean. Particularly, phosphites highly induced the expression of the enzyme phenylalanine ammonia-lyase in soybean plants<sup>116</sup> which was also up-regulated in a SDS-partially resistant soybean cultivar,<sup>117</sup> and thus, they could be promising compounds for the activation of defense responses against FSSC. Additionally, manganese phosphite inhibited *F.v.* and *F.t.* mycelial growth *in vitro*.<sup>118</sup>

## HOST-INDUCED GENE SILENCING

Host-induced gene silencing (HIGS) refers to the strategy that employs plant produced sRNAs (small RNAs) to silence specific genes in a non-viral pathogen to generate protection against pathogens. Hairpin RNA structures can be introduced into the plant genome by transgenesis to trigger the silencing mechanism. Alternatively, a non-transgenic strategy, named spray-induced gene silencing (SIGS), involves the exogenous application of dsRNAs (double-stranded RNAs) or siRNAs (small interfering RNAs) to plants. In recent years, HIGS and SIGS have been proposed as an emerging approach to control filamentous fungi such as *F. graminearum*, *F. oxysporum*, *F. verticilloides*, and *F. culmorum*.<sup>119-120</sup> However, this approach has not yet been exploited against SDS-causing *Fusarium* species. Considering the potential of this technique against filamentous fungi, here we briefly emphasize some relevant aspects for the selection of putative targets that could confer resistance against members of the FSSC. Also, we discuss the advantages of this technique in the light of sustainable management.

To design an effective strategy for SDS control by HIGS, the selection of a suitable pathogen gene that will be targeted by sRNAs must be very well established. Its role in pathogenesis must be considered to avoid those target genes that are redundant for the infection process. Also, it is important to prevent the off-target silencing of host plant or beneficial microorganism genes.<sup>119</sup> Genes involved in protein transport, cell differentiation, conidiation, regulation of primary metabolism, and modulation of plant-hormone pathways are generally essential for pathogenicity making them good target for silencing. Moreover, cell wall-degrading enzymes (CWDEs) are induced during the *F.v.* late infection phase<sup>92</sup> and are required for the invasion of plant tissues,<sup>121</sup> and hence, could be interesting targets for HIGS. Despite the impact of SDS on soybean yield, only a few *F.v.* genes involved in the pathogenesis process have been characterized to date. *FvSTR1*, a striatin orthologue in *F.v.*, plays a role in asexual development and virulence.<sup>122</sup> Interestingly, the disruption of *FvSTR1* resulted in complete loss of virulence in *F.v.* in a greenhouse experiment, showing that this gene could be a promising target for the HIGS strategy against SDS-causing *Fusarium* species. The sucrose non-fermenting protein kinase 1 gene (*FvSNF1*) could also be a target for HIGS against SDS. This gene codifies for a key component of the glucose de-repression pathway and its disruption abolishes the expression of the galactose oxidase gene. Moreover, *Fvsnf1* mutants exhibit a reduced expression of CWDE-coding genes in contrast to the wild-type strain and are severely impaired in their ability to cause SDS on challenged soybean.<sup>123</sup> Additional identification of new fungal genes involved in pathogenesis will increase the repertoire of HIGS targets, and therefore allow the successful implementation of this strategy against SDS.

HIGS has some advantages over the use of fungicides for SDS management since it can avoid the off-target effects against beneficial microorganisms through the careful selection and design of the



Accepted Article  
fungal targets. Furthermore, the scarce availability of fungicide target molecules increases the risk of selecting resistant strains. The HIGS strategy is advantageous since it is highly versatile in terms of the availability of target RNA molecules in the pathogen, hence, the appearance of HIGS-resistant strains could be easily overcome.<sup>119</sup> Finally, the use of SIGS can replace the employment of transgenic crops and, in this way, bypass the regulatory process required for their approval.

## BREEDING FOR SDS RESISTANCE

Tolerant cultivars are the most effective way of managing SDS. Despite the efforts of breeding programs, to date, only moderately resistant genotypes have been identified which can exhibit SDS foliar symptoms under favorable environmental conditions. In recent years, several *loci* and SNPs involved in tolerance against SDS have been identified through quantitative trait loci (QTL) and genotyping-by-sequencing (GBS) analysis, contributing to marker-assisted breeding programs.<sup>26</sup> These findings have been extensively summarized in a previously published review.<sup>124</sup> Considering that the development of improved cultivars by conventional breeding methods takes several years, new breeding techniques (NBTs) have recently appeared as a faster strategy that allows breeders to generate improved varieties in a few generations. NBTs comprise a wide range of techniques that include genome editing, cisgenesis, and intragenesis. Occasionally, changes to DNA sequences introduced by these techniques can be indistinguishable from those that can occur by conventional breeding and, therefore, the regulatory process may be short compared to the time required for the approval of transgene breeding cultivars.<sup>125</sup> The use of NBTs involves the identification of single major resistance genes or the discovery of disease-susceptibility *loci*. Here, we review some recent works that characterize soybean genes whose modification may enable the development of novel SDS resistance cultivars through NBTs.

Two single major genes conferring SDS resistance have been characterized.<sup>126-127</sup> These authors found a set of soybean genes whose expression is downregulated following *F.v.* infection and hypothesized that *F.v.* suppresses the transcription of these genes to induce host susceptibility.<sup>126</sup> In agreement with this, the overexpression of two of these genes, ankyrin repeat-containing protein (GmARP1) and a plasma membrane protein named *Glycine max* disease resistance 1 (GmDR1), were shown to enhance resistance against *F.v.* in transgenic soybean lines. Additionally, the overexpression of GmDR1 confers resistance against a broad spectrum of pathogens such as the soybean cyst nematode, spider mites, and soybean aphids, presumably through the recognition of PAMPs.<sup>127</sup> Interestingly, both defense genes, GmARP1 and GmDR1, were overexpressed under root-specific and infection inducible soybean promoters and, thus, these cultivars could be considered intragenic since soybean genetic elements were employed for the design of expression constructs. Therefore, these findings show that the overexpression of single major genes under endogenous promoters can be an effective strategy to provide resistance against SDS. Nevertheless, more research is needed at the field level to evaluate the durability and applicability of resistant-modified plants.

In recent years, the appearance of genome editing (GE) techniques such as the CRISPR-Cas system has made it possible to improve plant tolerance to biotic stress. This system is more adaptable and cheaper than other GE techniques and enables either DNA mutation or DNA base editing, expanding its uses.<sup>121</sup> CRISPR/Cas can be employed to reduce the susceptibility to fungal pathogens by disrupting the expression of susceptible genes (*S* genes) that are involved in plant sensitivity to

Accepted Article

phytotoxins.<sup>128,129</sup> Based on this, the study of *S loci* against SDS has gained relevance in the last years. In the searching for *S loci*, two soybean STAY-GREEN genes (GmSGR1 and GmSGR2) were found to be associated with the resistance to SDS foliar chlorosis.<sup>130</sup> However, soybean plants with a double mutation of GmSGR1 and GmSGR2 stayed green but displayed necrosis and reduced photosynthesis in response to *F.v.* phytotoxins. These detrimental agronomic traits make it impractical to use these *S loci* for the improvement of soybean tolerance against SDS. Further characterization of not deleterious *S loci* may be an effective strategy in the control of SDS. In this regard, the analysis of FvTox1-induced transcriptional changes in soybean leaves may help characterize additional *S loci* involved in FvTox1 recognition and to develop new SDS tolerant cultivars.<sup>131</sup> Future research on this topic, combined with the development of precise and reproducible SDS phenotyping methods,<sup>124</sup> will favor the generation of new tools for the management of SDS.

## CONCLUSIONS AND FUTURE OUTLOOK

Innovative technologies that emerged in recent years, such as gene editing and HIGS, together with a better understanding of the FSSC/soybean interaction, can be exploited to develop new sustainable management strategies for the control of SDS. The adoption of these tools is driven by the lack of efficient management options for the control of this disease. In particular, the current management options for SDS are limited because no completely resistant cultivars are available. Additionally, only two fungicide seed treatments, fluopyram and pydiflumetofen, are commercially available for the control of SDS. There is also an increased risk of fungicide resistance as these two molecules have the same specific target site. Moreover, fungicides generally impact both pathogenic and nonpathogenic fungi, and consequently, tend to reduce the population of beneficial microorganisms that help to avoid soil-borne pathogen epidemics. While efforts to develop resistant and tolerant soybean varieties continue, all available management measures should be applied in an integrated approach to control the incidence of SDS. In the present review, we discuss currently used strategies for SDS control. Besides, we propose new potential approaches that can be implemented. It is expected that in the coming years, new strategies based on recent technologies (HIGS, CRISPR, the use of nano micronutrients) will expand the repertoire of effective tools for the management of SDS. Finally, the implementation of each of these tools should be part of an integrated management strategy for the control of SDS.

## ACKNOWLEDGMENTS

The authors would like to thank Dr. Gabriela Conti (IABIMO, INTA) for the assistance with English-language editing. This work was supported by the National Institute of Agriculture Technology (INTA, INTA PD1116), the National Scientific and Technical Research Council-Argentina (CONICET) and the University of Buenos Aires (UBA, University of Buenos Aires Project UBACyT 20020170100147BA).

## DISCLOSURE

All authors declare that they have no competing interests.

1. Aoki T, O'Donnell K, Homma Y, Lattanzi AR. Sudden-death syndrome of soybean is caused by two morphologically and phylogenetically distinct species within the *Fusarium solani* species complex - *F. virguliforme* in North America and *F. tucumaniae* in South America. *Mycologia* **95**(4):660–684 (2003).
2. Aoki T, O'Donnell K, Scandiani MM. Sudden death syndrome of soybean in South America is caused by four species of *Fusarium*: *Fusarium brasiliense* sp. nov., *F. cuneirostrum* sp. nov., *F. tucumaniae*, and *F. virguliforme*. *Mycoscience*. **46**(3):162–183 (2005).
3. O'Donnell K, Sink S, Scandiani M, Luque A, Colletto A, Biasoli M, et al. Soybean sudden death syndrome species diversity within north and South America revealed by multilocus genotyping. *Phytopathology*. **100**(1):58–71 (2010).

- Accepted Article
4. Aoki T, Scandiani MM, O'Donnell K. Phenotypic, molecular phylogenetic, and pathogenetic characterization of *Fusarium crassistipitatum* sp. nov., a novel soybean sudden death syndrome pathogen from Argentina and Brazil. *Mycoscience*. **53**(3):167–186 (2012).
  5. Tewoldemedhin YT, Lamprecht SC, Vaughan MM, Doehring G, O'Donnell K. Soybean SDS in South Africa is caused by *Fusarium brasiliense* and a novel undescribed *Fusarium* sp. *Plant Dis*. **101**(1):150–157 (2017).
  6. Wang J (2016) Molecular diagnostics, epidemiology, and population genetics of the soybean sudden death syndrome pathogen, *Fusarium virguliforme*. PhD Thesis, Michigan State University.
  7. Geiser DM, Al-Hatmi AMS, Aoki T, Arie T, Balmas V, Barnes I, et al. Phylogenomic analysis of a 55.1 kb 19-gene dataset resolves a monophyletic *Fusarium* that includes the *Fusarium solani* Species Complex. *Phytopathology* (2020). DOI: 10.1094/PHYTO-08-20-0330-LE
  8. Wang J, Sang H, Jacobs JL, Oudman KA, Hanson LE, Chilvers MI. Soybean sudden death syndrome causal agent *Fusarium brasiliense* present in Michigan. *Plant Dis*. **103**(6):1234–1243 (2019).
  9. Srivastava SK, Huang X, Brar HK, Fakhoury AM, Bluhm BH, Bhattacharyya MK. The genome sequence of the fungal pathogen *Fusarium virguliforme* that causes sudden death syndrome in soybean. *PLoS One*. **9**(1)(2014). e81832. doi.org/10.1371/journal.pone.0081832
  10. Hughes TJ, O'Donnell K, Sink S, Rooney AP, Scandiani MM, Luque A, et al. Genetic architecture and evolution of the mating type locus in fusaria that cause soybean sudden death syndrome and bean root rot. *Mycologia*. **106**(4):686–697 (2014).
  11. Sang H, Witte A, Jacobs JL, Chang HX, Wang J, Roth MG, et al. Fluopyram sensitivity and functional characterization of sdhb in the *Fusarium solani* species complex causing soybean sudden death syndrome. *Front Microbiol*. **9**:1–12 (2018).
  12. Covert SF, Aoki T, O'Donnell K, Starkey D, Holliday A, Geiser DM, et al. Sexual reproduction in the soybean sudden death syndrome pathogen *Fusarium tucumaniae*. *Fungal Genet Biol*. **44**(8):799–807 (2007).
  13. Scandiani MM, Aoki T, Luque AG, Carmona MA, O'Donnell K. First report of sexual reproduction by the soybean sudden death syndrome pathogen *Fusarium tucumaniae* in Nature. *Plant Dis*. **94**(12):1411–1416 (2010).
  14. Brar HK, Swaminathan S, Bhattacharyya MK. The *Fusarium virguliforme* toxin FvTox1 causes foliar sudden death syndrome-like symptoms in soybean. *Mol Plant-Microbe Interact*. **24**(10):1179–1188 (2011).
  15. Abeysekara NS, Bhattacharyya MK. Analyses of the xylem sap proteomes identified candidate *Fusarium virguliforme* proteinaceous toxins. *PLoS One*. **9**(5)(2014).
  16. Chang HX, Domier LL, Radwan O, Yendrek CR, Hudson ME, Hartman GL. Identification of multiple phytotoxins produced by *Fusarium virguliforme* including a phytotoxic effector (fvnis1) associated with sudden death syndrome foliar symptoms. *Mol Plant-Microbe Interact*. **29**(2):96–108 (2016).



- Accepted Article
17. Pudake RN, Swaminathan S, Sahu BB, Leandro LF, Bhattacharyya MK. Investigation of the *Fusarium virguliforme* fvtox1 mutants revealed that the FvTox1 toxin is involved in foliar sudden death syndrome development in soybean. *Curr Genet.* **59**(3):107–117 (2013).
  18. Rosati RG, Lario LD, Hourcade ME, Cervigni GDL, Luque AG, Scandiani MM, Spampinato, CP. Primary metabolism changes triggered in soybean leaves by *Fusarium tucumaniae* infection. *Plant Sci.* **274**:91–100 (2018).
  19. Hartman GL, Huang YH, Li S. Phytotoxicity of *Fusarium solani* culture filtrates from soybeans and other hosts assayed by stem cuttings. *Australasian Plant Pathology.* **33**: 9–15 (2004).
  20. Xiang Y, Scandiani MM, Herman TK, Hartman G. Optimizing production and evaluation conditions for *Fusarium virguliforme* cell-free toxic filtrates to evaluate soybean genotype responses to isolates causing sudden death syndrome. *Plant Dis.* **99**:502–507 (2015).
  21. Tang E, Hill CB, Hartman GL. Carbon utilization profiles of *Fusarium virguliforme* isolates. *Can J Microbiol.* **56**(12):979–986 (2010).
  22. Roth MG, Webster RW, Mueller DS, Chilvers MI, Faske TR, Mathew FM, et al. Integrated Management of Important Soybean Pathogens of the United States in Changing Climate. *J Integr Pest Manag.* **11**(1) (2020).
  23. Roy KW, State M, Rupe JC, Hershman DE, Abney TS, Lafayette W, et al. Sudden Death Syndrome of Soybean. *Plant Dis.* **81**(10):1100–1111 (1997).
  24. Gongora-Canul C, Nutter FW, Leandro LFS. Temporal dynamics of root and foliar severity of soybean sudden death syndrome at different inoculum densities. *Eur J Plant Pathol.* **132**(1):71–79 (2012).
  25. Kandel Y, Bradley C, Chilvers M, Mathew F, Tenuta A, Smith D, et al. Relationship Between Sudden Death Syndrome caused by *Fusarium virguliforme* and Soybean Yield: A Meta-Analysis. *Plant Dis.* **104**:1736–1743 (2020).
  26. Hartman GL, Chang HX, Leandro LF. Research advances and management of soybean sudden death syndrome. *Crop Prot.* **73**:60–66 (2015).
  27. Rupe JC. Frequency and pathogenicity of *Fusarium solani* recovered from soybeans with Sudden Death Syndrome. *Plant Dis.* **73**:581–584 (1989).
  28. Sanogo S, Yang XB. Relation of sand content, pH, and potassium and phosphorus nutrition to the development of sudden death syndrome in soybean. *Can J Plant Pathol.* **23**(2):174–180 (2001).
  29. Chong SK, Hildebrand KK, Luo Y, Myers O, Indorante SJ, Kazakevicius A, et al. Mapping soybean sudden death syndrome as related to yield and soil/site properties. *Soil Tillage Res.* **84**(1):101–107 (2005).
  30. Srour AY, Gibson DJ, Leandro LFS, Malvick DK, Bond JP, Fakhoury AM. Unraveling microbial and edaphic factors affecting the development of sudden death syndrome in soybean. *Phytobiomes J.* **1**(2):91–101 (2017).

- Accepted Article
31. Bandara AY, Weerasooriya DK, Bradley CA, Allen TW, Esker PD. Dissecting the economic impact of soybean diseases in the United States over two decades. *PLoS One*. **15**(4):1–28 (2020). <http://dx.doi.org/10.1371/journal.pone.0231141>
  32. Koenning SR, Wrather JA. Suppression of Soybean Yield Potential in the Continental United States by Plant Diseases from 2006 to 2009. *Plant Heal Prog*. **11**(1):5 (2010).
  33. Wrather A, Shannon G, Balardin R, Carregal L, Escobar R, Gupta GK, et al. Effect of Diseases on Soybean Yield in the Top Eight Producing Countries in 2006. *Plant Heal Prog*. **11**(1):29 (2010).
  34. Scandiani MM, Carmona MA, Luque AG, Matos K da S, Lenzi L, Formento ÁN, et al. Aislamiento, identificación y daños asociados al síndrome de la muerte súbita en el cultivo de soja en Argentina. *Trop Plant Pathol*. **37**(5):358–362 (2012).
  35. McLean KS, Lawrence GW. Interrelationship of *Heterodera glycines* and *Fusarium solani* in Sudden Death Syndrome of Soybean. *J Nematol*. **25**(3):434–439 (1993).
  36. Xing L, Westphal A. Synergism in the interaction of *Fusarium virguliforme* with *Heterodera glycines* in sudden death syndrome of soybean. *J Plant Dis Prot*. **120**(5–6):209–217 (2013).
  37. Westphal A, Li C, Xing L, McKay A, Malvick D. Contributions of *Fusarium virguliforme* and *Heterodera glycines* to the disease complex of sudden death syndrome of soybean. *PLoS One*. **9**(6):1–13 (2014).
  38. Roth MG, Noel ZA, Wang J, Warner F, Byrne AM, Chilvers MI. Predicting soybean yield and sudden death syndrome development using at-planting risk factors. *Phytopathology*. **109**(10):1710–1719 (2019).
  39. Doucet ME, Lax P, Coronel N. The Soybean Cyst Nematode *Heterodera glycines* Ichinohe, 1952 in Argentina. In: Ciancio A, Mukerji K., editors. Integrated Management and Biocontrol of Vegetable and Grain Crops Nematodes Integrated Management of Plant Pests and Diseases, vol 2. Springer, Dordrecht, pp 127-148 (2008).
  40. Juliatti BCM. Análise de genótipos de soja quanto à resistência ao nematoide do cisto. Universidade federal de Uberlandia; 2015.
  41. Matsuo É, Sedyama T, Oliveira R, Cruz CD, Oliveira RCT. Avaliação de genótipos de soja em relação ao nematoide de cisto. *Bragantia*. **71**:173–181 (2012).
  42. Hunter MC, Smith RG, Schipanski ME, Atwood LW, Mortensen DA. Agriculture in 2050: Recalibrating targets for sustainable intensification. *Bioscience*. **67**(4):386–391 (2017).
  43. Pérez-Brandán C, Huidobro J, Grümberg B, Scandiani MM, Luque AG, Meriles JM, et al. Soybean fungal soil-borne diseases: A parameter for measuring the effect of agricultural intensification on soil health. *Can J Microbiol*. **60**(2):73–84 (2014).
  44. Hollomon DW. Fungicide Resistance: 40 Years on and Still a Major Problem. In: Ishii H, Hollomon D, editors. Fungicide Resistance in Plant Pathogens. Springer, Tokyo, pp 3-11. (2015).

- Accepted Article
45. Bockus WW, Shroyer JP. The impact of reduced tillage on soilborne plant pathogens. *Annu Rev Phytopathol.* **36**:485–500 (1998).
  46. Spampinato CP, Scandiani MM, Luque AG. Soybean sudden death syndrome: Fungal pathogenesis and plant response. *Plant Pathol.* **70**: 3– 12 (2021). doi.org/10.1111/ppa.13275
  47. Navi SS, Yang XB. Foliar Symptom Expression in Association with Early Infection and Xylem Colonization by *Fusarium virguliforme* (formerly *F. solani* f. sp. *glycines*), the Causal Agent of Soybean Sudden Death Syndrome. *Plant Heal Prog.* **9**(1):24 (2008).
  48. Sjarpe DA, Kandel YR, Chilvers MI, Giesler LJ, Malvick DK, McCarville MT, et al. Multi-location evaluation of fluopyram seed treatment and cultivar on root infection by *Fusarium virguliforme*, foliar symptom development, and yield of soybean. *Can J Plant Pathol.* **42**(2):192–202 (2020). doi.org/10.1080/07060661.2019.1666166
  49. Kandel YR, Bradley CA, Chilvers MI, Mathew FM, Tenuta AU, Smith DL, et al. Effect of seed treatment and foliar crop protection products on sudden death syndrome and yield of soybean. *Plant Dis.* **103**(7):1712–1720 (2019).
  50. Kandel YR, Wise KA, Bradley CA, Chilvers MI, Tenuta AU, Mueller DS. Fungicide and cultivar effects on sudden death syndrome and yield of soybean. *Plant Dis.* **100**(7):1339–1350 (2016).
  51. Adee EA. Effect of Saltro Soybean Seed Treatment on Sudden Death Syndrome in Kansas in 2019. *Kansas Agric Exp Stn Res Reports.* **6**(5) (2020). DOI: 10.4148/2378-5977.7932
  52. Umetsu N, Shirai Y. Development of novel pesticides in the 21st century. *J Pestic Sci.* **45**(2):54–74 (2020).
  53. Asociación de Laboratorios Agropecuarios Privados, editors. Lenzi L., Couretot L., A.; C, Gadbán L., A.; S, Russian H. Evaluación de tratamiento de semillas con Tymirium™ para el control del síndrome de la muerte súbita de la soja. In: 1er Congreso Argentino de Semillas. Argentina; 2020.
  54. Syngenta. Syngenta announces TYMIRIUM™ technology brand. <https://www.syngenta.com/pt-br/node/2198>
  55. Brent KJ, Hollomon DW. Fungicide Resistance in Plant Management: How can it be managed? FRAC Monograph No. 1 (second, revised edition). Fungicide resistance action committee, Brussels, 57 p. (2007)
  56. Lamichhane JR, You MP, Laudinot V, Barbetti MJ, Aubertot JN. Revisiting sustainability of fungicide seed treatments for field crops. *Plant Dis.* **104**(3):610–623 (2020).
  57. Wang J, Bradley CA, Stenzel O, Pedersen DK, Reuter-Carlson U, Chilvers MI. Baseline sensitivity of *Fusarium virguliforme* to fluopyram fungicide. *Plant Dis.* **101**(4):576–582 (2017).
  58. Scherm H, Yang XB. Development of sudden death syndrome of soybean in relation to soil temperature and soil water matric potential. *Phytopathology.* **86**: 642–649 (1996).

- Accepted Article
59. Leandro LFS, Robertson AE, Mueller DS, Yang X-B. Climatic and Environmental Trends Observed During Epidemic and Non-epidemic Years of Soybean Sudden Death Syndrome in Iowa. *Plant Heal Prog.* **14**(1):18 (2013).
  60. Hershman DE. Influence of Planting Date and Cultivar on Soybean Sudden Death Syndrome in Kentucky. *Plant Dis.* **74**:761-766 (1990).
  61. Marburger DA, Smith DL, Conley SP. Revisiting planting date and cultivar effects on soybean sudden death syndrome development and yield loss. *Plant Dis.* **100**(10):2152–2157 (2016).
  62. Kandel YR, Wise KA, Bradley CA, Tenuta AU, Mueller DS. Effect of planting date, seed treatment, and cultivar on plant population, sudden death syndrome, and yield of Soybean. *Plant Dis.* **100**(8):1735–1743 (2016).
  63. Rupe, J. C., Sabbe W. F., Robbins R. T., and Gbur, E. E., Jr. 1993. Soil and plant factors associated with sudden death syndrome of soybean. *J. Prod. Agric.* 6:218-221.
  64. Scherm, H., Yang, X. B., and Lundeen, P. 1998. Soil variables associated with sudden death syndrome in soybean fields in Iowa. *Plant Dis.* 82:1152-1157.
  65. Vick CM, Chong SK, Bond JP, Russin JS. Response of soybean sudden death syndrome to subsoil tillage. *Plant Dis.* **87**:629–632 (2003).
  66. Vick, C.M., Bond, J.P., Chong, S.K. and Russin, J.S. (2006) Response of soybean sudden death syndrome to tillage and cultivar. *Can. J. Plant Pathol.*, 28, 77–83.
  67. Kandel YR, Leandro LFS, Mueller DS. Effect of tillage and cultivar on plant population, sudden death syndrome, and yield of soybean in Iowa. *Plant Heal Prog.* **20**(1):29–34 (2019).
  68. Coupe RH, Capel PD. Trends in pesticide use on soybean, corn and cotton since the introduction of major genetically modified crops in the United States. *Pest Manag Sci.* **72**(5):1013–22 (2016).
  69. Sanogo S, Yang XB, Scherm H. Effects of herbicides on *Fusarium solani* f. sp. *glycines* and Development of sudden death syndrome in glyphosate-tolerant soybean. *Phytopathology.* **90**(1):57–66 (2000).
  70. Sanogo S, Yang XB, Lundeen P. Field response of glyphosate-tolerant soybean to herbicides and sudden death syndrome. *Plant Dis.* **85**(7):773–9 (2001).
  71. Njiti VN, Myers O, Schroeder D, Lightfoot DA. Roundup Ready Soybean: Glyphosate Effects on *Fusarium solani* Root Colonization and Sudden Death Syndrome. *Agron J.* **95**(5):1140–1145 (2003).
  72. Kandel YR, Bradley CA, Wise KA, Chilvers MI, Tenuta AU, Davis VM, et al. Effect of glyphosate application on sudden death syndrome of glyphosate-resistant soybean under field conditions. *Plant Dis.* **99**(3):347–354 (2015).
  73. Vargas Gil S, Meriles JM, Haro R, Casini C, March GJ. Crop rotation and tillage systems as a proactive strategy in the control of peanut fungal soilborne diseases. *BioControl.* **53**(4):685–698 (2008).



- Accepted Article
74. Mueller DS, Pedersen WL, Hartman GL. Effect of crop rotation and tillage system on sclerotinia stem rot on soybean. *Can J Plant Pathol.* **24**(4):450–456 (2002).
  75. Kolander TM, Bienapfl JC, Kurle JE, Malvick DK. Symptomatic and asymptomatic host range of *Fusarium virguliforme*, the causal agent of soybean sudden death syndrome. *Plant Dis.* **96**(8):1148–53 (2012).
  76. Couteaudier Y, Alabouvette C. Survival and inoculum potential of conidia and chlamydospores of *Fusarium oxysporum f.sp. lini* in soil. *Can J Microbiol.* **36**(8):551–556 (1990).
  77. Panth M, Hassler SC, Baysal-Gurel F. Methods for management of soilborne diseases in crop production. *Agriculture.* 10,16 (2020).
  78. Pérez-Brandán, C. Arzeno JL, Huidobro J, Conforto C, Grümberg B, Hilton S, Bending GD, et al. The effect of crop sequences on soil microbial, chemical and physical indicators and its relationship with soybean sudden death syndrome (complex of *Fusarium species*). *Spanish J Agric Res.* **12**:252–264 (2014).
  79. Xing L, Westphal A. Effects of crop rotation of soybean with corn on severity of sudden death syndrome and population densities of *Heterodera glycines* in naturally infested soil. *F CropRes.* **112**(1):107–117 (2009).
  80. Marburger DA, Venkateshwaran M, Conley SP, Esker PD, Lauer JG, Ané JM. Crop Rotation and Management Effect on *Fusarium spp.* Populations. *Crop Sci.* **55**(1):365–376 (2015).
  81. Baetsen-Young A, Wai CM, VanBuren R, Day B. *Fusarium virguliforme* transcriptional plasticity is revealed by host colonization of maize versus soybean. *Plant Cell.* **32**(2):336–51 (2020).
  82. Leandro LFS, Eggenberger S, Chen C, Williams J, Beattie GA, Liebman M, et al. Cropping system diversification reduces severity and incidence of soybean sudden death syndrome caused by *Fusarium virguliforme*. *Plant Dis.* **102**(9):1748–58 (2018).
  83. Rupe JC, Robbins RT, Gbur EE. Effect of crop rotation on soil population densities of *Fusarium solani* and *Heterodera glycines* and on the development of sudden death syndrome of soybean. *Crop Prot.* **16**(6):575–80 (1997).
  84. Westphal A, Xing L. Soil suppressiveness against the disease complex of the soybean cyst nematode and sudden death syndrome of soybean. *Phytopathology.* **101**(7):878–86 (2011).
  85. Kobayashi-Leonel R, Mueller D, Harbach C, Tylka G, Leandro L. Susceptibility of cover crop plants to *Fusarium virguliforme*, causal agent of soybean sudden death syndrome, and *Heterodera glycines*, the soybean cyst nematode. *J Soil Water Conserv.* **72**(6):575–583 (2017).
  86. Budi JW. Approaches to sustainable management of sudden death syndrome of soybean. (2020). Graduate Theses and Dissertations. 18058. <https://lib.dr.iastate.edu/etd/18058>
  87. Yang S, Li X, Chen C, Kyveryga P, Yang XB. Assessing field-specific risk of soybean sudden death syndrome using satellite imagery in Iowa. *Phytopathology.* **106**(8):842–853 (2016).

88. Chilvers M. Risk prediction tools and the interaction of soybean sudden death syndrome and soybean cyst nematode. *Crops & Soils*. 52: 18-19 (2019).
89. Herrmann I, Vosberg SK, Ravindran P, Singh A, Chang HX, Chilvers MI, et al. Leaf and canopy level detection of *Fusarium virguliforme* (sudden death syndrome) in soybean. *Remote Sens*. **10**(3), 426 (2018).
90. Hatton NM, Menke E, Sharda A, van der Merwe D, Schapaugh W. Assessment of sudden death syndrome in soybean through multispectral broadband remote sensing aboard small unmanned aerial systems. *Comput Electron Agric*. **167**:105094 (2019). doi.org/10.1016/j.compag.2019.105094
91. Raza MM, Harding C, Liebman M, Leandro LF. Exploring the potential of high-resolution satellite imagery for the detection of soybean sudden death syndrome. *Remote Sens*. **12**(7), 1213 (2020) doi.org/10.3390/rs12071213
92. Sahu BB, Baumbach JL, Singh P, Srivastava SK, Yi X, Bhattacharyya MK. Investigation of the *Fusarium virguliforme* transcriptomes induced during infection of soybean roots suggests that enzymes with hydrolytic activities could play a major role in root necrosis. *PLoS One*. **12**(1):1–22 (2017).
93. Dordas C. Role of nutrients in controlling plant diseases in sustainable agriculture: A review. *Agron. Sustain. Dev*. **28**, 33–46 (2008).
94. Huber DM, Haneklaus S. Managing nutrition to control plant disease. *Landbauforsch Volkenrode*. **57**(4):313–322 (2007).
95. Bloem E, Haneklaus S, Schnug E. Milestones in plant sulfur research on sulfur-induced-resistance (SIR) in Europe. *Front Plant Sci*. **5**:1–12 (2015).
96. Pérez CDP, De La Torre Roche R, Zuverza-Mena N, Ma C, Shen Y, White JC, et al. Metalloid and Metal Oxide Nanoparticles Suppress Sudden Death Syndrome of Soybean. *J Agric Food Chem*. **68**(1):77–87 (2020).
97. Ma C, Borgatta J, Hudson BG, Tamijani AA, De La Torre-Roche R, Zuverza-Mena N, et al. Advanced material modulation of nutritional and phytohormone status alleviates damage from soybean sudden death syndrome. *Nat. Nanotechnol*. **15**:1033–1042 (2020). doi.org/10.1038/s41565-020-00776-1
98. Köhl J, Kolnaar R, Ravensberg WJ. Mode of action of microbial biological control agents against plant diseases: Relevance beyond efficacy. *Front Plant Sci*. **10**, 845 (2019). doi.org/10.3389/fpls.2019.00845
99. Gupta S, Dikshit AK. Biopesticides: An ecofriendly approach for pest control. *J Biopestic*. **3**(1):186–8 (2010).
100. Pimentel MF, Arnão E, Warner AJ, Subedi A, F. Rocha L, Srour A, et al. *Trichoderma* isolates inhibit *Fusarium virguliforme* growth, reduce root rot, and induce defense-related genes on soybean seedlings. *Plant Dis*. **104**(7) (2020).

- Accepted Article
101. Pawlowski ML, Hartman GL. Reduction of Sudden death syndrome foliar symptoms and *Fusarium virguliforme* DNA in roots inoculated with *Rhizophagus intraradices*. *Plant Dis.* **104**(5):1415–20 (2020).
  102. Giachero ML, Marquez N, Gallou A, Luna CM, Declerck S, Ducasse DA. An in vitro method for studying the three-way interaction between soybean, *Rhizophagus irregularis* and the soil-borne pathogen *Fusarium virguliforme*. *Front Plant Sci.* **8**, 1033 (2017). <https://doi.org/10.3389/fpls.2017.01033>
  103. Pin Viso N, Sautua F, Scandiani M, Luque A, Simonetti E, Carmona M. In vitro antagonistic activity of native bacteria isolated from soils of the Argentine Pampas against *Fusarium tucumaniae* and *Fusarium virguliforme*. *African J Microbiol Res.* **10**(27):1031–5 (2016).
  104. Nian J, Yu M, Bradley CA, Zhao Y. Lysobacter enzymogenes strain C3 suppresses mycelium growth and spore germination of eight soybean fungal and oomycete pathogens and decreases disease incidences. *Biol Control.* **152**:104424 (2021). [doi.org/10.1016/j.biocontrol.2020.104424](https://doi.org/10.1016/j.biocontrol.2020.104424)
  105. Marquez N, Giachero ML, Gallou A, Debat HJ, Declerck S, Ducasse D. Transcriptome analysis of mycorrhizal and nonmycorrhizal soybean plantlets upon infection with *Fusarium virguliforme*, one causal agent of sudden death syndrome. *Plant Pathol.* **68**:470–480 (2019).
  106. Mauch-Mani B, Baccelli I, Luna E, Flors V. Defense Priming: An Adaptive Part of Induced Resistance. *Annu Rev Plant Biol.* **68**:485–512 (2017).
  107. Kachroo A, Kachroo P. Mobile signals in systemic acquired resistance. *Curr Opin Plant Biol.* **58**:41–47 (2020).
  108. Gaffney T, Friedrich L, Vernooij B, Negrotto D, Gaffney T, Friedrich L, et al. Requirement of Salicylic Acid for the Induction of Systemic Acquired Resistance. *Science.* **261** (5122): 754–756 (1993).
  109. Latgé JP. The cell wall: A carbohydrate armour for the fungal cell. *Mol Microbiol.* **66**(2):279–290 (2007).
  110. Narula K, Elagamey E, Abdellatef M, Sinha A, Ghosh S, Chakraborty, N CS. Chitosan-triggered immunity to *Fusarium* in chickpea is associated with changes in the plant extracellular matrix architecture, stomatal closure and remodeling of the plant metabolome and proteome. *Plant J.* **103**(2):561–83 (2020).
  111. Prapagdee B, Kotchadat K, Kumsopa A, Visarathanonth N. The role of chitosan in protection of soybean from sudden death syndrome caused by *Fusarium solani* f. sp. *glycines*. *Bioresour Technol.* **98**(7):1353–8 (2007).
  112. Bawa G, Feng L, Yan L, Du Y, Shang J, Sun X, et al. Pre-treatment of salicylic acid enhances resistance of soybean seedlings to *Fusarium solani*. *Plant Mol Biol.* **101**(3):315–23 (2019).
  113. Abdelsamad NA, MacIntosh GC, Leandro LFS. Induction of ethylene inhibits development of soybean sudden death syndrome by inducing defense-related genes and reducing *Fusarium virguliforme* growth. *PLoS One.* **14**(5):e0215653 (2019). doi: 10.1371/journal.pone.0215653.

- Accepted Article
114. Schnake A, Hartmann M, Schreiber S, Malik J, Brahmman L, Yildiz I, et al. Inducible biosynthesis and immune function of the systemic acquired resistance inducer N-hydroxypipelicolic acid in monocotyledonous and dicotyledonous plants. *J Exp Bot.* **71**(20):6444–59 (2020).
  115. Abeysekara NS, Swaminathan S, Desai N, Guo L, Bhattacharyya MK. The plant immunity inducer pipelicolic acid accumulates in the xylem sap and leaves of soybean seedlings following *Fusarium virguliforme* infection. *Plant Sci.* **243**:105–14 (2016).
  116. Gill US, Sun L, Rustgi S, Tang Y, von Wettstein D, Mysore KS. Transcriptome-based analyses of phosphite-mediated suppression of rust pathogens *Puccinia emaculata* and *Phakopsora pachyrhizi* and functional characterization of selected fungal target genes. *Plant J.* **93**(5):894–904 (2018).
  117. Iqbal MJ, Yaegashi S, Ahsan R, Shopinski KL, Lightfoot DA. Root response to *Fusarium solani* f. sp. *glycines*: Temporal accumulation of transcripts in partially resistant and susceptible soybean. *Theor Appl Genet.* **110**(8):1429–1438 (2005).
  118. Carmona M, Simonetti E, Ravotti ME, Scandiani M, Luque A, Formento N., et al. In vitro antifungal / fungistatic activity of manganese phosphite against soybean soil- borne pathogens. *Rev Int Bot Exp.* 9457:265–269 (2017).
  119. Machado AK, Brown NA, Urban M, Kanyuka K, Hammond-Kosack KE. RNAi as an emerging approach to control *Fusarium* head blight disease and mycotoxin contamination in cereals. *Pest Manag Sci.* **74**(4):790–799 (2018).
  120. Rampersad SN. Pathogenomics and management of fusarium diseases in plants. *Pathogens.* **9**(5) (2020).
  121. Tyagi S, Kumar R, Kumar V, Won SY, Shukla P. Engineering disease resistant plants through CRISPR-Cas9 technology. *GM Crop Food.* **12**(1):125–144 (2021).
  122. Islam KT, Bond JP, Fakhoury AM. FvSTR1, a striatin orthologue in *Fusarium virguliforme*, is required for asexual development and virulence. *Appl Microbiol Biotechnol.* **101**(16):6431–6445 (2017).
  123. Islam KT, Bond JP, Fakhoury AM. FvSNF1, the sucrose non-fermenting protein kinase gene of *Fusarium virguliforme*, is required for cell-wall-degrading enzymes expression and sudden death syndrome development in soybean. *Curr Genet.* **63**(4):723–38 (2017).
  124. Chang HX, Roth MG, Wang D, Cianzio SR, Lightfoot DA, Hartman GL, et al. Integration of sudden death syndrome resistance loci in the soybean genome. *Theor Appl Genet.* **131**(4):757–773 (2018). doi.org/10.1007/s00122-018-3063-0
  125. Lassoued R, Macall DM, Smyth SJ, Phillips PWB, Hesseln H. How should we regulate products of new breeding techniques? Opinion of surveyed experts in plant biotechnology. *Biotechnol Reports.* **26**:e00460. (2020) doi.org/10.1016/j.btre.2020.e00460
  126. Ngaki MN, Wang B, Srivastava SK, Farooqi MS, Kambakam S, Swaminathan S, et al. Transcriptomic Study of the Soybean- *Fusarium virguliforme* Interaction Revealed a Novel Ankyrin-Repeat Containing Defense Gene, Expression of Whose during Infection Led to



Enhanced Resistance to the Fungal Pathogen in Transgenic Soybean Plants. *PLoS One*. **19**;11(10):e0163106 (2016). doi: 10.1371/journal.pone.0163106.

127. Ngaki MN, Sahoo DK, Wang B, Bhattacharyya MK. Overexpression of a plasma membrane protein generated broad-spectrum immunity in soybean. *Plant Biotechnol J*. (2020). <https://doi.org/10.1111/pbi.13479>
128. Pavan S, Jacobsen E, Visser RGF, Bai Y. Loss of susceptibility as a novel breeding strategy for durable and broad-spectrum resistance. *Mol Breed*. **25**(1):1–12 (2010).
129. Vleeshouwers VGAA, Oliver RP. Effectors as tools in disease resistance breeding against biotrophic, hemibiotrophic, and necrotrophic plant pathogens. *Mol Plant-Microbe Interact*. **27**(3):196–206 (2014).
130. Chang HX, Tan R, Hartman GL, Wen Z, Sang H, Domier LL, et al. Characterization of soybean *STAY-GREEN* genes in susceptibility to foliar chlorosis of sudden death syndrome. *Plant Physiol*. **180**(2):711–717 (2019).
131. Wang B, Swaminathan S, Bhattacharyya MK. Identification of *Fusarium virguliforme* FvTox1-interacting synthetic peptides for enhancing foliar sudden death syndrome resistance in soybean. *PLoS One*. **10**(12):e0145156 (2015). doi: 10.1371/journal.pone.0145156
132. Rupe JC, Gbur JR EE. Effect of plant age, maturity group and environmental on disease progress of Sudden Death Syndrome of Soybean. *Plant Dis*. **79**:139–43 (1995).
133. Wrather JA, Anderson TR, Arsyad DM, Gai J, Ploper LD, Porta-Puglia A, et al. Soybean disease loss estimates for the top 10 soybean producing countries in 1994. *Plant Dis*. **81**(1):107–10 (1997).
134. Njiti VN, Shenaut MA, Suttner RJ, Schmidt ME, Gibson PT. Relationship between soybean sudden death syndrome disease measures and yield components in F6 – derived lines. *Crop Sci*. **38**(3):673–8 (1998).
135. Freitas TM de Q, Meneghetti RC, Balardin RS. Dano devido à podridão vermelha da raiz na cultura da soja. *Ciência Rural*. **34**(4):991–6 (2004).
136. Luo Y, Hildebrand K, Chong SK, Myers O, Russin JS. Soybean yield loss to sudden death syndrome in relation to symptom expression and root colonization by *Fusarium solani* f. sp. *glycines*. *Plant Dis*. **84**(8):914–20 (2000).
137. Costa SS, Matos KS, Tessmann DJ, Seixas CDS, Pfenning LH. *Fusarium paranaense* sp. nov., a member of the *Fusarium solani* species complex causes root rot on soybean in Brazil. *Fungal Biol*. **120**(1):51–60 (2016).
138. Aoki T, Tanaka F, Suga H, Hyakumachi M, Scandiani MM, O'Donnell K. *Fusarium azukicola* sp. nov., an exotic azuki bean root-rot pathogen in Hokkaido, Japan. *Mycologia*. **104**(5):1068–1084 (2012).
139. Kandel YR, McCarville MT, Adey EA, Bond JP, Chilvers MI, Conley SP, et al. Benefits and profitability of fluopyram-amended seed treatments for suppressing sudden death syndrome and protecting soybean yield a meta-analysis. *Plant Dis*. **102**(6):1093–1100 (2018).

**Table 1. Differential aspects of SDS in the three main soybean producing countries**

Feature	Argentina	Brazil	USA	Reference
Soybean area in 2020 (million hectares)	16.7	36.9	30.33	Global Market Analysis, FAS, USDA, November 2020
Soybean production in 2020 (million metric tons)	49	126	96.67	
Yield losses caused by SDS	22% (4-46) <sup>34</sup> and up to 90% <sup>136</sup>	38% (30-40) <sup>135,137</sup>	5 to 15% on average, 20 to 80% according to cultivar reaction, weather conditions and growth stage at the time of infection <sup>23,132-134</sup>	23, 34, <sup>□□</sup> 132-134, 136 <sup>□</sup>
SDS and cyst nematode association importance	not important / unknown	not important / unknown	Important	37 <sup>□</sup>
Prevalent <i>Fusarium</i> species causing SDS	<i>F. tucumaniae</i> <sup>1,2</sup>	<i>F. brasiliense</i> <sup>2</sup>	<i>F. virguliforme</i> <sup>1</sup>	1-4,8
Other <i>Fusarium</i> species reported to cause SDS <sup>(a)</sup>	<i>F. virguliforme</i> <sup>3</sup> <i>F. brasiliense</i> <sup>2</sup> <i>F. crassistipitatum</i> <sup>4</sup>	<i>F. tucumaniae</i> <sup>1,2</sup> <i>F. crassistipitatum</i> <sup>4</sup>	<i>F. brasiliense</i> <sup>8</sup>	
Other <i>Fusarium</i> species associated with SDS <sup>(b)</sup>	-	<i>F. paranaense</i> sp. nov. <sup>137</sup>	<i>F. phaseoli</i> <sup>8</sup> <i>Fusarium</i> sp. <sup>8(c)</sup>	
<i>Fusarium</i> sexual stage	Present	not reported	not reported	3,12

<sup>(a)</sup> In South Africa, a novel undescribed *Fusarium* sp. was isolated from soybean and caused root-rot and foliar SDS symptoms on soybean.<sup>5</sup> <sup>□</sup> In Japan, *Fusarium azukicola* sp. nov. was isolated from azuki bean (*Vigna angularis*) and caused root-rot and typical SDS foliar symptoms when inoculated on soybean<sup>138</sup> <sup>□</sup>.

<sup>(b)</sup> *Fusarium* species isolated from soybean and other Fabaceae causing only root rot or root rot and foliar chlorosis, but not typical SDS symptoms on soybean.

<sup>(c)</sup> undescribed *Fusarium* sp. from FSSC clade 2 and strains in FSSC clade 5 and FSSC clade 11.

**Table 2. Efficacy of the management tools reported for SDS control**

SDS management practices	Efficacy (a)	Availability (b)	Reference
Fungicide seed treatments			
-fluopyram	+	USA	11, 48-50, 57, □ □62, 139 □
-pydiflumetofen	+	USA	51
-cyclobutrifluram	?	(c)	53-54 □
Tillage	?	ARG, BRA, USA	65-66 □
Late-planting (d)	+	ARG, BRA, USA	60
Short-term crop rotations	-	ARG, BRA, USA	78-80 □
Long-term crop rotations	+	ARG, BRA, USA	82-□ 83
Soil suppressiveness development	+	ARG, BRA, USA	30
Legume cover crops	-	ARG, BRA, USA	85 □
Other cover crops (grasses, Brassicaceae)	+	ARG, BRA, USA	85
Green manures	+	ARG, BRA, USA	86
Remote sensing methods	+	USA	84, □ 86-88
Disease risk prediction models	+	USA	38, 88 □
Plant nutrition	+	ARG, BRA, USA	28, □ 96, 97
Biological control	+	ARG, USA	100-105
Plant defense induction	+	under research	111-113 □
Host-induced gene silencing	*	under research	119-120 □
Genetic resistance			
	+	ARG, BRA, USA	124, 126-127 □

(a) — = not effective, + = effective, ? = mixed results, \* = expected results in the coming years.

(b) Management practices that are being used or are available.

(c) Registration and commercial release are planned for Latin America in 2021 / 2022.

(d) There is no conclusive evidence to recommend growers to sacrifice yield at late planting to combat SDS regarding early planting.

**Figure 1. Soybean sudden death syndrome disease cycle.** *Fusarium* overwinters in soybean and corn residues as macroconidia (stained with cotton blue) in sporodochia on root surface (A), early symptoms of root infection (B), late symptoms of root infection (C), leaf and pod symptoms caused by fungal toxins (D), reddish pigmentation is observed in basal stems at the end of the disease cycle (E), *F. tucumaniae* can overwinter as perithecia (F).





