



## Okara: A Nutritionally Valuable By-product Able to Stabilize *Lactobacillus plantarum* during Freeze-drying, Spray-drying, and Storage

#### Gabriel Quintana, Esteban Gerbino and Andrea Gómez-Zavaglia\*

Center for Research and Development in Food Cryotechnology (CCT-CONICET La Plata), Buenos Aires, Argentina

#### **OPEN ACCESS**

Edited by:

Juan Aguirre, Universidad de Chile, Chile

#### Reviewed by:

Evandro De Souza, Federal University of Paraíba, Brazil Vittorio Capozzi, University of Foggia, Italy

> \*Correspondence: Andrea Gómez-Zavaglia angoza@qui.uc.pt

#### Specialty section:

This article was submitted to Food Microbiology, a section of the journal Frontiers in Microbiology

Received: 24 September 2016 Accepted: 29 March 2017 Published: 12 April 2017

#### Citation:

Quintana G, Gerbino E and Gómez-Zavaglia A (2017) Okara: A Nutritionally Valuable By-product Able to Stabilize Lactobacillus plantarum during Freeze-drying, Spray-drying, and Storage. Front. Microbiol. 8:641. doi: 10.3389/fmicb.2017.00641 Okara is a nutritionally valuable by-product produced in large quantities as result of soymilk elaboration. This work proposes its use as both culture and dehydration medium during freeze-drying, spray-drying, and storage of Lactobacillus plantarum CIDCA 83114. Whole and defatted okara were employed as culture media for L. plantarum CIDCA 83114. The growth kinetics were followed by plate counting and compared with those of bacteria grown in MRS broth (control). No significant differences in plate counting were observed in the three media. The fatty acid composition of bacteria grown in whole and defatted okara showed a noticeable increase in the unsaturated/saturated (U/S) fatty acid ratio, with regard to bacteria grown in MRS. This change was mainly due to the increase in polyunsaturated fatty acids, namely C18:2. For dehydration assays, cultures in the stationary phase were neutralized and freeze-dried (with or without the addition of 250 mM sucrose) or spray-dried. Bacteria were plate counted immediately after freeze-drying or spray-drying and during storage at 4°C for 90 days. Freeze-drying in whole okara conducted to the highest bacterial recovery. Regarding storage, spraydried bacteria previously grown in whole and defatted okara showed higher plate counts than those grown in MRS. On the contrary, freeze-dried bacteria previously grown in all the three culture media were those with the lowest plate counts. The addition of sucrose to the dehydration media improved their recovery. The higher recovery of microorganisms grown in okara after freeze-drying and spray-drying processes and during storage was ascribed to both the presence of fiber and proteins in the dehydration media, and the increase in U/S fatty acids ratio in bacterial membranes. The obtained results support for the first time the use of okara as an innovative matrix to deliver L. plantarum. Considering that okara is an agro-waste obtained in large quantities, these results represent an innovative strategy to add it value, providing a symbiotic ingredient with promising industrial applications in the development of novel functional foods and feeds.

Keywords: okara, Lactobacillus plantarum, fatty acid composition, preservation process, storage

## INTRODUCTION

Okara is a white-yellowish puree remaining after filtration of the smashed soybeans seeds used for the production of soymilk (Stanojevic et al., 2013), and is produced in large quantities (about 1.1 kg per kilogram of soybean processed for soymilk production) (Li Y. et al., 2012). It is rich in high quality proteins, unsaturated lipids and dietary fiber, and also contains isoflavones, minerals and oligosaccharides (i.e., raffinose, stachyose) (Jimenez-Escrig et al., 2008; Song et al., 2009). Because of this nutritional richness, and the high water activity, okara is very prone to putrefaction. Although this agro-waste is generally used for animal feed, surplus is still burned as waste or dumped as landfill (Vong and Liu, 2016), which entails a significant disposal problem. For this reason, it is important to find novel strategies to add value to this material.

Several health benefits have been ascribed to okara, including antioxidant capacity, prevention of obesity and liver fat accumulation, decrease of cholesterol plasma levels and prevention of cardiovascular diseases (Jimenez-Escrig et al., 2008; Bedani et al., 2015; Vong and Liu, 2016). These properties make it a source of functional ingredients at a low cost (O'Toole, 1999; Li Y. et al., 2012). Okara was used as fermentation substrate to produce a variety of products for human consumption, including  $\alpha$ -glucosidase inhibitor, β-fructofuranosidase, fibrinolytic enzymes, iturin A, edible fungi, chitosan, alcohol, biosurfactants, among others (Kitawaki et al., 2007; Li B. et al., 2012; Li Y. et al., 2012; Villanueva-Suárez et al., 2013; Zhu et al., 2014; Vong and Liu, 2016). It was also used to supplement culture media or probiotic-containing products (Bedani et al., 2013, 2014). Inoculation of okara with lactic acid bacteria has been reported as an efficient approach to inhibit the growth of spoilage microorganisms (O'Toole, 1999) and to add value to this by-product (Lê et al., 2004; Espinosa-Martos and Rupérez, 2009; Villanueva-Suárez et al., 2013; Vong and Liu, 2016). Considering this, okara could be used as a cost-effective culture medium for the production of large quantities of starters at an industrial level.

Among lactic acid bacteria, *Lactobacillus plantarum* is a flexible and versatile species that is encountered in a variety of environmental niches, including some dairy, meat, and many vegetable or plant fermentations (Kleerebezem et al., 2003). Several strains of *L. plantarum* have demonstrated probiotic and technological properties. For this reason, the strains of this species have been used for a large number of applications, namely the formulation of fortified bread and pasta (Capozzi et al., 2016), for the fermentation starter cultures (Berbegal et al., 2016), for the fermentation of regional beverages (Yadav et al., 2016), as antibiotic replacement (Al-Madboly and Abdullah, 2015), and for the elimination of the

undesirable biogenic amines in wines (Capozzi et al., 2012), among others. In particular, *L. plantarum* CIDCA 83114, a strain isolated from kefir grains, have demonstrated both probiotic and technological properties, including the inhibition of growth and/or activity of *Escherichia coli* O157:H7 and *Salmonella* (Golowczyc et al., 2011b; Kakisu et al., 2013), and the overcoming of thermal and dehydrating processes (Romano et al., 2014; Tavera-Quiroz et al., 2015). These properties make it a good candidate to grow in okara and add value to this by-product.

The production of lactic acid bacteria as industrial starters requires adequate stabilization processes to avoid bacterial damage or death. In general, starters are delivered to food companies, or directly to consumers, under the form of ready to use and highly concentrated products. The commercialization of these concentrated cultures requires the application of preservation (or stabilization) processes to increase their shelflife. Among them, freeze-drying has been the most widely used process (Fonseca et al., 2015). Spray-drying is an alternative technique that conducts to the transformation of liquid systems into dry particulate powders when they come into direct contact with a drying medium (air) at high temperatures. It has been increasingly used to dehydrate lactic acid bacteria (Golowczyc et al., 2010, 2011a,b, 2013; Sosa et al., 2016). It is a cost-effective process, especially suitable for the production of large quantities of microorganisms (Golowczyc et al., 2010, 2011a,b, 2013; Sosa et al., 2016).

During the preservation processes, the decrease of water activity produces damages on the bacterial structures, decreasing their viability (Tymczyszyn et al., 2007a). To prevent these damages, sugars are generally used as protective agents (Carvalho et al., 2004), sucrose and trehalose being the most widely used among lactic acid bacteria. More recently more complex sugars, namely fructo and galacto-oligosaccharides, inuline, fiber, have been successfully used (Santos et al., 2014; Romano et al., 2015). Considering that okara includes about 50% of oligo and polysaccharides (Vong and Liu, 2016), its use as dehydration medium during preservation processes could provide protective compounds, avoiding the addition of external protectants.

Taking this background into account, the aim of this work was twofold: to use okara as culture medium for *L. plantarum* CIDCA 83114; and to evaluate the stabilizing capacity of okara during bacterial freeze-drying, spray-drying and storage at  $4^{\circ}$ C for 90 days.

#### MATERIALS AND METHODS

## Raw Material, Centesimal Composition, and Fourier Transform Infrared assays (FTIR)

Okara was obtained from Soyana S. H. (San Martín, Argentina). After being received, it was centrifuged five times to remove the excess of water. The sediment was frozen at  $-80^{\circ}$ C for 48 h and then, freeze-dried on a Heto FD4 equipment (Heto Lab Equipment, Denmark) for 48 h (temperature of condenser:  $-50^{\circ}$ C; chamber pressure: 0.04 mbar).

**Abbreviations:** a<sub>w</sub>, water activity; CFUs, cell forming units; cyc19, 0 cyclopropane fatty acid (methyleneoctadecanoic acid); d.b., dry basis; EDTA, ethylenediaminetetraacetic acid; FAMEs, fatty acids methyl esters; FTIR, Fourier transform infrared spectroscopy; MUFAs, monounsaturated fatty acids; PBS, phosphate saline buffer; PUFAs, polyunsaturated fatty acids; U/S, unsaturated/saturated.

Moisture content of the freeze-dried okara was determined by measuring the weight loss upon drying in a vacuum oven at 70°C until constant weight (Association of Official Analytical Chemists [AOAC], 1980). Total nitrogen was determined using the Kjeldahl method (Association of Official Analytical Chemists [AOAC], 1995) (conversion factor to transform nitrogen into protein: 6.25). Lipids were assessed by extraction with diethyl ether solvent in a Soxhlet system (Association of Official Analytical Chemists [AOAC], 1995). Ash content was determined by carbonization of the dried samples followed by incineration in a muffle furnace at 550°C. Dietary fiber was assessed using Megazyme testkit K-ACHDF 11/08 (Megazyme International Ireland Limited, Bray, Ireland) according to the manufacturer's procedure (McCleary, 2007). The centesimal composition was expressed in grams per 100 g of dry basis. The centesimal composition of defatted okara was determined after lipids' extraction (Association of Official Analytical Chemists [AOAC], 1995).

Fourier transform infrared spectroscopy spectra of freezedried whole and defatted okara were registered in the  $4000-500 \text{ cm}^{-1}$  range on KBr pellets, prepared in a 1:200 okara-KBr ratio. They were recorded in a transmission mode by co-adding 64 scans with 4 cm<sup>-1</sup> spectral resolution, on a Thermo Nicolet iS10 spectrometer (Thermo Scientific, Waltham, MA, USA).

Whole and defatted okara were suspended in distilled water to obtain 5% w/v suspensions. The suspensions were homogenized in an Ultra Turrax T25 (IKA, Staufen im Breisgau, Germany) and autoclaved for 15 min at 121°C. They were then used as culture media.

# Bacterial Strain, Culture Conditions, and Growth Kinetics

*Lactobacillus plantarum* CIDCA 83114 was isolated from kefir grains (Garrote et al., 2001) maintained frozen at  $-80^{\circ}$ C in 120 g/L non-fat milk solids (Difco, USA). Microbial cells were activated for 24 h in MRS broth (de Man et al., 1960) at 37°C in aerobic conditions. The resulting culture was inoculated (1% v/v) in fresh MRS broth and incubated in the same conditions. Cultures in the stationary phase [18 h incubation and  $\sim 1 \times 10^{13}$ colony forming units per mL (CFU/mL)] were used to inoculate 100 mL whole and defatted okara (inoculum concentration 2% v/v). They were then incubated at 37°C with shaking. In a parallel assay, microorganisms were grown in MRS broth in the same conditions (control). The three growth kinetics were followed by plate counting in MRS agar every 1 h, and results were expressed in CFU/mL.

#### **Extraction of Lipids from Bacteria**

Microorganisms were grown in okara and defatted okara at 37°C. Once attained the stationary phase, cultures were treated with 2% w/v EDTA pH 12 for 10 min to sequester calcium from okara and partly solubilize insoluble structures (Holt, 1982). After the treatment with EDTA, cultures were centrifuged, supernatants were discarded and bacterial pellets were used for lipids' extraction. Controls: bacteria in the stationary phase grown

in MRS, harvested by centrifugation, and washed with PBS [(K<sub>2</sub>HPO<sub>4</sub> 0.144 g/L; NaCl 9.00 g/L; Na<sub>2</sub>HPO<sub>4</sub> g/L), pH 7].

Lipids were extracted according to the modified Bligh and Dyer method (Marinetti, 1993). Briefly, bacterial pellets from microorganisms grown in whole and defatted okara and in MRS were suspended in chloroform-methanol-water (1:2:0.8 v/v/v) (4.75 mL per g of cells) for 12 h at 4°C, and then centrifuged at 8000 g for 10 min at 10°C. The supernatants were collected, and a second extraction on bacterial pellets was carried out. Both supernatants were mixed and chloroform-water (1:1) was added (12.5 mL per g of bacteria). The final mixture was centrifuged at 8000 g for 20 min. The chloroform phase was collected and dried under vacuum (Rotavapor<sup>®</sup> RE 120 – Büchi, Flawil, Switzerland). Lipids were dissolved in chloroform (final concentration 3 mg/mL) and stored at  $-20^{\circ}$ C for up to 2 weeks.

## **Characterization of Fatty Acids**

The FAME were prepared by adding 2 mL sulfuric acid (20 g/L in methanol) to 3 mg bacterial lipids, heating at 60°C for 2 h, extracting the esters with 1 mL chloroform-water (2:0.7 v/v) and washing twice with 0.7 mL water. The obtained FAME were analyzed on a gas chromatograph interfaced with a mass spectrometer detector (Shimadzu QP 5050A, Tokyo, Japan) using capillary column ZB-5 (30 m × 0.25 mm). The analysis conditions were: injection temperature 250°C, detector temperature 280°C, and column temperature initially 100°C increased to 280°C at 6°C min<sup>-1</sup>. The FAME were identified by mass spectrometry (GC–MS). The fatty acid composition was determined by considering the relative contribution of each peak area with regard to the sum of the areas corresponding to all peaks. Results were expressed as % of the total area.

## **Freeze-drying**

Microorganisms were grown in okara and defatted okara at 37°C. Aliquots of 1 mL containing L. plantarum CIDCA 83114 were collected in the stationary phase and neutralized with NaOH to pH 7. Then, they were transferred into 5 mL glass vials under aseptic conditions and frozen at -80°C for 48 h. The freeze-drying process was carried out at -50°C using a Heto FD4 freeze drier (Heto Lab Equipment, Denmark) and lasted 48 h. In parallel assays, sucrose was added to the neutralized okara (whole and defatted) containing L. plantarum CIDCA 83114 (final concentration of sucrose: 250 mM). Freeze-drying conditions were the same as those described above. In turn, microorganisms grown in MRS broth were harvested, neutralized and resuspended in 250 mM sucrose or in PBS. Freeze-drying process was carried out as described for bacteria grown in okara. Results were expressed as log N/N<sub>0</sub>, where N and N<sub>0</sub> were the CFU/mL after and before freeze-drying, respectively.

#### Spray-drying

In a parallel experiment, 100 mL of okara suspensions (whole and defatted) containing *L. plantarum* CIDCA 83114 in the stationary phase were neutralized and spray-dried in a laboratory-scale spray-dryer (Büchi B290 mini spray-dryer, Flawil, Switzerland) at a constant air inlet temperature of 180°C and an outlet temperature of 65–70°C. Microorganisms grown in MRS broth

were harvested neutralized and resuspended in PBS prior spraydrying. Results were expressed as  $\log N/N_0$ , where N and  $N_0$  were the CFU/mL after and before spray-drying, respectively.

#### **Storage Conditions**

After freeze-drying or spray-drying, samples were stored in silica gel desiccators at 4°C for 90 days. During storage, and at regular intervals, freeze-dried and spray-dried samples were rehydrated in PBS, homogenized for 1 min in a vortex mixer and maintained at room temperature for 30 min. Bacterial suspensions were serially diluted and plate counted on MRS agar, after incubation at  $37^{\circ}$ C for 48 h in aerobic conditions.

#### **Mathematical Modeling**

Storage plots were fitted using lineal and non-lineal mathematical regressions according to Eqs 1 and 2, respectively:

$$\log N/N_0 = -kt$$

where N is the CFU/mL at a given time of storage, N<sub>0</sub> is the CFU/mL at time equal to 0 (immediately after the preservation process), t is the time of storage (expressed in days) and k is the constant of bacterial inactivation (expressed in days<sup>-1</sup>).

Equation 2 describes non-lineal decays as:

$$\log N/N_0 = -A e^{-k-t} + A$$

where N is the CFU/mL at a given time of storage, N<sub>0</sub> is the CFU/mL at time equal to 0 (immediately after the process), t is the time of storage (expressed in days), k is the constant of bacterial inactivation (expressed in days<sup>-1</sup>) and A is the value of Log N/N<sub>0</sub> at the *plateau*.

#### Water Activity Measurements

The  $a_w$  of freeze-dried and spray-dried samples was determined using an Aqualab water activity instrument (Aqualab, Model Series 3TE, USA). The equipment was calibrated using standard salt solutions provided by the manufacturer.

#### Statistical Analysis and Reproducibility

All experiments were carried out on duplicate samples using three independent cultures of bacteria. The relative differences were reproducible independently on the cultures used. Analysis of variance (ANOVA) of cultivable bacteria corresponding to the different treatments was performed using the statistical program GraphPad Prism version 5.01 for Windows (GraphPad Software, Inc., San Diego, CA, USA 2007). Comparison of means by Tukey methods were tested, and if p < 0.05 the difference was considered statistically significant.

#### **RESULTS AND DISCUSSION**

The centesimal composition of whole and defatted okara is presented in **Table 1**, and was comparable to that reported for okara in other works (Vong and Liu, 2016). The FTIR spectra corresponding to whole and defatted okara are shown in **Figure 1**.



The main differences observed in the spectra are related with the lipid and carbohydrate bands. In fact, a clear decrease of the typical narrow  $\nu$ CH<sub>2</sub> bands corresponding to hydrocarbon lipid chains (2931 and 2860 cm<sup>-1</sup>) and the  $\nu$ C = O band at 1750 cm<sup>-1</sup> ascribed to the carbonyl groups of fatty acids, was observed in defatted okara. In addition, a noticeable increase of the relative intensity of the bands in the 1200–900 cm<sup>-1</sup> region, ascribed to the absorption of the C-O-C glycosidic linkage, was observed (Romano et al., 2014) (**Figure 1**). This reflects the higher contribution of fiber in defatted okara, which is consistent with the information shown in **Table 1**.

**Figure 2** depicts the growth kinetics of *L. plantarum* CIDCA 83114 in okara, defatted okara and MRS. No significant differences were observed for microorganisms grown in whole and defatted okara with regard to the control grown in MRS (p > 0.05), demonstrating that both whole and defatted okara are

TABLE 1   Centesimal composition of freeze-dried whole and defatted	
okara.	

Centesimal composition	Whole okara (g/100 g d.b.)	Defatted okara (g/100 g d.b.)	
Protein	$20.90 \pm 1.88$	$23.25 \pm 2.09$	
Lipids	13.40 ± 1.24 (a)	$3.67 \pm 0.34$ (b)	
Whole fiber	$54.51 \pm 2.43$	$60.64 \pm 2.70$	
Ash	$1.54 \pm 0.67$	$1.71 \pm 0.75$	
Humidity	$9.64\pm2.30$	$10.73\pm2.69$	

Different letters (a, b) denote statistically significant differences (p < 0.05).



as adequate as MRS to grow *L. plantarum* CIDCA 83114. Because of the large availability of okara, using okara as culture medium represents a clear advantage for the large scale production of starters, as costs can be substantially reduced. It has been reported that certain strains of lactobacilli and bifidobacteria ferment okara, and for this reason, okara has been proposed as a potential source of prebiotic fiber (Espinosa-Martos and Rupérez, 2009).

The composition of culture media strongly determines the composition of bacterial lipid membranes (Smittle et al., 1974; Tymczyszyn et al., 2005), and this latter is in turn related with their stability during preservation processes (Gomez-Zavaglia et al., 2000). For this reason, efforts were made to determine the fatty acid composition of okara lipids, which included C18:2 (54.61%), C18:1 (21.34%), C16:0 (11.57%), and C18:3 (7.97%) fatty acids, in agreement with the information published by Peñalvo et al. (2004). Taking into account the high contribution of C18:2 and C18:1, okara can be considered as a culture medium enriched in unsaturated fatty acids. Because of the high nutritional value and the technological properties, we have foreseen that okara oil can be used for different applications, thus remaining defatted okara as a final by-product. For this reason, and considering that defatted okara may be a real by-product in case okara oil is used as food ingredient, we decided to include it in this work.

**Table 2** shows the fatty acid composition of *L. plantarum* CIDCA 83114 grown whole and defatted okara and in MRS (control). *L. plantarum* CIDCA 83114 grown in MRS had mainly C16:0, C18:1 and cycC19:0, with an U/S ratio of 0.63. Growing microorganisms in okara induced noticeable changes in the fatty acid composition, mainly due to the increase in the polyunsaturated fatty acids (i.e., C18:2) (**Table 2**). The contribution of other fatty acids, like C16:1 and cycC19:0, strongly dropped for bacteria grown in okara. A moderate

TABLE 2 | Fatty acid composition of *Lactobacillus plantarum* CIDCA 83114 grown in MRS, whole and defatted okara.

Ratio	MRS	Whole okara	Defatted okara	
C14:0	$4.68 \pm 0.23$ (a1) <sup>a</sup>	$2.01 \pm 0.10$ (a1)	$1.00 \pm 0.05$ (a1)	
C16:0	31.75 ± 1.59 (a2)	$25.83 \pm 1.29$ (b2)	$21.10 \pm 1.06$ (b2)	
C16:1	$6.91 \pm 0.35$ (a3)	n.d. <sup>b</sup>	$0.63 \pm 0.03$ (b3)	
C18:0	$4.83 \pm 0.24$ (a4)	$11.08 \pm 0.56$ (b4)	$6.93 \pm 0.35$ (a4)	
C18:1	$30.09 \pm 1.52$ (a5)	$16.94 \pm 0.85$ (b5)	$20.00 \pm 1.00$ (b5)	
C18:2	0.67 ± 0.03 (a6)	44.14 ± 2.21 (b6)	$47.32 \pm 2.37$ (c6)	
cycC19:0	18.78 ± 0.94 (a7)	n.d.	$0.49 \pm 0.02$ (b7)	
U/S <sup>c</sup>	0.63	1.57	2.30	
PUFA/MUFA <sup>d</sup>	0.02	2.61	2.29	

<sup>a</sup>Analysis of variance has been carried out on rows. They were indicated as: a1, a2, a3, b1, b2, b3, etc. <sup>b</sup>n.d, not detected. <sup>c</sup>U/S: unsaturated/saturated ratio. Unsaturated fatty acids include C16:1, C18:1, and C18:2. Saturated fatty acids include C14:0, C16:0, C18:0, and cycC19:0. <sup>d</sup>PUFA/MUFA: polyunsaturated/monounsaturated ratio. Polyunsaturated fatty acids include C18:2 and monounsaturated fatty acids include C16:1 and C18:1.

decrease of C16:0 and C18:1 was also observed. As result of these changes, an inversion in the U/S ratio was observed for microorganisms grown in whole and defatted okara. Moreover, due to the high contribution of C18:2, the PUFA/MUFA ratio also increased with regard to the controls grown in MRS (Table 2). It has been reported that growing bacteria in media enriched with unsaturated fatty acids induces an increase in the U/S fatty acids ratio, which prevents lipids from being closely packed, and thus increases membrane fluidity (Muller et al., 2011; Hansen et al., 2015). Hansen et al. (2015) reported that the addition of 10 mg/L C18:2 to the growth medium promotes its incorporation into bacterial membranes. If considered that: (i) defatted okara had a lipid concentration of  $3.67 \pm 0.34$  g/100 g d.b. (Table 1), (ii) bacteria were grown in 5% w/v okara suspensions (see Materials and Methods) and (iii) okara fatty acids included 54.61% C18:2, it can be calculated that microorganisms grown in defatted okara were cultured in the presence of 1 mg/mL C18:2. This concentration was higher than that reported by Hansen et al. (2015), thus explaining the increase of PUFA/MUFA ratio in microorganisms grown in defatted okara (Table 2). The same rationalization can be carried out for microorganisms grown in whole okara, which explains the changes observed in the fatty acid composition (Table 2).

The effect of the growth and dehydration conditions on bacterial recovery was further analyzed. Freeze-dried microorganisms (with and without sucrose) had lower  $a_w$  than spray-dried ones, and bacteria grown in okara (whole and defatted), lower values than bacteria grown in MRS (**Table 3**). Bacteria grown in whole okara were those showing the lowest loss of viability after freeze-drying (left groups of bars in **Figure 3**). The log N/N<sub>0</sub> value for freeze-dried bacteria grown in defatted okara was significantly lower (p < 0.05), and comparable with that of bacteria grown in MRS and suspended in PBS (p > 0.05). The addition of sucrose in the dehydration media improved the recovery of microorganisms grown in defatted okara and MRS after freeze-drying (p < 0.05) and did not have a significant effect on bacteria grown in whole okara (p > 0.05) (middle group

Growth medium	Dehydration condition	a <sub>w</sub> a	Fitting		
			No significant decay	Linear <sup>b</sup>	Non-linear <sup>c</sup>
Whole okara	Freeze-drying	0.2544		k: 0.053 days <sup>-1</sup> ( <i>R</i> <sup>2</sup> : 0.9618)	
	Freeze-drying with sucrose	0.2466	X <sup>d</sup>		
	Spray-drying	0.3754	Х		
Defatted okara	Freeze-drying	0.2635			k: 0.04254 days <sup>-1</sup> A: 2.994 ( <i>R</i> <sup>2</sup> : 0.9239)
	Freeze-drying with sucrose	0.2525	Х		
	Spray-drying	0.3904			k: 0.03561 days <sup>-1</sup> A: 2.881 ( <i>R</i> <sup>2</sup> : 0.9536)
MRS	Freeze-drying	0.4141		k: 0.0780 days <sup>-1</sup> ( <i>R</i> <sup>2</sup> : 0.9419)	
	Freeze-drying with sucrose	0.2884		k: 0.035 days <sup>-1</sup> ( <i>R</i> <sup>2</sup> : 0.9622)	
	Spray-drying	0.4772			k: 0.1216 days <sup>-1</sup> A: 4.082 ( <i>R</i> <sup>2</sup> : 0.9193)

TABLE 3 | Water activity and inactivation constants of *L. plantarum* CIDCA 83114 grown in okara (whole and defatted) and MRS, and then freeze-dried (with and without 250 mM sucrose) or spray-dried.

<sup>a</sup>Water activity after the treatment. <sup>b</sup>Linear fitting is described by Eq. 1: Log N/N<sub>0</sub> = -kt. N = CFU/mL, where N<sub>0</sub> = CFU/mL at time equal to, t = time of storage in days;  $k = \text{constant of bacterial inactivation, expressed in days}^{-1}$  (see text). <sup>c</sup>Non-linear fitting is described by Eq. 2:  $y = -A e^{-k-t} + A$ , where N = CFU/mL at a given time of storage; N<sub>0</sub> = CFU/mL at time equal to 0; t = time of storage expressed in days,  $k = \text{constant of bacterial inactivation expressed in days}^{-1}$ ;  $A = \text{value of Log N/N<sub>0</sub> once attained the plateau. <sup>d</sup>X denote no significant decay during storage.$ 



of bars). It is interesting to note that the growth medium did not induce significant differences after freeze-drying with sucrose (p > 0.05) (middle group of bars). For the spray-drying process, the loss of viability for microorganisms grown in whole and defatted okara was significantly lower than that of bacteria grown in MRS and suspended in PBS (p < 0.05) (right group of bars).

The effect of the growth medium on the preservation processes was also analyzed. Microorganisms grown in whole okara showed

no significant differences in the Log N/N<sub>0</sub> after the three treatments (p > 0.05) (**Figure 3**). In turn, for bacteria grown in defatted okara and in MRS, spray-drying and freeze-drying with sucrose were significantly less harmful processes than freeze-drying without sucrose (p < 0.05).

The viability loss of L. plantarum CIDCA 83114 during storage at 4°C is shown in Figure 4. Freeze-drying was the most harmful process for L. plantarum CIDCA 83114 grown in the three culture media (Figure 4A). After 90 days of storage at 4°C, the viability of microorganisms grown in whole and defatted okara decayed  $4.62 \pm 0.25$  and  $2.65 \pm 0.06$  logarithmic units, respectively with regard to the corresponding viability obtained immediately after freeze-drying, considered as N<sub>0</sub> (time equal to 0). A significantly more deleterious effect was observed for microorganisms grown in MRS (<0.05), as a decrease of 7.10  $\pm$  0.14 logarithmic units was observed after 90 days of storage at 4°C. The addition of sucrose to the dehydration media had a protective effect during storage in all the three conditions assayed (Figure 4B). In this condition, microorganisms grown in whole and defatted okara did not show a significant decrease of viability with regard of the corresponding values obtained immediately after freeze-drying (p > 0.05) (Figure 4B). Although viability of microorganisms grown in MRS significantly decreased during storage (p < 0.05), this decrease was lower than that observed for bacteria grown in MRS and freeze-dried without sucrose (Figure 4A). In turn, spray-drying resulted an adequate process to stabilize bacteria grown in whole and defatted okara for 90 days at 4°C (Figure 4C). In this condition, no significant viability loss was observed for bacteria grown in whole okara (p < 0.05), and a loss of  $2.71 \pm 0.22$ logarithmic units, for bacteria grown in defatted okara. Growing bacteria in MRS before spray-drying induced the highest drop of viability (4.15  $\pm$  0.66 logarithmic units after 90 days of storage).

Polyhydroxy compounds, namely sugars and fiber, are generally used as protective compounds during freeze-drying, spray-drying and storage of lactic acid bacteria (Tymczyszyn et al., 2008; Santos et al., 2014; Romano et al., 2015). Fiber and oligosaccharides present in okara could improve the bacterial



FIGURE 4 | Log N/N<sub>0</sub> of *L. plantarum* CIDCA 83114 during storage. N<sub>0</sub>: viability obtained immediately after each preservation process (in log CFU/mL) and correspond to the values of N used in Figure 3. N: viability after each time of storage (log CFU/mL). Full black circles: bacteria grown in whole okara; gray squares: bacteria grown in defatted okara; white triangles: bacteria grown in MRS. (A) Freeze-drying; (B) Freeze-drying in the presence of sucrose; (C) Spray-drying. Dash lines show the mathematical regressions. In the cases of no significant decays (see Table 3), experimental values were just connected.

capacity to overcome freeze-drying and spray-drying processes and also stabilize microorganisms during storage (**Figures 3**, **4**) (Hincha et al., 2003; Sosa et al., 2016). The addition of sucrose in the dehydration medium improved the recovery of microorganisms grown in defatted okara and in MRS (**Figure 3**), which was an expected result considering the reported protective effect of sucrose during preservation of lactic acid bacteria (Tymczyszyn et al., 2007a,b). In this regard, it is worth to mention that although there are recent articles reporting the sucrose metabolism in *L. plantarum* strains (Chen et al., 2014; Mendoza-Llerenas et al., 2016), in this work, *L. plantarum* CIDCA 83114 is exposed to sucrose in non-growing conditions. Thus, sucrose is not hydrolyzed and acts as a protective agent.

The mathematical fitting of the storage graphs allowed the description of three different groups:

(a) no significant viability decays after 90 days of storage: this group included bacteria grown in whole okara and freeze-dried with sucrose or spray-dried, and microorganisms grown in defatted okara and freeze-dried with sucrose (**Figure 4** and **Table 3**).

(*b*) *linear decay after 90 days of storage:* this group included bacteria grown in okara and freeze-dried without sucrose, and those grown in MRS, freeze-dried with or without sucrose (**Table 3**). Eq. 1 describes the behavior of this group.

(c) non-linear decay after 90 days of storage: this group is composed of bacteria grown in defatted okara freeze-dried without sucrose or spray-dried, and bacteria grown in MRS and then, spray-dried (**Table 3**). Eq. 2 describes the behavior of this group.

For probiotic products, the European Food Safety Agency (EFSA) requires a bacterial viability of at least  $10^6-10^7$  CFU/g at the moment of being consumed (Hill et al., 2014). The *k* parameters obtained from Eqs 1 and 2 (**Table 3**) represent important tools to determine these values and in consequence, the shelf-life of *L. plantarum* CIDCA 83114 stabilized in okara matrices at the storage conditions.

#### CONCLUSION

Fermenting okara with *L. plantarum* CIDCA 83114 appears as an interesting strategy to deal with okara spoilage and disposal. In addition, dehydrating microorganisms in okara resulted in an adequate approach to stabilize *L. plantarum* CIDCA 83114. In particular, the successful stabilization of fermented okara by spray-drying appears as a cost-effective solution for the production of starters at an industrial level with a lowenergy consumption. The composition of okara had a strong influence on bacterial recovery after freeze-drying, spray-drying and storage: the presence of PUFAs induced an increase in the U/S and PUFA/MUFA ratios of lipid membranes, increasing membrane fluidity, and the fiber and oligosaccharides present in okara acted as protectants during dehydration and storage.

Using a by-product produced in large quantities as culture medium provides a cost-effective alternative for the production of starters at large scale. In addition, it provides a valuable solution for industrials, as they can solve two problems: okara disposal and availability of low-cost culture media. Regarding a nutritional perspective, okara provides high quality nutrients as vehicles of potentially probiotic microorganisms, whose importance in human and animal health is out of question. Taking this into account, the results support the use of okara matrices as suitable vehicles for probiotic strains, with promising applications in the development of novel functional foods and feeds.

## **AUTHOR CONTRIBUTIONS**

GQ and EG did the experimental work. EG and AG-Z conceived the work, analyzed and discussed results and wrote the manuscript.

#### REFERENCES

- Al-Madboly, L. A., and Abdullah, A. K. (2015). Potent antagonistic activity of Egyptian *Lactobacillus plantarum* against multiresistant and virulent foodassociated pathogens. *Front. Microbiol.* 6:347. doi: 10.3389/fmicb.2015.00347
- Association of Official Analytical Chemists [AOAC] (1980). Official Methods of Analysis of Association of Official Analytical Chemists, 13th Edn. Washington, DC: Association of Official Analytical Chemists.
- Association of Official Analytical Chemists [AOAC] (1995). Official Methods of Analysis Association of Official Analytical Chemists, 2 Vol. 16th Edn. Washington, DC: Association of Official Analytical Chemists.
- Bedani, R., Campos, M. M., Castro, I. A., Rossi, E. A., and Saad, S. M. I. (2014). Incorporation of soybean by-product okara and inulin in a probiotic soy yoghurt: texture profile and sensory acceptance. J. Sci. Food Agric. 94, 119–125. doi: 10.1002/jsfa.6212
- Bedani, R., Rossi, E. A., Cavallini, D. C. U., Pinto, R. A., Vendramini, R. C., Augusto, E. M., et al. (2015). Influence of daily consumption of synbiotic soybased product supplemented with okara soybean by-product on risk factors for cardiovascular diseases. *Food Res. Int.* 73, 142–148. doi: 10.1016/j.foodres.2014. 11.006
- Bedani, R., Rossi, E. A., and Isay Saad, S. M. (2013). Impact of inulin and okara on *Lactobacillus acidophilus* La-5 and *Bifidobacterium animalis* Bb-12 viability in a fermented soy product and probiotic survival under in vitro simulated gastrointestinal conditions. *Food Microbiol.* 34, 382–389. doi: 10.1016/j.fm. 2013.01.012
- Berbegal, C., Peña, N., Russo, P., Grieco, F., Pardo, I., Ferrer, S., et al. (2016). Technological properties of *Lactobacillus plantarum* strains isolated from grape must fermentation. *Food Microbiol.* 57, 187–194. doi: 10.1016/j.fm.2016. 03.002
- Capozzi, V., Menga, V., Digessù, A. M., De Vita, P., van Sinderen, D., Cattivelli, L., et al. (2011). Biotechnological production of vitamin B2-enriched bread and pasta. J. Agric. Food Chem. 59, 8013–8020. doi: 10.1021/jf201519h
- Capozzi, V., Russo, P., Ladero, V., Fernández, M., Fiocco, D., Alvarez, M. A., et al. (2012). Biogenic amines degradation by *Lactobacillus plantarum*: toward a potential application in wine. *Front. Microbiol.* 3:122. doi: 10.3389/fmicb.2012. 00122
- Carvalho, A., Silva, J., Hob, P., Teixeira, P., Malcata, F. X., and Gibbs, P. (2004). Relevant factors for the preparation of freeze-dried lactic acid bacteria. *Int. Dairy J.* 14, 835–847. doi: 10.1016/j.idairyj.2004.02.001
- Chen, C., Zhoua, F., Rena, J., Aia, L., Donga, Y., Wua, Z., et al. (2014). Cloning, expression and functional validation of a  $\beta$ -fructofuranosidase from *Lactobacillus plantarum. Proc. Biochem.* 49, 758–767. doi: 10.1016/j.procbio. 2014.02.013
- de Man, J. O., Rogosa, M., and Sharpe, M. E. (1960). A medium for the cultivation of lactobacilli. J. Appl. Bacteriol. 23, 130–135. doi: 10.1111/j.1365-2672.1960. tb00188.x
- Espinosa-Martos, I., and Rupérez, P. (2009). Indigestible fraction of okara from soybean: composition, physicochemical properties and in vitro fermentability

## FUNDING

This work was supported by the Argentinean Agency for the Scientific and Technological Promotion (ANPCyT) (Projects PICT/2014/0912 and PICT/2013/1258) and the Argentinean National Research Council (CONICET) (PIP2012-2014114-201101-00024).

#### ACKNOWLEDGMENTS

EG and AG-Z are members of the research career CONICET. GQ is doctoral fellow from CONICET. Dr. G. Kociubinski is especially acknowledged for his helpful suggestions and discussions.

by pure cultures of *Lactobacillus acidophilus* and *Bifidobacterium bifidum*. *Eur. Food Res. Technol.* 228, 685–693. doi: 10.1007/s00217-008-0979-7

- Fonseca, F., Cenard, S., and Passot, S. (2015). "Freeze-drying of lactic acid bacteria," in *Cryopreservation and Freeze-drying Protocols*, eds W. F. Wolkers and H. Oldenhof (New York, NY: Springer Science+Business Media). doi: 10.1007/ 978-1-4939-2193-5\_24
- Garrote, G. L., Abraham, A. G., and De Antoni, G. L. (2001). Chemical and microbiological characterisation of kefir grains. *J. Dairy Res.* 68, 639–652.
- Golowczyc, M. A., Gerez, C. L., Silva, J., Abraham, A. G., De Antoni, G. L., and Teixeira, P. (2011a). Survival of spray-dried *Lactobacillus kefir* is affected by different protectants and storage conditions. *Biotechnol. Lett.* 33, 681–686. doi: 10.1007/s10529-010-0491-6
- Golowczyc, M. A., Silva, J., Abraham, A. G., De Antoni, G. L., and Teixeira, P. (2010). Preservation of probiotic strains isolated from kefir by spray drying. *Lett. Appl. Microbiol.* 50, 7–12. doi: 10.1111/j.1472-765X.2009.02759.x
- Golowczyc, M. A., Silva, J., Teixeira, P., De Antoni, G. L., and Abraham, A. G. (2011b). Cellular injuries of spray-dried *Lactobacillus* spp. isolated from kefir and their impact on probiotic properties. *Int. J. Food Microbiol.* 144, 556–560. doi: 10.1016/j.ijfoodmicro.2010.11.005
- Golowczyc, M. A., Vera, C., Santos, M., Guerrero, C., Carasi, P., Illanes, A., et al. (2013). Use of whey permeate containing *in situ* synthesised galactooligosaccharides for the growth and preservation of *Lactobacillus plantarum*. *J. Dairy Res.* 80, 374–381. doi: 10.1017/S0022029913000356
- Gomez-Zavaglia, A., Disalvo, E. A., and De Antoni, G. L. (2000). Fatty acid composition and freeze-thaw resistance in lactobacilli. J. Dairy Res. 67, 241–247. doi: 10.1017/S0022029900004179
- Hansen, M. L. R. W., Petersen, M. A., Risbo, J., Hümmer, M., and Clausen, A. (2015). Implications of modifying membrane fatty acid composition on membrane oxidation, integrity, and storage viability of freeze-dried probiotic, *Lactobacillus acidophilus* La-5. *Biotechnol. Prog.* 31, 799–807. doi: 10.1002/btpr. 2074
- Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., et al. (2014). Expert consensus document: The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat. Rev. Gastroenterol.* 11, 506–514. doi: 10.1038/ nrgastro.2014.66
- Hincha, D. K., Zuther, E., and Heyer, A. G. (2003). The preservation of liposomes by raffinose family oligosaccharides during drying is mediated by effects on fusion and lipid phase transitions. *Biochim. Biophys. Acta* 1612, 172–177. doi: 10.1016/S0005-2736(03)00116-0
- Holt, C. (1982). Inorganic constituents of milk III. The colloidal calcium phosphate of cow's milk. J. Dairy Res. 49, 29–38. doi: 10.1017/S002202990002210X
- Jimenez-Escrig, A., Tenorio, M. D., Espinosa-Martos, I., and Ruperez, P. (2008). Health-promoting effects of a dietary fiber concentrate from the soybean byproduct okara in rats. J. Agric. Food Chem. 56, 7495–7501. doi: 10.1021/ jf800792y
- Kakisu, E., Abraham, A. G., Tironi Farinati, C., Ibarra, C., and De Antoni, G. L. (2013). Lactobacillus plantarum isolated from kefir protects vero cells from

cytotoxicity by type-II Shiga toxin from *Escherichia coli* O157:H7. *J. Dairy Res.* 80, 64–71. doi: 10.1017/S0022029912000659

- Kitawaki, R., Takagi, N., Iwasaki, M., Asao, H., Okada, S., and Fukuda, M. (2007). Plasma cholesterol-lowering effects of soymilk and okara treated by lactic acid fermentation in rats. J. Jpn. Soc. Food Sci. 54, 379–382. doi: 10.3136/nskkk. 54.379
- Kleerebezem, M., Boekhorst, J., van Kranenburg, R., Molenaar, W., Kuipers, O. P., Leer, R., et al. (2003). Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. U.S.A.* 100, 1990–1995. doi: 10.1073/pnas. 0337704100
- Lê, N. T., Champagne, C. P., Lee, B. H., and Goulet, J. (2004). Growth of Lactobacillus paracasei ssp. paracasei on tofu whey. Int. J. Food Microbiol. 89, 67–75. doi: 10.1016/S0168-1605(03)00109-0
- Li, B., Qiao, M., and Lu, F. (2012). Composition, nutrition, and utilization of okara (soybean residue). *Food Rev. Int.* 28, 231–252. doi: 10.1080/87559129. 2011.595023
- Li, Y., Hao, J., Cheng, Y., Zhao, R., Yin, L., and Li, L. (2012). Improvement of okara mouthfeel by *Aspergillus niger* and *Aspergillus oryzae* fermentation. *Trans. Chin. Soc. Agric. Eng.* 28, 248–253. doi: 10.3969/j.issn.1002-6819.2012. 07.041
- Marinetti, G. V. (1993). "Isolation, analysis and identification of lipids," in *Techniques of Lipidology*, eds M. Kates, H. R. Burdon, and P. H. Knippemberg (Amsterdam: Elsevier), 220–227. doi: 10.1016/S0075-7535(08) 70544-8
- McCleary, B. V. (2007). An integrated procedure for the measurement of total dietary fibre (including resistant starch), non-digestible oligosaccharides and available carbohydrates. *Anal. Bioanal. Chem.* 389, 291–308. doi: 10.1007/ s00216-007-1389-6
- Mendoza-Llerenas, E. O., Pérez, D. J., Gómez-Sandoval, Z., Escalante-Minakata, P., Ibarra-Junquera, V., Razo-Hernández, R. S., et al. (2016). Lactobacillus plantarum WCFS1 β-fructosidase: evidence for an open funnel-like channel through the catalytic domain with importance for the substrate selectivity. Appl. Biochem. Biotechnol. 180, 1056–1075. doi: 10.1007/s12010-016-2152-2
- Muller, J. A., Ross, R. P., Sybesma, W. F. H., Fitzgerald, G. F., and Stanton, C. (2011). Modification of the technical properties of *Lactobacillus johnsonii* NCC 533 by supplementing the growth medium with unsaturated fatty acids. *Appl. Environ. Microbiol.* 77, 6889–6898. doi: 10.1128/AEM.05213-11
- O'Toole, D. K. (1999). Characteristics and use of okara, the soybean residue from soy milk production: a review. J. Agric. Food Chem. 47, 363–371. doi: 10.1021/ jf9807541
- Peňalvo, J. L., Castilho, M. C., Silveira, M. I. N., Matallana, M. C., and Torija, M. E. (2004). Fatty acid profile of traditional soymilk. *Eur. Food Res. Technol.* 219, 251–253. doi: 10.1007/s00217-004-0945-y
- Romano, N., Tavera-Quiroz, M. J., Bertola, N., Mobili, P., Pinotti, A., and Gómez-Zavaglia, A. (2014). Edible methylcellulose-based films containing fructooligosaccharides as vehicles for lactic acid bacteria. *Food Res. Int.* 64, 560–566. doi: 10.1016/j.foodres.2014.07.018
- Romano, N., Tymczyszyn, E., Mobili, A., and Gómez-Zavaglia, A. (2015). "Chapter 10: Prebiotics as protectants of lactic acid bacteria," in *Probiotics, Prebiotics, and Synbiotics: Bioactive Foods in Health Promotion, Part 1: Prebiotics in Health Promotion,* 2nd Edn, eds R. R. Watson and V. R. Preedy (Cambridge, MA: Academic Press), 155–164. doi: 10.1016/B978-0-12-802189-7.00010-1
- Santos, M., Gerbino, E., Araujo-Andrade, C., Tymczyszyn, E. E., and Gomez-Zavaglia, A. (2014). Stability of freeze-dried *Lactobacillus delbrueckii* subsp. *bulgaricus* in the presence of galacto-oligosaccharides and lactulose as determined by near infrared spectroscopy. *Food Res. Int.* 59, 53–60. doi: 10. 1016/j.foodres.2014.01.054

- Smittle, R. B., Gilliland, S. E., Speck, M. L., and Walter, W. M. Jr. (1974). Relationship of cellular fatty acid composition to survival of *Lactobacillus bulgaricus* in liquid nitrogen. *Appl. Microbiol.* 27, 738–743.
- Song, D., Chang, S. K. C., and Ibrahim, A. A. (2009). Effect of fermentation substrates on enzyme production and degradation of oligosaccharides in pinto bean flour as affected by particle size. *J. Food Process. Pres.* 33, 527–546. doi: 10.1111/j.1745-4549.2008.00304.x
- Sosa, N., Gerbino, E., Golowczyc, M. A., Schebor, C., Gómez-Zavaglia, A., and Tymczyszyn, E. E. (2016). Effect of galacto-oligosaccharides: maltodextrin matrices on the recovery of *Lactobacillus plantarum* after spray-drying. *Front. Microbiol.* 7:584. doi: 10.3389/fmicb.2016.00584
- Stanojevic, S. P., Barac, M. B., Pesic, M. B., Jankovic, V. S., and Vucelic-Radovic, B. V. (2013). Bioactive proteins and energy value of okara as a byproduct in hydrothermal processing of soy milk. *J. Agric. Food Chem.* 61, 9210–9219. doi: 10.1021/jf4012196
- Tavera-Quiroz, M. J., Romano, N., Mobili, P., Pinotti, A., Gomez-Zavaglia, A., and Bertola, N. (2015). Green apple baked snacks functionalized with edible coatings of methylcellulose containing *Lactobacillus plantarum*. J. Funct. Foods 16, 164–173. doi: 10.1016/j.jff.2015.04.024
- Tymczyszyn, E. E., Díaz, M. R., Gómez-Zavaglia, A., and Disalvo, E. A. (2007a). Volume recovery, surface properties and membrane integrity of *Lactobacillus delbrueckii* subsp. *bulgaricus* dehydrated in the presence of trehalose or sucrose. J. Appl. Microbiol. 103, 2410–2419. doi: 10.1111/j.1365-2672.2007.03482.x
- Tymczyszyn, E. E., Díaz, R., Pataro, A., Sandonato, N., Gómez-Zavaglia, A., and Disalvo, E. A. (2008). Critical water activity for the preservation of *Lactobacillus bulgaricus* by vacuum drying. *Int. J. Food Microbiol.* 128, 342–347. doi: 10.1016/ j.ijfoodmicro.2008.09.009
- Tymczyszyn, E. E., Gomez-Zavaglia, A., and Disalvo, E. A. (2005). Influence of the growth at high osmolality on the lipid composition, water permeability and osmotic response of *Lactobacillus bulgaricus*. Arch. Biochem. Biophys. 443, 66–73. doi: 10.1016/j.abb.2005.09.004
- Tymczyszyn, E. E., Gómez-Zavaglia, A., and Disalvo, E. A. (2007b). Effect of sugars and growth media on the dehydration of *Lactobacillus delbrueckii* ssp. *bulgaricus. J. Appl. Microbiol.* 102, 845–851. doi: 10.1111/j.1365-2672.2006. 03108.x
- Villanueva-Suárez, M. J., Pérez-Cózar, M. L., and Redondo-Cuenca, A. (2013). Sequential extraction of polysaccharides from enzymatically hydrolyzed okara byproduct: physicochemical properties and in vitro fermentability. *Food Chem.* 141, 1114–1119. doi: 10.1016/j.foodchem.2013.03.066
- Vong, W. C., and Liu, S. Q. (2016). Biovalorisation of okara (soybean residue) for food and nutrition. *Trends Food Sci. Technol.* 52, 139–147. doi: 10.1016/j.tifs. 2016.04.011
- Yadav, R., Puniya, A. K., and Shukla, P. (2016). Probiotic properties of *Lactobacillus plantarum* RYPR1 from an indigenous fermented beverage Raabadi. *Front. Microbiol.* 7:1683. doi: 10.3389/fmicb.2016.01683
- Zhu, Y., Li, X., Liu, R., and Li, H. (2014). Antioxidant activity and nutritional characterization of okara fermented by various microorganisms. J. Chin. Inst. Food Sci. Tech. 14, 20–27. doi: 10.1080/87559129.2011.595023

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Quintana, Gerbino and Gómez-Zavaglia. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.