



First record of *Neotomys ebriosus* Thomas, 1894 (Mammalia: Rodentia: Sigmodontinae) from Tucuman province, northwestern Argentina

Ignacio Ferro^{1, 2, 3} & Ruben M. Barquez²

¹ CONICET – Universidad Nacional de Jujuy, Instituto de Ecorregiones Andinas (INECOA), Av. Bolivia 1239, CP: 4600, San Salvador de Jujuy, Jujuy, Argentina

² CONICET– Universidad Nacional de Tucumán, Programa de Investigaciones de Biodiversidad Argentina (PIDBA), Facultad de Ciencias Naturales e Instituto Miguel Lillo, Miguel Lillo 205, CP: 4000, San Miguel de Tucumán, Tucumán, Argentina

³ Corresponding author. E-mail: ignacioferro@gmail.com

Abstract. *Neotomys ebriosus* Thomas, 1894 (Mammalia: Rodentia: Sigmodontinae) is a monotypic genus of sigmodontine rodent restricted to the Central Andes of South America. Its distribution is inferred from a few localities; although some maps include Tucumán province, in northwestern Argentina, within the range of this species, previous records are not known. Here, we report from Cerro Muñoz (Tafí de Valle Department) the first documented record for the province.

Key words. Andean Swamp Rat; cloud grasslands; distribution; Puna; Aconquija

The Andean Swamp Rat, *Neotomys ebriosus* Thomas, 1894 (Mammalia: Rodentia: Cricetidae) (Fig. 1), is a rodent endemic to the Central Andean highlands, ranging from about 10° S in central Peru southward throughout western Bolivia and northern Chile to northwestern Argentina at about 29° S (ORTIZ & JAYAT 2015). However, it is only known from a few specimens and localities, and thus, much about its distribution, natural history, and patterns of geographic variation are still poorly understood (ORTIZ & JAYAT 2015). Until recently, this monotypic genus was considered as “incertae sedis” in the suprageneric classification of sigmodontinae rodents (D’ELÍA et al. 2006, 2007, 2015), but they are now known to constitute a well-supported “Andean clade”, together with *Euneomys* and *Irenomys* according to molecular data (MARTÍNEZ et al. 2012). Thus, this clade was included in a new tribe, Euneomyini (PARDIÑAS et al. 2015). In Argentina *N. ebriosus* is known from Jujuy, Salta, Catamarca, La Rioja and San Juan provinces (BARQUEZ 1989, JAYAT et al. 2011). Herein, we provide the first reliable record from Tucumán province of this enigmatic rodent and refine the known distribution of this species.

Our new record is from Cañada del Fuerte Viejo, Cerro Muñoz, (26°53'41.00" S, 065°48'40.32" W; 3300 m above sea level), Tafí del Valle Department, Tucumán province, Argentina (Fig. 2). The specimen was captured during a field trip on 14 July 2016 in a Sherman-like trap set in a 5 × 5 trap grid (25 traps, each separated by 15 m), located on a cushion peat-

land covered by bunch grass. The specimen was preserved as a skin and skeleton under the field number LIF 950 (Luis Ignacio Ferro field catalogue) and deposited at the Colección Mamíferos Lillo (CML 10859), Facultad de Ciencias Naturales e Instituto Miguel Lillo, University of Tucumán.

Neotomys ebriosus is distinctive because of its strongly rufous muzzle, the presence of hairs of the same color at the base of the ears and on the rump, the length of the tail shorter than the head and body combined, and by the characteristic longitudinal grooves along the labial borders of the upper incisors (Fig. 1). Another species of rodent having a rufous muzzle and short tail is *Abrothirx jelskii* (Thomas, 1894), which is sympatric with *N. ebriosus* at several localities along their ranges. However, *A. jelskii* is much smaller, has a marked contrast between dorsal (gray) and ventral (white) pelage, and has white postauricular patches which are absent in *N. ebriosus*. In *Neotomys* the ventral hairs are plumbeous at the bases and white at the tips and thus conferring an overall



Figure 1. Photograph of the living specimen of *Neotomys ebriosus* (CML 10859), showing some distinctive characters of the species, as the rufous muzzle and ears, and the short tail. Inserted is a photograph of the distinctive upper incisors grooves.




Figure 2. Localities of occurrence for *Neotomys ebriosus* in northwestern Argentina. Dots indicate previous published records. 1: Yavi, Jujuy province (PARDIÑAS & ORTIZ 2001); 2: Sierra de Tilcara, Jujuy province (DÍAZ & BARQUEZ 2007); 3: Valle Encantado, Salta province (PARDIÑAS & ORTIZ 2001); 4: Paycuqui, Catamarca province (specimen revised by the authors CML 6893); 5: Barranca Larga, Catamarca province (PARDIÑAS & ORTIZ 2001); 6: Famatina range, La Rioja province (JAYAT et al. 2011); 7: San Guillermo National Park, San Juan province (BARQUEZ 1983). The star indicates the new record from Tucumán province. The gray area in the insert map shows the entire distributional range of *Neotomys ebriosus*.

grizzled greyish color with a gradual contrast between dorsal and ventral regions. Among cranio-dental characters, a longitudinal groove along the labial border of the upper incisors is characteristic. For further information on this species, see THOMAS (1894), SANBORN (1947), STEPPAN (1995), PARDIÑAS & ORTIZ (2001), and ORTIZ & JAYAT (2015).

The presence of *N. ebriosus* in Tucumán province was known only by a fossil mandible exhumed from Pleistocene sediments in La Angotura (Tafi del Valle department), 1900

m (PARDIÑAS & ORTIZ 2001). During the Pleistocene glacial maximum, down-slope migration of the vegetation belt would have allowed for the presence of *N. ebriosus* at lower elevations. However, high-altitude habitats in Tucumán are inaccessible by road, hindering knowledge of the sigmodontine fauna in the highlands of the province (FERRO & BARQUEZ 2008). Our new record is especially significant because high-altitude environments and the sigmodontine fauna are among the least studied in northwestern Argentina and particularly in Tucumán province. The nearest known occurrence of *N. ebriosus* is probably that reported by THOMAS (1926) from the “Aconquija Mountain, Catamarca”, but the exact location of this record remains unknown and could be anywhere along the approximately 150 km long mountain range. Our new data adds *N. ebriosus* to the list of living sigmodontine rodent species of Tucumán province; it refines this species’ known distribution with an extension of 100 km to the east from the closest known record (Barranca Larga, Belén Department, Catamarca). The specimen we report here was captured in a cloud grassland belonging to the Yungas ecoregion, together with tree other sigmodontine rodents: *Akodon spegazzinii* THOMAS, 1897, *Phyllotis tucumanus* THOMAS, 1912 and *Andinomys edax* THOMAS, 1902

ACKNOWLEDGEMENTS

We appreciate the support from CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina)—for supporting grants to RMB and LIF and several fellowships to LIF—as well as the Agencia Nacional de Promoción Científica y Tecnológica for a grant (PICT 2011 N°0322) and SAREM (Sociedad Argentina para el Estudio de los Mamíferos) for “the Osvaldo Reig Award 2007” given to LIF. We also thank the PIDBA team, particularly Mónica Díaz, Fernanda López Berrizbeitia, and Pablo Gaudioso for logistical support and curation of our specimen at the CML. We are grateful to an anonymous reviewer. Finally, we appreciate the fundamental help of the Nieva family and Pocho Roldán from Tafi del Valle.



Figure 3. Photograph of the collecting area of *Neotomys ebriosus* (CML 10859) in Tucumán province. Biogeographically corresponds to the uppermost vegetation formation of the yungas forest: the cloud grasslands on the eastern slopes of the central Andes. The specimen was captured among the bunch grasses next to the riverside.

LITERATURE CITED

- BARQUEZ, R.M. 1983. La distribución de *Neotomys ebriosus* Thomas en la Argentina y su presencia en la provincia de San Juan (Mammalia, Rodentia, Cricetidae). *Historia Natural* 3: 189–191.
- DÍAZ, M.M. & BARQUEZ, R.M. 2007. The wild mammals of Jujuy province, Argentina: systematics and distribution; pp. 417–578, in: KELT, D.A., LESSA, E., SALAZAR-BRAVO, J.A. & PATTON, J.L. (eds.). *The quintessential naturalist: honoring the life and legacy of Oliver P. Pearson*. University of California Publications, Zoology 134.
- D'ELÍA, G. 2015. Sigmodontinae incertae sedis; pp. 70–73, in: PATTON, J.L., PARDIÑAS, U.F.J. & D'ELÍA, G. (eds.). *Mammals of South America, Volume 2: rodents*. Chicago/London: The University of Chicago Press.
- D'ELÍA, G., TETA, P. & PARDIÑAS, U.F.J. 2006. Incertae sedis; pp. 197–202, in: Barquez, R.M., Díaz, M.M. & Ojeda, R.A. (eds.). *Mamíferos de Argentina: sistemática y distribución*. Tucumán: Sociedad Argentina para el Estudio de los Mamíferos.
- D'ELÍA, G., Pardiñas, U.F.J., Teta, P. & Patton, J.L. 2007. Definition and diagnosis of a new tribe of sigmodontine rodents (Cricetidae: Sigmodontinae), and a revised classification of the subfamily. *Gayana* 71: 187–194. <https://doi.org/10.4067/S0717-65382007000200007>
- FERRO, L.I. & BARQUEZ, R.M. 2008. Comentarios sobre la distribución de *Abrothrix andinus* y *Calomys lepidus* (Rodentia: Cricetidae) en la provincia de Tucumán, Argentina. *Mastozoología Neotropical* 15: 197–201.
- JAYAT, J.P., ORTIZ, P.E., GONZÁLEZ, R., LOBO ALLENDE, R. & MADOZZO JAÉN, M.C. 2011. Mammalia, Rodentia, Sigmodontinae Wagner, 1843: new locality records, filling gaps and geographic distribution maps from La Rioja province, northwestern Argentina. *Check List* 7: 614–618. <https://doi.org/10.15560/7.5.614>
- MARTINEZ, J.J., FERRO, L.I., MOLLERACH, M.I. & BARQUEZ, R.M. 2012. The phylogenetic relationships of the Andean swamp rat genus *Neotomys* (Rodentia, Cricetidae, Sigmodontinae) based on mitochondrial and nuclear markers. *Acta Theriologica* 55: 277–287. <https://doi.org/10.1007/s13364-011-0070-9>
- ORTIZ, P.E. & JAYAT, J.P. 2015. Genus *Neotomys*; pp. 99–101, in: PATTON, J.L. PARDIÑAS, U.F.J. & D'ELÍA, G. (eds.). *Mammals of South America. Volume 2: rodents*. Chicago/London: The University of Chicago Press.
- PARDIÑAS, U.F.J., TETA, P. AND SALAZAR-BRAVO, J. 2015. A new tribe of sigmodontinae rodents (Cricetidae). *Mastozoología Neotropical* 22: 171–186.
- PARDIÑAS, U.F.J. & ORTIZ, P.E. 2001. *Neotomys ebriosus*, an enigmatic South American rodent (Muridae, Sigmodontinae): its fossil record and present distribution in Argentina. *Mammalia* 65: 244–250.
- SANBORN, C.C. 1947. The South American rodents of the genus *Neotomys*. *Fieldiana Zoology* 11(7): 51–57. <https://doi.org/10.5962/bhl.title.5567>
- STEBBINS, S.J. 1995. Revision of the tribe Phyllotini (Rodentia: Sigmodontinae), with a phylogenetic hypothesis for the Sigmodontinae. *Fieldiana, Zoology, New Series* 80: 1–112. <https://doi.org/10.5962/bhl.title.3336>
- THOMAS, O. 1894. Descriptions of some new Neotropical Muridae. *Annals and Magazine of Natural History, including Zoology, Botany and Geology* (ser. 6) 14: 346–366. <https://doi.org/10.1080/00222939408677813>
- THOMAS, O. 1926. The Spedan Lewis South American Exploration.— IV. List of mammals obtained by Sr. Budin on the boundary between Jujuy and Bolivia. *Annals and Magazine of Natural History, including Zoology, Botany and Geology* (ser. 9) 18: 193–195. <https://doi.org/10.1080/00222932608633494>

Authors' contributions. IF collected the data, revised specimens in collections, and wrote the first version of the manuscript RMB revised specimens in collections, read, made suggestions and accepted the last version of the manuscript.

Received: 27 August 2016

Accepted: 24 March 2017

Academic editor: Terrance C. Demos