

Research Article

Fermented Milks from Small Ruminant: Effect on Metabolism and Immune Status of Mice Fed Mild Caloric Restricted Diet

Fabersani E¹, Torres S^{2,3}, Valdez C², González S^{1,2} and Gauffin Cano P^{1,4*}

¹Centro de Referencia Para Lactobacilos (CERELA) – CONICET, Argentina

²National University of Tucumán, Argentina

³Institute of Chemistry NOA – CONICET, Argentina

⁴Universidad Del Norte Santo Tomás de Aquino, Argentina

*Corresponding author: Gauffin Cano P, Universidad Del Norte Santo Tomás de Aquino, CERELA, Chacabuco 145 (4000) Tucumán, Argentina, Tel: 054-381-4310465; Fax: 054-381-4310465; Email: pgauffin@cerela.org.ar

Received: August 14, 2014; Accepted: September 16, 2014; Published: September 18, 2014

Abstract

The purpose of this study was to evaluate the effect of the administration of Fermented Goat's Milks (FGMs) in mice fed mild caloric restricted diet. The ability of these FGMs to ameliorate immune and metabolic parameters related to mild caloric restriction was studied. We also analyzed if the time (45 or 90 days) of caloric restriction have influence on leptin secretion, IgA+ cells number and phagocytic activity of peritoneal macrophages, and also test the correlation between these parameters. Mice were fed with a mild caloric restricted diet during 45 or 90 days. After these periods of caloric restriction, mice were refed with balanced conventional diet (BCD) plus goat's milks or BCD plus goat's milks fermented with *Lactobacillus rhamnosus* CRL1425 (FGM-Lr) or *Lactobacillus casei* CRL431 (FGM-Lc). All renutrition diets induced an increase of serum glucose, triglycerides, total proteins and leucocytes cells. A decrease in cholesterol levels was observed after the diets. FGM-Lr induced lower triglycerides values than others diets. All FGMs restore IgA+ cells in intestinal mucosa, and the FGM-Lc group had higher positive cells number than the *ad libitum* control. Furthermore, phagocytic activity of peritoneal macrophages only increased in mice fed with FGM-Lc. Mild caloric restriction induced a decrease on leptin circulating values at 90 days. The renutrition with FGM-Lr lead to lower leptin levels than mild caloric restriction controls, while FGM-Lc induced higher leptin levels. Positive correlation between serum leptin concentration and immune parameters was observed in all groups under study. We showed that leptin levels could positively predict the immune mucosal competence in mild caloric restricted mice. These results suggest that the FGMs are able to modulate, in different way, serum leptin levels, IgA+ cells number and phagocytic activity. Furthermore, FGM-Lc could be more effective for nutritional treatment in malnutrition status.

Keywords: Mild caloric restriction; Leptin renutrition; Goat's milk; Probiotic

Abbreviations

GM: Goat's Milk; FGM: Fermented Goat's Milk; FGM-Lr: Fermented Goat's Milk with *Lactobacillus Rhamnosus* CRL1425; FGM-Lc: Fermented Goat's Milk with *Lactobacillus casei* CRL431; BCD: Balanced Conventional Diet; AD: *Ad libitum* control group;

CR: Caloric Restricted group. Group of mice which after diet adaptation period (2days) received 75% of BCD; AD45: *Ad libitum* group of 45 days; AD90: *Ad libitum* group of 90 days; CR45: group of mice which after diet adaptation period (2days) received 75% of BCD during 45 days; CR90: group of mice which after diet adaptation period (2days) received 75% of BCD during 90 days; CR/Re: Caloric Restricted / Renourished group; CR45/Re-Lr: Caloric Restricted mice during 45 days were renourished with *ad libitum* BCD plus goat's milk during 7 days followed by 5 days in which animals were fed with BCD plus FGM with *Lactobacillus rhamnosus* CRL1425; CR45/Re-Lr: Caloric Restricted mice during 45 days were renourished with *ad libitum* BCD plus goat's milk during 7 days followed by 5 days in which animals were fed with BCD plus FGM with *Lactobacillus rhamnosus* CRL1425; CR90/Re+Lr: Caloric Restricted mice during 90

days were renourished with *ad libitum* BCD plus goat's milk during 7 days followed by 5 days in which animals were fed with BCD plus FGM with *Lactobacillus rhamnosus* CRL1425; CR45/Re-Lc: Caloric Restricted mice during 45 days were renourished with *ad libitum* BCD plus goat's milk during 7 days followed by 5 days in which animals were fed with BCD plus FGM with *Lactobacillus casei* CRL431; CR90/Re-Lc: Caloric Restricted mice during 90 days were renourished with *ad libitum* BCD plus goat's milk during 7 days followed by 5 days in which animals were fed with BCD plus FGM with *Lactobacillus casei* CRL431; CR45/Re-GM: Caloric Restricted mice during 45 days were renourished with *ad libitum* BCD plus whole goat's milk during 12 days; T: effect of time determinate by ANOVA test; D: effect of diet determinate by ANOVA test

Introduction

Clinically, malnutrition is characterized by inadequate intake of protein, energy and micronutrients. Generally, it is associated with frequent infections and disorders. Currently, nearly 12% of the global population is estimated to be undernourished, and the vast majority of these live in developing countries [1], therefore, the management

of undernourishment has continued being a public health priority.

According to the WHO child growth standards, underweight is defined as weight-for-age below 2 SD the median [2]. Actually, there is a prevalence of stunting and underweight among children under-five years of age worldwide. This situation leads to millions of children still remain at risk [2]. Children with mild malnutrition have an increased risk of mortality. If some of these children do not receive adequate support, they may progress towards a severe acute malnutrition or a severe stunting.

The adaptation to caloric restriction diet is characterized by metabolic, endocrine, and immunologic changes. There is robust evidence that leptin could be a sensitive marker of nutritional status [3]. It is directly correlated with several biochemical and anthropometric parameters [3]. Leptin, the product of the obesity gene, is a 16-kDa circulating hormone that has been recognized to have a major influence on energy balance [4]. Leptin is a mediator of long-term regulation of energy balance via hypothalamic-mediated effects. This hormone suppresses food intake, by inhibiting orexigenic neuropeptides and stimulating anorexigenic ones, as well as increases energy expenditure [5,6]. The presence of leptin receptor in multiple biologic systems, as well as in various immune cells, suggest this hormone also intervene in the modulation of immune function [7]. During caloric restriction the loss of body fat results in profound reductions in circulating levels of leptin, amongst other adipokines [8]. However, the decline in leptin in response to caloric restriction appears to be partially dependent on other factors, such as the composition of the diet [9,10]. Also, intensity and duration of caloric restriction seems to affect leptin drop. In a previous study we showed that a short-term (12 days) mild calorie-restricted diet (set at around a 25 % reduction in energy consumption) does not significantly modify serum leptin levels in mice [11].

It is known that nutritional deficiencies can adversely affect the number and activity of immune cells leading to a depression of the immune system [12,13]. Reduced body fat or nutritional deprivation, typically associated with hypoleptinemia, is a direct cause of secondary immunodeficiency and increased susceptibility to infection [4]. Matarese *et al.* suggested that leptin plays an important role in the regulation of the immune system in energy- or leptin-deficient states [14]. The malnutrition could be reversed with an appropriate renutrition [12,15,16]. In this regard, fermented dairy products could be useful in malnourished groups [17]. Probiotic dairy products are widely accepted as health products and included within functional foods [17]. Probiotics defined as live microorganisms that, when provided in adequate amounts, confer a health benefit on the host, are included as important components of the daily diet [17]. In this context, there is sufficient scientific evidence demonstrating that probiotic fermented milks are responsible for restoring gastrointestinal and immune functions in moderate malnutrition [12,15,16].

Milks from small ruminants, such as goat's milk, are an important and cheap source of milk and especially in tropical countries, and could be an alternative for nutritional treatment [11,18,19]. The nutritive quality of milk is determined by its composition. These conventional milks differ in lipid, protein, carbohydrate, vitamin and mineral contents, and thus in their nutritional properties,

compared with cow's milk [2,11]. Investigations have shown that small ruminant milks, and particularly goat's milk, have functional benefits for the treatment of malnutrition [11,18]. However, little is known regarding the potential role of fermented goat's milks on metabolism and immune status during mild malnutrition recovery. Therefore, in this study we evaluate the effect of the administration of Fermented Goat's Milks (FGMs) in mice fed caloric restricted diet. The ability of these fermented milks to ameliorate immune and metabolic parameters related to mild caloric restriction was studied. We also analyzed if the time (45 or 90 days) of caloric restriction have influence on leptin secretion.

Material and Methods

Bacterial strains, media and culture conditions


Two lactic acid bacteria strains were selected as probiotic cultures for fermentation of goat's milk due to their source or ability to stimulate the immune system. *Lactobacillus rhamnosus* CRL1425 and *Lactobacillus casei* CRL431 were obtained from Centro de Referencia para Lactobacilos (CERELA, Tucumán, Argentina) culture collection. *L. rhamnosus* CRL1425 is a microorganism with proved technological properties which was isolated from goat's cheese [20]. *L. casei* CRL431 was isolated from intestine of healthy children and has extensively showed immunomodulatory properties in malnourished hosts [12,15,16]. The cultures were stored at -70°C and activated in Man-Rogosa-Sharpe (MRS) broth (1% v/v inoculums). The cells were harvested by centrifugation at 3,000 g for 10 min and washed three times with sterile 0.01 mol/L Phosphate-Buffered Saline (PBS), pH 7.2 for development of fermented goat's milks (FGMs).

Preparation of goat's milks fermented with *Lactobacillus rhamnosus* CRL1425 or *Lactobacillus casei*

CRL431: Fresh, non-fortified, whole goat's milk was provided by INTA "Santa Cruz" (Catamarca, Argentina). The microbiological quality of the milk was assessed according to Salva *et al.* [18]. Goat's milk fat content, protein and non-fat solids were determined with a milk analyzer (Ekomilk M). Before use, the goat's milk was pasteurized (85°C , 15 min) and cooled to 42°C . Goat's milk was fermented with *Lactobacillus rhamnosus* CRL1425 (named as FGM-Lr) or *Lactobacillus casei* CRL431 (named as FGM-Lc). FGMs were obtained by the inoculation (1%) of a fresh (overnight 18 h) culture of these strains into goat's milk followed by 8 h of incubation (37°C , aerobiosis). FGMs were microbiologically analyzed at the end of milk fermentation. The samples were diluted in sterile peptone water (0.1% v/v) and subsequently plated in duplicate onto the selective media. For total viable counts of bacteria, spread plating was performed in duplicate on MRS agar. The plates were incubated at 37°C for 48–72 h. The counts of *Lactobacillus casei* CRL431 and *L. rhamnosus* CRL1425 reached a maximum of 1×10^9 cfu/ml and 7.21×10^8 , respectively, after fermentation process. These FGMs were prepared daily for the feeding trials.

Animals and experimental design

The experimental protocol was approved by the Institutional of Laboratory Animals Care and Use of CERELA, and experimental procedures were carried out in accordance with the present laws of Argentina, according to Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) requirements (Resolution D, N° 1047/05) for laboratory animal research.



The overall experimental protocol is summarized in Figure 1. Weanling (21 d) male albino Swiss mice were supplied by CERELA Institute (Tucumán-Argentina). All animals were housed in individual metabolic cages and acclimated to 22°C with a 12 h light - 12 h dark cycle. Mice were fed with Balanced Conventional Diet (BCD) providing 21% calories as protein (casein), 66% as carbohydrates (corn starch) and 13% as fat (soy bean oil), supplemented with vitamin mixture (2.2%) (ICN, Biomedicals, Inc., Ohio, USA), and salt mixture (4%) (ICN, Biomedicals, Inc., Ohio, USA) and given water *ad libitum*. After 2 day diet adaptation period, mice were matched by weight and assigned to one of three main groups where the effects of mild caloric restriction (during 45 or 90 days) and renutrition diets were studied:

(1) *Ad libitum* control group (AD): after a period of adaptation to the diet (2 days), mice (n=30) were provided with *ad libitum* access to BCD for all assay period. Six mice were killed at day 0 (baseline values). Other 6 mice were killed at day 45 or 90 (in coincidence with

the end of the period of caloric restriction). Other 6 mice were killed after 12 days (in coincidence with the end of the period of renutrition). All these mice constituted the AD45 and AD90 groups.

(2) Caloric restricted group (CR): after diet adaptation period (2 days), one animal group (n=24) received food in an amount equivalent to 75% of the *ad libitum* diet, to match 10-25% weight loss compared with the AD group (as was previously set) (11). Six (6) mice were killed at day 45 or 90 (end of the period of caloric restriction). Other 6 mice were killed after 12 days (in coincidence with the end of the period of renutrition). All these mice constituted the CR45 and CR90 groups.

(3) Caloric Restricted / Renourished group (CR/Re): after caloric restriction period, mice (n=36) were separated into three feeding groups for each period of caloric restriction. Animals of these test groups were refed with *ad libitum* BCD plus whole goat's milk during 7 days followed by 5 days in which animals were fed with BCD plus (a) FGM-Lr or (b) FGM-Lc or (c) goat's milk (goat's milk control = GM). Previous studies showed that treatment with fermented milks for 5 days is the optimum time for a better recovery of metabolic and immune status (12,15,16). These mice constituted the CR45/Re-Lr or CR90/Re-Lr; CR45/Re-Lc or CR90/Re-Lc and CR45/Re-GM or CR90/Re-GM groups, respectively. Nutrient composition of FGMs is presented in Table 1. FGMs and GM were given twice daily (about 4 ml/mouse) during the renutrition period in replacement of drinking water. FGMs and GM were entirely consumed, ensuring the same energy consumption in all groups.

Throughout the trial, mice were sacrificed at beginning of treatment period (0 day), at the end of the two caloric restriction periods (45 or 90 days), and at the end of renutrition periods (after 12 days, that is to say at day 57 or 102, respectively) (Figure 1). Animals were fasted (12 hours) before killing. Blood samples were collected from cardiac puncture, stored at room temperature for 1 h and kept overnight at 4 °C. Then, it was centrifuged at 1,000 xg for 10 min to collect the serum and stored at -70 °C until analysis. The small intestines were removed for immunological studies. Macrophages were isolated from peritoneum to evaluate the phagocytic activity.

Biochemical measurements

Serum glucose, cholesterol and triglycerides were determined by oxidase/peroxidase method according to Trinder [21] (GT Lab, Rosario, Argentina). Total protein concentration was determined using Bradford technique [22]. Haematocrit (HTO), and the number of leukocytes were determined by the haematocytometric methods. Haemoglobin concentration was determined by colorimetric assays (GT Lab, Rosario, Argentina). Serum leptin concentration was assayed using a sensitive commercially available ELISA kit according

Table 1: Composition of goat's milk and fermented goat's milks with *Lactobacillus rhamnosus* CRL1425 or *Lactobacillus casei* CRL431.

Components	Goat's milk	FGM-Lr	FGM-Lc
Proteins (%; w/v)	3.28	3.4	3.33
Total lipid (%; w/v)	4.16	4.15	4.19
Carbohydrate (%; w/v)	4.45	4.5	4.43
Calories (KJ/ml)	286.21	288.34	285.88

FGM-Lr: Fermented goat's milk with *L. rhamnosus* CRL1425 ; FGM-Lc: Fermented goat's milk with *L. casei* CRL431.

Table 2: Effect of goat's milk and fermented goat's milks by *L. rhamnosus* CRL1425 (FGM-Lr) and *L. casei* CRL 431 (FGM-Lc) on metabolic and haematologic parameters in CR45 mice under feeding conditions.

Experimental Groups													
Variables	AD		CR45		CR/Re		CR45/Re		CR45/Re		P (A_OVA)		
	Mean	SD	Mean	SD	FGM-Lr		FGM-Lc		GM				
					Mean	SD	Mean	SD	Mean	SD	T	D	T*D
Glucose (g/l)	1.63 ^a	0.06	0.40 ^b	0.03	1.59 ^a	0.27	1.58 ^a	0.09	1.53 ^a	0.06	NS	NS	NS
Cholesterol (mg/dl)	127.50 ^a	8.94	137.00 ^a	6.32	104.00 ^b	3.79	101.00 ^b	10.12	115.00 ^c	8.05	<0.001	<0.001	<0.001
Triglycerides (mg/dl)	109.00 ^a	1.26	95.00 ^b	1.26	102.00 ^d	8.94	115.50 ^d	4.43	117.5 ^d	13.28	NS	<0.001	NS
Total Protein (g/l)	4.60 ^a	0.20	4.05 ^b	0.35	4.64 ^a	0.80	4.70 ^a	0.63	4.8 ^a	0.80	NS	NS	NS
Albumin (g/l)	3.12 ^a	0.48	2.88 ^a	0.40	2.99	0.63	3.09	0.27	3.15 ^a	0.41	NS	NS	NS
Haematocrit (%)	45.75 ^a	2.48	43.70 ^a	2.14	43.15 ^b	1.79	42.90 ^b	1.07	43.30 ^b	1.39	NS	NS	NS
Haemoglobin (g/dl)	14.02 ^a	2.15	14.80 ^a	1.88	13.8 ^a	1.07	13.75 ^a	0.44	13.60 ^a	0.80	NS	NS	NS

*Animals were fed with a caloric restriction diet during 45 days (caloric restriction period), after which received *ad libitum* BCD plus goat's milk (GM) during 7 days and then mice were fed with *ad libitum*

BCD plus fermented goat's milk (FGM) with *L. rhamnosus* CRL1425 (CR/Re45-Lr) or *L. casei* CRL431 (CR/Re45-Lc), or GM (CR/Re45-GM) during 5 days (renutrition period).

Values are means ± standard deviation (n = 6 for all groups). Means for each row without a common superscript letter differ significantly (P<0.05, Student's t test). T: effect of time of caloric restriction; D: effect of different diet, T*D: interaction between time of caloric restriction and diet (ANOVA test). NS: no significant.

Table 3: Effect of goat's milk (GM) and fermented goat's milks by *L. rhamnosus* CRL1425 (FGM-Lr) and *L. casei* CRL 431 (FGM-Lc) on metabolic and haematologic parameters in CR90 mice under feeding conditions.

Experimental groups													
Variables	AD		CR90		CR/Re		CR90/Re		CR90/Re		P (A_OVA)		
	Mean	SD	Mean	SD	FGM-Lr		FGM-Lc		GM				
					Mean	SD	Mean	SD	Mean	SD	T	D	T*D
Glucose (g/l)	1.64 ^a	0.10	0.48 ^b	0.04	1.52 ^a	0.27	1.55 ^a	0.13	1.56 ^a	0.07	NS	NS	NS
Cholesterol (mg/dl)	133.00 ^a	14.10	110.00 ^b	6.32	100.00 ^c	7.49	115.00 ^b	6.96	120.00 ^b	5.06	<0.001	<0.001	<0.001
Triglycerides (mg/dl)	121.00 ^a	11.50	102.00 ^b	7.16	130.00 ^c	6.05	140.00 ^d	8.94	148.00 ^d	9.84	NS	<0.001	<0.001
Total Protein (g/l)	4.63 ^a	0.89	4.09 ^b	0.54	4.80 ^a	0.36	4.70 ^a	0.63	4.95 ^a	0.49	NS	NS	NS
Albumin (g/l)	3.12 ^a	0.45	2.80 ^a	0.20	3.10 ^a	0.15	2.90 ^a	0.30	3.16 ^a	0.18	NS	NS	NS
Haematocrit (%)	45.75 ^a	2.77	42.00 ^a	2.58	44.24 ^a	2.29	41.35 ^a	3.47	43.68 ^a	2.24	NS	NS	NS
Haemoglobin (g/dl)	14.02 ^a	2.40	13.98 ^a	1.16	13.96 ^a	1.30	13.30 ^a	1.10	13.87 ^a	1.30	NS	NS	NS

* Animals were fed with a caloric restriction (CR) diet during 90 days (caloric restriction period), after which received *ad libitum* BCD plus goat's milk (GM) during 7 days and then mice were fed with *ad libitum* BCD plus fermented goat's milk (FGM) with *L. rhamnosus* CRL1425 (CR/Re90-Lr) or *L. casei* CRL431 (CR/Re90-Lc), or GM (CR/Re90-GM) during 5 days (renutrition period).

Values are means ± standard deviation (n = 6 for all groups). Means for each row without a common superscript letter differ significantly (P<0.05, Student's t test). T: effect of time of caloric restriction; D: effect of different diet, T*D: interaction between time of caloric restriction and diet (ANOVA test). NS: no significant.

to the manufacturer's instructions (R&D Systems, Minneapolis, MN).

Immunofluorescence assay for IgA secreting cells


The small intestines were removed after 12 days of renutrition period and washed with saline solution (NaCl 0.15 M). The tissues were prepared using the modified method described by Sainte-Marie [16]. Serial paraffin sections (4 µm) were made and used for direct immunofluorescence assays to determine the number of IgA+ cells in the lamina propria. After deparaffinization using xylene and rehydration in a decreasing gradient of ethanol, slides were incubated with α-chain monospecific antibody conjugated with fluorescein isothiocyanate (FITC, Sigma, St. Louis, USA) for IgA+ cells. The results are expressed as the number of positive cells per 10 fields of vision (magnification 100×) using a fluorescent light microscope.

Ex vivo phagocytosis assay of peritoneal macrophages

After 12 days of renutrition period, peritoneal macrophages were obtained according to De Moreno de LeBlanc et al. [23] and the cell concentration were adjusted at 1 × 10⁶ cells/ml. Phagocytosis assay was performed using a *Saccharomyces boulardii* suspension at a concentration of 10⁷ cells/ml. Opsonized yeast in mouse autologous serum (10%) were added to 0.2 ml of macrophage suspension. The mixture was incubated for 30 min at 37°C. The percentage of phagocytosis was expressed as the percentage of phagocytosing macrophages in 200 cells counted in an optical microscope.

Statistical analysis

Statistical analyses were carried out using SPSS 11.0 software (SPSS Inc., Chicago, IL, USA). The experimental data were expressed



as mean ± SD and were statistically evaluated by two-way ANOVA. Tukey's test (for pairwise comparisons of the means of the different groups) was used to test for differences between the groups. Single comparisons between groups were assessed by Student's t test. Differences were considered significant at P < 0.05. As an exploratory step, a principal component analysis was performed, including IgA cells number, leukocytes number, phagocytic activity and leptin concentration. Pearson correlation coefficients were used to detect correlations, where appropriate. P values of <0.05 were considered to be significant.

Results

Effects of mild caloric restriction/renitration on body weight


Body weight was followed throughout the trials (Figure 2). As expected, mild caloric restriction diet resulted in lower body weight in mice of both CR groups (CR45 and CR90), with a decrease of approximately 21% in CR45 and 22% in CR90 group. At the end of renitration period, all diets in both CR groups, caused a significant increase in body weights in comparison with the respective CR group of the same age. ANOVA test showed that there was no effect of Time (T) of caloric restriction diet on the body weight in both CR groups and there was a significant effect of diet (D) only in CR45 groups. Student's t test revealed that the FGM-Lc induced a higher increase in body weight than GM.

Effects of mild caloric restriction/renitration on serum biochemical variables

Table 2 and 3 shows serum glucose, cholesterol, triglycerides,

total proteins, and haematological parameters (haematocrit, haemoglobin) values of CR animals. As it was expected, in both CR groups, the caloric restriction diet induced a significant decrease in serum glucose, triglycerides and total proteins (Table 2 and 3). All renitration diets induced a significant increase of glucose, triglycerides and total proteins values in comparison with CR45 or CR90 controls (Table 2 and 3). In CR90/Re animals, an influence of diet (D) on the triglyceride levels were observed; FGM-Lr induced lower triglyceride values than FGM-Lc and GM administration (ANOVA test) (table 3). Cholesterol values not changed significantly at 45 days of caloric restriction diet. But, a significant fall was observed at 90 days of diet (ANOVA test) (Table 2 and 3). Both, FGM-Lr and FGM-Lc induced a hypocholesterolemic effect compared with CR45/Re-GM mice, while in CR90/Re group, only FGM-Lr administration was able to keep lower values of cholesterol (Table 2 and 3). Serum albumin modifications were not observed in any groups (Table 2 and 3). Also, the haematocrit percentage and haemoglobin concentration did not modify with any treatment in comparison with control groups (Table 2 and 3).

The results of serum leptin levels are shown in Figure 3. The 45 days of caloric restriction induced a slight increase of serum leptin, while 90 days of caloric restriction diet, produced a significant descent (ANOVA test). In the renitration period, effect of time (T) of caloric restriction on leptin levels was not observed. However, influence of diet (D) was demonstrated (ANOVA test). Serum leptin values decreased after administration of FGM-Lr compared to CR and AD control, while FGM-Lc induced a significant increase on serum leptin when it was administered to both CR groups. The values observed with GM were similar to AD control mice.



Impact of mild caloric restriction/renutrition on immune response. Effect on IgA secreting cells of the small intestine and phagocytic activity of macrophages isolated from peritoneum

Table 4 shows the number of leukocytes, the number of IgA secreting cells and the percentage of phagocytic activity of peritoneal macrophages in mice fed with caloric restriction diet and renourished with FGMs. Effect of time (T) of caloric restriction was observed in leukocytes number (ANOVA test). The administration of different renutrition diets (D) has influence on the studied immune parameters (ANOVA test). In both CR groups, leukocytes number diminished significantly. The administration of the FGMs during the renutrition period significantly improved the number of leukocytes, but without retrieving a value similar to the controls. The number of IgA+ cells significantly decreased in both CR group and FGMs feeding restore the count of these cells, and exceeding in the case of FGM-Lc administration, the AD control values. The percentage of phagocytic activity of peritoneal macrophages, markedly affected due to caloric restriction diet, only increased significantly after renutrition period with FGM-Lc in both CR/Re mice.

Figure 4 shows the principal component analysis (PCA) with focus on groupings of diets with respect to leptin and immune parameters levels. The analysis revealed that two components can be extracted, which together accounted for 82.6% of the variability in the grouping of diets. The first principal component discriminated better the animals according to their diets, specially AD animals and those renourished with FGM+Lc, from the other diets, and was more influenced by leukocytes number. The second principal component singled out AD animals and was more influenced by serum leptin levels. The loading plot showed that all the animals fed with the same diet formed a group well separated from other groups. Only CR groups were discriminated between animals fed during 45 days and




Figure 4: Biplot of the principal component analysis with focus on groupings of diets with respect to leptin and immune parameters level (IgA+ cells number; leukocytes number; peritoneal phagocytic activity) using the first and second principal components. Animals were fed with a caloric restriction (CR) diet during 45 or 90 days (caloric restriction period), after which received *ad libitum* BCD plus whole goat’s milk (GM) during 7 days followed by 5 days in which animals were fed with BCD plus fermented goat’s milk (FGM) with *L. rhammnosus* CRL1425 (CR/Re45-Lr, CR/Re90-Lr) or *L. casei* CRL431 (CR/Re45-Lc, CR/Re90-Lc), or GM (CR/Re45-GM, CR/Re90-GM) (renutrition period).

those fed during 90 days, which demonstrates the influence of the time of caloric restriction with respect to immune parameters and leptin levels.

Correlations of serum leptin levels with IgA+ cells number, leukocytes number and peritoneal phagocytic activity are illustrated in Table 5. Significant and positive strong correlation between leptin concentration and immune parameters were found in all studied groups when they were analyzed individually. Thus, serum leptin

Table 4: Effect of goat’s milk and fermented goat’s milks by *L. rhammnosus* CRL1425 (FGM-Lr) and *L. casei* CRL 431 (FGM-Lc) on leukocytes number, IgA-secreting cells on the small intestine and phagocytic activity of peritoneal macrophages in CR45 or CR90 mice under feeding conditions*.

Experimental Groups		Leukocytes (cells/∞1)	RC45/Re		Leukocytes (cells/∞1)	RC90/Re	
			IgA+ cells _umber (positive cells per 10 villi)	Peritoneal phagocytic activity (%)		IgA+ cells _umber (positive cells per 10 villi)	Peritoneal phagocytic activity (%)
Control	AD	6.83 ^a ± 0.84	108.25 ^a ± 7.89	46.45 ^a ± 7.11	6.80 ^a ± 0.66	108.25 ^a ± 7.89	46.45 ^a ± 7.11
	CR	3.75 ^b ± 0.62	73.75 ^b ± 7.54	24.95 ^b ± 5.02	2.55 ^b ± 0.66	66.50 ^b ± 6.45	20.20 ^b ± 4.10
	A_OVA	T				T	
CR/Re	FGM-Lr	4.45 ^c ± 0.76	118.75 ^{a,c} ± 4.35	26.25 ^b ± 4.86	4.47 ^c ±0.86	117.00 ^{a,c} ± 5.29	25.25 ^b ± 2.87
	A_OVA		D	D		D	D
	FGM-Lc	5.00 ^c ± 0.66	139.75 ^d ± 5.62	37.40 ^c ± 4.50	5.00 ^c ±0.73	135.50 ^d ± 4.93	36.00 ^c ± 4.69
	A_OVA		D	D		D	
	GM	5.10 ^c ± 0.84	113.25 ^{a,c} ± 2.36	24.50 ^b ± 4.65	4.99 ^c ± 0.51	115.75 ^{a,c} ± 4.35	25.25 ^b ± 5.74
	A_OVA		D	D		D	D

* Animals were fed with a caloric restriction (CR) diet during 45 or 90 days (caloric restriction period) after which received *ad libitum* BCD plus goat’s milk (GM) during 7 days and then mice were fed with *ad libitum* BCD plus fermented goat’s milk (FGM) with *L. rhammnosus* CRL1425 (CR/Re45- Lr) or *L. casei* CRL431 (CR/Re45-Lc), or GM (CR/Re45-GM) during 5 days (renutrition period). The number of IgA+ cells was determined by direct immunofluorescence on the small intestine tissue slides from mice of different experimental groups. The results were expressed as the number of positive cells per ten fields of vision (1000X). Mice were sacrificed at the end of the study period to isolate the macrophages from peritoneum. The phagocytic activity of these cells was determined using *Saccharomyces boulardii*. The values are expressed as mean of percentage of phagocytosis expressed as the percentage of phagocytic macrophages in 200 cells counted. Values are means ± standard deviation (n = 6 for all groups). Means for each column without a common superscript letter differ significantly (P<0.05, Student’s t test). T: effect of time of caloric restriction; D: effect of different diet, T*D: interaction between time of caloric restriction and diet (ANOVA test).

Table 5: Relationship between serum leptin levels and immune parameters (IgA-secreting cells on the small intestine, phagocytic activity of peritoneal macrophages and blood leukocytes number) in *Ad-libitum*, CR and CR/Re mice under feeding conditions*.

Leptin	IgA+ cells number	Peritoneal phagocytic activity	Leukocytes number
	<i>r</i>	<i>r</i>	<i>r</i>
AD	0.880 ^a	0.885 ^a	0.711 ^a
CR45	0.870 ^a	0.961 ^a	0.753 ^b
CR90	0.923 ^a	0.944 ^a	0.781 ^b
FGM-Lr45	0.927 ^a	0.951 ^a	0.758 ^b
FGM-Lr90	0.975 ^a	0.944 ^a	0.899 ^a
FGM-Lc45	0.976 ^a	0.882 ^a	0.909 ^a
FGM-Lc90	0.890 ^a	0.902 ^a	0.911 ^a
GM45	0.822 ^b	0.810 ^b	0.865 ^a
GM90	0.990 ^a	0.912 ^a	0.869 ^a
Total#	0.405^a	0.382^a	0.389^a

* Animals were fed with a caloric restriction (CR) diet during 45 or 90 days (caloric restriction period) after which received *ad libitum* BCD plus goat's milk (GM) during 7 days and then mice were fed with *ad libitum* BCD plus fermented goat's milk (FGM) with *L. rhamnosus* CRL1425 (FGM-Lr) or *L. casei* CRL431 (FGM-Lc), or GM during 5 days (renutrition period).

r, Pearson correlation; ^a Correlation is significant at the 0.01 level; ^b Correlation is significant at the 0.05 level.

#, Relation between leptin and immune parameters regardless of the type of diet.

levels were positive predictors of the number of IgA+ cells, leukocytes and phagocytic activity, with Pearson correlation coefficients ranging from 0.711 and 0.990 ($P < 0.05$). Even when the data were analyzed regardless of the type of diet, leptin concentration still significantly and positively correlated with IgA+ cells number ($r = 0.405$; $P < 0.01$), leukocytes number ($r = 0.389$; $P < 0.01$) and peritoneal phagocytic activity ($r = 0.382$; $P < 0.01$).

Discussion

The results obtained in this work showed some beneficial effects of mild caloric restriction on health that are comparable to previous reports, such as decline in serum triglycerides, cholesterol, glucose and leptin levels [9,24]. However, long term mild caloric restriction adversely affects other parameters and may compromised immune function.

Fermented dairy products are widely accepted as health products and actually, the development of new functional foods are required by specific benefits on health. The goat's milk represented an alternative option for food industry for development of different functional food with important impact on malnutrition therapy. We selected goat's milk for nutritional rehabilitation because in a previous study, and in agreement with other authors, we showed that it could be effectively used, as an alternative milk source, for gaining weight and for improving the altered metabolic parameters due to malnutrition [11]. In accordance with our previous results obtained in a mild caloric restriction model we also demonstrate here the efficiency of FGMs to restore (during refeeding period) metabolic parameters values, such as serum glucose and triglycerides. Lower levels of triglycerides obtained with FGM-Lr administration could be related to liporegulatory effects of leptin. Previous studies have already demonstrated a strain-dependent effect on cholesterol and triglycerides levels in rodents and also in humans [25,26].

It is well known that leptin could be used as a sensitive marker of nutritional status [27]. In a previous work we showed that the short-term (12 days) mild caloric restriction does not significantly modify

serum leptin levels in mice [11]. In the current work, mice were maintained in this nutritional condition during 45 and 90 days. We observed that mice subject to mild caloric restriction diet for a period of 45 days showed a slight increase on serum leptin concentration. Those leptin values suggested that this hormone could play a role in a metabolic adaptive phenomenon necessary in the active growth period of weaned mice. Gat-Yablonsky *et al.* reported a significant stimulatory effect of leptin on longitudinal growth in a mouse model, even in the presence of low caloric intake [28,29]. The leptin levels decreased significantly only after implementation of 90 days of mild caloric restriction diet, probably related to a decrease in total white adipose tissue mass associated to the drop in the body weight of CR90 mice. Dagon *et al.* analyzed leptin functionality under different degrees of restriction diet in rats; and these authors observed that serum leptin concentrations decreased promptly after implementation of both, 40% or 60% of caloric restriction diets [30]. The discrepancies between our results (decrease of leptin concentration only after 90 days of caloric restriction diet) and those of Dagon *et al.*, could be attributed to the mild caloric restriction animal model used in this work, where animals only suffer 25% of caloric restriction.

Some strains of lactobacilli are known to modulate mucosal immune functions when were administered orally [13,15,16,31]. Several studies showed variable effects of *Lactobacillus* on the production of leptin. The main reason for the discrepancy is the route of administration of *Lactobacillus* strain and/or animal species or breed used. Bleau *et al.* [34] described *Lactobacillus*-induced decrease in leptin release by adipocyte derived from SJL mice, whereas the same *Lactobacillus* preparation led to an increase in leptin release when used to treat C57BL/6 adipocytes. In another report, probiotic capsules containing *L. acidophilus* had no effect on plasma leptin in a group of men even after 2 months of oral intake [33]. Oral administration of *L. plantarum* to smokers led to a decrease on leptin concentrations in plasma and other parameters which were attributed to their anti-inflammatory properties [35]. In other study, it was shown that direct administration of *Lactobacillus* supernatant into the brains of rats resulted in weight loss without a decrease in food

consumption and this was accompanied with an increase in leptin expression in neurons and peripheral adipose tissues as intestine [36]. This modulatory effect on leptin secretion was also observed with other probiotics, such as *Bifidobacterium* genus. In previous studies we demonstrated that administration of *B. pseudocatenulatum* CECT 7765 significantly reduced leptin levels in high-fat-diet fed mice [37]. In the current work we proved that fermented milks impact on leptin circulating levels and this effect is strain dependent. Despite body weight recovery was similar after renutrition with both FGMs, we showed that FGM-*L. rhammnosus* ETC14 and FGM-*L. casei* CRL431 were able to induce changes in serum leptin levels differently. The leptin concentration was significantly decreased with FGM-*L. rhammnosus* ETC14 administration, while FGM-*L. casei* CRL431 displayed higher values than AD and CR controls.

Principal component analysis clearly discriminated between diets, which were grouped with respect to serum leptin concentration and immune parameters, regardless of the time of caloric restriction, except for CR controls. Specially, serum leptin levels and immune parameters positively influenced the differences between animals renourished with FGM-Lc and the others groups. However, PCA revealed a scarce impact of leptin and immune response in FGM-Lr groups.

According to previous studies, IgA+ cells number and phagocytic capacity of peritoneal macrophages can be used as immunologic parameters to evaluate probiotic properties of fermented milks [9,16,38]. In this work, we demonstrated that *L. casei* CRL431 was able to improve immune competence in mild calorie restricted mice [15]. This result is in concordance with other studies in malnourished and immunosuppressed mice [39–41]. Since the administration of FGM-Lc also increases serum leptin levels, it could be expected that this hormone intervene in the immunomodulatory mechanisms triggered by this microorganism. It is known that leptin is able to set up macrophages to be more responsive to lipopolysaccharide [42] and exerts direct effects on CD4+ T lymphocyte proliferation, macrophage phagocytosis, and secretion of inflammatory cytokines such as IL-1 and tumor necrosis factor (TNF) α [31]. In line with this agreement, La Cava *et al.* showed that the restoration of leptin concentration in *ob/ob* mice might boost immune responses, via increase of thymic T cell output and cell mediated Th1 immune responses [4,44]. In addition, leptin-deficient mice have been reported to display reduced cellularity in spleen and thymus, in cell-mediated proinflammatory immune response, and in antibody production [13,45].

L. rhammnosus ETC14 is a microorganism with proved technological properties [20]. However, it showed low probiotic effects (up-regulation of immune response) because it produce a slight increase of IgA producing cells without change on phagocyte activity. The immune response triggered by *L. rhammnosus* ETC14 is accompanied by a decrease in serum leptin concentration. According to Bleau *et al.* the inhibition of leptin secretion may thus be a new way by which some lactobacilli strains may reduce inflammatory diseases [34]. Leptin is involved in several autoimmune conditions such as diabetes [46] or arthritis [47]. La Cava *et al.* suggested that modulation of circulating leptin levels may be considered as a newer possible strategy to intervene on some inflammatory and autoimmune conditions [44]. In this context, our results suggested that FGM-Lr could be used as downregulator of leptin secretion, which

could be helpful in inflammatory diseases. Studies of leptin (*OB*) mRNA expression levels in white adipose tissue should be performed to test this hypothesis.

Considering this background it could be assumed that leptin may contribute to improved macrophage phagocytosis and adaptative response (production of IgA-antibody-secreting cells) in malnourished individuals. As expected, leptin levels positively correlate with IgA+ cells number and peritoneal phagocytic activity, but also with leukocytes number, in all the groups studied. Furthermore, when the data were analyzed together, regardless of the type of diet, the correlation between leptin and the immune parameters remained positive. As was previously mentioned, numerous studies have shown a multifunctional role of leptin in modulating immune and inflammatory reactions, both in humans and experimental animals [42,48–51]. Since probiotic bacteria improve the function of immune system in malnourished humans and experimental animals, and as we demonstrated, these bacteria can regulate leptin levels in mice, it could be expected that probiotic bacteria produce the same response in humans and could be part of the mechanisms by which probiotics modulate the immune system.

Conclusion

In conclusion, the regulation of serum leptin secretion during mild caloric restriction appears to depend on food deprivation period. Long time food restriction could lead to a decrease in leptin secretion. Whereas, during short-time moderate food restriction, leptin levels could be sustained as a way to regulate the energy homeostasis. On the other hand, not traditional fermented milk products could modify circulating leptin levels in a strain-dependent way, feature that could be exploited in hosts with different nutritional status or in inflammatory diseases. Thus, renutrition diets with both FGMs are able to modulate in different way, leptin secretion, IgA+ cells and phagocytic activity, and FGM-Lc could be useful in malnutrition status.

A developing area of research is the study of the complex link between leptin, nutrition, immune status and probiotics. We suggested that caloric restriction represents an ideal model to investigate how changes in the nutritional status influence neuroendocrine, reproductive, and immune functions.

Therefore, we aimed to gain further insight into the mechanisms that could underlie the link between nutritional and immune status. One way to take into account is performing a more extensive analysis of leptin levels and their relationship with other immune parameters in animals exposed to caloric restriction and renourished with probiotics.

Acknowledgment

We are grateful to R. Pivotto (INTA “Santa Cruz”- Catamarca) for supplying goat’s milks used in these studies. We also thank Jose Luis Alvarado for technical assistance in animal care during feeding phases of the experiments. This work was supported by PICT 2004 N° 21447, PIP0010, PICT2011-0804 and UNSTA 2011.

References

1. McGuire S. WHO, World Food Programme, and International Fund for Agricultural Development. 2012. The State of Food Insecurity in the World

2012. Economic growth is necessary but not sufficient to accelerate reduction of hunger and malnutrition. Rome, FAO. *Adv Nutr.* 2013; 4: 126-127.
2. UNICEF, WHO, WB. Levels and trends in child malnutrition: UNICEF-WHO-World Bank Joint Child Malnutrition Estimates. UNICEF, New York; WHO, Geneva; the World Bank, Washington, DC: UNICEF, WHO, WB. 2012.
3. Kilic M, Taskin E, Ustundag B, Aygun AD. The evaluation of serum leptin level and other hormonal parameters in children with severe malnutrition. *Clin Biochem.* 2004; 37: 382-387.
4. Cai C, Shi FD, Matarese G, La Cava A. Leptin as clinical target. *Recent Pat Inflamm Allergy Drug Discov.* 2009; 3: 160-166.
5. Klok MD, Jakobsdottir S, Drent ML. The role of leptin and ghrelin in the regulation of food intake and body weight in humans: a review. *Obes Rev.* 2007; 8: 21-34.
6. Konieczna J, García AP, Sánchez J, Palou M, Palou A, Picó C. Oral leptin treatment in suckling rats ameliorates detrimental effects in hypothalamic structure and function caused by maternal caloric restriction during gestation. *PLoS One.* 2013; 8: e81906.
7. Barrenetxe J, Villaro AC, Guembe L, Pascual I, Muñoz-Navas M, Barber A, et al. Distribution of the long leptin receptor isoform in brush border, basolateral membrane, and cytoplasm of enterocytes. *Gut.* 2002; 50: 797-802.
8. Speakman JR, Mitchell SE. Caloric restriction. *Mol Aspects Med.* 2011; 32: 159-221.
9. Duivenvoorde LP, van Schothorst EM, Bunschoten A, Keijer J. Dietary restriction of mice on a high-fat diet induces substrate efficiency and improves metabolic health. *J Mol Endocrinol.* 2011; 47: 81-97.
10. Pallaro AN, Roux ME, Slobodianik NH. Nutrition disorders and immunologic parameters: study of the thymus in growing rats. *Nutrition.* 2001; 17: 724-728.
11. Gauffin Cano MP, Van Nieuwenhove C, Chaila Z, Bazán C, González S. Effects of short-term mild calorie restriction diet and renutrition with ruminant milks on leptin levels and other metabolic parameters in mice. *Nutrition.* 2009; 25: 322-329.
12. Cano PG, Perdígón G. Probiotics induce resistance to enteropathogens in a re-nourished mouse model. *J Dairy Res.* 2003; 70: 433-440.
13. Maldonado Galdeano C, Novotny Nunez I, de Moreno de LeBlanc A, Carmuega E, Weill R, Perdigon G. Impact of a probiotic fermented milk in the gut ecosystem and in the systemic immunity using a non-severe protein-energy-malnutrition model in mice. *BMC Gastroenterol.* 2011; 11:64.
14. Matarese G. Leptin and the immune system: how nutritional status influences the immune response. *Eur Cytokine Netw.* 2000; 11: 7-14.
15. Gauffin CP, Agüero G, Perdigon G. Adjuvant effects of *Lactobacillus casei* added to a renutrition diet in a malnourished mouse model. *Biocell.* 2002; 26: 35-48.
16. Cano PG, Agüero G, Perdígón G. Immunological effects of yogurt addition to a re-nutrition diet in a malnutrition experimental model. *J Dairy Res.* 2002; 69: 303-316.
17. FAO, WHO. Joint FAO/WHO Expert Consultation on Evaluation of Health and Nutritional Properties of Probiotics in Food including Powder Milk with Live Lactic Acid Bacteria. 2001.
18. Salva S, Nuñez M, Villena J, Ramón A, Font G, Alvarez S. Development of a fermented goats' milk containing *Lactobacillus rhamnosus*: in vivo study of health benefits. *J Sci Food Agric.* 2011; 91: 2355-2362.
19. Haldar C, Ghosh S. Immune Modulation in Goats by Melatonin and Other Hormones: A Novel Horizon of Research. *J Immun Res.* 2014; 1: 7.
20. Katz M, Medina R, Gonzalez S, Oliver G. Esterolytic and lipolytic activities of lactic acid bacteria isolated from ewe's milk and cheese. *J Food Prot.* 2002; 65: 1997-2001.
21. Trinder P. Determination of blood glucose using an oxidase-peroxidase system with a non-carcinogenic chromogen. *J Clin Pathol.* 1969; 22: 158-161.
22. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976; 72: 248-254.
23. De LeBlanc Ade M, Castillo NA, Perdigon G. Anti-infective mechanisms induced by a probiotic *Lactobacillus* strain against *Salmonella enterica* serovar Typhimurium infection. *Int J Food Microbiol.* 2010; 138: 223-231.
24. Reimer RA, Maurer AD, Lau DC, Auer RN. Long-term dietary restriction influences plasma ghrelin and GOAT mRNA level in rats. *Physiol Behav.* 2010; 99: 605-610.
25. Kekkonen RA, Sysi-Aho M, Seppanen-Laakso T, Julkunen I, Vapaatalo H, Oresic M, et al. Effect of probiotic *Lactobacillus rhamnosus* GG intervention on global serum lipidomic profiles in healthy adults. *World J Gastroenterol.* 2008; 14: 3188-3194.
26. Yin YN, Yu QF, Fu N, Liu XW, Lu FG. Effects of four Bifidobacteria on obesity in high-fat diet induced rats. *World J Gastroenterol.* 2010; 16: 3394-3401.
27. Procaccini C, Jirillo E, Matarese G. Leptin as an immunomodulator. *Mol Aspects Med.* 2012; 33: 35-45.
28. Gat-Yablonski G, Ben-Ari T, Shtaf B, Potievsky O, Moran O, Eshet R, et al. Leptin reverses the inhibitory effect of caloric restriction on longitudinal growth. *Endocrinology.* 2004; 145: 343-350.
29. Gat-Yablonski G, Phillip M. Leptin and regulation of linear growth. *Curr Opin Clin Nutr Metab Care.* 2008; 11: 303-308.
30. Dagon Y, Avraham Y, Magen I, Gertler A, Ben-Hur T, Berry EM. Nutritional status, cognition, and survival: a new role for leptin and AMP kinase. *J Biol Chem.* 2005; 280: 42142-42148.
31. Galdeano CM, de Leblanc Ade M, Carmuega E, Weill R, Perdígón G. Mechanisms involved in the immunostimulation by probiotic fermented milk. *J Dairy Res.* 2009; 76: 446-454.
32. Gauffin Cano P, Chaila, Z, González, S, P. de Ruiz Holgado, A. Characterization of metabolic effects of energy mal-nutrition. An experimental model for «in vivo» studies in weaned and adult mice. *J Food Technol.* 2005; 3: 196-203.
33. McMullen MH, Hamilton-Reeves JM, Bonorden MJ, Wangen KE, Phipps WR, Feirtag JM, et al. Consumption of *Lactobacillus acidophilus* and *Bifidobacterium longum* does not alter phytoestrogen metabolism and plasma hormones in men: a pilot study. *J Altern Complement Med.* 2006; 12: 887-894.
34. Bleau C, Lamontagne L, Savard R. New *Lactobacillus acidophilus* isolates reduce the release of leptin by murine adipocytes leading to lower interferon-gamma production. *Clin Exp Immunol.* 2005; 140: 427-435.
35. Naruszewicz M, Johansson ML, Zapolska-Downar D, Bukowska H. Effect of *Lactobacillus plantarum* 299v on cardiovascular disease risk factors in smokers. *Am J Clin Nutr.* 2002; 76: 1249-1255.
36. Sousa R, Halper J, Zhang J, Lewis SJ, Li Wl. Effect of *Lactobacillus acidophilus* supernatants on body weight and leptin expression in rats. *BMC Complement Altern Med.* 2008; 8: 5.
37. Cano PG, Santacruz A, Trejo FM, Sanz Y. *Bifidobacterium* CECT 7765 improves metabolic and immunological alterations associated with obesity in high-fat diet-fed mice. *Obesity (Silver Spring).* 2013; 21: 2310-2321.
38. Matarese G, Carrieri PB, La Cava A, Perna F, Sanna V, De Rosa V, et al. Leptin increase in multiple sclerosis associates with reduced number of CD4(+)CD25+ regulatory T cells. *Proc Natl Acad Sci U S A.* 2005; 102: 5150-5155.
39. Agüero G, Villena J, Racedo S, Haro C, Alvarez S. Beneficial immunomodulatory activity of *Lactobacillus casei* in malnourished mice pneumonia: effect on inflammation and coagulation. *Nutrition.* 2006; 22: 810-819.
40. Racedo S, Villena J, Medina M, Agüero G, Rodríguez V, Alvarez S. *Lactobacillus casei* administration reduces lung injuries in a *Streptococcus pneumoniae* infection in mice. *Microbes Infect.* 2006; 8: 2359-2366.
41. Haro C, Villena J, Zelaya H, Alvarez S, Agüero G. *Lactobacillus casei* modulates the inflammation-coagulation interaction in a pneumococcal pneumonia experimental model. *J Inflamm.* 2009; 6: 28.
42. Lago R, Gómez R, Lago F, Gómez-Reino J, Gualillo O. Leptin beyond body

- weight regulation--current concepts concerning its role in immune function and inflammation. *Cell Immunol.* 2008; 252: 139-145.
43. Vaughan T, Li L. Molecular mechanism underlying the inflammatory complication of leptin in macrophages. *Mol Immunol.* 2010; 47: 2515-2518.
44. La Cava A, Matarese G. The weight of leptin in immunity. *Nat Rev Immunol.* 2004; 4: 371-379.
45. Núñez IN, Galdeano CM, Carmuega E, Weill R, de Moreno de LeBlanc A, Perdígón G. Effect of a probiotic fermented milk on the thymus in Balb/c mice under non-severe protein-energy malnutrition. *Br J Nutr.* 2013; 110: 500-508.
46. Matarese G, Castelli-Gattinara G, Cancrini C, Bernardi S, Romiti ML, Savarese C, et al. Serum leptin and CD4+ T lymphocytes in HIV+ children during highly active antiretroviral therapy. *Clin Endocrinol (Oxf).* 2002; 57: 643-646.
47. Conde J, Scotece M, López V, Gómez R, Lago F, Pino J, et al. Adiponectin and leptin induce VCAM-1 expression in human and murine chondrocytes. *PLoS One.* 2012; 7: e52533.
48. Meyers J, McTiernana, A, Ulrich C. Leptin and immune function: integrating the evidence. *Nutr Res.* 2005; 25: 791-803.
49. Meyers JA, Liu AY, McTiernan A, Wener MH, Wood B, Weigle DS, et al. Serum leptin concentrations and markers of immune function in overweight or obese postmenopausal women. *J Endocrinol.* 2008; 199: 51-60.
50. Fantuzzi G. Three questions about leptin and immunity. *Brain Behav Immun.* 2009; 23: 405-410.
51. Fernández-Riejos P, Najib S, Santos-Alvarez J, Martín-Romero C, Pérez-Pérez A, González-Yanes C, et al. Role of leptin in the activation of immune cells. *Mediators Inflamm.* 2010; 2010: 568343.