



Study on the efficacy of edible coatings on quality of blueberry fruits during shelf-life



C. Mannozi^{a, *}, J.P. Cecchini^b, U. Tylewicz^a, L. Siroli^a, F. Patrignani^{a, c}, R. Lanciotti^{a, c}, P. Rocculi^{a, c}, M. Dalla Rosa^{a, c}, S. Romani^{a, c}

^a Department of Agricultural and Food Sciences, University of Bologna, Cesena, Italy

^b Instituto de Tecnología de Alimentos, Universidad Nacional del Litoral, Santa Fe, Argentina

^c Interdepartmental Centre for Agri-Food Industrial Research, University of Bologna, Cesena, Italy

ARTICLE INFO

Article history:

Received 30 June 2016

Received in revised form

27 December 2016

Accepted 29 December 2016

Available online 29 December 2016

Keywords:

Fruit

Quality

Storage

Alginate

Pectin

ABSTRACT

Edible films or coatings could be used as an alternative way of conservation, because of their ability to reduce respiration and transpiration rate, maintain firmness and generally delay fruit senescence. The aim of this research was to evaluate the influence of different types of coating: sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe), on some blueberries quality characteristics, cell viability and microbial growth during 14 days of storage at 4 °C.

Blueberry samples differently coated did not show significant differences in weight loss, pH, soluble solid and dry matter content. However, the application of Al, Pe and Al + Pe improved the firmness of blueberry samples as compared to the uncoated one. Changes in the surface reflection properties in the coated blueberries induced a general lower lightness and a more intense blue hue colour than the control sample. The microbiological results indicated that the coating of blueberry, in particular with Al or Pe, significantly reduced the growth kinetics of yeasts and mesophilic aerobic bacteria.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Blueberries are appreciated for their rich composition in bioactive compounds such as flavonoids, phenolic acids, tannins and anthocyanins giving them nutraceutical properties. However, fresh fruit deteriorate rapidly due to loss of water and juice (product of superficial lesions), mould and/or putrefaction (Yang et al., 2014). The shelf-life of fresh blueberries usually is in the range of 10–40 days depending on different factors such as fruit maturity, cultivar, harvest method and storage conditions (Abugoch et al., 2016). Various technologies are used to reduce spoilage, extend the shelf-life and retain the nutritional value of fruit products; among this group particular attention can be given to refrigeration, UV irradiation, ozonation and modified packaging atmosphere (Duan, Wu, Strik, & Zhao, 2011). The use of edible films or coatings represents an alternative way of preservation because of their ability to reduce moisture, solute migration, respiration and transpiration rate, to maintain firmness and generally delay

senescence (Tezotto-Uliana, Fargoni, Geerdink, & Kluge, 2014). The efficiency and stability of edible coatings or films depend on their compositions. Edible films and coatings are generally based on biological materials such as proteins, lipids and polysaccharides, alone or, more often, in combination.

Sodium alginate is a natural linear polysaccharide obtained from brown seaweeds and has many important physical and biological properties, such as moisture retention, gel-forming capability, good biocompatibility, low price and high availability (Pei, Chen, Li, & Zhou, 2008).

Pectin is a complex of acidic polysaccharides that form an interpenetrating network in the plant cell wall; it is one of the most important citrus by-products that are industrially extracted from apple pomace and citrus peels. Generally it is used to increase viscosity and gel strength of food products (Krochta & Mulder-Johnston, 1997).

Some studies confirm that the application of edible coatings on fruit surface can increase the shelf-life of different fruits, for example raspberries (Tezotto-Uliana et al., 2014) and tropical fruits (Cerqueira, Lima, Teixeira, Moreira, & Vicente, 2009). However, there are few works about coatings effects on blueberries (Chiabrando & Giacalone, 2015; Duan et al., 2011). In both papers,

* Corresponding author. University of Bologna, Department of Agricultural and Food Sciences, p.zza Goidanich 60, 47521 Cesena (FC), Italy.

E-mail address: cinzia.mannozi2@unibo.it (C. Mannozi).

the authors showed that the use of alginate coating on berries had a positive effect on firmness, titratable acidity and maintained surface lightness of coated fruit products. However, to the best of our knowledge there are no papers presented in the literature on the effect of pectin-based coating on blueberries.

Although edible films are not intended to completely replace conventional packages, the efficiency of food protection can be improved by combining both actions. The objectives of this study were to investigate the effectiveness of sodium alginate, pectin and both of these polysaccharides based coatings in improving some qualitative characteristics of blueberry fruits during shelf-life.

2. Material and methods

2.1. Fruit material

Organic blueberries were purchased once from local market. Berry fruits were kept at 0 ± 1 °C until they were used, for no longer than one week, as suggested by Perkins-Veazie, Clark, Collins, & Magee, 1995 and Jackson, Sanford, Lawrence, McRae, & Stark, 1999. Fresh blueberries with the same colour and size and no damages were selected for the experiments.

2.2. Preparation of coating solutions

Three different coating solutions were prepared, each of them contained 15 g/kg of glycerol ($\geq 99.5\%$ Sigma-Aldrich, St. Louis, MO USA) and 2 g/kg of Tween[®] 20 (Sigma-Aldrich, St. Louis, MO USA) and solved in distilled water. In a first solution, sodium alginate (Al) (Sigma-Aldrich, St. Louis, MO USA) was added in the quantity of 20 g/kg. The second one was enriched by 20 g/kg of pectin (Pe) from citrus peel (Galacturonic acid $\geq 74.0\%$ Sigma, St. Louis, MO USA), and the third one was prepared by combination of Sodium Alginate and Pectin (Al + Pe) in equals amounts of 10 g/kg + 10 g/kg. Afterwards, all coating solutions were homogenised at 5000 rpm for 2 min in order to remove air bubbles.

2.3. Sample preparation

Blueberry fruits were sanitized with sodium hypochlorite water solution (0.2 g/kg), rinsed in distilled water and dried with absorbing paper. Whole fruits were dipped in coating solutions, in two process steps, each one of 30 s duration. The berry samples were drained in a ventilated oven at 25 ± 1 °C for 30 min following the first step dipping, and for 60 min following the second step dipping. Blueberries dipped in distilled water with the same procedures were used as control. Coated berry samples were then placed in plastic trays (PET) closed in micro-perforated bags (PLA) and stored at 4 °C for 14 days. Coated samples and control ones were analysed at 0, 2, 4, 6, 10 and 14 days of storage. Totally 4 samples were obtained: 3 differently coated blueberry samples (Al, Pe, Al + Pe) and 1 not coated control sample. For each sample 540 blueberries were used. Three trays for every sampling time were made, containing 30 blueberries each, from which fruits were taken randomly from the three trays and used for analytical determinations.

2.4. Quality determinations

2.4.1. Weight loss, dry matter, pH and soluble solid content

Weight loss (WL) of blueberry samples during storage was measured by weighting fruits in the trays before storage and at every day of analysis, following the standard method of AOAC (1994).

Dry matter content was determined gravimetrically by

difference in weight before and after drying at 70 °C, until a constant weight was reached (AOAC International, 2002).

pH was determined at 20 °C with a pH meter CRISON GLP21 (Shinghai Shilu-Instruments, China).

Soluble solid content (SSC) analysis were performed at 20 °C by measuring the refractive index of blueberry juice with digital hand refraktometer mod. DR301-95 (Kruess, Germany).

For each treatment-time condition, dry matter was determined in triplicate from 8 blueberries from each tray; pH and SSC were determined also in triplicate on three different juice samples each obtained from 10 berries from each tray, after filtering through Whatman #1 filter paper.

2.4.2. Colour and texture

Surface colour of blueberry was measured using spectrophotometer HUNTERLAB ColorFlexTM, mod. A60-1010-615 (Reston, Virginia). For each sample L^* , a^* and b^* parameters from CIELAB scale were measured and Hue angles (h°) index was calculated.

Penetration test was performed with a Texture Analyser mod. TA-HDi500 (Stable Micro Systems, Godalming, UK) equipped with a 50N load cell and a 2 mm diameter stainless steel probe. Penetration test speed was 0.5 mm s^{-1} , the test ended when a maximum deformation of 80% was reached. Results were expressed as average of 12 measurements carried out on 12 blueberries for each treatment-time condition.

2.4.3. Cell viability

The cell viability test was performed on blueberries slices obtained from 9 different blueberries using fluorescein diacetate (FDA, Sigma-Aldrich, USA, $\lambda_{\text{ex}} = 495 \text{ nm}$, $\lambda_{\text{em}} = 518 \text{ nm}$), as described by Tylewicz, Romani, Widell, and Galindo (2013). Viable cells could be easily identified by a bright fluorescence. Observations were performed under a fluorescent light in a Nikon upright microscope (Eclipse Ti-U, Nikon Co, Tokyo, Japan) equipped with a Nikon digital video camera (digital sight DS-Qi1Mc, Nikon Co, Tokyo, Japan) at a magnification of 4 \times .

2.4.4. Microbial growth

The total loads of mesophylic aerobic bacteria, lactic acid bacteria, yeasts, moulds and total coliforms were evaluated according to the methods reported by Siroli et al. (2015). Briefly, 10 g portion of each sample were used (around 6 berries), suspended in 90 ml of sterile saline solution (9 g/l NaCl) and homogenized using a Stomacher for 2 min at room temperature; serial dilutions were made. The microbiological analyses were performed in triplicate immediately after treatments and during storage.

2.4.5. Data analyses

Analysis of variance (ANOVA) and the test of mean comparison, according to Fisher's least significant difference (LSD) were applied on all obtained data. Level of significance was $p < 0.05$.

The statistical software used was STATISTICA, v 8.0 (StatSoft, Tulsa, Oklahoma).

3. Results and discussion

3.1. Weight loss, dry matter, pH and soluble solid content

The fruits weight loss during storage usually is caused by the migration of the water from the fruit to the surrounding environment. As reported in Table 1, all samples underwent a slight loss of weight during 14 days of storage. Coated samples did not show any significant differences in weight loss as compared to the control. These results are probably due to a slight loss of water undergone

Table 1
Weight loss (%) of control (Control) and sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe) coated blueberry samples during 14 days of storage at 4 °C.

Weight loss (%)					
	T2	T4	T6	T10	T14
Control	-1.1 ± 0.1 ^a	-1.2 ± 0.3 ^a	-2.3 ± 0.3 ^a	-3.9 ± 0.2 ^a	-5.9 ± 0.8 ^a
Al	-1.05 ± 0.05 ^a	-1.8 ± 0.4 ^a	-2.34 ± 0.05 ^a	-4.2 ± 0.2 ^a	-6 ± 1 ^a
Pe	-0.83 ± 0.07 ^a	-1.5 ± 0.2 ^a	-2.2 ± 0.3 ^a	-4.0 ± 0.5 ^a	-5.5 ± 0.2 ^a
Al + Pe	-2 ± 1 ^a	-2.2 ± 0.1 ^a	-2.3 ± 0.5 ^a	-4.1 ± 0.6 ^a	-5.6 ± 0.3 ^a

Data are reported as average values and standard deviations obtained from three replicates for each treatment-time conditions. Values with different letter within the column are significantly different ($p < 0.05$).

by samples. The moisture loss of fresh fruit and vegetables is due to the gradient of water vapor pressure that occurs from different locations in the cell tissues (Yaman & Bayoındurlu, 2002). The cold storage conditions (temperature and relative humidity) could have an effect on the difference of vapor pressure between blueberries and the environment resulting in non-significant weight losses. In fact, as reported by Nunes (2015) the weight loss up to 4–5% does not significantly influence the freshness of the fruit.

As reported in Table 2, no significant differences ($p < 0.05$) on dry matter and pH were detected, among control and differently coated samples at each considered storage time. Concerning the SSC, significant differences ($p < 0.05$), even if slight, were observed only at 10 days of storage; in particular Al and Al + Pe presented higher SSC values as compared to the control and Pe coated blueberry fruits. As a general trend dry matter, pH and SSC tended to increase during storage in both control and coated fruit samples. pH and SSC showed the same behaviour increasing with longer storage time, similar results have been provided by Duan et al. (2011). The increase of pH and SSC is probably due to metabolic processes and reactions during post-harvest storage, which continue to convert starch and acids into the sugar.

3.2. Colour and texture

In Table 3 colour data (lightness - L^* and Hue angles - h°) of blueberry samples during 14 days of storage at 4 °C are reported. Coating induced a general lower lightness and a more intense blue hue colour in blueberry samples as compared with the control one

($p < 0.05$), probably due to the glossy effect of coating. The observed lower luminosity value of coated samples could be caused by the modifications in the surface reflection properties (Hoagland & Parris, 1996). L^* values of control and coated samples tended to increase during the first days of storage, then remained relatively stable and decreased after the sixth storage day.

The visual perception of the intensity of blue colour was always more intense in the coated than in the control samples, as indicated from their highest hue values. Moreover, the blueberry samples showed a general decrease in hue values from 0 to 10 days that tended to increase on 14th day. The h° decrease of blueberries during the first period of storage is probably caused by oxidation or condensation reactions of phenolic compounds resulting in loss of anthocyanins during cold storage (Reque et al., 2014). Moreover, the increase of hue values at the end of storage might be caused by a possible anthocyanins synthesis during ripening as also observed by the higher pH and SSC values (Table 2).

As shown in Fig. 1 blueberry coated samples presented a significantly ($p < 0.05$) higher firmness compared to control sample until the first 10 days of storage. After this period, texture of blueberries coated samples decreased, reaching the same value of control one (1.75N). The higher firmness values of coated samples are probably due to the presence of coating that provide a structural rigidity to the surface of fruit (Duan et al., 2011). Pe and Al + Pe showed the same behaviour of the Al based coating. This result of Al coating was in agreement with Rojas-Graü, Tapia, & Martín-Belloso, (2008) on fresh-cut apple and Fan et al. (2009) on strawberry fruits. Moreover, the retention of firmness could be explained by the delay

Table 2
Dry matter (g/kg) pH and soluble solid content (°Bx) of control (Control) and sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe) coated blueberry samples during 14 days of storage at 4 °C.

Dry Matter (g/kg)						
	T0	T2	T4	T6	T10	T14
Control	178.2 ± 0.4 ^a	193.0 ± 0.6 ^a	183 ± 2 ^a	204.5 ± 0.1 ^a	199 ± 1 ^a	198 ± 1 ^a
Al	177.8 ± 0.8 ^a	180 ± 2 ^a	194.8 ± 0.7 ^a	194.30 ± 0.02 ^a	183.0 ± 0.1 ^a	202 ± 1 ^a
Pe	185.8 ± 0.6 ^a	179.9 ± 0.7 ^a	195.9 ± 0.2 ^a	204.9 ± 0.4 ^a	196.2 ± 0.2 ^a	194.3 ± 0.6 ^a
Al + Pe	185.4 ± 0.2 ^a	186.5 ± 0.8 ^a	190.7 ± 0.5 ^a	184.32 ± 0.06 ^a	193.5 ± 0.4 ^a	188.0 ± 0.6 ^a
pH						
	T0	T2	T4	T6	T10	T14
Control	3.49 ± 0.00 ^a	4.09 ± 0.03 ^a	3.5 ± 0.2 ^a	3.7 ± 0.5 ^a	3.7 ± 0.2 ^a	4.1 ± 0.1 ^a
Al	3.47 ± 0.07 ^a	3.9 ± 0.2 ^a	3.8 ± 0.8 ^a	3.35 ± 0.08 ^a	3.4 ± 0.2 ^a	4.03 ± 0.05 ^a
Pe	3.28 ± 0.04 ^a	3.8 ± 0.1 ^a	3.4 ± 0.2 ^a	3.52 ± 0.00 ^a	3.38 ± 0.07 ^a	4.0 ± 0.1 ^a
Al + Pe	3.55 ± 0.04 ^a	3.8 ± 0.2 ^a	3.5 ± 0.2 ^a	3.31 ± 0.02 ^a	3.6 ± 0.3 ^a	3.58 ± 0.02 ^a
SSC						
	T0	T2	T4	T6	T10	T14
Control	13.4 ± 0.7 ^a	13 ± 2 ^a	15.0 ± 0.2 ^a	15.2 ± 0.2 ^a	12.7 ± 0.9 ^b	15 ± 2 ^a
Al	12.6 ± 0.7 ^a	15 ± 2 ^a	15 ± 3 ^a	14.6 ± 0.1 ^a	15.1 ± 0.9 ^a	15 ± 2 ^a
Pe	13 ± 2 ^a	15 ± 1 ^a	13 ± 1 ^a	13.1 ± 0.5 ^a	11.6 ± 0.4 ^b	18 ± 1 ^a
Al + Pe	13 ± 2 ^a	14 ± 1 ^a	14 ± 1 ^a	15.6 ± 0.2 ^a	15.0 ± 0.4 ^a	17 ± 1 ^a

Data are reported as average values and standard deviations obtained from three replicates for each treatment-time conditions. Values with different letter within the column are significantly different ($p < 0.05$).

Table 3

Lightness- L^* and Hue angles- h° of control (Control) and sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe) coated blueberry samples during 14 days of storage at 4 °C.

L^*						
	T0	T2	T4	T6	T10	T14
Control	21 ± 1 ^a	28.4 ± 0.1 ^a	31.5 ± 0.8 ^a	30.5 ± 0.5 ^a	28.5 ± 0.6 ^a	29 ± 1 ^a
Al	19.33 ± 0.07 ^a	18.9 ± 0.1 ^b	22.74 ± 0.05 ^c	22.2 ± 0.6 ^c	19.4 ± 0.6 ^b	16.48 ± 0.00 ^c
Pe	14 ± 2 ^b	19.5 ± 0.5 ^b	23.2 ± 0.4 ^{bc}	26.0 ± 0.2 ^b	19.3 ± 0.2 ^b	19.59 ± 0.02 ^b
Al + Pe	15.3 ± 0.6 ^b	15.9 ± 0.8 ^c	24.9 ± 0.2 ^b	25.6 ± 0.5 ^b	17.6 ± 1.4 ^b	19.9 ± 0.4 ^b
h°						
	T0	T2	T4	T6	T10	T14
Control	100 ± 11 ^b	90 ± 3 ^c	97 ± 5 ^c	93 ± 4 ^b	72 ± 6 ^c	89 ± 6 ^c
Al	140 ± 11 ^a	126 ± 10 ^{ab}	117 ± 7 ^b	102 ± 9 ^b	75 ± 6 ^b	145 ± 11 ^b
Pe	145 ± 11 ^a	139 ± 7 ^a	128 ± 5 ^a	134 ± 6 ^a	87 ± 6 ^a	151 ± 11 ^b
Al + Pe	154 ± 11 ^a	123 ± 9 ^b	111 ± 7 ^b	126 ± 5 ^a	85 ± 5 ^{ab}	179 ± 11 ^a

Data are reported as average values and standard deviations obtained from twelve replicates for each treatment-time conditions. Values with different letter within the column are significantly different ($p < 0.05$).




Fig. 1. Firmness (N) of control (Control ○) and differently coated blueberry samples (sodium alginate - Al ◇; pectin - Pe □; sodium alginate plus pectin - Al + Pe △) during 14 days of storage at 4 °C.

of pectin and proto-pectin degradation, involved in maintaining the structural integrity of the fruits (Thompson, 1996).

3.3. Cell viability

Fig 2 shows the micrographs resulted from microscopic observations of control (a) and Al + Pe coated blueberry samples (b) after 14 days of storage. The pictures demonstrate that cell viability in all tissues is preserved until 14 days of storage both in case of control and coated samples. The results provide evidence that cell viability (viable cells could be identified by a bright fluorescence on the Figure) can be preserved in blueberries also after the application of coating. If the protoplasts of the cells did not retain the FDA, this means disruption of the plasma membrane (cell lysis) or loss of membrane semi permeability (Halperin & Koster, 2006). These results provide versatile tool to conduct study of the metabolism of blueberry tissues that was maintained despite storage and the application of different types of coatings.

3.4. Microbial growth

As reported in Table 4, yeasts were detected only in control sample after 2 and 4 days of storage while in all the coated samples




Fig. 2. Cell viability for (a) control (Control) and sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe), coated blueberry samples (b) at 14 days of storage after treatment using fluorescein diacetate (FDA) marker. Bar = 100 μm.

Table 4

Yeast count of control (Control) and sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe) coated blueberry samples.

	T0	T2	T4	T6	T10	T14
Control	nd*	2.2 ± 0.3 ^a	3.2 ± 0.2 ^a	3.5 ± 0.3 ^a	3.6 ± 0.3 ^a	3.3 ± 0.3 ^a
Al	nd*	nd*	nd*	nd*	2.1 ± 0.2 ^b	2.0 ± 0.2 ^b
Pe	nd*	nd*	nd*	nd*	1.8 ± 0.3 ^b	1.9 ± 0.2 ^b
Al + Pe	nd*	nd*	nd*	2.2 ± 0.2 ^b	2.2 ± 0.2 ^b	2.2 ± 0.2 ^b

Counts are expressed in log₁₀ cfu/g (±standard deviation). Means followed by different letters are significantly different (p < 0.05) and are obtained from three replicates for each treatment-time conditions.

* Under the detection limit (1 log₁₀ cfu/g).

Table 5

Mesophylic aerobic bacteria of control (Control) and sodium alginate (Al), pectin (Pe) and sodium alginate plus pectin (Al + Pe) coated blueberry samples.

	T0	T2	T4	T6	T10	T14
Control	nd*	2.3 ± 0.3 ^a	2.9 ± 0.3 ^a	3.1 ± 0.3 ^a	4.1 ± 0.3 ^a	4.5 ± 0.3 ^a
Al	nd*	nd*	1.5 ± 0.3 ^b	1.9 ± 0.3 ^b	2.0 ± 0.2 ^b	2.6 ± 0.3 ^b
Pe	nd*	nd*	nd*	2.1 ± 0.3 ^b	2.2 ± 0.3 ^b	2.7 ± 0.2 ^b
Al + Pe	nd*	nd*	1.5 ± 0.3 ^b	3.0 ± 0.3 ^a	3.6 ± 0.4 ^a	4.2 ± 0.4 ^a

Counts are expressed in log₁₀ cfu/g (±standard deviation). Means followed by different letters are significantly different (p < 0.05) and are obtained from three replicates for each treatment-time conditions.

* Under the detection limit (1 log₁₀ cfu/g).

the yeast cell loads were below the detection limit. In samples coated with Al or Pe yeasts were detected only after 10 days of storage. However, at the end of storage the yeast loads of coated samples were 1.09–1.38 logarithmic cycles lower than control samples.

Significant differences were also evidenced in the total aerobic mesophilic cell loads among the samples during the storage period (Table 5). In fact, in this case only the control sample showed mesophilic cell loads above the detection limit after 2 days of storage. Samples coated with Al or Pe showed mesophilic cell loads from the sixth days of storage, significantly lower than the controls and samples coated with Al + Pe. Finally, no significant differences were found for lactic acid bacteria and total coliform cell loads in relation to the coating adopted, whose loads resulted below 2.0 log CFU/g, during the whole period of storage. The microbiological results indicate that the coating of blueberry, in particular with Al or Pe, significantly reduce the growth kinetics of yeasts and mesophilic aerobic bacteria that play a dominant role in the spoilage of minimally processed fruits (Siroli et al., 2014).

4. Conclusions

The use of coating showed a positive effect mainly on firmness and microbial growth of treated blueberry samples. The firmness was maintained until 10 storage days also for the Pe and Al + Pe coated blueberries. Furthermore, the application of coatings reduced the growth kinetics of yeasts and mesophilic aerobic bacteria, in particular with the application of Al and Pe based coatings. Results from this study indicate the possibility of using edible coatings to develop ready-to-eat fresh blueberries with no reduction in their shelf-life. Further researches will focus on the effect of these edible coatings on blueberry bioactive compounds and sensorial properties.

Acknowledgements

Financial support for this project is provided by funding bodies within the FP7 ERA-Net CORE Organic Plus, and with cofunds from the European Commission (No 618107).

References

- Abugoch, L., Tapia, C., Plasencia, D., Pastor, A., Castro-Mandujano, O., López, L., et al. (2016). Shelf-life of fresh blueberries coated with quinoa protein/chitosan/sunflower oil edible film. *Journal of the Science of Food and Agriculture*, 96, 619–626.
- AOAC. (1994). *Official Methods of Analysis* (p. 22209). 1111 North 19th Street, Suite 20, 16th Edi. Arlington, Virginia USA: Association of Official Analytical Chemists.
- AOAC International. (2002). *Official methods of analysis (OMA) of AOAC International, 17th Edition, USA*. Method number: 920.15. Available at: <http://www.eoma.aoc.org/>.
- Cerqueira, M. A., Lima, Á. M., Teixeira, J. A., Moreira, R. A., & Vicente, A. A. (2009). Suitability of novel galactomannans as edible coatings for tropical fruits. *Journal of Food Engineering*, 94, 372–378.
- Chiabrando, V., & Giacalone, G. (2015). Anthocyanins, phenolics and antioxidant capacity after fresh storage of blueberry treated with edible coatings. *International Journal of Food Sciences and Nutrition*, 66, 248–253.
- Duan, J., Wu, R., Strik, B. C., & Zhao, Y. (2011). Effect of edible coatings on the quality of fresh blueberries (Duke and Elliott) under commercial storage conditions. *Postharvest Biology and Technology*, 59, 71–79.
- Fan, Y., Xu, Y., Wang, D., Zhang, L., Sun, J., Sun, L., et al. (2009). Effect of alginate coating combined with yeast antagonist on strawberry (*Fragaria × ananassa*) preservation quality. *Postharvest Biology and Technology*, 53, 84–90.
- Halperin, S. J., & Koster, K. L. (2006). Sugar effects on membrane damage during desiccation of pea embryo protoplasts. *Journal of Experimental Botany*, 57, 2303–2311.
- Hoagland, P. D., & Parris, N. (1996). Chitosan/pectin laminated films. *Journal of Agricultural and Food Chemistry*, 44, 1915–1919.
- Jackson, E., Sanford, K., Lawrence, R., McRae, K., & Stark, R. (1999). Lowbush blueberry quality changes in response to prepacking delays and holding temperatures. *Postharvest Biology and Technology*, 15, 117–126.
- Krochta, J. M., & Mulder-Johnston, D. (1997). Edible and biodegradable polymer films: Challenges and opportunities. *Food technology (USA)*.
- Nunes, M. C. N. (2015). Correlations between subjective quality and physicochemical attributes of fresh fruits and vegetables. *Postharvest Biology and Technology*, 107, 43–54.
- Pei, H. N., Chen, X. G., Li, Y., & Zhou, H. Y. (2008). Characterization and ormidazole release in vitro of a novel composite film prepared with chitosan/poly (vinyl alcohol)/alginate. *Journal of Biomedical Materials Research Part A*, 85, 566–572.
- Perkins-Veazie, P., Clark, J., Collins, J., & Magee, J. (1995). Southern highbush blueberry clones differ in postharvest fruit quality. *Fruit varieties journal (USA)*, 49, 46–52.
- Reque, P. M., Steffens, R. S., Jablonski, A., Flores, S. H., Rios, A. d. O., & de Jong, E. V. (2014). Cold storage of blueberry (*Vaccinium* spp.) fruits and juice: Anthocyanin stability and antioxidant activity. *Journal of Food Composition and Analysis*, 33, 111–116.
- Rojas-Graü, M. A., Tapia, M. S., & Martín-Belloso, O. (2008). Using polysaccharide-based edible coatings to maintain quality of fresh-cut Fuji apples. *LWT-Food Science and Technology*, 41, 139–147.
- Siroli, L., Patrignani, F., Serrazanetti, D. I., Tabanelli, G., Montanari, C., Gardini, F., et al. (2015). Lactic acid bacteria and natural antimicrobials to improve the safety and shelf-life of minimally processed sliced apples and lamb's lettuce. *Food microbiology*, 47, 74–84.
- Siroli, L., Patrignani, F., Serrazanetti, D. I., Tabanelli, G., Montanari, C., Tappi, S., et al. (2014). Efficacy of natural antimicrobials to prolong the shelf-life of minimally processed apples packaged in modified atmosphere. *Food Control*, 46, 403–411.
- Tezotto-Uliana, J. V., Fargoni, G. P., Geerdink, G. M., & Kluge, R. A. (2014). Chitosan applications pre-or postharvest prolong raspberry shelf-life quality. *Postharvest Biology and Technology*, 91, 72–77.
- Thompson, A. K. (1996). *Postharvest technology of fruit and vegetables: Harlow*. Oxford: Blackwell Science Ltd.
- Tylewicz, U., Romani, S., Widell, S., & Galindo, F. G. (2013). Induction of vesicle formation by exposing apple tissue to vacuum impregnation. *Food and Bioprocess Technology*, 6, 1099–1104.
- Yaman, Ö., & Bayoındırlı, L. (2002). Effects of an edible coating and cold storage on shelf-life and quality of cherries. *LWT-Food science and Technology*, 35(2), 146–150.
- Yang, G., Yue, J., Gong, X., Qian, B., Wang, H., Deng, Y., et al. (2014). Blueberry leaf extracts incorporated chitosan coatings for preserving postharvest quality of fresh blueberries. *Postharvest Biology and Technology*, 92, 46–53.