

## Prognostic Implications of Cytogenetic Features in Myelodysplastic Syndromes

Carolina B Belli, PhD<sup>1#a</sup>, Yesica Bestach, MSc<sup>1b</sup>, Laura Kornblihtt, PhD<sup>2#c</sup>, Irene B Larripa, PhD<sup>1,3d</sup>

1. Laboratory of Hematological Genetics, Institute of Experimental Medicine (IMEX-CONICET)/ National Academy of Medicine (ANM), Buenos Aires, 2. Division of Hematology, Clinic Hospital, 'J de San Martín', University of Buenos Aires, Buenos Aires, 3. Department of Genetics, Institute of Hematological Research (IIHEMA)/ANM, Buenos Aires, # = on behalf of the Argentine MDS Study Group, Argentine Society of Hematology, Argentine.

a. Assistant Researcher, b. Doctoral Fellow, c. Hematologist, d. Superior Researcher.

### Abstract

Myelodysplastic Syndromes (MDS) are a heterogeneous group of hematological diseases characterized by refractory cytopenia(s) and variable risk of leukemic progression. Cytogenetic analysis is important in day-to-day clinical practice helping to define subgroups of MDS patients who share similarities in the course of the disease. There are recurring aberrations affecting chromosomes 5, 7, 8, and 20. While all of them do not suggest a therapeutic approach, their presence has been considered as a risk indicator since the original international prognostic scoring system (IPSS) was published. The most recent cytogenetic stratifications tried to find the prognostic significance of less frequent alterations which have been longer included in the intermediate group. Moreover, monitoring of karyotype changes is suggested to evaluate cytogenetic response to treatments and the acquisition of new aberrations associated to an unfavorable outcome. This review focuses on different cytogenetic risk stratifications that have been published during the past twenty years and the molecular background of the most relevant chromosomal findings.

### Keywords

Myelodysplastic Syndromes, cytogenetic analysis, karyotype, monosomal karyotype, risk stratifications, scoring systems, prognostic, survival, cytogenetic aberrations, IPSS, risk assessment.

**Disclosure:** The authors have no conflicts of interest to declare.

**Acknowledgements:** This work was supported by Argentine grants from the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and the Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT).

**Received:** November 12, 2012 **Accepted:** January 10, 2013 **Citation:** *Oncology & Hematology Review*, 2013;9(1):60–7

**Correspondence:** Carolina B. Belli, PhD, Laboratorio de Genética Hematológica, Instituto de Medicina Experimental – Consejo Nacional de Investigaciones Científicas y Técnicas, (IMEX - CONICET)/ Academia Nacional de Medicina (ANM), Pacheco de Melo 3081, 1425, Buenos Aires, Argentina. E: cbelli@hematologia.anm.edu.ar

Myelodysplastic Syndromes (MDS) are a heterogeneous group of clonal hematopoietic stem cell diseases characterized by morphological changes in the bone marrow, peripheral cytopenia(s) and susceptibility to bone marrow failure with or without progression to Acute Myeloid Leukemia (AML). Although there is an inherited predisposition to develop MDS/AML in patients with congenital bone marrow failure syndromes, most of MDS are idiopathic and develop *de novo* MDS without any pre-existing hematologic abnormalities. In secondary MDS, cytotoxic chemotherapy and radiation exposure are major factors in the induction of marrow failure and subsequent malignant transformation.<sup>1-5</sup>

Cytogenetic analyses have proven an extremely valuable diagnostic tools in MDS, involving the whole genome, supporting the cytomorphologic diagnosis, confirming disease clonality, providing prognostic information and, in some cases, helping with selection of treatment. The frequency of clonal chromosomal abnormalities in *de novo* MDS varies between 14–65 % (see *Table 1*) increasing to nearly 80 % in secondary MDS.<sup>6-31</sup> Cytogenetic findings are not specific for the disease because they can be observed in other oncohematological disorders and none of them is specifically associated with any morphologic subtype [except for the isolated del(5q) in the presence of blast count less than 5 %, which is considered as distinct entity by the World Health Organization (WHO)

classification]. However, the proportion of normal karyotypes decreases according to worsening prognosis from 54–59 % in refractory anemia (RA) and RA with ringed sideroblasts (RS) to 41–34 % in RA with excess of blasts (RAEB) and RAEB in transformation (RAEBt) patients (see *Table 2*).<sup>7-8,10-11,16,21-22,24,27,32-34</sup> The wide range of percentage of observed cytogenetic alterations in published MDS series may be related to difficulties in assessing and discriminating between regional variations, differences in classification, sizes of the population based-studies and the inclusion/exclusion of therapy-related MDS or AML cases (see *Table 1*).<sup>6-31</sup>

MDS are not associated with any specific chromosomal alteration and can vary from simple to complex.<sup>6,9,11-12,34-35</sup> Although all chromosomes may be involved in a variety of cytogenetic aberrations, chromosomes 5, 7, 11, 17, 20 and Y are the most commonly affected, mainly by partial or total loss; followed by trisomies of chromosomes 8 and 21, and rare translocations or other structural alterations predominantly implicating chromosomes 1, 2, 3 and 11.<sup>34</sup> Deletions are more often interstitials and display variable sizes; however 'common deleted regions' have been described for most of these.<sup>36-37</sup> The predominance of partial or total chromosomal losses suggests involvement of tumor suppressor genes. One of both alleles is deleted and the remaining may also be deleted, mutated or silenced by aberrant hypermethylation. However, other relevant mechanisms have been recently

**Table 1: Cytogenetic Alterations in Different Published Myelodysplastic Syndromes Series**

Series	Country	Normal (%)	5q- (%)	7/7q- (%)	+ 8 (%)	20q- (%)	-Y (%)	Others (%)	Complex Karyo- types (%)	Total
Morel et al., 1993 <sup>6</sup> \$	France	259 (63)	19 (5)	11 (3)	12 (3)	9 (3)	8 (2)	23 (6)	67 (16)	408
Paquette et al., 1993 <sup>7</sup>	USA	124 (60)	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	208
Parlier et al., 1994 <sup>8</sup> \$	Switzerland	48 (44)	9 (8)	5 (4)	4 (4)	5 (5)	5 (4)	17 (16)	16 (15)	109
Greenberg et al., 1997 <sup>9</sup> \$	International	489 (60)	48 (6)	10 (1)	38 (5)	16 (2)	17 (2)	117 (14)	81 (10)	816
de Souza Fernandez, et al., 2000 <sup>10</sup> \$	Brazil	76 (61)	9 (7)	6 (5)	3 (2)	2 (2)	3 (2)	25 (20)	1 (1)	125
Solé et al., 2000 <sup>11</sup> \$	Spain	313 (49)	32 (5)	24 (4)	31 (5)	8 (1)	8 (1)	136 (21)	88 (14)	640
Belli et al., 2002 <sup>12</sup> \$	Argentina	116 (59)	5 (3)	9 (5)	6 (3)	4 (2)	1 (1)	35 (17)	22 (11)	198
Lee et al., 2003 <sup>13</sup> \$	Korea	67 (56)	2 (2)	2 (2)	7 (6)	0 (0)	3 (3)	20 (17)	18 (15)	119
Lau et al., 2004 <sup>14</sup> \$	Singapore	21 (53)	0 (0)	n.s.	n.s.	n.s.	n.s.	n.s.	8 (20)	40
Lorand-Metze et al., 2004 <sup>15</sup> &	Brazil	74 (86)	1 (1)	1 (1)	1 (1)	0 (0)	2 (2)	0 (0)	7 (8)	86
Solé et al., 2005 <sup>16</sup> \$	Spain	514 (53)	55 (6)	43 (4)	56 (6)	13(1)	17(2)	163 (17)	107 (11)	968
Malcovati et al., 2005 <sup>17</sup> &	Italy	222 (58)	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	386
Müller-Berndorff et al., 2006 <sup>18</sup> \$	Germany	46 (58)	5 (6)	3 (4)	n.s.	n.s.	1 (1)	13 (16)	11 (14)	79
Irons et al., 2006 <sup>19</sup> &	China	121 (74)	0 (0)	n.s.	n.s.	n.s.	n.s.	n.s.	8 (5)	164
Sakuma et al., 2006 <sup>20</sup> &#	Japan	43 (43)	0 (0)	1 (1)	6 (6)	5 (5)	6 (6)	12 (12)	27 (27)	100
Bernasconi et al., 2007 <sup>21</sup> \$	Italy	197 (40)	43 (9)	33 (7)	22 (4)	15 (3)	5 (1)	135 (27)	41 (8)	491
Haase et al., 2007, 2008 <sup>22-23</sup> \$ # *	Germany- Austria	988 (48)	146 (7)	86 (4)	81 (4)	36 (2)	41 (2)	395 (19)	299 (14)	2072
Gmidene et al., 2008 <sup>24</sup> \$	Tunisia	110 (49)	30 (13)	17 (8)	7 (3)	7 (3)	1 (1)	34 (15)	18 (8)	224
Poznyakova et al., 2008 <sup>25</sup> &	USA	571 (55)	61 (6)	17 (2)	38 (4)	28 (3)	22 (2)	111 (11)	181 (18)	1029
Kantarjian et al., 2008 <sup>26</sup> \$ *	USA	440 (46)	27 (3)	n.s.	n.s.	21 (2)	14 (1)	123 (13)	n.s.	958
Vundinti et al., 2009 <sup>27</sup> \$ @ *	India	66 (45) £	5 (3)	11 (8)	4 (3)	1 (1)	0 (0)	58 (40)	0 (0)	145
Schanz 2011 <sup>28</sup> \$	Germany-Austria-USA	1156 (49)	146 (6)	99 (4)	132 (6)	52 (2)	42 (2)	266 (11)	458 (20)	2351
Qu et al., 2012 <sup>29</sup> &	China	186 (35)	8 (2)	n.s.	67 (13)	n.s.	n.s.	n.s.	85 (16)	532
Belli et al., 2012 <sup>30</sup> \$	Argentina	296 (57)	19 (4)	15 (3)	26 (5)	14 (3)	14 (3)	93 (18)	41 (8)	518
Schanz et al., 2012 <sup>31</sup> \$	International	1543 (55)	180 (6)	59 (2)	133 (5)	48 (2)	63 (2)	519 (19)	256 (9)	2801
Mean (Range)		52 (35-86)%	6 (0-13)%	3 (1-8)%	5 (1-13)%	2 (0-5)%	2 (1-6)%	16 (0-40)%	12 (0-27)%	

\$ = FAB classified population, # = including patients with AML, \* = including patients with secondary MDS & WHO classified population, @ = including pediatric cases, £ = including FISH results; ns = non specified, ¥ = excluding patients with Chronic Myelomonocytic Leukemia.

added to this list: segments of acquired uniparental disomy (injury resulting in loss of heterozygosity that does not alter the number of copies) that probably represent an attempt to repair chromosomal deletions using the remaining chromosome copy as template<sup>38</sup> and haploinsufficiency (when the protein produced by a normal gene copy is not sufficient to fulfil its function).<sup>39</sup>

### Risk Stratification of Cytogenetic Findings in Myelodysplastic Syndromes

Clinical course of MDS is highly variable, ranging from stable disease over 10 or more years to death within a few months due to cytopenias or transformation to acute myeloid leukemia. Since the development of

**Table 2: Normal Karyotype According to Each French-American-British Subgroup**

Series	Country	FAB Categories				
		RA (%)	RARS (%)	RAEB (%)	RAEBt (%)	CMMML (%)
Paquette et al., 1993 <sup>7</sup>	USA	53/71 (75)	30/43 (70)	23/ 54(43)	4/21 (19)	14/19 (74)
Parlier et al., 1994 <sup>8</sup>	Switzerland	10/21 (48)	13/26 (50)	7/16 (44)	3/10 (30)	14/25 (56)
de Souza Fernandez, et al., 2000 <sup>10</sup>	Brazil	59/74 (80)	8/9 (89)	6/21 (29)	0/14 (0)	3/7 (43)
Solé et al., 2000 <sup>11</sup>	Spain	104/184 (57)	51/76 (67)	80/208 (39)	24/79 (30)	54/93 (58)
Nösslinger et al., 2001 <sup>32</sup>	Austria	(45)	(55)	(48)	(38)	(52)
Matsuda et al, 2005 <sup>33</sup>	Japan	72/102 (71)				
	Germany	94/199 (47)				
Solé et al., 2005 <sup>16</sup>	Spain	151/264 (57)	106/160 (66)	109/ 269 (41)	31/ 105 (30)	117/170 (69)
Bernasconi et al., 2007 <sup>21</sup>	Italy	99/209 (47)	45/79 (57)	44/152 (29)	9/51 (18)	Not included
Haase et al., 2007 <sup>22</sup>	Germany-Austria	267/573 (47)	147/252 (58)	179/415 (43)	132/305 (43)	170/272 (63)
Gmidene et al., 2008 <sup>24</sup>	Tunisia	36/78 (46)	10/21 (48)	29/ 57 (51)	6/16 (38)	20/25 (80)
Vandinti et al., 2009 <sup>27</sup>	India	29/55 (53)	7/9 (78)	17/41 (41)	6/30 (20)	7/10 (70)
Belli et al., 2011 <sup>34</sup>	Argentina	133/208 (64)	22/36 (61)	51/103 (50)	9/25 (36)	30/49 (61)
Mean % (Range)		54 (45-80)	59 (48-89)	41 (29-51)	34 (0-43)	64 (43-80)

RA = Refractory Anemia, RARS = RA with ringed sideroblast, RAEB = RA with excess of blasts, RAEBt = RAEB in transformation, CMMML = Chronic Myelomonocytic Leukemia

the Bournemouth index in 1985<sup>40</sup>, various scoring systems have been designed, based on clinical characteristics at presentation, in order to define prognostic subgroups.

The karyotype was recognized as an independent prognostic factor in 1993, and its inclusion among different prognostic systems has contributed to improve the assessment of prognosis.<sup>6</sup> Morel et al., 1993, confirmed that the presence of complex karyotypes ( $\geq 3$  altered chromosomes) was associated with short survival and leukemic evolution,<sup>6</sup> whilst Lausanne-Bournemouth index included the presence of 2 aberrations among the worst prognostic karyotypes<sup>35</sup> (see Table 3).

In 1997, the International Prognostic Scoring System (IPSS)<sup>9</sup> established that the most frequent cytogenetic findings were associated with specific prognosis and clearly differentiated three cytogenetic risk categories: good, intermediate and poor with median survivals of 46, 29 and 10 months, respectively. This modality was later adopted by the WHO classification-based Prognostic Scoring System (WPSS) in 2005.<sup>41</sup> Complex karyotypes ( $\geq 3$  cytogenetic alterations) and -7/del(7q) were associated with worse prognosis and unfavorable outcome after allogeneic bone marrow transplant.<sup>42</sup> Schanz et al. showed that the independent prognostic impact of poor risk cytogenetics on the overall survival was equivalent to the impact of high blast counts and its predictive power was unaffected by type of therapy given.<sup>28</sup> Normal karyotype, Y chromosome loss, isolated del(5q) or del(20q) constituted good prognostic findings that were associated with longer survival and reduced risk of progression. However, the intermediate group, including trisomy 8 and other miscellaneous, less frequent single and double chromosome defects, showed considerable heterogeneity.<sup>9</sup> Recent evidence suggests that some of these less common abnormalities may play a significant role in the MDS clinical outcome when larger patient series are examined (see Table 3).<sup>6,9,11,16,21-23,25,30-31,34-35,43-46</sup>

The 'prognostic index cytogenetics' (pi score)<sup>44</sup>, published in 1999, differentiated four risk groups according to cytogenetic aberrations (see Table 3). Pfeilstöcker et al., 1999, found that patients with isolated del(5q), del(20q), Y chromosome loss or other single miscellaneous findings show an intermediate-I outcome compared with normal karyotype. In addition, they described no statistically differences between patients that present -7/del(7q) or +8.<sup>44</sup>

In 2005, the Spanish Cytogenetic Working Group (GCECGH) defined four cytogenetic prognostic subgroups based on 968 patients. They included isolated del(11q) and del(12p) into the good prognosis group; rearrangements of 3q21q26, +8, +19, t(11q) and del(17p) into the intermediate-I; a new intermediate-II group of miscellaneous single or double cytogenetic abnormalities and the i(17q) into the worst prognostic group. These risk groups showed median survivals of 4.3, 2.7, 1.0 and 0.7 years, respectively.<sup>16</sup>

Bernasconi et al., 2007<sup>21</sup>, redefined the IPSS cytogenetic categories. In terms of survival, they included del(11q) and del(12p) within the good risk, del(7q) within the intermediate risk and rearrangements of 3q within the high-risk category. Patients classified accordingly showed 2-years overall survival of 90, 69 and 26 %, respectively. In terms of the risk of AML progression, the isolated del(11q) and del(12p) were associated with good outcome, del(20q) with intermediate risk; whereas, +8 with higher risk to develop AML.<sup>21</sup>

In 2007 and 2008, the German-Austrian MDS Group analyzed a series of 1,202 MDS patients treated only with supportive care and defined four different cytogenetic groups with median survival of 55, 29, 15 and 9 months, respectively. In this study, the abnormalities associated with a favorable clinical course, with median survivals longer than 32 months, were: normal karyotype, t(1q), del(5q), t(7q), del(9q), del(12p), del(15q)/t(15q), t(17q), del(20q), +21, -21, -X, and -Y. The intermediate-I alterations

**Table 3: Cytogenetic Risk Groups according to different Prognostic Scoring Systems**

Cytogenetic Groups of Risk						
Prognostic Systems		Very Good	Good	Intermediate		Very Poor
				Intermediate I	Intermediate II	
Lille (Morel et al., 1993) <sup>6</sup>			Normal Karyotype 1 or 2 altered chromosomes			≥ 3 altered chromosomes
Lausanne-Bournemouth (Parlier et al., 1995) <sup>35</sup>			Normal Karyotype 1 cytogenetic alteration			≥ 2 alterations
IPSS (Greenberg et al., 1997) <sup>9</sup>			Normal Karyotype -Y, del(5q), del(20q)	+8, 1 or 2 cytogenetic alterations (others)		≥ 3 alterations -7/del(7q)
Cytogenetic Prognostic Index (Pi) (Pfeilstöcker et al., 1999) <sup>§44</sup>			Normal Karyotype	1 point	2 o 3 points	≥ 4 points
GCECGH (Solé et al., 2005) <sup>16</sup>			Normal Karyotype, -Y, del(5q), del(20q), del(11q), del(12p)	+8, Rearr. 3(q21q26), +9, +18, +19, t(11q), del(17p)	cytogenetic alterations (others)	≥ 3 alterations; -7/del(7q), i(17q)
WPSS (Malcovati et al., 2005) <sup>17</sup>			Normal Karyotype -Y, del(5q), del(20q)	+8, 1 or 2 cytogenetic alterations (others)		≥ 3 alterations -7/del(7q)
(Bernasconi et al., 2007) <sup>21</sup>	SV		Normal Karyotype, -Y, del(5q), del(20q), del(11q), del(12p)	+8, del(7q), 1 or 2 cytogenetic alterations (others)		Rearr. 3(q21q26), -7, ≥ 3 alterations
	PFI		Normal Karyotype, -Y, del(5q), del(11q), del(12p)	Rearr. 3(q21q26), del(20q), 1 or 2 cytogenetic alterations (others)		-7/del(7q), +8, ≥ 3 alterations
German- Austrian Group (Haase et al., 2007, 2008) <sup>22-23</sup> IPSS-R (Schanz et al., 2010) <sup>46</sup>			Normal Karyotype; -Y, -X, del(5q), del(20q), +1/+1q, t(1q), t(7q), del(9q), del(12p), chr. 15 abn., t(17q), +/-21	+8 del(11q)	= 3 cytogenetic alterations; -7/del(7q); 3q rearr., +19, t(11)(q23)	> 3 alterations t(5q)
MDACC (Kantarjian et al., 2008) <sup>26</sup>			Normal Karyotype, 1 o 2 cytogenetic alterations (others)			≥ 3 alterations -7/del(7q)
German- Austrian Group (Haase et al., 2009) <sup>45</sup>			Normal Karyotype, -Y, del(5q), del(20q), del(11q), t(11)(q23), del(12p), +21 and 2 cytogenetic alterations including chr. 5	+8, +1q, rearr. 3q, t(7q), +19, -21, 2 cytogenetic alterations excluding chr. 5 or 7	-X -7/del(7q) simplex o double = 3 alterations	> 3 alterations
IPSS-R (Schanz et al., 2010) <sup>46</sup>	del(11q), -Y		Normal karyotype, der(1;7), del(5q), del(12p), del(20q), or double incl. del(5q)	-7/7q-, +8, i(17q), +19, +21, any other single, or any other double, independent clones		der(3)(q21)/der(3)(q26), double incl. -7/del(7q), complex 3 abnormalities
IPSS-MK (Belli et al., 2011, 2012) <sup>30,34</sup>			Normal Karyotype, -Y, del(5q), del(20q), other isolated deletions	+8, 1 o 2 cytogenetic alterations (others)		≥ 3 alterations -7/del(7q), MK
IPSS-R (Schanz et al., 2012) <sup>31</sup>	del(11q), -Y		Normal Karyotype, del(5q), del(12p), del(20q), 2 cytogenetic alterations including chr. 5	del(7q), +8, i(17q), +19, any other single or double, independent clones		der(3)(q21)/der(3)(q26), -7, double incl. -7/del(7q), complex 3 abnormalities

§ 1 point: alterations chr 5, non complex alterations excluding chr. 5, 7 and 8; 3 points: chr 7 or 8 alterations; 4 points: complex alterations. SV: Survival; PFI: progression free interval; MK: Monosomal Karyotype.

were isolated del(11q) and +8, with median survival of 26 and 23 months, respectively. The intermediate-II alterations were t(11q23), rearrangements of 3q, +19, -7/del(7q) and complex (= 3 anomalies), with median survivals ranging from 20 to 14 months. In the very poor prognostic group were complex (> 3 anomalies) and t(5q) with median survival of 9 and 4 months, respectively.<sup>22-23</sup> In 2009, the same group presented a new stratification for some of above described alterations<sup>45</sup> (see *Table 3*).

In 2010, the German–Austrian group, the International MDS risk analysis workshop (IMRAW), the GCECGH, and the International Cytogenetics Working Group of the MDS Foundation (ICWG) and the M.D. Anderson Cancer Center (MDACC), proposed five cytogenetic prognostic subgroups for the IPSS-revised (IPSS-R) stratification.<sup>46</sup> In 2012, after some modifications, the final published categories, based on 2801 patients, were defined as follows: very good [del(11q) or -Y], good [normal, del(5q), del(12p), del(20q), or double abnormalities including del(5q)], intermediate [del(7q), +8, i(17q), +19, any other single or double, independent clones], poor [der(3)(q21)/ der(3)(q26), -7, double incl. -7/del(7q), or complex (= 3 abnormalities)], and very poor [complex (>3 abnormalities)] with median overall survivals of 61, 49, 26, 16 and 6 months, respectively.<sup>31</sup>

Recently we have found that the presence of an isolated deletion, excluding del(7q), is a good prognostic finding, while the presence of a monosomal karyotype (MK) is a high risk marker (IPSS-MK). This alternative but complementary strategy to the original IPSS system showed an independent prognostic impact and a better discriminating power than not only the original IPSS categories<sup>34</sup> but also the initially proposed IPSS-R ones in our series.<sup>30</sup> MK refers to the presence of two or more distinct autosomal monosomies or a single monosomy in the presence of structural abnormalities.<sup>47</sup> New published data supported the adverse prognosis associated with MK in MDS.<sup>48</sup>

## Cytogenetic Alterations Isolated Y Chromosome Loss

Loss of Y chromosome is relatively common with a reported frequency of 2 % (range 1–6 %) for the overall MDS population (see *Table 1*). Y chromosome loss is dependent on age, diagnosis, and appears to be neutral or favorable for survival or therapy with no consistent relationship between changes in the percentage and disease progression or remission.<sup>49</sup> Although the United Kingdom Cancer Cytogenetics Group study sustain that Y loss should not be interpreted as a marker of the malignant clone in elderly<sup>50</sup>, Wiktor et al. found that if the clone is present in >75 % of metaphase cells, it probably represents a disease-associated clonal population. In addition, Y chromosome loss appeared to be more frequent in AML and MDS when compared to a control group or to those with Myeloproliferative Disorders or B-cell diseases.<sup>49</sup>

## Isolated deletion of the long arm of chromosome 5

The isolated del(5q) is observed with a reported frequency of 6 % (range 0–13 %) in different published series (see *Table 1*). The 5q-syndrome is characterized by the presence of a del(5q) as the sole karyotypic abnormality, female preponderance, macrocytic anemia, normal to high platelet counts, less than 5 % of blasts in bone marrow, dismegakaryopoiesis with hypolobulated megakaryocytes, long survival with low risk of leukemic evolution<sup>51</sup> and good response to lenalidomide treatment.<sup>52</sup> The clinical outcome of del(5q) patients is

not determined by the different breakpoints of the interstitial deletion, but instead by marrow blast cell percentage, being more evident for blast cell counts more than 10 %<sup>21</sup> and by a therapy-related etiology. In addition, the good prognosis is significantly modified by additional cytogenetic changes or mutations that can transform the clone allowing its expansion.<sup>22-23,53-56</sup> These adverse marrow subclones might be present in patients with early stage disease and may expand due to the acquisition of new genetic alterations or because the clones are insensitive to lenalidomide<sup>57</sup>, therefore monitoring of cytogenetic response is mandatory.<sup>58</sup> TP53 mutations were present in almost one fifth of low-risk MDS patients with del(5q) and were associated with lower complete cytogenetic response and subsequent leukemic evolution.<sup>56</sup> In addition, TP53 mutations were associated with del(5q) in the context of complex karyotype and lenalidomide treatment had no effect on the majority of these patients.<sup>59</sup>

The deleted region varies in size: the proximal endpoints vary from 5q11 to 5q23 while the distal breaks from 5q31 to 5qter.<sup>60</sup> The common deleted region contains more than 24 genes.<sup>37</sup> A recent study implicated haploinsufficiency of multiple genes as the relevant genetic consequence of this deletion.<sup>39</sup> The decreased expression of these genes, including the ribosomal protein 14 gene (RPS14), miR-145, miR-146a and the tumor suppressor gene secreted protein acidic and rich in cysteine (SPARC), among others, may cooperate to cause several of the key features of the 5q- syndrome and may be related to lenalidomide response in these patients.<sup>37,55-56,61-65</sup>

## Partial or total loss of chromosome 7

Isolated -7/del(7q) is observed with a reported frequency of 3 % (1–8 %) (see *Table 1*). Several studies suggest ‘critical regions’ with a marked heterogeneity in the breakpoints involving 7q22, 7q32-34 and 7q35-36 bands.<sup>66-69</sup> These distinct critical loci may contribute alone or in combination to the evolution of MDS and AML.<sup>70</sup> A correlation between survival and deletion limits in del(7q) myeloid neoplasms suggests that the size of the del(7q) clone, presence of complex aberrations, refractoriness to therapy, survival and clinical progression depend on the deleted region on the del(7q) marker.<sup>71</sup> The MLL5 is a MLL family member located within the human chromosome band 7q22 and seems to play an essential role in regulating proliferation and functional integrity of hematopoietic stem/progenitor cells.<sup>72</sup>

The number of apoptotic and Fas-expressing CD34 cells was decreased in MDS patients with monosomy 7 (-7)<sup>73</sup>, and the gene expression pattern in their progenitor cells was consistent with functional characteristics of high proliferation and malignant potential.<sup>74</sup> In addition, monosomy 7 CD34 cells showed increased expression of the Granulocyte Colony Stimulating Factor receptor (GCSFR) class IV mRNA isoform which is defective in signaling cell maturation and differentiation.<sup>75</sup>

The presence of an isolated -7/del(7q) was considered as an unfavorable finding by the ICWG<sup>76</sup> and also by the IPSS<sup>9</sup>, the GCECGH<sup>16</sup>, WPSS, MDACC<sup>26</sup> and the IPSS-MK<sup>30,34</sup> systems, among others. However, other risk stratifications assigned an intermediate-I or-II risk to -7/del(7q).<sup>22-23,44,46</sup> Despite the different risk categorization assigned to these alterations median survivals show good consistency, which vary from 14 to 16 months, when both alterations are plotted together.<sup>16,23,29-30</sup> However, when these alterations were separately analyzed, del(7q) was more



favorable as compared with the loss of the whole chromosome 7 with regard to overall survival, with median survivals ranging from 16–26 and 9–16 months, respectively<sup>16,21,23,25,31</sup>, as well as risk of AML transformation.<sup>31</sup>

## Trisomy 8

The trisomy 8 (+8) is a unique cytogenetic abnormality in 5 % (1–13 %) of MDS patients (see *Table 1*). Up-regulated genes in patients with trisomy 8 were primarily involved in immune and inflammatory responses.<sup>74,77</sup> All patients had significant CD8+ T-cell expansions of one or more T-cell receptor VB subfamilies. These findings are consistent with an immune response where activated T cells, in proximity to trisomy 8 cells which would express neoantigen, would release cytokines up-regulating Fas expression on the surface of hematopoietic cells.<sup>73</sup> MDS patients with +8 respond to immunosuppressive therapies with durable reversal of cytopenias and restoration of transfusion independence.<sup>78</sup>

## Deletion (20q)

The isolated del(20q) is observed in 2 % (range 0–5 %) of MDS patients (see *Table 1*). These patients present with higher reticulocyte, lower platelet, and marrow blast counts.<sup>79</sup> The deletion is interstitial with heterogeneity of both centromeric and telomeric breakpoints including a commonly deleted region corresponding to the 20q11.2–q12 bands where several genes are located.<sup>36</sup> The most attractive candidate is the human homologue of a Drosophila tumor suppressor gene (L3MBTL1) which regulates chromatin structure during mitosis.<sup>80</sup>

## Complex Karyotypes

Multiple chromosome rearrangements ( $\geq 3$  cytogenetic alterations) are found in 12 % (range 0–27 %) (see *Table 1*) and 50 % of patients with *de novo* and secondary MDS, respectively. They often result in chromosome loss of 5q (50 %) (as interstitial deletion or as a result of unbalanced translocations), del(7q) (40 %) and del(17p) [frequently accompanied by del(5q)], gain of chromosome 8 and rare specific translocations.<sup>16,54,81–82</sup> Characteristics associated with complex karyotypes include old age, short survival time<sup>82</sup>, and its prognostic value has been shown to be independent of blast cell percentage.<sup>43</sup> The median survival seems to depend on the number of observed aberrations, and ranges from 16–17 and 9–6 months for patients with 3 and  $>3$  abnormalities, respectively.<sup>22–23,31</sup> Among complex karyotypes, around 80 % fulfilled the monosomal karyotype (MK) criteria associated with significant inferior survival.<sup>48</sup>

## Less frequent cytogenetic alterations

Different series include a wide range of less frequent ( $<1$  %) alteration, for example: rearrangements (3q), +9/del(9q), +11/del(11q), del(12p)/t(12p), -13/del(13q), del(17p)/i(17q)/t(17p) idic(Xq) and several balanced translocations such as t(6;9)(p23;q34), resulting in a fusion of DEK and CAN/NUP214; t(11;16)(q23;p13), leading to a chimeric MLL/CREBBP gene; t(2;11)(p21;q23), which leads to a strong up-regulation of microRNA miR-125b-1, among others.<sup>9–10,16,11,21,46</sup>

Major abnormalities of 3q, including t(1;3)(p36;q21), inv(3)(q21q26), t(3;3)(q21;q26), t(3;5)(q25;q34) or t(3;21)(q26;q22), are observed in 0.5–2 % of MDS and AML patients, and 30–50% of them are therapy related. Patients with rearrangements of 3q26 usually present with excess of blast, abnormal multilineage hematopoiesis, frequent dismegakaryopoiesis, normal or elevated

platelet counts and inappropriate expression of the zinc finger transcription factor oncogene EVI1 located at 3q26. The patients show a particularly aggressive course of disease and unfavorable treatment outcome characterized by short median survival and high risk of disease progression to AML.<sup>83–85</sup>

Deletions of chromosome 11q [del(11q)] as part of a non-complex karyotype are present predominately in *de novo* MDS with an overall 0.6% frequency. Wang et al. described 32 MDS cases that were characterized by lack of cryptic MLL rearrangements, transfusion-dependent anemia (65 %), frequent ring sideroblasts (59 %), bone marrow hypocellularity (22 %) and less severe thrombocytopenia, with an overall survival of 35 months, and 38 % cases progressed to AML.<sup>86</sup> However, a study of 20 (0.7 %) *de novo* MDS patients, found an overall median survival of 141 months. As a result, the IPSS-R classified del(11q) as a very good prognostic finding.<sup>31</sup>

Trisomy 11 (+11) is observed rarely in MDS and its presence correlates with clinical aggressiveness, significantly inferior survival to patients in the IPSS intermediate-risk cytogenetic group and would be best considered a high-risk cytogenetic abnormality in MDS prognosis. Moreover, MLL partial tandem duplication is observed in 50 % of MDS patients.<sup>87</sup>

Cytogenetic abnormalities of 12p in hematologic malignancies result in different molecular changes: deletions, amplifications and structural rearrangements. FISH analysis of patients with del(12p) demonstrated that the deletions are interstitial,<sup>88</sup> and overlapped deleted region might be localized between CDKN1B (KIP7, p27) and ETV6 (TEL) genes.<sup>89</sup> Recently, in a study, by single nucleotide polymorphism microarray analysis, of seven patients with del(12p) encompassing only the region centromeric of ETV6 it was observed that the minimally deleted region displays a size of 815 kb and nine genes are localized, including CDKN1B.<sup>90</sup> The clinical outcome of del(12p) patients seems to be determined by the different breakpoints of the interstitial deletion and the presence of the small deletion del(12)(p11.2p13) has been associated with a mild clinical course.<sup>91</sup> Latest reports are in agreement regarding the good outcome of MDS patients with del(12p) who show a median overall survival longer than 59 months.<sup>16,22–23,25,31,34</sup>

Alterations resulting in 17p deletion include -17, del(17p), i(17q) and unbalanced translocations that mainly involve chromosome five. They are observed in 3 % to 5 % of AML and MDS cases (most of them are therapy-related), usually as a part of complex karyotypes where del(5q) is frequently accompanying.<sup>54,81</sup> The 17p- syndrome has been described as a morphologic-cytogenetic-molecular entity based on a strong correlation between cytogenetic rearrangements leading to 17p loss, a typical form of dysgranulopoiesis combining pseudo-Pelger-Huët hypolobulation, small vacuoles in neutrophils, loss of one copy of TP53 (17p13) and mutation of the remaining TP53 allele.<sup>81</sup>

Patients with i(17q) are characterized by male predominance, severe anemia, prominent pseudo-Pelger-Huët neutrophils, granulocytic hyperplasia, increased micromegakaryocytes and poor clinical prognosis.<sup>92</sup> An association between these patients and wild-type TP53 allele was first described by Fioretos in 1999<sup>93</sup>, and recently confirmed.<sup>94</sup> The selective advantage may be conferred by gene dosage imbalances resulting from loss of 17p and gain of 17q material.

Most recent cytogenetic stratifications tried to find the prognostic significance of rare alterations [(i.e. der(1;7), rearrangements of 3q, del(11q), del(12p), i(17q), +19 and +21, among others]. The frequencies of these alterations ranged among 0.2–0.7 %, and the respective percentages represent a low number of patients which may be reflected in differences in the assessed prognostic implications. However seems to be an apparent coincidence regarding the good prognostic associated with del(12p) and del(11q).<sup>16,21-23,25,31,34,45-46</sup> Large study groups are important to obtain the minimum of 10 cases required to properly evaluate rare alterations in order to fulfill the ICWG criteria.<sup>76,95</sup>

## Discussion and Conclusion

Cytogenetic information, in addition to the development of new technologies, has contributed an additional level of biological understanding related to MDS clone. And far from becoming less relevant following the introduction of molecular, immunophenotypic, and histochemical staining techniques, the standard cytogenetic analysis is currently recommended for all MDS patients, retaining and increasing its importance in day-to-day clinical practice strategies.

Although MDS are still heterogeneous, cytogenetic findings help to define subgroups of patients who share similarities in the course of

the disease. There are recurring, non random cytogenetic alterations specially affecting chromosomes 5, 7, 8 and 20. While they do not suggest a therapeutic approach, except for the isolated del(5q), these aberrations have been considered as risk indicators from the time the original IPSS was published.<sup>9</sup> Most recent cytogenetic stratifications tried to find the prognostic significance of less frequent alterations which has been longer included in the intermediate group.<sup>16,21-23,25,30-31,34,45-46</sup> Moreover, monitoring of karyotype changes is suggested, not only to evaluate cytogenetic response to treatment but also to evaluate the acquisition of new cytogenetic alterations that are related to an unfavorable clinical outcome.<sup>96-97</sup>

The next step will be to search for unifying molecular abnormalities across the syndromes which would further refine classification, risk stratification, and, at the same time, provide potential and novel therapeutic targets. The use of approved therapeutic agents (lenalidomide, decitabine and azacitidine) in MDS patients during the past decade has provided more information as well as more questions regarding this heterogeneous pathology because only a subgroup of patients will respond. Further insights into the biology of responding versus non responding patients are required to develop new adapted scoring systems based on large available information. ■

- Bennett J, Catovsky D, Daniel M, et al., Proposals for the classification of the myelodysplastic Syndromes, *Br J Haematol*, 1982;51:189–99.
- Sanz GF, Sanz MA, Vallespi T, Etiopathogenesis, prognosis and therapy of myelodysplastic syndromes, *Hematol Cell Ther*, 1997;39:277–94.
- Heaney M, Golde G, Myelodysplasia, *N Engl J Med*, 1999;340:1649–60.
- Vardiman J, Thiele J, Arber D, et al., The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes, *Blood*, 2009;114:937–51.
- Tefferi A, Vardiman J, Myelodysplastic Syndromes, *N Engl J Med*, 2009;361:1872–85.
- Morel P, Hebbat M, Lai JL, et al., Cytogenetic analysis has strong independent prognostic value in the novo myelodysplastic syndromes and can be incorporated in a new scoring system: a report on 408 cases, *Leukemia*, 1993;7:1315–23.
- Paquette R, Landaw E, Pierre R, et al., N-RAS mutations are associated with poor prognoses and increased risk of leukaemia in myelodysplastic syndrome, *Blood*, 1993;82:590–6.
- Parlier V, van Melle G, Beris P, et al., Hematologic, Clinical, and Cytogenetic Analysis in 109 Patients with Primary Myelodysplastic Syndrome. Prognostic Significance of Morphology and Chromosome Findings, *Cancer Genet Cytogenet*, 1994;78:219–31.
- Greenberg P, Cox C, LeBeau MM, et al., International Scoring System for evaluating Prognosis in Myelodysplastic Syndromes, *Blood*, 1997;89:2079–88.
- de Souza Fernández T, Ornellas MD, Otero de Carvalho L, et al., Chromosomal alterations associated with evolution from myelodysplastic syndrome to acute myeloid leukemia, *Leuk Res*, 2000;24:839–48.
- Solé F, Espinet B, Sanz G, et al., Grupo Cooperativo Español de Citogenética Hematológica. Incidence, characterization and prognostic significance of cromosomal abnormalities in 640 patients with primary myelodysplastic syndromes, *Br J Haematol*, 2000;108:346–56.
- Belli C, Acevedo S, Bengió R, et al., Detection of risk groups in myelodysplastic syndrome. A multicenter study, *Haematologica*, 2002;87:9–16.
- Lee JH, Lee JH, Shin YR, et al., Application of different prognostic scoring systems and comparison of FAB and WHO classification in Korean patients with myelodysplastic syndrome, *Leukemia*, 2003;17:305–13.
- Lau L, Chng WJ, Liu T, et al., Clinico-pathological analysis of Myelodysplastic Syndromes according to French–American–British classification and International Prognostic Scoring System, *Ann Acad Med Singapore*, 2004;33:589–95.
- Lorand-Metze I, Pinheiro M, Ribeiro E, et al., Factors influencing survival in myelodysplastic syndromes in Brazilian population: comparison of FAB and WHO classifications, *Leuk Res*, 2004;28:587–94.
- Solé F, Luño E, Sanzo C, et al., Identification of novel cytogenetic markers with prognostic significance in a series of 968 patients with primary myelodysplastic syndromes, *Haematologica*, 2005;90:1168–78.
- Malcovati L, Porta MG, Pascutto C, et al., Prognostic factors and life expectancy in myelodysplastic syndromes classified according to WHO criteria: a basis for clinical decision making, *J Clin Oncol*, 2005;23:7594–603.
- Müller-Berndorff H, Haas P, Kunzmann R, et al., Comparison of five prognostic scoring systems, the French–American–British (FAB) and World Health Organization (WHO) classifications in patients with myelodysplastic syndromes: results of a single-center analysis, *Ann Hematol*, 2006;85:502–13.
- Irons RD, Wang X, Gross SA, et al., Prevalence of MDS subtypes in Shanghai, China: A comparison of the World Health Organization and French American British Classifications, *Leuk Res*, 2006;30:769–75.
- Sakuma T, Hayashi Y, Kanomata N, et al., Histological and cytogenetic characterization of bone marrow in relation to prognosis and diagnosis of myelodysplastic syndromes, *Pathol Int*, 2006;56:191–9.
- Bernasconi P, Klersy C, Boni M, et al., World Health Organization classification in combination with cytogenetic markers improves the prognostic stratification of patients with de novo primary myelodysplastic syndromes, *Br J Haematol*, 2007;137:193–205.
- Haase D, Germing U, Schanz J, et al., New insights into the prognostic impact of the karyotype in MDS and correlation with subtypes: evidence from a core dataset of 2124 patients, *Blood*, 2007;110:4385–95.
- Haase D, Cytogenetic features in myelodysplastic syndromes, *Ann Hematol*, 2008;87:515–26.
- Gmidène A, Sennana H, Fenaux P, et al., Cytogenetic abnormalities in Tunisian de novo myelodysplastic syndrome: a comparison with other populations, *Leuk Res*, 2008;32:1824–9.
- Pozdnyakova O, Miron PM, Tang G, et al., Cytogenetic abnormalities in a series of 1,029 patients with primary myelodysplastic syndromes: a report from the US with a focus on some undefined single chromosomal abnormalities, *Cancer*, 2008;113:3331–40.
- Kantarjian H, O'Brien S, Ravandi F, et al., Proposal for a New Risk Model in Myelodysplastic Syndrome That Accounts for Events Not Considered in the Original International Prognostic Scoring System, *Cancer*, 2008;113:1351–61.
- Vundinti B, Kerketta L, Jijina F, Ghosh K, Cytogenetic study of myelodysplastic syndrome from India, *Indian J Med Res*, 2009;130:155–9.
- Schanz J, Steidl C, Fonatsch C, et al., Coalesced multicentric analysis of 2,351 patients with myelodysplastic syndromes indicates an underestimation of poor-risk cytogenetics of myelodysplastic syndromes in the international prognostic scoring system, *J Clin Oncol*, 2011;29:1963–70.
- Qu S, Xu Z, Zhang Y, et al., Impacts of the cytogenetic categories in the Revised International Prognostic Scoring System on the prognosis of primary myelodysplastic syndromes: results of a single-center study, *Leuk Lymphoma*, 2012;53:940–6.
- Belli C, Bestach Y, Correa W, et al., Prognostic relevance of cytogenetic systems in Myelodysplastic Syndromes, *Leuk Lymphoma*, 2012;53:1640–2.
- Schanz J, Tüchler H, Solé F, et al., New comprehensive cytogenetic scoring system for primary myelodysplastic syndromes (MDS) and oligoblastic acute myeloid leukemia after MDS derived from an international database merge, *J Clin Oncol*, 2012;30:820–9.
- Nösslinger T, Reisner R, Koller E, et al., Myelodysplastic syndromes, from French–American–British to World Health Organization: comparison of classifications on 431 unselected patients from a single institution, *Blood*, 2001;98:2935–41.
- Matsuda A, Germing U, Jinnai I, et al., Difference in clinical features between Japanese and German patients with refractory anemia in myelodysplastic syndromes, *Blood*, 2005;106:2633–40.
- Belli C, Bengió R, Negri Aranguren P, et al., Partial and total monosomal karyotypes in Myelodysplastic Syndromes: comparative prognostic relevance among 421 patients, *Am J Hematol*, 2011;86:540–5.
- Parlier V, van Melle G, Beris PH, et al., Prediction of 18–Month Survival in Patients with Primary Myelodysplastic Syndrome. A regression Model and Scoring System Based on the Combination of chromosome Findings and Bournemouth Score, *Cancer Genet Cytogenet*, 1995;81:158–65.
- Bench AJ, Nacheva EP, Hood TL, et al., Chromosome 20 deletions in myeloid malignancies: reduction of the common deleted region, generation of a PAC/BAC contig and identification of candidate genes. UK Cancer Cytogenetics Group (UKCCG), *Oncogene*, 2000;19:3902–13.
- Boulwood J, Fidler C, Strickson A, et al., Narrowing and genomic annotation of the commonly deleted region of the 5q– syndrome, *Blood*, 2002;99:4638–41.
- Gondek L, Tiu R, O'Keefe C, et al., Chromosomal lesions and uniparental disomy detected by SNP arrays in MDS, MDS/MPD, and MDS-derived AML, *Blood*, 2008;111:1534–42.
- Graubert TA, Payton MA, Shao J, et al., Integrated genomic analysis implicates haploinsufficiency of multiple chromosome 5q31.2 genes in de novo myelodysplastic syndromes pathogenesis, *PLoS One*, 2009;4:e4583.
- Mufti G, Stevens J, Oscier D, et al., Myelodysplastic syndromes: a scoring system with prognostic significance, *Br J Haematol*, 1985;59:425–33.
- Malcovati L, Germing U, Kuendgen A, et al., Time-Dependent Prognostic Scoring System for Predicting Survival and Leukemic Evolution in Myelodysplastic Syndromes, *J Clin Oncol*, 2007;25:3503–10.
- Nevill TJ, Shepherd JD, Sutherland HJ, et al., IPSS poor-risk karyotype as a predictor of outcome for patients with myelodysplastic syndrome following myeloablative

stem cell transplantation, *Biol Blood Marrow Transplant*, 2009;15:205–13.

43. Bernasconi P, Klersy C, Boni M, et al., Incidence and prognostic significance of karyotype abnormalities in de novo primary myelodysplastic syndromes: a study on 331 patients from a single institution, *Leukemia*, 2005;19:1424–31.

44. Pfeilstöcker M, Reisner R, Nösslinger T, et al., Cross-validation of prognostic scores in myelodysplastic syndromes on 386 patients from a single institution confirms importance of cytogenetics, *Br J Haematol*, 1999;106:455–63.

45. Haase D, Schanz J, Tuechler H, et al., Updated cytogenetic risk features in MDS – present state, *Leuk Res*, 2009;33:S9–S10.

46. Schanz J, Tuechler H, Sole F, et al., Proposal of a new, comprehensive cytogenetic scoring system for primary MDS, *Haematologica*, 2010, 95 (Suppl.2), Abstract 535.

47. Breems DA, Van Putten WL, De Greef GE, et al., Monosomal karyotype in acute myeloid leukemia: a better indicator of poor prognosis than a complex karyotype, *J Clin Oncol*, 2008;26:4791–7.

48. Patnaik MM, Hanson CA, Hodnefield JM, et al., Monosomal karyotype in myelodysplastic syndromes, with or without monosomy 7 or 5, is prognostically worse than an otherwise complex karyotype, *Leukemia*, 2011;25:266–70.

49. Wiktor A, Rybicki BA, Piao ZS, et al., Clinical significance of Y chromosome loss in hematologic disease, *Genes Chrom Cancer*, 2000;27:11–6.

50. United Kingdom Cancer Cytogenetics Group (UKCCG), Loss of the Y chromosome from normal and neoplastic bone marrows, *Genes Chrom Cancer*, 1992;5:83–8.

51. Van der Berghe H, Vermaelen K, Mecucci C, et al., The 5q– anomaly, *Cancer Genet Cytogenet*, 1985;17:189–255.

52. List A, Kurtin S, Roe DJ, et al., Efficacy of lenalidomide in myelodysplastic syndromes, *N Engl J Med*, 2005;352:549–57.

53. Larripa J, Acevedo S, Palau V, et al., Leukaemic transformation in patients with 5q– and additional abnormalities, *Haematologica*, 1991;76:363–7.

54. Castro PD, Liang JC, Nagarajan L, Deletions of chromosome 5q13.3 and 17p loci cooperate in myeloid neoplasms, *Blood*, 2000;95:2138–43.

55. Jädersten M, Karsan A, Clonal evolution in myelodysplastic syndromes with isolated del(5q): the importance of genetic monitoring, *Haematologica*, 2011;96:177–80.

56. Jädersten M, Saft L, Smith A, et al., TP53 mutations in low-risk myelodysplastic syndromes with del(5q) predict disease progression, *J Clin Oncol*, 2011;29:1971–9.

57. Jädersten M, Saft L, Pellagatti A, et al., Clonal heterogeneity in the 5q– syndrome: p53 expressing progenitors prevail during lenalidomide treatment and expand at disease progression, *Haematologica*, 2009;94:1762–6.

58. Göhring G, Giagounidis A, Büsche G, et al., Cytogenetic follow-up by karyotyping and fluorescence in situ hybridization: implications for monitoring patients with myelodysplastic syndrome and deletion 5q treated with lenalidomide, *Haematologica*, 2011;96:319–22.

59. Möllgård L, Saft L, Treppendahl MB, et al., Clinical effect of increasing doses of lenalidomide in high-risk myelodysplastic syndrome and acute myeloid leukemia with chromosome 5 abnormalities, *Haematologica*, 2011;96:963–71.

60. Pedersen B, Jensen I, Clinical and prognostic implications of chromosome 5q deletions: 96 high resolution studied patients, *Leukemia*, 1991;5:556–73.

61. Pellagatti A, Jädersten M, Forsblom AM, et al., Lenalidomide inhibits the malignant clone and up-regulates the SPARC gene mapping to the commonly deleted region in 5q– syndrome patients, *Proc Natl Acad Sci USA*, 2007;104:11406–11.

62. Barlow JL, Drynan LF, Hewett DR, et al., A p53-dependent mechanism underlies macrocytic anemia in a mouse model of human 5q– syndrome, *Nat Med*, 2010;16:59–66.

63. Pellagatti A, Hellström-Lindberg E, Giagounidis A, et al., Haploinsufficiency of RPS14 in 5q– syndrome is associated with deregulation of ribosomal– and translation-related genes, *Br J Haematol*, 2008;142:57–64.

64. Pellagatti A, Marafioti T, Paterson JC, et al., Induction of p53 and up-regulation of the p53 pathway in the human 5q– syndrome, *Blood*, 2010;115:2721–3.

65. Wei S, Chen X, Rocha K, et al., A critical role for phosphatase haploinsufficiency in the selective suppression of deletion 5q MDS by lenalidomide, *Proc Natl Acad Sci USA*, 2009;106:12974–9.

66. Le Beau M, Espinosa R, Davis E, et al., Cytogenetic and molecular delineation of a region of chromosome 7 commonly deleted in malignant myeloid diseases, *Blood*, 1996;88:1930–5.

67. Lewis S, Abrahamson G, Boulwood J, et al., Molecular characterization of the 7q deletion in myeloid disorders, *Br J Haematol*, 1996;93:75–80.

68. Fischer K, Fröhling S, Scherer S, et al., Molecular Cytogenetic Delineation of Deletions and Translocations Involving Chromosome Band 7q22 in Myeloid Leukemias, *Blood*, 1997;89:2036–41.

69. Döhner K, Brown J, Hehmann U, et al., Molecular Cytogenetic Characterization of a Critical Region in Bands 7q35–q36 Commonly Deleted in Malignant Myeloid Disorders, *Blood*, 1998;92:4031–6.

70. Liang H, Fairman J, Claxton D, et al., Molecular anatomy of chromosome 7q deletions in myeloid neoplasms: Evidence for multiple critical loci, *Proc Natl Acad Sci USA*, 1998;95:3781–5.

71. Pedersen B, Ellegaard J, A factor encoded by 7q31 suppresses expansion of the 7q– clone and delays cytogenetic progression, *Cancer Genet Cytogenet*, 1994;78:181–8.

72. Madan V, Madan B, Brykczynska U, et al., Impaired function of primitive hematopoietic cells in mice lacking the Mixed-Lineage–Leukemia homolog MLL5, *Blood*, 2009;113:1444–54.

73. Chen G, Zeng W, Miyazato A, et al., Distinctive gene expression profiles of CD34 cells from patients with myelodysplastic syndrome characterized by specific chromosomal abnormalities, *Blood*, 2004;104:4210–8.

74. Sloand E, Kim S, Fuhrer M, et al., Fas-mediated apoptosis is important in regulating cell replication and death in trisomy 8 hematopoietic cells but not in cells with other cytogenetic abnormalities, *Blood*, 2002;100:4427–32.

75. Sloand E, Yong A, Ramkissoon S, et al., Granulocyte colony-stimulating factor preferentially stimulates proliferation of monosomy 7 cells bearing the isoform IV receptor, *Proc Natl Acad Sci USA*, 2006;103:14483–8.

76. Chun K, Hagemeljer A, Iqbal A, Slovak ML, Implementation of standardized international karyotype scoring practices is needed to provide uniform and systematic evaluation for patients with myelodysplastic syndrome using IPSS criteria: An International Working Group on MDS Cytogenetics Study, *Leuk Res*, 2010;34:160–5.

77. Sloand EM, Pfannes L, Chen G, et al., CD34 cells from patients with trisomy 8 myelodysplastic syndrome (MDS) express early apoptotic markers but avoid programmed cell death by up-regulation of antiapoptotic proteins, *Blood*, 2007;109:2399–405.

78. Sloand EM, Mainwaring L, Fuhrer M, et al., Preferential suppression of trisomy 8 compared with normal hematopoietic cell growth by autologous lymphocytes in patients with trisomy 8 myelodysplastic syndrome, *Blood*, 2005;106:841–51.

79. Braun T, de Botton S, Taksin AL, et al., Characteristics and outcome of myelodysplastic syndromes (MDS) with isolated 20q deletion: a report on 62 cases, *Leuk Res*, 2011;35:863–7.

80. MacGrogan D, Alvarez S, DeBlasio T, Jhanwar SC, Nimer SD, Identification of candidate genes on chromosome band 20q12 by physical mapping of translocation breakpoints found in myeloid leukemia cell lines, *Oncogene*, 2001;20:4150–60.

81. Soenen V, Preudhomme C, Roumier C, et al., 17p Deletion in Acute Myeloid Leukemia and Myelodysplastic Syndrome. Analysis of Breakpoints and Deleted Segments by Fluorescence In Situ, *Blood*, 1998;91:1008–15.

82. Lindvall C, Nordenskjöld M, Porwit A, et al., Molecular cytogenetic characterization of acute myeloid leukemia and myelodysplastic syndromes with multiple chromosome rearrangements, *Haematologica*, 2001;86:1158–64.

83. Testoni N, Borsaru G, Martinelli G, et al., 3q21 and 3q26 cytogenetic abnormalities in acute myeloblastic leukemia: biological and clinical features, *Haematologica*, 1999;84:690–4.

84. Cui W, Sun J, Cotta CV, et al., Myelodysplastic syndrome with inv(3)(q21q26.2) or t(3;3)(q21;q26.2) has a high risk for progression to acute myeloid leukemia, *Am J Clin Pathol*, 2011;136:282–8.

85. Martinelli G, Ottaviani E, Buonamici S, et al., Association of 3q21q26 syndrome with different RPN1/EV11 fusion transcripts, *Haematologica*, 2003;88:1221–8.

86. Wang SA, Abruzzo LV, Hasserjian RP, et al., Myelodysplastic syndromes with deletions of chromosome 11q lack cryptic MLL rearrangement and exhibit characteristic clinicopathologic features, *Leuk Res*, 2011;35:351–7.

87. Wang SA, Jabbar K, Lu G, et al., Trisomy 11 in myelodysplastic syndromes defines a unique group of disease with aggressive clinicopathologic features, *Leukemia*, 2010;24:740–7.

88. Kobayashi H, Montgomery KT, Bohlander SK, et al., Fluorescence in situ hybridization mapping of translocations and deletions involving the short arm of human chromosome 12 in malignant hematologic diseases, *Blood*, 1994;84:3473–82.

89. Sato Y, Suto Y, Pietenpol J, et al., TEL and KIP1 define the smallest region of deletions on 12p13 in hematopoietic malignancies, *Blood*, 1995;86:1525–33.

90. Haeflrich C, Bacher U, Kohlmann A, et al., CDKN1B, encoding the cyclin-dependent kinase inhibitor 1B (p27), is located in the minimally deleted region of 12p abnormalities in myeloid malignancies and its low expression is a favorable prognostic marker in acute myeloid leukemia, *Haematologica*, 2011;96:829–36.

91. Streubel B, Sauerland C, Heil G, et al., Correlation of cytogenetic, molecular cytogenetic, and clinical findings in 59 patients with ANLL or MDS and abnormalities of the short arm of chromosome 12, *Br J Haematol*, 1998;100:521–33.

92. Xiao Z, Liu S, Yu M, et al., Isochromosome 17q in patients with Myelodysplastic Syndrome: six new cases, *Haematologica*, 2003;88:714–5.

93. Fioretos T, Strömbeck B, Sandberg T, et al., Isochromosome 17q in blast crisis of chronic myeloid leukemia and in other hematologic malignancies is the result of clustered breakpoints in 17p11 and is not associated with coding TP53 mutations, *Blood*, 1999;94:225–32.

94. Kanagal-Shamanna R, Bueso-Ramos CE, Barkoh B, et al., Myeloid neoplasms with isolated isochromosome 17q represent a clinicopathologic entity associated with myelodysplastic/myeloproliferative features, a high risk of leukemic transformation, and wild-type TP53, *Cancer*, 2011, Oct 28. [Epub ahead of print].

95. Slovak M, Dewald G, International Working Group on MDS cytogenetics: October 2007 meeting report, *Leuk Res*, 2008;32:1329–32.

96. Bernasconi P, Klersy C, Boni M, et al., Does cytogenetic evolution have any prognostic relevance in myelodysplastic syndromes? A study on 153 patients from a single institution, *Ann Hematol*, 2010;89:545–51.

97. Wang H, Wang XQ, Xu XP, Lin GW, Cytogenetic evolution correlates with poor prognosis in myelodysplastic syndrome, *Cancer Genet Cytogenet*, 2010;196:159–66.