

Hyperfine interaction of individual atoms on a surface

Science 19 Oct 2018:

Vol. 362, Issue 6412, pp. 336-339

DOI: 10.1126/science.aat7047

1. Philip Willke^{1,2,3},
2. Yujeong Bae^{1,2,3},
3. Kai Yang¹,
4. Jose L. Lado^{4,5},
5. Alejandro Ferrón⁶,
6. Taeyoung Choi^{2,3},
7. Arzhang Ardavan⁷,
8. Joaquín Fernández-Rossier^{4,*},
9. Andreas J. Heinrich^{2,3,†},
10. Christopher P. Lutz^{1,†}

1. ¹IBM Almaden Research Center, San Jose, CA 95120, USA.

2. ²Center for Quantum Nanoscience, Institute for Basic Science (IBS), Seoul 03760, Republic of Korea.

3. ³Department of Physics, Ewha Womans University, Seoul 03760, Republic of Korea.

4. ⁴QuantaLab, International Iberian Nanotechnology Laboratory (INL), 4715-310 Braga, Portugal.

5. ⁵Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland.

6. ⁶Instituto de Modelado e Innovación Tecnológica (CONICET-UNNE), Facultad de Ciencias Exactas, Naturales y Agrimensura, Universidad Nacional del Nordeste, W3404AAS Corrientes, Argentina.

7. ⁷Centre for Advanced Electron Spin Resonance, Clarendon Laboratory, Department of Physics, University of Oxford, Oxford OX1 3PU, UK.

Bajo Embargo