

Domination parameters with number 2: interrelations and algorithmic consequences

Flavia Bonomo* Boštjan Brešar† Luciano N. Grippo‡ Martin Milanič§
Martín D. Safe¶

November 7, 2016

Abstract

In this paper, we study the most basic domination invariants in graphs, in which number 2 is intrinsic part of their definitions. We classify them upon three criteria, two of which give the following previously studied invariants: the weak 2-domination number, $\gamma_{w2}(G)$, the 2-domination number, $\gamma_2(G)$, the $\{2\}$ -domination number, $\gamma_{\{2\}}(G)$, the double domination number, $\gamma_{\times 2}(G)$, the total $\{2\}$ -domination number, $\gamma_{t\{2\}}(G)$, and the total double domination number, $\gamma_{t\times 2}(G)$, where G is a graph in which a corresponding invariant is well defined. The third criterion yields rainbow versions of the mentioned six parameters, one of which has already been well studied, and three other give new interesting parameters. Together with a special, extensively studied Roman domination, $\gamma_R(G)$, and two classical parameters, the domination number, $\gamma(G)$, and the total domination number, $\gamma_t(G)$, we consider 13 domination invariants in graphs. In the main result of the paper we present sharp upper and lower bounds of each of the invariants in terms of every other invariant, a large majority of which are new results proven in this paper. As a consequence of the main theorem we obtain new complexity results regarding the existence of approximation algorithms for the studied invariants, matched with tight or almost tight inapproximability bounds, which hold even in the class of split graphs.

Keywords: graph domination, total domination, rainbow domination, 2-domination, integer domination, double domination, split graph, approximation algorithm, inapproximability

AMS subject classification (2010): 05C69, 05C85, 68R10, 68W25, 94C15

*Departamento de Computación, FCEyN, Universidad de Buenos Aires, and CONICET, Argentina. E-mail address: fbonomo@dc.uba.ar

†Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia, and Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia. E-mail address: bostjan.bresar@um.si

‡Instituto de Ciencias, Universidad Nacional de General Sarmiento, Los Polvorines, Buenos Aires, Argentina. E-mail address: lgrippo@ungs.edu.ar

§University of Primorska, UP IAM, Muzejski trg 2, SI-6000 Koper, Slovenia, and University of Primorska, UP FAMNIT, Glagoljaška 8, SI-6000 Koper, Slovenia. E-mail address: martin.milanic@upr.si

¶Departamento de Matemática, Universidad Nacional del Sur, Bahía Blanca, Buenos Aires, Argentina, and Instituto de Ciencias, Universidad Nacional de General Sarmiento, Los Polvorines, Buenos Aires, Argentina. E-mail address: msafe@uns.edu.ar

1 Introduction

1.1 Prologue

A continuously growing interest in the area of graph domination, which arises from both practical applications and combinatorial challenges, has made the theory rather incoherent; two monographs surveying domination theory were published almost twenty years ago [46,47]. Due to a large number of domination-type concepts, it is not always easy to notice and appreciate some deep results that capture a broad aspect of the theory. Several results in domination theory have been in some sense rediscovered, because an approach that works for one concept can often be used with some slight adjustment for several other related concepts. We wish to make a step in the direction of making the situation more transparent, by classifying some of the most basic domination invariants, in which number 2 is involved in the definition. We make a comparison of their values in graphs between each pair of them, and as a consequence, since the discovered translations between parameters can be efficiently constructed, a general approach that joins some algorithmic and complexity issues on all of these concepts is established. In many cases our results imply that an algorithm for one invariant gives a good approximation algorithm for some other invariant; in addition, strong inapproximability results are inferred for almost all considered parameters, which hold even in the class of split graphs. (Let us mention that in [7] some connections between a (different and smaller) group of domination parameters has been established, yet the main focus was on claw-free graphs.)

1.2 Classification of parameters

The central focus of the paper is on several domination invariants of graphs, which have number 2 appearing in their definition (in particular, vertices must be dominated twice or using the sum of weights 2), and we can classify them upon three different criteria. The first criterion is the set of weights that are allowed to be assigned to vertices, which can be either $\{0, 1, 2\}$ or only $\{0, 1\}$ (in rainbow versions, which we will consider in parallel, these weights can be either $\{\emptyset, \{a\}, \{b\}, \{a, b\}\}$ or only $\{\emptyset, \{a\}, \{b\}\}$). The second criterion distinguishes three possibilities with respect to the set of vertices that need to be dominated, and at the same time the type of neighborhoods, which are considered in domination. The possibilities are as follows: only vertices with weight 0 need to be dominated ('outer domination'), all vertices need to be dominated and vertices with a positive weight dominate their closed neighborhoods ('closed domination'), and finally all vertices need to be dominated and only open neighborhoods are dominated by vertices with positive weight ('open domination'). The following table shows the six concepts that arise from these two criteria, all of which have already been studied in the literature (in parenthesis a standard symbol of the corresponding graph invariant is written¹):

	$\{0, 1, 2\}$	$\{0, 1\}$
outer	weak 2-domination (γ_{w2})	2-domination (γ_2)
closed	$\{2\}$ -domination ($\gamma_{\{2\}}$)	double domination ($\gamma_{\times 2}$)
open	total $\{2\}$ -domination ($\gamma_{t\{2\}}$)	total double domination ($\gamma_{t\times 2}$)

¹The total double domination was also denoted by $\gamma_{\times 2, t}$ in the literature, and was also called the double total domination.

The third criterion is based on the so-called rainbow variations of these parameters, and thus distinguishes domination parameters as being rainbow or not. This criterion is motivated by the concept known as k -rainbow domination introduced in [9]; in the case $k = 2$ the corresponding graph invariant was denoted by γ_{r2} , see, e.g., [10]. Note that in this paper the concept will be called rainbow weak 2-domination, and the invariant will be denoted by $\tilde{\gamma}_{w2}$, suggesting that it is the rainbow counterpart of the concept of weak 2-domination, whose graph invariant is denoted by γ_{w2} . The k -rainbow domination (and 2-rainbow domination, in particular) has been considered in several papers [12–14, 68, 70, 74, 76], and is interesting also because of its strong connection with the domination of Cartesian products of graphs; in fact, some initial results on the 2-rainbow domination number in [45] were expressed in the terminology of domination of prisms. In this paper we are mainly concerned with its conceptual features, which initiates several other rainbow domination parameters. Intuitively speaking they are obtained as follows: weight 0 is replaced by the label \emptyset , weight 1 by labels $\{a\}$ and $\{b\}$, and weight 2 by the label $\{a, b\}$, while the conditions imposed by each parameter are meaningfully adjusted to the rainbow version. The main difference is that instead of the sum of values of weights, in a rainbow version one considers the union of labels, and also the condition of having weight 2 in a neighborhood corresponds to having label $\{a, b\}$.

Given a graph G its weak 2-domination number is denoted by $\gamma_{w2}(G)$, its 2-domination number by $\gamma_2(G)$, its $\{2\}$ -domination number by $\gamma_{\{2\}}(G)$, its double domination number by $\gamma_{\times 2}(G)$, its total $\{2\}$ -domination number by $\gamma_{t\{2\}}(G)$ and its total double domination number by $\gamma_{t\times 2}(G)$. (We remark that the notion of *weak 2-domination* appeared in the literature also under the name “weak 2-rainbow domination” [10].) By the above reasoning each of these parameters has its rainbow counter-part, which we will denote in a systematic way, by putting the symbol \sim above γ , indicating that we are considering the rainbow version of the known concept. Two of the parameters among $\tilde{\gamma}_2(G)$, $\tilde{\gamma}_{w2}(G)$, $\tilde{\gamma}_{\{2\}}(G)$, $\tilde{\gamma}_{\times 2}(G)$, $\tilde{\gamma}_{t\{2\}}(G)$ and $\tilde{\gamma}_{t\times 2}(G)$ (namely $\tilde{\gamma}_{\{2\}}(G)$ and $\tilde{\gamma}_{t\{2\}}(G)$) turn out to be easily expressible by the known graph invariants, and we have thus not studied them any further. We believe that other four rainbow domination parameters are worth of consideration.

There is yet another well studied domination parameter, which involves number 2, but does not directly fit into the above frame. Nevertheless, the so-called *Roman domination*, introduced in [75] (see also [20, 64]) has been considered in a number of papers, and is conceptually relevant also to our study. In the condition of the Roman dominating function, only the vertices with weight 0 must have in the neighborhood a vertex with weight 2, while there is no such restriction for the vertices with weight 1 and 2. Beside Roman domination, whose parameter is denoted by γ_R , we decided to include in our study also the two classical domination concepts, i.e., the *domination* and the *total domination*, denoted by γ and γ_t , respectively. Hence in our main result, see Table 2 (on p. 14), thirteen domination parameters are mutually compared. To stay within a reasonable length of the paper (and to stay in line with the basic classification presented in this paper) we do not consider other variations that also involve number 2 in their definitions. In particular, we do not consider the concepts that arise from basic parameters by imposing additional restrictions (such as paired domination [48], independent Roman domination [2], exact double domination [16], etc.).

1.3 Algorithmic complexity

The main result of this paper is the list of the sharp upper and lower bounds for each of the parameters, expressed in terms of any other parameter. The comparison is not only interesting in its own right, but also has several consequences regarding algorithmic and complexity properties of the invariants involved.

For some of the invariants studied in this paper NP-completeness of their decision problems was known in the literature. In addition, for $\gamma, \gamma_t, \gamma_2, \gamma_{\times 2}, \gamma_{t \times 2}$ it was known that any polynomial time approximation of these values to within a multiplicative factor of $(1 - \epsilon) \ln n$ is very unlikely even when restricted to n -vertex split graphs: it would imply $P = NP$. (See Section 6 for details.) By using the main result of this paper we are able to infer such theorems about inapproximability in split graphs for all but three considered invariants. For two of the remaining invariants (namely, for rainbow 2-domination, $\tilde{\gamma}_2$, and rainbow double domination, $\tilde{\gamma}_{\times 2}$) we obtain the same result using a direct reduction from the SET COVER problem. The only exception to the inapproximability bounds is the rainbow total double domination number, $\tilde{\gamma}_{t \times 2}$, for which we prove that there is no polynomially computable function f such that there exists an $f(n)$ -approximation algorithm for this invariant in an n -vertex split graphs for which this parameter is finite, unless $P = NP$. We prove this using a reduction from the NP-complete HYPERGRAPH 2-COLORABILITY problem.

On a positive side, for all of the invariants studied in this paper we prove the existence of approximation algorithms matching the logarithmic lower bound up to a constant factor, with an obvious exception of $\tilde{\gamma}_{t \times 2}$ and two other parameters, $\tilde{\gamma}_2$ and $\tilde{\gamma}_{\times 2}$, for which this is still open.

1.4 Organization of the paper

In Section 2 we state the definitions of the parameters studied in this paper as well as some preliminaries on three covering parameters in graphs, and summarize the definitions in Table 1. In Section 3 we present the main results, expressed in the 13×13 table (Table 2), in which rows and columns represent the considered domination parameters, and each entry contains the upper bound of the row-parameter with respect to the column-parameter in the family of all graphs for which both parameters are finite. Since the diagonal elements are trivially just the equalities, this means that altogether we have $13 \cdot 12 = 156$ sharp upper bounds between all pairs of parameters. Table 3 in the same section gives a road map for deduction of proofs, either by references to results in one of the next sections, or by references to the papers in which the results were proven, or (in many cases) by using transitivity.

In Section 4 we make the comparison of the parameters, by proving the upper bounds, if they exist, of parameters expressed as functions of other parameters. We omit the proofs of most of such bounds that can be found in the literature, as well as of those that follow by transitivity from other bounds in Table 2. Having in mind this optimization of the proofs, we only need to prove 17 propositions in this section. Then, in Section 5, we present the values of the parameters in different families of graphs, some showing the sharpness of the bounds in Table 2, and some other showing that a particular parameter is not bounded by a function of another parameter.

In Section 6 we discuss the algorithmic and complexity consequences of the bounds obtained in Section 3, proving new lower and upper bounds regarding the (in-)approximability of the corresponding optimization problems, subject to the $P \neq NP$ assumption. We combine this with a survey on previously known (in-)approximability and NP-hardness results on these parameters.

2 Definitions and preliminaries

Unless stated otherwise, we consider finite, undirected, simple graphs. Given a vertex $x \in V(G)$, $N(x) = \{v \in V(G) \mid xv \in E(G)\}$ denotes its (*open*) *neighborhood*, and $N[x] := N(x) \cup \{x\}$ is the *closed neighborhood* of x . For a graph G and $X \subseteq V(G)$, we write $N(X)$ for $(\bigcup_{v \in X} N(v)) \setminus X$. As usual, $\Delta(G)$ and $\delta(G)$ stand for the maximum, resp. the minimum degree of vertices in G .

Let $f : V(G) \rightarrow X$ be a function such that X is either a set of real numbers or a set of finite sets. For an arbitrary subset $W \subseteq V(G)$, we denote its *weight with respect to f* (or just *weight* when f is clear from the context) by

$$f(W) = \sum_{w \in W} |f(w)|,$$

where the notation $|r|$ denotes either the cardinality of r (if r is a set), or the absolute value of r (if r is a real number). For a function $f : V(G) \rightarrow X$, where X is an arbitrary set of finite sets, we denote

$$f_{\cup}(W) = \bigcup_{w \in W} f(w).$$

Next, we present definitions of all the invariants studied in the paper. Whenever an invariant is not defined for all graphs, i.e., if there is a graph G for which no function satisfying the corresponding constraints exists, we use the convention of stating that the value of the invariant in G is infinite.

Domination, total domination.

Definition 2.1. (γ , row/column 1 in Table 2)

Let $G = (V, E)$ be a graph. A dominating function of G is a function $f : V \rightarrow \{0, 1\}$ such that for all $v \in V(G)$ it holds that

$$f(N[v]) \geq 1.$$

Equivalently,

$$f(v) = 0 \implies f(N(v)) \geq 1.$$

The domination number of G is denoted by $\gamma(G)$ and equals the minimum weight $f(V)$ over all dominating functions f of G .

Any set of the form $D = \{v \in V \mid f(v) = 1\}$ where f is a dominating function of G is said to be a dominating set of G . Note that the minimum size of a dominating set equals $\gamma(G)$.

Domination number is one of the classical graphs invariants; together with several of its variations it was surveyed in two monographs [46, 47].

Definition 2.2. (γ_t , row/column 2 in Table 2)

Let $G = (V, E)$ be a graph. A total dominating function of G is a function $f : V \rightarrow \{0, 1\}$ such that for all $v \in V(G)$ it holds that

$$f(N(v)) \geq 1.$$

The total domination number of G is denoted by $\gamma_t(G)$ and equals the minimum weight $f(V)$ over all total dominating functions f of G .

Any set of the form $D = \{v \in V \mid f(v) = 1\}$ where f is a total dominating function of G is said to be a total dominating set of G . Note that the minimum size of a total dominating set equals $\gamma_t(G)$.

Clearly, the total domination number is well-defined (i.e. is finite) in graphs with no isolated vertices. The recent monograph [56] presents a thorough survey on total domination theory.

Weak 2-domination, rainbow weak 2-domination.

Definition 2.3. (γ_{w2} , row/column 3 in Table 2)

Let $G = (V, E)$ be a graph. A weak 2-dominating function of G is a function $f : V \rightarrow \{0, 1, 2\}$ such that for all $v \in V(G)$ it holds that

$$f(v) = 0 \implies f(N(v)) \geq 2.$$

The weak 2-domination number of G is denoted by $\gamma_{w2}(G)$ and equals the minimum weight $f(V)$ over all weak 2-dominating functions f of G .

Definition 2.4. ($\tilde{\gamma}_{w2}$, row/column 9 in Table 2)

Let $G = (V, E)$ be a graph. A rainbow weak 2-dominating function of G is a function $f : V \rightarrow \mathcal{P}(\{a, b\})$ such that for all $v \in V(G)$ it holds that

$$f(v) = \emptyset \implies |f_{\cup}(N(v))| \geq 2.$$

Equivalently, for all $v \in V(G)$ with $f(v) = \emptyset$, it holds that $f_{\cup}(N(v)) = \{a, b\}$. The rainbow weak 2-domination number of G is denoted by $\tilde{\gamma}_{w2}(G)$ and equals the minimum weight $f(V)$ over all rainbow weak 2-dominating functions f of G .

Rainbow weak 2-domination was introduced less than 10 years ago in [9], under the name *2-rainbow domination*; it has already been considered in a number of papers. Weak 2-domination was studied in [10] with the aim to give more insight in the (weak) 2-rainbow domination. It has probably been known before, although we were unable to find a reference confirming it. Weak 2-domination should not be confused with the concept of weak domination, as introduced in [71].

{2}-domination, rainbow {2}-domination.

Definition 2.5. ($\gamma_{\{2\}}$, row/column 4 in Table 2)

Let $G = (V, E)$ be a graph. A {2}-dominating function of G is a function $f : V \rightarrow \{0, 1, 2\}$ such that for all $v \in V(G)$ it holds that

$$f(N[v]) \geq 2.$$

The {2}-domination number of G is denoted by $\gamma_{\{2\}}(G)$ and equals the minimum weight $f(V)$ over all {2}-dominating functions f of G .

The concept of {2}-domination was introduced in 1991 [26], and considered later on in several papers. In particular, several recent papers consider the variation of Vizing's conjecture on the domination number of Cartesian products of graphs with respect to this domination invariant, see [8, 18, 57, 58].

Definition 2.6. Let $G = (V, E)$ be a graph. A rainbow {2}-dominating function of G is a function $f : V \rightarrow \mathcal{P}(\{a, b\})$ such that for all $v \in V(G)$ it holds that

$$|f_{\cup}(N[v])| \geq 2.$$

Equivalently, $f_{\cup}(N[v]) = \{a, b\}$ holds for all $v \in V(G)$. The rainbow $\{2\}$ -domination number of G is denoted by $\tilde{\gamma}_{\{2\}}(\mathbf{G})$ and equals the minimum weight $f(V)$ over all rainbow $\{2\}$ -dominating functions f of G .

It is easy to see that the rainbow $\{2\}$ -domination number is closely related to the domination number. Indeed, for every graph G , it holds that $\tilde{\gamma}_{\{2\}}(\mathbf{G}) = 2\gamma(\mathbf{G})$. Hence, we will not discuss this parameter any further in the rest of the paper, except briefly in Sections 3 and 6 (in Theorems 6.7 and 6.11).

Total $\{2\}$ -domination, rainbow total $\{2\}$ -domination.

Definition 2.7. ($\gamma_{t\{2\}}$, row/column 5 in Table 2)

Let $G = (V, E)$ be a graph. A total $\{2\}$ -dominating function of G is a function $f : V \rightarrow \{0, 1, 2\}$ such that for all $v \in V(G)$ it holds that

$$f(N(v)) \geq 2.$$

The total $\{2\}$ -domination number of G is denoted by $\gamma_{t\{2\}}(\mathbf{G})$ and equals the minimum weight $f(V)$ over all total $\{2\}$ -dominating functions f of G .

Clearly, the total $\{2\}$ -domination number is finite precisely in graphs with no isolated vertices. While the concept has been known for some time, see two recent papers on the total $\{k\}$ -domination [4, 63].

Definition 2.8. Let $G = (V, E)$ be a graph. A rainbow total $\{2\}$ -dominating function of G is a function $f : V \rightarrow \mathcal{P}(\{a, b\})$ such that for all $v \in V(G)$ it holds that

$$|f_{\cup}(N(v))| \geq 2.$$

Equivalently, $f_{\cup}(N(v)) = \{a, b\}$ holds for all $v \in V(G)$. The rainbow total $\{2\}$ -domination number of G is denoted by $\tilde{\gamma}_{t\{2\}}(\mathbf{G})$ and equals the minimum weight $f(V)$ over all rainbow total $\{2\}$ -dominating functions f of G .

Similarly as the rainbow $\{2\}$ -domination number is related to the domination number via the relation $\tilde{\gamma}_{\{2\}}(\mathbf{G}) = 2\gamma(\mathbf{G})$, the rainbow total $\{2\}$ -domination number is related to the total domination number via the relation $\tilde{\gamma}_{t\{2\}}(\mathbf{G}) = 2\gamma_t(\mathbf{G})$. Hence, we will not discuss this parameter any further in the rest of the paper, except briefly in Sections 3 and 6 (in Theorems 6.7 and 6.11).

2-domination, rainbow 2-domination.

Definition 2.9. (γ_2 , row/column 6 in Table 2)

Let $G = (V, E)$ be a graph. A 2-dominating function of G is a function $f : V \rightarrow \{0, 1\}$ such that for all $v \in V(G)$ it holds that

$$f(v) = 0 \implies f(N(v)) \geq 2.$$

The 2-domination number of G is denoted by $\gamma_2(\mathbf{G})$ and equals the minimum weight $f(V)$ over all 2-dominating functions f of G .

Any set of the form $D = \{v \in V \mid f(v) = 1\}$ where f is a 2-dominating function of G is said to be a 2-dominating set of G . Note that the minimum size of a 2-dominating set equals $\gamma_2(\mathbf{G})$.

The concept of k -domination (and 2-domination in particular) was introduced back in 1985 [32], and was later studied quite extensively, see some recent papers [11, 22, 29, 42].

Definition 2.10. ($\tilde{\gamma}_2$, row/column 10 in Table 2)

Let $G = (V, E)$ be a graph. A rainbow 2-dominating function of G is a function $f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$ such that

$$f(v) = \emptyset \implies |f_{\cup}(N(v))| \geq 2.$$

Equivalently, for all $v \in V(G)$ with $f(v) = \emptyset$, it holds that $f_{\cup}(N(v)) = \{a, b\}$. The rainbow 2-domination number of G is denoted by $\tilde{\gamma}_2(G)$ and equals the minimum weight $f(V)$ over all rainbow 2-dominating functions f of G .

Double domination, rainbow double domination.

Definition 2.11. ($\gamma_{\times 2}$, row/column 7 in Table 2)

Let $G = (V, E)$ be a graph. A double dominating function of G is a function $f : V \rightarrow \{0, 1\}$ such that for all $v \in V(G)$ it holds that

$$f(N[v]) \geq 2.$$

The double domination number of G is denoted by $\gamma_{\times 2}(G)$ and equals the minimum weight $f(V)$ over all double dominating functions f of G .

Any set of the form $D = \{v \in V \mid f(v) = 1\}$ where f is a double dominating function of G is said to be a double dominating set of G . Note that the minimum size of a double dominating set equals $\gamma_{\times 2}(G)$.

Double domination number is finite in graphs without isolated vertices. It was introduced in [44] (see also [43]), and was studied by a number of authors; consider for instance some recent papers [6, 23, 27, 61].

Definition 2.12. ($\tilde{\gamma}_{\times 2}$, row/column 11 in Table 2)

Let $G = (V, E)$ be a graph. A rainbow double dominating function of G is a function $f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$ such that for all $v \in V(G)$ it holds that

$$|f_{\cup}(N[v])| \geq 2.$$

Equivalently, $f_{\cup}(N[v]) = \{a, b\}$ holds for all $v \in V(G)$. The rainbow double domination number of G is denoted by $\tilde{\gamma}_{\times 2}(G)$ and equals the minimum weight $f(V)$ over all rainbow double dominating functions f of G .

Note that every graph without isolated vertices has domatic number at least 2, which means that it admits a domatic 2-partition, that is, a partition of its vertex set into two dominating sets. (The domatic number was introduced in a paper from 1970's [21], extensively studied afterwards, and surveyed in [81].) To see this, note that for any maximal independent set S in the graph, the pair $(S, V \setminus S)$ is a domatic 2-partition. Given a domatic 2-partition (A, B) , setting $f(v) = \{a\}$ for all $v \in A$ and $f(v) = \{b\}$ for all $v \in B$ results in a rainbow double dominating function of G , which

shows that the rainbow double domination number is well defined for all graphs without isolated vertices.

The above observation can be strengthened as follows: The rainbow double domination number of a graph G without isolated vertices equals the so-called *disjoint domination number* of G , defined in [49] as the minimum value of $|A| + |B|$ over all pairs (A, B) of disjoint dominating sets of G , and denoted by $\gamma\gamma(G)$. The disjoint domination number was studied in several papers, [1, 3, 53, 59, 66].

Proposition 2.1. *For every graph G without isolated vertices, we have $\tilde{\gamma}_{\times 2}(G) = \gamma\gamma(G)$.*

Proof. Suppose that A and B are two disjoint dominating sets in G . Then setting $f(v) = \{a\}$ for all $v \in A$, $f(v) = \{b\}$ for all $v \in B$, and $f(v) = \emptyset$ for all $v \in V \setminus (A \cup B)$ results in a rainbow double dominating function of G with total weight $|A| + |B|$. Conversely, if $f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$ is a rainbow double dominating function of G , then $f^{-1}(\{a\})$ and $f^{-1}(\{b\})$ form a pair of disjoint dominating sets of G of total size equal to the total weight of f . \square

Total double domination, rainbow total double domination.

Definition 2.13. ($\gamma_{t \times 2}$, row/column 8 in Table 2)

Let $G = (V, E)$ be a graph. A total double dominating function of G is a function $f : V \rightarrow \{0, 1\}$ such that for all $v \in V(G)$ it holds that

$$f(N(v)) \geq 2.$$

The total double domination number of G is denoted by $\gamma_{t \times 2}(G)$ and equals the minimum weight $f(V)$ over all total double dominating functions f of G .

Any set of the form $D = \{v \in V \mid f(v) = 1\}$ where f is a total double dominating function of G is said to be a total double dominating set of G . Note that the minimum size of a total double dominating set equals $\gamma_{t \times 2}(G)$.

The total double domination number is finite precisely in graphs G with $\delta(G) \geq 2$. The invariant, which is in some papers called *double total domination*, was studied for instance in [52, 54].

Definition 2.14. ($\tilde{\gamma}_{t \times 2}$, row/column 12 in Table 2)

Let $G = (V, E)$ be a graph. A rainbow total double dominating function of G is a function $f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$ such that for all $v \in V(G)$ it holds that

$$|f_{\cup}(N(v))| \geq 2.$$

Equivalently, $f_{\cup}(N(v)) = \{a, b\}$ holds for all $v \in V(G)$. The rainbow total double domination number of G is denoted by $\tilde{\gamma}_{t \times 2}(G)$ and equals the minimum weight $f(V)$ over all rainbow total double dominating functions f of G .

The *total domatic number* of a graph G without isolated vertices is the maximum number of total dominating sets of G that form a partition of its vertex set, cf. [80]. Analogously to the disjoint domination number of a graph, we define the *disjoint total domination number* of a graph G as the minimum value of $|A| + |B|$ over all pairs (A, B) of disjoint total dominating sets of G , and denote it by $\gamma_t \gamma_t(G)$. Note that this parameter is finite if and only if G admits a partition of its vertex set into two total dominating sets, that is, if its total domatic number is at least 2 (this is the case, for instance, for all k -regular graphs with $k \geq 4$ [55]). A similar parameter for digraphs was recently considered in [62].

Proposition 2.2. *For every graph G , we have $\tilde{\gamma}_{t \times 2}(G) = \gamma_t \gamma_t(G)$. In particular, the rainbow total double domination number of G is finite if and only if $V(G)$ can be partitioned into two total dominating sets.*

Proof. Let $f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$ be a minimum rainbow total double dominating function of G . Since $f \cup (N(v)) = \{a, b\}$ for all $v \in V(G)$, the set of vertices $f^{-1}(\{a\})$, that is, the set of vertices labeled by $\{a\}$, is a total dominating set in G , and, similarly, so is $f^{-1}(\{b\})$. Since these two sets are disjoint, we have $\gamma_t \gamma_t(G) \leq \tilde{\gamma}_{t \times 2}(G)$. Note that in this case, $V(G)$ can be partitioned into two total dominating sets, namely $f^{-1}(\{\emptyset, \{a\}\})$ and $f^{-1}(\{b\})$.

Conversely, suppose that $\gamma_t \gamma_t(G)$ is finite, and take a pair A, B of disjoint total dominating sets A and B such that $|A| + |B| = \gamma_t \gamma_t(G)$. Then, the function $f : V \rightarrow \{\{a\}, \{b\}\}$, defined by $f(v) = \{a\}$ for all $v \in A$, $f(v) = \{b\}$ for all $v \in B$, and $f(v) = \emptyset$ for all $v \in V \setminus (A \cup B)$, is a rainbow total double dominating function of G with $f(V) = \gamma_t \gamma_t(G)$. This implies that $\tilde{\gamma}_{t \times 2}(G) \leq \gamma_t \gamma_t(G)$, and consequently $\tilde{\gamma}_{t \times 2}(G) = \gamma_t \gamma_t(G)$. The above argument also shows that if $V(G)$ can be partitioned into two total dominating sets, then G has a rainbow total double dominating function. \square

Roman domination.

Definition 2.15. (γ_R , row/column 13 in Table 2)

Let $G = (V, E)$ be a graph. A Roman dominating function of G is a function $f : V \rightarrow \{0, 1, 2\}$ such that for all $v \in V(G)$ it holds that

$$f(v) = 0 \implies (\exists w \in N(v) \text{ such that } f(w) = 2).$$

The Roman domination number of G is denoted by $\gamma_R(G)$ and equals the minimum weight $f(V)$ over all Roman dominating functions f of G .

As already mentioned in the introduction, the concept of Roman domination was introduced by Stewart in [75], see also [20]. It was studied also in the PhD thesis of Dreyer [28] and in a series of papers, see, e.g., [31, 64, 65, 67, 73] for some recent references.

Defining the rainbow Roman domination number in the obvious way does not lead to a new graph parameter: it coincides with the Roman domination number.

For each of above defined domination parameters, given a weight function f , if the defining condition is satisfied for a vertex $v \in V(G)$, we say that v is dominated (with respect to f).

For later use in Section 4, we now recall also the definitions of three covering parameters in graphs.

Edge covers, 2-edge covers, and 2-vertex covers.

An *edge cover* of G is a function $f : E \rightarrow \{0, 1\}$ such that for all $v \in V$, it holds that

$$\sum_{w \in V, vw \in E} f(vw) \geq 1.$$

If G is a graph with no isolated vertices, the *edge cover number* of G is denoted by $\rho(G)$ and equals the minimum weight $f(E)$ over all edge covers f of G .

A *2-edge cover* of G is a function $f : E \rightarrow \{0, 1, 2\}$ such that for all $v \in V$, it holds that

$$\sum_{w \in V, vw \in E} f(vw) \geq 2.$$

If G is a graph with no isolated vertices, the *2-edge cover number* of G is denoted by $\rho_2(G)$ and equals the minimum weight $f(E)$ over all 2-edge covers f of G .

A *2-vertex cover* of G is a function $f : V \rightarrow \{0, 1, 2\}$ such that for all $vw \in E$, it holds that

$$f(v) + f(w) \geq 2.$$

The *2-vertex cover number* of G is denoted by $\tau_2(G)$ and equals the minimum weight $f(V)$ over all 2-vertex covers f of G .

The following result is a consequence of several works by Gallai [37–40] (cf. [72, Chapter 30]), and will be used to prove the bound of $\gamma_{t\{2\}}$ in terms of γ_{w2} in Proposition 4.6.

Theorem 2.3. *For every graph $G = (V, E)$ with no isolated vertices, $\rho_2(G) + \tau_2(G) = 2|V|$.*

Name	Notion	Function	Condition
domination	γ	$f : V \rightarrow \{0, 1\}$	$f(N[v]) \geq 1 \quad \forall v$
total domination	γ_t	$f : V \rightarrow \{0, 1\}$	$f(N(v)) \geq 1 \quad \forall v$
weak 2-domination	γ_{w2}	$f : V \rightarrow \{0, 1, 2\}$	$f(N(v)) \geq 2$ if $f(v) = 0$
rainbow weak 2-domination	$\tilde{\gamma}_{w2}$	$f : V \rightarrow 2^{\{a,b\}}$	$ f_{\cup}(N(v)) \geq 2$ if $f(v) = \emptyset$
$\{2\}$ -domination	$\gamma_{\{2\}}$	$f : V \rightarrow \{0, 1, 2\}$	$f(N[v]) \geq 2 \quad \forall v$
rainbow $\{2\}$ -domination	$\tilde{\gamma}_{\{2\}}$	$f : V \rightarrow 2^{\{a,b\}}$	$ f_{\cup}(N[v]) \geq 2 \quad \forall v$
total $\{2\}$ -domination	$\gamma_{t\{2\}}$	$f : V \rightarrow \{0, 1, 2\}$	$f(N(v)) \geq 2 \quad \forall v$
rainbow total $\{2\}$ -domination	$\tilde{\gamma}_{t\{2\}}$	$f : V \rightarrow 2^{\{a,b\}}$	$ f_{\cup}(N(v)) \geq 2 \quad \forall v$
2-domination	γ_2	$f : V \rightarrow \{0, 1\}$	$f(N(v)) \geq 2$ if $f(v) = 0$
rainbow 2-domination	$\tilde{\gamma}_2$	$f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$	$ f_{\cup}(N(v)) \geq 2$ if $f(v) = \emptyset$
double domination	$\gamma_{\times 2}$	$f : V \rightarrow \{0, 1\}$	$f(N[v]) \geq 2 \quad \forall v$
rainbow double domination	$\tilde{\gamma}_{\times 2}$	$f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$	$ f_{\cup}(N[v]) \geq 2 \quad \forall v$
total double domination	$\gamma_{t \times 2}$	$f : V \rightarrow \{0, 1\}$	$f(N(v)) \geq 2 \quad \forall v$
rainbow total double domination	$\tilde{\gamma}_{t \times 2}$	$f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$	$ f_{\cup}(N(v)) \geq 2 \quad \forall v$
Roman domination	γ_R	$f : V \rightarrow \{0, 1, 2\}$	$\exists w \sim v : f(w) = 2$ if $f(v) = 0$
edge cover	ρ	$f : E \rightarrow \{0, 1\}$	$f(E(v)) \geq 1 \quad \forall v$
2-edge cover	ρ_2	$f : E \rightarrow \{0, 1, 2\}$	$f(E(v)) \geq 2 \quad \forall v$
vertex cover	τ	$f : V \rightarrow \{0, 1\}$	$f(v) + f(w) \geq 1 \quad \forall vw \in E$
2-vertex cover	τ_2	$f : V \rightarrow \{0, 1, 2\}$	$f(v) + f(w) \geq 2 \quad \forall vw \in E$

Table 1: Summary of definitions of the parameters under study.

3 Comparison of parameters

In this section, we state our main result: the comparison of the values of 13 domination parameters in graphs for each pair of them (Table 2). We start with describing, in Fig. 1, the Hasse diagram of the relation \leq on the 15 graph parameters defined in Section 2. Given two of these parameters, say ρ and ρ' , we write $\rho \leq \rho'$ if and only if for every graph G for which both $\rho(G)$ and $\rho'(G)$ are well defined, it holds that $\rho(G) \leq \rho'(G)$. The relations represented in this figure will be used often in our proofs of upper bounds for the parameters in terms of functions of other parameters, and in the proofs that these bounds are sharp.




Figure 1: Hasse diagram of the relation \leq among various domination parameters.

Proposition 3.1. *For any two parameters ρ and ρ' in the Hasse diagram on Fig. 1, ρ is below ρ' in the diagram if and only if $\rho \leq \rho'$.*

Proof. Here, we will only argue the ‘only if’ direction of the proof, that is, if ρ is below ρ' in the diagram then $\rho \leq \rho'$. The other direction will follow from results in Section 5.

Clearly, it suffices to verify the statement only for the ‘covering’ pairs (ρ, ρ') in the diagram, that is, pairs such that ρ is immediately below ρ' in the diagram – the inequalities for all the remaining pairs follow by transitivity. The inequality $\gamma \leq \gamma_{w2}$ can be proved by observing that

if $f : V(G) \rightarrow \{0, 1, 2\}$ is a minimum weight weak 2-dominating function of G , then the set $f^{-1}(\{1, 2\})$ is a dominating set of G of total size $|f^{-1}(1)| + |f^{-1}(2)| \leq f(V) = \gamma_{w2}(G)$. Similarly, if $f : V(G) \rightarrow \{0, 1, 2\}$ is a minimum weight $\{2\}$ -dominating function of G , then the set $f^{-1}(\{1, 2\})$ is a total dominating set of G of total size $|f^{-1}(1)| + |f^{-1}(2)| \leq f(V) = \gamma_{\{2\}}(G)$. This implies the inequality $\gamma_t \leq \gamma_{\{2\}}$. The inequality $\gamma_t \leq \tilde{\gamma}_{w2}$ was proved in [35] and the inequality $\tilde{\gamma}_{w2} \leq \gamma_R$ in [79].

If f is a dominating function of G , then $2f$ is a Roman dominating function of G (cf. [20, 51]). Hence, for every graph G , it holds $\gamma_R(G) \leq 2\gamma(G) = \tilde{\gamma}_{\{2\}}(G)$, which establishes the relation $\gamma_R \leq \tilde{\gamma}_{\{2\}}$ in the diagram.

That every rainbow parameter is above its original counterpart is a direct consequence of definitions. Similarly, all other inequalities represented in the diagram \mathcal{H} can be easily derived just by looking at the definitions of parameters. \square

In Table 2 we summarize the bounds relating any two of the 13 considered parameters, or the fact that there is no bound. We will give the necessary proofs of upper bounds in Section 4 and summarize them in Table 3. All the bounds are sharp, as will be shown in Section 5, which will also contain the proofs of nonexistence of bounds between certain pairs of parameters. The families proving the nonexistence of a bound and the examples showing sharpness are summarized in Tables 4 and 5 (on p. 27 and 29), respectively.

We can also consider the following weaker version of the \leq relation on the 15 graph parameters defined in Section 2. The relation \preceq is defined by $\rho \preceq \rho'$ if and only if there exists a function f such that for every graph G for which both $\rho(G)$ and $\rho'(G)$ are well defined, it holds that $\rho(G) \leq f(\rho'(G))$. This relation is reflexive and transitive, but not antisymmetric. It induces an equivalence relation \approx , defined by $\rho \approx \rho'$ if and only if $\rho \preceq \rho'$ and $\rho' \preceq \rho$. The results summarized in Table 2 imply that the relation \approx has exactly four equivalence classes, which are linearly ordered by the quotient partial order obtained from \preceq by collapsing each equivalence class of \approx into a single element. See Fig. 2 for a depiction of these four equivalence classes and the Hasse diagram of the corresponding linear order.




Figure 2: The Hasse diagram representing the preorder \preceq on the considered domination parameters.

$\rho \leq f(\rho')$	1	2	3	4	5	6	7	8	9	10	11	12	13
γ	γ	γ_t	γ_{w2}	$\gamma_{\{2\}}$	$\gamma_{t\{2\}}$	γ_2	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$\tilde{\gamma}_{w2}$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	γ_R
$=$	γ	γ_t	γ_{w2}	$\gamma_{\{2\}} - 1$	$\gamma_{t\{2\}} - 1$	γ_2	$\gamma_{\times 2} - 1$	$\gamma_{t \times 2} - 1$	$\tilde{\gamma}_{w2}$	$\tilde{\gamma}_2$	$\frac{1}{2} \tilde{\gamma}_{\times 2}$	$\frac{1}{2} \tilde{\gamma}_{t \times 2}$	$\gamma_R - 1^*$
2γ	2γ	$=$	$\frac{3\gamma_{w2} - 1}{2}$	$\gamma_{\{2\}}$	$\gamma_{t\{2\}} - 1$	$\frac{3\gamma_2 - 1}{2}$	$\gamma_{\times 2}$	$\gamma_{t \times 2} - 1$	$\tilde{\gamma}_{w2}$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\frac{1}{2} \tilde{\gamma}_{t \times 2}$	γ_R
2γ	2γ	$2\gamma_t$	$=$	$\gamma_{\{2\}}$	$\gamma_{t\{2\}}$	γ_2	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$\tilde{\gamma}_{w2}$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	γ_R
2γ	2γ	$2\gamma_t$	$2\gamma_{w2} - 1^*$	$=$	$\gamma_{t\{2\}}$	$2\gamma_2 - 1^*$	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$2\tilde{\gamma}_{w2} - 1^*$	$2\tilde{\gamma}_2 - 1^*$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	$2\gamma_R - 2^*$
4γ	4γ	$2\gamma_t$	$2\gamma_{w2}$	$2\gamma_{\{2\}}$	$=$	$2\gamma_2$	$2\gamma_{\times 2}$	$\gamma_{t \times 2}$	$2\tilde{\gamma}_{w2}$	$2\tilde{\gamma}_2$	$2\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	$2\gamma_R$
$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$=$	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$\neg\exists f$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	$\neg\exists f$
$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$2\gamma_2 - 1$	$=$	$\gamma_{t \times 2}$	$\neg\exists f$	$2\tilde{\gamma}_2 - 1$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	$\neg\exists f$
$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$3\gamma_2 - 2$	$2\gamma_{\times 2} - 1$	$=$	$\neg\exists f$	$3\tilde{\gamma}_2 - 2$	$2\tilde{\gamma}_{\times 2} - 1$	$\tilde{\gamma}_{t \times 2}$	$\neg\exists f$
2γ	2γ	$2\gamma_t$	$2\gamma_{w2} - 2$	$2\gamma_{\{2\}} - 2$	$2\gamma_{t\{2\}} - 2$	$2\gamma_2 - 2$	$2\gamma_{\times 2} - 2$	$2\gamma_{t \times 2} - 2$	$=$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	γ_R
$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$=$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	$\neg\exists f$
$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$2\tilde{\gamma}_2$	$=$	$\tilde{\gamma}_{t \times 2}$	$\neg\exists f$
$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$\neg\exists f$	$=$	$\neg\exists f$
2γ	2γ	$2\gamma_t$	$2\gamma_{w2} - 1$	$2\gamma_{\{2\}} - 2$	$2\gamma_{t\{2\}} - 2$	$2\gamma_2 - 1$	$2\gamma_{\times 2} - 2$	$2\gamma_{t \times 2} - 2$	$\frac{3}{2} \tilde{\gamma}_{w2}$	$\frac{3}{2} \tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	$=$

Table 2: Upper bounds for several domination parameters in terms of others. The entry in row indexed by parameter ρ and in column indexed by parameter ρ' represents either that ρ is not bounded from above by any function of ρ' (in this case the entry is “ $\neg\exists f$ ”), or gives an upper bound for ρ in terms of a function of ρ' . The bound $\rho \leq f(\rho')$ should be interpreted so that it holds for graphs G such that both $\rho(G)$ and $\rho'(G)$ are defined. For instance, the entry in row γ_t and column γ is 2γ , indicating that for every graph G without isolated vertices, it holds that $\gamma_t(G) \leq 2\gamma(G)$. Bounds of the form $\rho \leq f(\rho')$ that are marked with an asterisk are only valid for graphs with at least one edge. The (sharp) bounds for graphs with no edges are, respectively, $\gamma = \gamma_R$, $\gamma_{\{2\}} = 2\gamma_{w2} = 2\gamma_2 = 2\tilde{\gamma}_{w2} = 2\tilde{\gamma}_2 = 2\gamma_R$, and can be easily verified. The bound in entry (9, 3) holds if $G \neq K_1$. The correctness of the entries in the table is proved or referenced in Section 4 (proofs of upper bounds) and in Section 5 (proofs of nonexistence of upper bounds), while the references to proofs are summarized in Table 3. All the bounds are sharp. The examples for sharpness and for the unboundedness are summarized in Table 5.

$\rho \leq f(\rho')$	1	2	3	4	5	6	7	8	9	10	11	12	13
	γ	γ_t	γ_{u2}	$\gamma_{\{2\}}$	$\gamma_{t\{2\}}$	γ_2	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$\tilde{\gamma}_{u2}$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	γ_R
1	γ	\mathcal{H}	\mathcal{H}	\mathcal{H}	T4	\mathcal{H}	T4	T2	\mathcal{H}	\mathcal{H}	4.15	T11	4.17
2	γ_t	\mathcal{H} [48]	\mathcal{H}	4.4	\mathcal{H}	T3	\mathcal{H}	T5	\mathcal{H} [35]	\mathcal{H}	\mathcal{H}	4.16	\mathcal{H} [50]
3	γ_{u2}	T4	T1	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}
4	$\gamma_{\{2\}}$	4.2	T1	4.5	\mathcal{H}	T3	\mathcal{H}	\mathcal{H}	T3	T3	\mathcal{H}	\mathcal{H}	T1
5	$\gamma_{t\{2\}}$	T2	4.3	4.6	T2	T3	T3	\mathcal{H}	T3	T3	T3	\mathcal{H}	T3
6	γ_2	T5	T5	T5	T5	5.6	\mathcal{H}	\mathcal{H}	T5	\mathcal{H}	\mathcal{H}	\mathcal{H}	T5
7	$\gamma_{\times 2}$	T5	T5	T5	T6	4.11	\mathcal{H}	\mathcal{H}	T5	T6	\mathcal{H}	\mathcal{H}	T5
8	$\gamma_{t \times 2}$	T5	T5	T5	5.7	4.12	4.13	\mathcal{H}	T5	T6	T7	\mathcal{H}	T5
9	$\tilde{\gamma}_{u2}$	T13	T1	4.7	T3	T3	T3	T3	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H}	\mathcal{H} [79]
10	$\tilde{\gamma}_2$	T5	T5	T5	T6	T8	T8	5.8	T5	\mathcal{H}	\mathcal{H}	\mathcal{H}	T5
11	$\tilde{\gamma}_{\times 2}$	T5	T5	T5	T6	T8	T8	T10	T5	4.14	\mathcal{H}	\mathcal{H}	T5
12	$\tilde{\gamma}_{t \times 2}$	T11	T11	T11	T11	T11	T11	T11	T11	T11	5.9	\mathcal{H}	T11
13	γ_R	[20, 51]	T1	4.8	T4	T3	T4	T5	[15, 34]	T9	\mathcal{H}	\mathcal{H}	\mathcal{H}

Table 3: Summary of proofs for the entries of Table 2. Entries labeled \mathcal{H} refer to results that are already stated in the corresponding Hasse diagram of Fig. 1. Entries labeled “Tk” are proved by transitivity through parameter in row/column k . Details regarding the use of transitivity in proving nonexistence of bounds will be explained in Section 5.3. Entries labeled with numbers refer to propositions that directly prove the bound (or nonexistence of a bound). Finally, for some entries we give references to papers where a proof of the corresponding bound can be found.

4 Proofs of upper bounds in Table 2

In this section we prove the upper bounds from Table 2. All the bounds are sharp, which will be demonstrated in Section 5. In the first subsection we concentrate on the bounds that follow from the Hasse diagram in Fig. 1, while in the second subsection we give explicit proofs of the remaining bounds. Note that the bounds in the entries of Table 2 that are not proven directly in this section, follow by transitivity from other bounds, as shown in Table 3.

4.1 Upper bounds following from the Hasse diagram on Fig. 1

The first proposition of this subsection is a direct consequence of Proposition 3.1.

Proposition 4.1. *The upper bounds indicated by the following entries in Table 2 are correct: (1, 2), (1, 3), (1, 6), (1, 9), (1, 10) (2, 4) (2, 7) (2, 9), (2, 10) (2, 11) (2, 13), (3, 4), (3, 5), (3, 6), (3, 7), (3, 8), (3, 9), (3, 10), (3, 11), (3, 12), (3, 13), (4, 5), (4, 7), (4, 8), (4, 11), (4, 12), (5, 8), (5, 12), (6, 7), (6, 8), (6, 10), (6, 11), (6, 12), (7, 8), (7, 11), (7, 12), (8, 12), (9, 10), (9, 11), (9, 12), (9, 13), (10, 11), (10, 12), (11, 12), (13, 11), (13, 12). \square*

4.2 Other upper bounds in Table 2

In this subsection the remaining upper bounds are proved. To ease an examination the bounds are numbered by the ordered pairs (r, c) , where r stands for the row and c for the column in the table. Since a proof of the bound labeled by a pair (r, c) proceeds by taking an optimal solution for the parameter indexed by column c and modifying it into a feasible solution for the parameter indexed by row r , we group together the proofs of bounds sharing the same column coordinate. We proceed in increasing order of columns and, within the same column, in increasing order of rows.

The proofs of Propositions 4.5 and 4.6 below make use of two classical results due to Gallai: one on the structure of minimum edge covers in graphs, and one on the relation between the 2-edge-cover and the 2-vertex cover numbers of a graph (Theorem 2.3), respectively.

Bound (4,1)

Proposition 4.2. *For every graph G , $\gamma_{\{2\}}(G) \leq 2\gamma(G)$.*

Proof. The inequality $\gamma_{\{2\}}(G) \leq 2\gamma(G)$ follows from the fact that if $f : V(G) \rightarrow \{0, 1\}$ is a minimum weight dominating function of G , then $g = 2f : V(G) \rightarrow \{0, 1, 2\}$ is a $\{2\}$ -dominating function of G of weight exactly $2\gamma(G)$. \square

Bound (5,2)

Proposition 4.3. *For every graph G without isolated vertices, $\gamma_{t\{2\}}(G) \leq 2\gamma_t(G)$.*

Proof. The inequality $\gamma_{t\{2\}}(G) \leq 2\gamma_t(G)$ follows from the fact that if $f : V(G) \rightarrow \{0, 1\}$ is a minimum weight total dominating function of G , then $g = 2f : V(G) \rightarrow \{0, 1, 2\}$ is a total $\{2\}$ -dominating function of G of weight exactly $2\gamma_t(G)$. \square

Bound (2,3)

Proposition 4.4. *For every graph G without isolated vertices, $\gamma_t(G) \leq \frac{3\gamma_{w2}(G)-1}{2}$.*

Proof. Let G be a graph without isolated vertices, and let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum weight weak 2-dominating function of G . Let us define the following subsets of $V(G)$. Let $V_i = \{v \in V(G) \mid f(v) = i\}$, for $i \in \{0, 1, 2\}$. Let S be a maximal set of vertices in V_0 such that their neighborhoods intersected with V_1 are nonempty and pairwise disjoint. Let D_0 denote the set of vertices in V_1 that have no neighbor in S . For every vertex $v \in D_0 \cup V_2$, choose a vertex v' adjacent to it. (Notice that such a vertex exists, since G has no isolated vertices.) Let $D'_0 = \{v' \mid v \in D_0\}$ and $V'_2 = \{v' \mid v \in V_2\}$.

We claim that the set $D' = (V_1 \setminus D_0) \cup D'_0 \cup S \cup V_2 \cup V'_2$ is a total dominating set of G . If $v \in V_1 \setminus D_0$, then v has a neighbor in S . If $v \in D_0$, then v has a neighbor in D'_0 . If $v \in V_2$, then v has a neighbor in V'_2 . If $v \in V_0$, then v has either a neighbor in V_2 , or it has at least two neighbors in V_1 , and thus by the definition of S , v has a neighbor in $N(S) \cap V_1 \subseteq V_1 \setminus D_0$.

Since every vertex in S has at least two neighbors in V_1 , we have $|N(S) \cap V_1| \geq 2|S|$. Therefore, we can bound the size of D' from above as follows:

$$|D'| = (|V_1 \setminus D_0| + |D'_0|) + |S| + (|V_2| + |V'_2|) \leq |V_1| + \frac{|N(S) \cap V_1|}{2} + 2|V_2|. \quad (1)$$

If $D_0 \neq \emptyset$, then $|N(S) \cap V_1| \leq |V_1| - 1$ and hence by (1), we have

$$|D'| \leq |V_1| + \frac{|V_1| - 1}{2} + 2|V_2| \leq \frac{3(|V_1| + 2|V_2|) - 1}{2} = \frac{3\gamma_{w2}(G) - 1}{2}.$$

If $V_2 \neq \emptyset$, then $2|V_2| \leq 3|V_2| - 1$ and by (1) we obtain

$$|D'| \leq |V_1| + \frac{|N(S) \cap V_1|}{2} + 3|V_2| - 1 < \frac{3(|V_1| + 2|V_2|) - 1}{2} = \frac{3\gamma_{w2}(G) - 1}{2}.$$

Finally, if $D_0 = \emptyset$ and $V_2 = \emptyset$, then $D' = V_1 \cup S$, and we can obtain a smaller total dominating set D'' by deleting from D' an arbitrary vertex of V_1 . Indeed, every vertex in V_0 has at least two neighbors in V_1 , and hence it has a neighbor in D'' . We can bound the size of D'' from above as follows:

$$|D''| = |V_1| + |S| - 1 \leq |V_1| + \frac{|N(S) \cap V_1|}{2} - 1 \leq \frac{3|V_1|}{2} - 1 < \frac{3\gamma_{w2}(G) - 1}{2}.$$

In either case, we obtain $\gamma_t(G) \leq \frac{3\gamma_{w2}(G)-1}{2}$. □

Bound (4,3)

Proposition 4.5. *For every graph G with at least one edge, $\gamma_{\{2\}}(G) \leq 2\gamma_{w2}(G) - 1$.*

Proof. Let G be a graph with at least one edge.

Let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum weight weak 2-dominating function of G . Let $V_i = \{v \in V(G) \mid f(v) = i\}$, for $i \in \{0, 1, 2\}$.

We first deal with the case when $V_2 \neq \emptyset$. The function $g : V(G) \rightarrow \{0, 1, 2\}$ that agrees with f on all vertices except on the vertices with f -value equal to 1, each of which g maps to 2, is a $\{2\}$ -dominating function of G . Since

$$g(V(G)) = 2|V_1| + 2|V_2| \leq 2(|V_1| + 2|V_2| - 1) = 2f(V(G)) - 2 = 2\gamma_{w2}(G) - 2,$$

we have the inequality $\gamma_{\{2\}}(G) \leq 2\gamma_{w2}(G) - 2$ in this case.

Assume now that $V_2 = \emptyset$. Note that the set I of isolated vertices in G is a subset of V_1 , and let $V'_1 = V_1 \setminus I$.

We may assume that every vertex u in V'_1 has a neighbor in V_0 . Otherwise, if all neighbors of u are in V'_1 , then we can obtain a weak 2-dominating function of G of the same weight as f by moving the weight 1 from u to one of its neighbors, and the previous case ($V_2 \neq \emptyset$) applies.

Now, let H be the graph with vertex set V'_1 in which two vertices u and v are adjacent if and only if they have a common neighbor $n(u, v)$ in V_0 . Let $h : E(H) \rightarrow \{0, 1\}$ be a minimum weight edge cover of H , let $C = \{e \in E(H) \mid h(e) = 1\}$ be the support of h , and let $N = \{n(u, v) \mid uv \in C\}$. Note that $|N| = |C| \leq |V'_1| - 1$ (recall that a minimum edge cover induces a spanning forest of stars [40]). Consider the function $g : V(G) \rightarrow \{0, 1, 2\}$, defined as follows:

$$g(v) = \begin{cases} 2, & \text{if } v \in I; \\ 1, & \text{if } v \in N \cup V'_1; \\ 0, & \text{otherwise.} \end{cases}$$

Then, g is a $\{2\}$ -dominating function of G : if $v \in I$, then clearly $g(N[v]) = 2$; if $v \in N \cup V'_1$, then v has a neighbor in $N \cup V'_1$, and hence $g(N[v]) \geq 2$; if $v \in V_0 \setminus N$, then v has at least two neighbors in V'_1 , and again $g(N[v]) \geq 2$ holds. Since

$$g(V(G)) = 2|I| + |V'_1| + |N| \leq 2|I| + |V'_1| + |V'_1| - 1 = 2|V_1| - 1 = 2f(V(G)) - 1 = 2\gamma_{w2}(G) - 1,$$

we have the desired inequality $\gamma_{\{2\}}(G) \leq 2\gamma_{w2}(G) - 1$. \square

Bound (5,3)

Proposition 4.6. *For every graph G with no isolated vertices, $\gamma_{t\{2\}}(G) \leq 2\gamma_{w2}(G)$.*

Proof. Let $f : V \rightarrow \{0, 1, 2\}$ be a minimum weight weak 2-dominating function of a graph $G = (V, E)$ with no isolated vertices. In the proof we will construct a total $\{2\}$ -dominating function g of G with weight less or equal to $2f(V)$, yielding the bound $\gamma_{t\{2\}}(G) \leq 2\gamma_{w2}(G)$.

Let $V_i = \{v \in V(G) \mid f(v) = i\}$, for $i \in \{0, 1, 2\}$. Note that for any vertex x in V_0 we already have $f(N(x)) \geq 2$, which is the condition imposed on vertices in the total $\{2\}$ -dominating set. On the other hand, $f(N(y))$ can be less than 2 for vertices $y \in V_1 \cup V_2$. Note that as f is minimum, each $y \in V_2$ is adjacent to a vertex in V_0 . Suppose that $V_2 \neq \emptyset$ and let Y be a minimum set of vertices from V_0 that dominate all vertices from V_2 ; i.e., $V_2 \subset N(Y)$ and Y is a smallest possible subset of V_0 with this property. Clearly, $|V_2| \geq |Y|$. Now, let $f_1 : V \rightarrow \{0, 1, 2\}$ be the function obtained from f by setting $f_1(y) = 2$ for all $y \in Y$ (changing f only in vertices of Y). Note that for any vertex $x \in V_0 \cup V_2$, we have $f_1(N(x)) \geq 2$. Moreover, when restricted to the subgraph G_1 of G induced by $V_2 \cup N(V_2)$, f_1 is a total $\{2\}$ -dominating function of G_1 such that $f_1(V(G_1)) \leq 2f(V(G_1))$. Thus it suffices to consider the remainder of the graph, i.e., $G - V(G_1)$; we remark that the function g , which we are constructing, coincides with f_1 on $V(G_1)$.

Consider the set Z of vertices $z \in V_1$ that are adjacent to some other vertex in V_1 (in the case when a vertex z from V_1 is adjacent to a vertex in V_2 , then it is already in G_1 , with $f_1(N(z)) \geq 2$, so this case need not be considered any more). If $Z \neq \emptyset$, then let $f_2 : V \rightarrow \{0, 1, 2\}$ be the function obtained from f_1 only by increasing the value of all vertices z from V_1 that have a neighbor in V_1 , by setting $f_2(z) = 2$. Denote by G_2 the subgraph of G , induced by $Z \cup N(Z)$. Clearly, $f_2(N(z)) \geq 2$ for all vertices $z \in Z \cup N(Z)$. Moreover, when restricted to G_2 , f_2 is a total $\{2\}$ -dominating function of G_2 such that $f_2(V(G_2)) \leq 2f(V(G_2))$. Thus it suffices to consider the remainder of the graph, i.e., $G - (V(G_1) \cup V(G_2))$; we remark that the function g , which we are constructing, coincides with f_2 on $V(G_1 \cup G_2)$.

Let $x \in V_1$, such that x is not adjacent to any other vertex in $V_1 \cup V_2$ (thus x lies in $G - (V(G_1) \cup V(G_2))$). Then x has a neighbor in V_0 , since G has no isolated vertices. It is possible that all neighbors of x are in $G_1 \cup G_2$. Let G_3 be the subgraph of G , induced by all vertices x in V_1 and not in $G_1 \cup G_2$, such that all their neighbors are in $G_1 \cup G_2$. To each vertex of G_3 , we set $f_3(x) = 0$, and to an arbitrary neighbor $y \in V_0$ of x , we set $f_3(y) = 2$ (and $f_3(u) = f_2(u)$ for all other vertices of G). Note that $f_3(N(x)) \geq 2$ for any $x \in G_3$, and f_3 restricted to $V(G_1) \cup V(G_2) \cup V(G_3)$ is a total $\{2\}$ -dominating function of the subgraph induced by $V(G_1) \cup V(G_2) \cup V(G_3)$. In addition, $f_3(V(G_1) \cup V(G_2) \cup V(G_3)) \leq 2f(V(G_1) \cup V(G_2) \cup V(G_3))$; we remark that the function g , which we are constructing, coincides with f_3 on $V(G_1) \cup V(G_2) \cup V(G_3)$.

Denote by H the remainder of the graph, i.e., $H = G - (V(G_1) \cup V(G_2) \cup V(G_3))$. Note that each vertex in H from V_1 has at least one neighbor in $V_0 \cap V(H)$, and also each vertex from $V_0 \cap V(H)$ has at least two neighbors from $V_1 \cap V(H)$ (the latter is because f is a weak 2-dominating function, and vertices from $V_0 \cap V(H)$ are not adjacent to any vertex from V_2 nor to any of the vertices of V_1 that were settled in the previous cases). For each $x \in V(H)$ with $f(x) = 0$, choose arbitrarily any of its two neighbors y, z in V_1 , and delete all other edges between x and its neighbors in $(V_0 \cup V_1) \setminus \{y, z\}$; call the resulting graph H' . Clearly, H' is a spanning subgraph of H , in which each vertex in V_0 has degree exactly 2, while vertices from V_1 can have an arbitrary degree, including 0. Remove all the isolated vertices from H' to obtain the graph H'' (we remark that the isolated vertices will be settled at the end of the proof). Let K be an arbitrary connected component of H'' . Since vertices from V_0 in K have degree 2 and are adjacent to two vertices from V_1 , K is a subdivision of a graph K' , whose vertices correspond to vertices of $V_1 \cap K$, and edges in K' correspond to vertices in $V_0 \cap K$.

Now, a function $h' : V(K') \cup E(K') \rightarrow \{0, 1, 2\}$ in a natural way corresponds to the function $h : V(K) \rightarrow \{0, 1, 2\}$, where $h'(x) = h(x)$ for any $x \in V_1$, while $h'(e)$ where $e \in E(K')$ coincides with $h(y)$, where $y \in V(K)$ is the subdivision vertex of e . In addition, the following conditions imposed to h' :

for every vertex $x \in V(K')$,

$$\sum_{xy \in E(K')} h'(xy) \geq 2$$

and for every edge $xy \in E(K')$,

$$h'(x) + h'(y) \geq 2,$$

are equivalent to the corresponding function h being a total $\{2\}$ -dominating function of K . Moreover, as $f(K) = f_3(K) = |V_1 \cap K| = |V(K')|$, it suffices to prove that there exists a function h' satisfying the above conditions such that the total sum of values of h' is at most $2|V(K')|$, to establish the desired bound of the theorem in the component K . Let $h'_1 : V(K') \rightarrow \{0, 1, 2\}$ and $h'_2 : E(K') \rightarrow \{0, 1, 2\}$ be minimum weight 2-vertex and 2-edge covers of K' , respectively. Define

$h'(v) = h'_1(v)$ for each $v \in V(K')$ and $h'(e) = h'_2(e)$ for each $e \in E(K')$. By definition of 2-vertex cover and 2-edge cover, h' satisfies the desired properties. Moreover, by Theorem 2.3, the total sum of values of h' is exactly $2|V(K')|$. By the observation above, the corresponding function $h : V(K) \rightarrow \{0, 1, 2\}$ is a total $\{2\}$ -dominating function of K of weight at most $2|V(K) \cap V_1|$.

Clearly, in the same way as K all connected components of H'' of order at least two can be analyzed, and so the function h is extended to all vertices of H'' . Now, H' is obtained from H'' by adding connected components with only one vertex, and they have not yet been considered in the proof. Recall that each such vertex $x \in V(H')$ is in V_1 , and we set $h(x) = 0$ and $h(y) = 2$ to any neighbor $y \in V(H')$, with which the value of h , which corresponds to x , is $2 \cdot f(x)$, as desired. Altogether we deduce that h is a total $\{2\}$ -dominating function of H' of weight at most $2|V(H') \cap V_1|$. As H' is a spanning subgraph of H , clearly h is also a total $\{2\}$ -dominating function of H of the same weight. Finally, we set $g(x) = f_3(x)$ for all vertices in $V(G) \setminus V(H)$ and $g(x) = h(x)$ for all vertices in $V(H)$. We conclude the main part of this proof by observing that g is a total $\{2\}$ -dominating function of G , with $g(V) \leq 2f(V)$. \square

Bound (9,3)

Proposition 4.7. *For every graph $G \neq K_1$, it holds that $\tilde{\gamma}_{w2}(G) \leq 2\gamma_{w2}(G) - 2$.*

Proof. Let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum weak 2-dominating function of G , and let $V_i = \{v \in V(G) \mid f(v) = i\}$, for $i \in \{0, 1, 2\}$. We consider two cases.

Case 1. $V_2 \neq \emptyset$. Define a function $g : V(G) \rightarrow \mathcal{P}(\{a, b\})$ as follows:

$$g(v) = \begin{cases} \{a, b\}, & \text{if } v \in V_1 \cup V_2; \\ \emptyset, & \text{otherwise} \end{cases}$$

Since f is weak 2-dominating function of G , every vertex $v \in V(G)$ with $f(v) = 0$ is adjacent to either two vertices of f -weight 1, or to one vertex of f -weight 2. Consequently, every vertex $v \in V(G)$ with $g(v) = \emptyset$ satisfies $g_{\cup}(N(v)) = \{a, b\}$. Thus, g is a rainbow weak 2-dominating function of G , and we have $\tilde{\gamma}_{w2}(G) \leq g(V(G)) = 2(|V_1| + |V_2|) \leq 2(|V_1| + 2|V_2|) - 2 = 2f(V(G)) - 2 = 2\gamma_{w2}(G) - 2$.

Case 2. $V_2 = \emptyset$. In this case, every vertex $v \in V(G)$ with $f(v) = 0$ is adjacent to two vertices of f -weight 1. If $V_1 = V(G)$, then the function $g : V(G) \rightarrow \mathcal{P}(\{a, b\})$ assigning $\{a\}$ to every vertex is a rainbow weak 2-dominating function of G , yielding in this case $\tilde{\gamma}_{w2}(G) \leq g(V(G)) = |V(G)| \leq 2|V(G)| - 2 = 2\gamma_{w2}(G) - 2$, where the inequality holds since $G \neq K_1$. If $V_1 \neq V(G)$, then there is a vertex $w \in V(G)$ with $f(w) = 0$ and consequently $|V_1| \geq 2$. Let $u, v \in V_1$ be two distinct vertices. Define a function $g : V(G) \rightarrow \mathcal{P}(\{a, b\})$ as follows:

$$g(w) = \begin{cases} \{a\}, & \text{if } w = u; \\ \{b\}, & \text{if } w = v; \\ \{a, b\}, & \text{if } w \in V_1 \setminus \{u, v\}; \\ \emptyset, & \text{otherwise.} \end{cases}$$

Since f is weak 2-dominating function of G , every vertex $v \in V(G)$ with $f(v) = 0$ is adjacent to either two vertices of f -weight 1, or to one vertex of f -weight 2. It follows that every vertex $w \in V(G)$ with $g(w) = \emptyset$ is adjacent either to a vertex with g -label $\{a, b\}$, or to vertices u and v ; in either case we have $g_{\cup}(N(w)) = \{a, b\}$. Thus, g is a rainbow weak 2-dominating function of G , and we have $\tilde{\gamma}_{w2}(G) \leq g(V(G)) = 2 + 2(|V_1| - 2) = 2|V_1| - 2 = 2f(V(G)) - 2 = 2\gamma_{w2}(G) - 2$. \square

Bound (13,3)

Proposition 4.8. *For every graph G , it holds that $\gamma_R(G) \leq 2\gamma_{w2}(G) - 1$.*

Proof. Let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum weight weak 2-dominating function of G .

Suppose first that there exists a vertex $u \in V(G)$ such that $f(u) = 2$. Then, let $g : V(G) \rightarrow \{0, 1, 2\}$ be defined as follows:

$$g(v) = \begin{cases} 2, & \text{if } f(v) \in \{1, 2\}; \\ 0, & \text{otherwise.} \end{cases}$$

Note that g is a Roman dominating function of G of total weight at most $2f(V(G)) - 2 = 2\gamma_{w2}(G) - 2$. On the other hand, if $f(v) \in \{0, 1\}$ for all $v \in V(G)$, then let $u \in V(G)$ be a vertex such that $f(u) = 1$, and let $g : V(G) \rightarrow \{0, 1, 2\}$ be defined as follows:

$$g(v) = \begin{cases} 2, & \text{if } v \neq u \text{ and } f(v) = 1; \\ 1, & \text{if } v = u; \\ 0, & \text{otherwise.} \end{cases}$$

Note that g is a Roman dominating function of G of total weight exactly $2f(V(G)) - 1 = 2\gamma_{w2}(G) - 1$. Hence, we have the desired inequality in each case. \square

Bound (1,4)

Proposition 4.9. *For every graph G , $\gamma(G) \leq \gamma_{\{2\}}(G) - 1$.*

Proof. Let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum $\{2\}$ -dominating function of G , and let $V_i = \{v \in V(G) \mid f(v) = i\}$, for $i \in \{0, 1, 2\}$. If $V_1 = \emptyset$, then V_2 is a dominating set of G and hence in this case $\gamma(G) \leq |V_2| \leq 2|V_2| - 1 = f(V(G)) - 1 = \gamma_{\{2\}}(G) - 1$. So we may assume that $V_1 \neq \emptyset$. Let $v_1 \in V_1$ and consider the set $D = (V_1 \cup V_2) \setminus \{v_1\}$. Then, $|D| = |V_1| + |V_2| - 1 \leq |V_1| + 2|V_2| - 1 = f(V(G)) - 1 = \gamma_{\{2\}}(G) - 1$. Hence, to show that $\gamma(G) \leq \gamma_{\{2\}}(G) - 1$, it suffices to argue that D is a dominating set of G , that is, that every $v \in V(G) \setminus D$ has a neighbor in D . If $v \in V(G) \setminus D$ and $v \neq v_1$, then v has a neighbor from V_2 or two neighbors from V_1 , because f is $\{2\}$ -dominating function of G . In the first case, this neighbor is clearly from D , while in the second case, there is at least one neighbor of v from V_1 , which is not equal to v_1 . Thus v has a neighbor from D . It remains to show that v_1 has a neighbor from D , which is also easy because $v_1 \in V_1$, hence it must have a neighbor from $V_1 \cup V_2$, which is thus from D . \square

Bound (2,5)

Proposition 4.10. *For every graph G without isolated vertices, $\gamma_t(G) \leq \gamma_{t\{2\}}(G) - 1$.*

Proof. Let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum total $\{2\}$ -dominating function of G , and let $V_i = \{v \in V(G) \mid f(v) = i\}$, for $i \in \{0, 1, 2\}$. If $V_1 = \emptyset$, then V_2 is a total dominating set of G and hence in this case $\gamma_t(G) \leq |V_2| \leq 2|V_2| - 1 = f(V(G)) - 1 = \gamma_{t\{2\}}(G) - 1$. So we may assume that $V_1 \neq \emptyset$. Let $v_1 \in V_1$ and consider the set $D = (V_1 \cup V_2) \setminus \{v_1\}$. Then, $|D| = |V_1| + |V_2| - 1 \leq |V_1| + 2|V_2| - 1 = f(V(G)) - 1 = \gamma_{t\{2\}}(G) - 1$. Hence, to show that $\gamma_t(G) \leq \gamma_{t\{2\}}(G) - 1$, it suffices to argue that D is a total dominating set of G , that is, that every $v \in V(G)$ has a neighbor in D . Since f is a total $\{2\}$ -dominating function of G , every vertex of v has either two neighbors in V_1 or one neighbor in V_2 . In particular, every vertex of v has a neighbor in either $V_1 \setminus \{v_1\}$, or in V_2 , and hence in D . \square

Bound (7,6)

Proposition 4.11. *For every graph G without isolated vertices, it holds that $\gamma_{\times 2}(G) \leq 2\gamma_2(G) - 1$.*

Proof. Let $f : V \rightarrow \{0, 1\}$ be a minimum 2-dominating function of G , and let $D = \{u \mid f(u) = 1\}$. For all $x \in V \setminus D$, we have $f(N[x]) \geq 2$, hence the condition imposed on a double dominating function is already fulfilled for these vertices. Let $y, w \in D$ be two neighbors of x with f -value positive. Let D' be a superset of D obtained from D by adding to it vertex x and for each vertex z from $D \setminus \{y, w\}$, adding an arbitrary vertex $u \in N(z)$. Clearly, for any $z \in D$, we have $|N[z] \cap D'| \geq 2$, where one of the vertices from $D' \cap N[z]$ is itself. Altogether we derive that D' is a double dominating set, that is, the function $f' : V \rightarrow \{0, 1\}$, which sets value 1 precisely to the vertices from D' , is a double dominating function of G , and its weight is $|D'| \leq 2|D| - 1 = 2\gamma_2(G) - 1$. \square

Bound (8,6)

Proposition 4.12. *For every graph G with $\delta(G) \geq 2$, $\gamma_{\text{tx}2}(G) \leq 3\gamma_2(G) - 2$.*

Proof. Let $f : V \rightarrow \{0, 1\}$ such that for all $v \in V$ it holds that $f(v) = 0 \implies f(N(v)) \geq 2$ and such that $|D| = \gamma_2(G)$, where $D = \{v \in V \mid f(v) = 1\}$. So, every vertex in $V \setminus D$ has at least two neighbors in D . Let D_2 be the set of vertices in D having at least two neighbors in D , let D_1 be the set of vertices in D having exactly one neighbor in D , and let D_0 be the set of vertices in D having no neighbors in D . Since $\delta(G) \geq 2$, we can always define a set D' of size at most $3|D_0| + 2|D_1| + |D_2|$ by adding to D one neighbor of v in $V \setminus D$ for each vertex v in D_1 , and two neighbors of v in $V \setminus D$ for each vertex v in D_0 . If $D_2 \neq \emptyset$, then $|D_2 \cup D_1| \geq 3$. Thus $|D'| \leq 3|D_0| + 2|D_1| + |D_2| \leq 3(|D_0| + |D_1| + |D_2|) - |D_1| - 2|D_2| \leq 3|D| - 2$. If $D_2 = \emptyset$ but $D_1 \neq \emptyset$, then $|D_1| \geq 2$. Again, $|D'| \leq 3|D| - 2$. In both cases, we can define $f' : V \rightarrow \{0, 1\}$ such that, for every vertex $v \in V$, $f'(v) = 1$ if and only if $v \in D'$. Function f' is a total double dominating function, thus $\gamma_{\text{tx}2}(G) \leq 3\gamma_2(G) - 2$.

Let now suppose $D_1 = D_2 = \emptyset$, so $D = D_0$ is an independent set, and consider the bipartite graph G' obtained from G by deleting all edges with both endpoints in $V \setminus D$. Since f is a 2-dominating function, $\delta(G') \geq 2$, and G' contains an even cycle C of length at least four. We can therefore define a set D' by adding to D the vertices of C and two neighbors of v in $V \setminus D$ for each vertex v in $D \setminus C$. Function $f' : V \rightarrow \{0, 1\}$ such that, for every vertex $v \in V$, $f'(v) = 1$ if and only if $v \in D'$ is a total double dominating function, and as $|D'| \leq 3|D| - 2$, we obtain $\gamma_{\text{tx}2}(G) \leq 3\gamma_2(G) - 2$. \square

Bound (8,7)

Proposition 4.13. *For every graph G with $\delta(G) \geq 2$, we have $\gamma_{\text{tx}2}(G) \leq 2\gamma_{\times 2}(G) - 1$.*

Proof. Let $D \subseteq V(G)$ be a minimum double dominating set of G . Then, every vertex in D has at least one neighbor in D , and every vertex in $V(G) \setminus D$ has at least two neighbors in D . Let $D_1 = \{v \in D \mid d_{G[D]}(v) = 1\}$. If $D_1 = \emptyset$, then D is also a total double dominating set, in which case $\gamma_{\text{tx}2}(G) \leq |D| = \gamma_{\times 2}(G) \leq 2\gamma_{\times 2}(G) - 1$.

Now let $D_1 \neq \emptyset$. Fix a vertex $v \in D_1$, and let $w \in N(v) \setminus D$. Note that such a vertex exists since $v \in D_1$ and $d_G(v) \geq 2$. Since $w \notin D$, there exists a neighbor of w in D , say v' , such that $v' \neq v$. For each vertex $x \in D_1 \setminus \{v, v'\}$, let x' denote an arbitrary neighbor of x outside D , and set $D' = D \cup \{w\} \cup X$, where $X = \{x' \mid x \in D_1 \setminus \{v, v'\}\}$.

We claim that D' is a total double dominating set of G . To see this, consider an arbitrary vertex $x \in V(G)$; we need to check that x has at least two neighbors in D' . If $x \in \{v, v'\}$, then $|N(x) \cap D| \geq 1$ and $N(x) \cap D' \supseteq (N(x) \cap D) \cup \{w\}$ (disjoint union). If $x \in D_1 \setminus \{v, v'\}$, then again $|N(x) \cap D| \geq 1$ and $N(x) \cap D' \supseteq (N(x) \cap D) \cup \{x'\}$ (disjoint union). If $x \in V(G) \setminus D_1$, then $|N(x) \cap D| \geq 2$ and $N(x) \cap D' \supseteq N(x) \cap D$. In either case, the conclusion follows.

The above implies that $\gamma_{t \times 2}(G) \leq |D'|$, hence it suffices to show that $|D'| \leq 2|D| - 1 = 2\gamma_{\times 2}(G) - 1$. If $v' \in D_1$, then $|X| \leq |D_1| - 2 \leq |D| - 2$, hence $|D'| \leq |D| + 1 + |X| \leq 2|D| - 1$. If $v' \notin D_1$, then $v' \in D \setminus D_1$, hence $|D_1| \leq |D| - 1$ and again we have $|X| \leq |D_1| - 1 \leq |D| - 2$, thus the same argument applies. \square

Bound (11,10)

Proposition 4.14. *For every graph G with no isolated vertices, $\tilde{\gamma}_{\times 2}(G) \leq 2\tilde{\gamma}_2(G)$.*

Proof. Let $f : V \rightarrow \{\emptyset, \{a\}, \{b\}\}$ such that for all $v \in V$ it holds that $f(v) = \emptyset \implies |f_{\cup}(N(v))| \geq 2$ and such that the total weight $f(V) = \tilde{\gamma}_2(G)$. Let $D_x = \{v \in V \mid f(v) = \{x\}\}$, for $x \in \{a, b\}$, and $D_0 = \{v \in V \mid f(v) = \emptyset\}$. Notice that $\tilde{\gamma}_2(G) = |D_a| + |D_b|$. Also, every vertex v in D_0 has a neighbor in D_a and a neighbor in D_b , in particular it satisfies $|f_{\cup}(N[v])| \geq 2$.

Vertices v of D_a having neighbors in D_b and vertices v of D_b having neighbors in D_a also satisfy $|f_{\cup}(N[v])| \geq 2$.

We will do the following process, successively, for all the vertices in D_a having neighbors only in $D_a \cup D_0$. Let v be such a vertex. If v has a neighbor w in D_0 , then we update $f(w) := \{b\}$, and update accordingly the sets D_b and D_0 . Now, $|f_{\cup}(N[v])| \geq 2$, and still $|f_{\cup}(N[z])| \geq 2$ for every z satisfying that before the update. If v has no neighbor in D_0 , then it has at least one neighbor in D_a . We update $f(v) := \{b\}$, and update accordingly the sets D_a and D_b . Again, $|f_{\cup}(N[v])| \geq 2$, and still $|f_{\cup}(N[z])| \geq 2$ for every z satisfying that before the update, because v had no neighbors in $D_b \cup D_0$. If, with the new definitions of D_a, D_b, D_0 , there are still vertices in D_a having neighbors only in $D_a \cup D_0$, we repeat the process. Notice that the size of this set strictly decreases on each step, while the size of vertices in D_b having neighbors only in $D_b \cup D_0$ never increases.

Once the set of vertices in D_a having neighbors only in $D_a \cup D_0$ is empty, we start processing vertices in D_b having neighbors only in $D_b \cup D_0$ analogously, interchanging the roles of a and b . Notice that the size of the set of vertices in D_b having neighbors only in $D_b \cup D_0$ strictly decreases on each step, while we never create vertices in D_a having neighbors only in $D_a \cup D_0$. So, once the former set is empty, all the vertices z in V satisfy $|f_{\cup}(N[z])| \geq 2$, hence f is a rainbow double dominating function of G . Notice that we have done at most $|D_a| + |D_b|$ steps and on each step we give a nonempty label to at most one vertex in D_0 . So the new weight $f(V)$ is at most $2\tilde{\gamma}_2(G)$. \square

Bound (1,11)

Proposition 4.15. *For every graph G without isolated vertices, $\gamma(G) \leq \frac{1}{2}\tilde{\gamma}_{\times 2}(G)$.*

Proof. Let G be a graph without isolated vertices, and let $f : V(G) \rightarrow \{\emptyset, \{a\}, \{b\}\}$ be a minimum weight rainbow double dominating function of G . Consider the sets $A = \{v \in V(G) \mid f(v) = \{a\}\}$ and $B = \{v \in V(G) \mid f(v) = \{b\}\}$. Without loss of generality, we may assume that $|A| \leq |B|$. Note that every vertex in $V(G) \setminus A$ has a neighbor in A , thus A is a dominating set of G , implying $\gamma(G) \leq |A| \leq \frac{1}{2}f(V(G)) = \frac{1}{2}\tilde{\gamma}_{\times 2}(G)$. \square

Bound (2,12)

Proposition 4.16. *For every graph G of total domatic number at least 2, $\gamma_t(G) \leq \frac{1}{2} \tilde{\gamma}_{t \times 2}(G)$.*

Proof. Let G be a graph with total domatic number at least 2, and let $f : V(G) \rightarrow \{\emptyset, \{a\}, \{b\}\}$ be a minimum weight rainbow total double dominating function of G . Consider the sets $A = \{v \in V(G) \mid f(v) = \{a\}\}$ and $B = \{v \in V(G) \mid f(v) = \{b\}\}$. Without loss of generality, we may assume that $|A| \leq |B|$. Note that every vertex in G has a neighbor in A . In particular, A is a total dominating set of G , implying $\gamma_t(G) \leq |A| \leq \frac{1}{2} f(V(G)) = \frac{1}{2} \tilde{\gamma}_{t \times 2}$. \square

Bound (1,13)

Proposition 4.17. *For every graph G with at least one edge, $\gamma(G) \leq \gamma_R(G) - 1$.*

Proof. Let $f : V(G) \rightarrow \{0, 1, 2\}$ be a minimum weight Roman dominating function of G , and let $V_i = \{v \in V(G) \mid f(v) = i\}$ for $i \in \{0, 1, 2\}$. We may assume that in addition f is chosen in such a way that $V_2 \neq \emptyset$. Indeed let $uv \in E(G)$, and assume that $f(u) = 1 = f(v)$. But then changing $f(u) = 1$ to $f(u) = 2$ and $f(v) = 1$ to $f(v) = 0$ yields a Roman dominating function of the same weight. Note that $V_1 \cup V_2$ is a dominating set, which implies $\gamma(G) \leq |V_1| + |V_2| < |V_1| + 2|V_2| = \gamma_R(G)$, hence $\gamma(G) \leq \gamma_R(G) - 1$. \square

5 Examples of sharpness and non-existence

5.1 Main families of graphs used in the proofs

In this section we present several families of graphs that will be used in the proofs of sharpness of the bounds from Table 2 or in the proofs of non-existence of such bounds indicated in the same table. Here we present the families that are used in several instances, while those that are used just once or twice will be presented along with Table 5, which summarizes the families used for each proof.

Given a graph G , by kG we define the disjoint union of k copies of G . Hence kK_2 , resp. kC_4 stands for the graphs on k components, each component being the connected graph on 2 vertices, resp. the *square*, that is, the cycle on four vertices.

Next, for a graph G , $S(G)$ denotes its subdivision graph, obtained from G by subdividing each of its edges exactly once. In $S(G)$ we distinguish between *original* and *subdivided* vertices that correspond to the vertices of G and to those internal vertices of the paths on three vertices replacing the edges of G to obtain $S(G)$ respectively. The stars with n leaves are denoted by $K_{1,n}$. We denote by $S(K_{1,n})^-$ the graph obtained from the subdivision graph $S(K_{1,n})$ of the n -star by deleting a leaf.

The graph H is the tree on 6 vertices, in which each of the two adjacent non-leaves is adjacent to two leaves.

We denote by F_n^3 the graph on $2n + 1$ vertices, obtained from nK_3 by identifying (gluing) one vertex of each triangle to a single vertex. Similarly, F_n^4 is the graph on $3n + 1$ vertices, obtained from nC_4 by identifying (gluing) one vertex of each square to a single vertex. See Fig. 3.

For $n \geq 2$, let K_n^{**} denote the graph obtained from the complete graph of order n by gluing two new triangles along each edge; see Fig. 4 for an example. In other words, for each pair x, y of vertices in the complete graph K_n two vertices are added, each of which is adjacent only to x and




Figure 3: Graphs F_n^3 and F_n^4

y . The added vertices in K_n^{**} (that is, those of degree 2) will be called *triangle vertices*; vertices that are not triangle will be called *original*.




Figure 4: Examples of graphs K_n^{**} and $S(K_n^{(2)})$

By $G^{(k)}$ we denote the multigraph obtained from a graph G by replacing each edge with k parallel edges. In particular, $K_n^{(2)}$ is the multigraph obtained from the complete graph of order n by duplicating each edge, and so $S(K_n^{(2)})$ is its subdivision graph; see Fig. 4 for an example. Note that $S(K_n^{(2)})$ is obtainable from K_n^{**} by deleting all edges joining pairs of original vertices. Next, the graph $S(K_3^{(n)})$ is the subdivision graph of the multigraph $K_3^{(n)}$ (i.e., the subdivision graph of the multigraph obtained from K_3 by adding $n - 1$ parallel edges between each pair of vertices). See Fig. 5 for an example.

We will denote by Q_n the graph that can be obtained from the multigraph $K_2^{(n)}$ (two vertices, connected by n parallel edges) by subdividing each edge twice, i.e., each edge is replaced by the path P_4 (the so-called double subdivision graphs of $K_2^{(n)}$). See Fig. 5 for an example.

Finally, graphs T_n are defined as follows. Let $V(T_n) = \{v_1, \dots, v_n, w_1, \dots, w_n, s_1, s_2, s_3, t_1, \dots, t_5\}$, so that s_1, s_2, s_3 induce a triangle, t_1, \dots, t_5 induce a C_5 , s_1 and s_3 are adjacent to v_i for every $i \in \{1, \dots, n\}$, t_1 and t_5 are adjacent to w_i for every $i \in \{1, \dots, n\}$, $v_i w_i \in E(T_n)$ for every $i \in \{1, \dots, n\}$, and there are no other edges; see Fig. 6.

It is for instance easy to see that for each n , the set $\{s_1, t_1, t_3\}$ is a minimum dominating set of T_n . We will discuss the values of several other parameters of T_n and the consequences for Table 4 in Subsection 5.3.

The sharpness of the bounds in Table 2 will be demonstrated using the values of relevant




Figure 5: Examples of graphs $S(K_3^{(n)})$ and Q_n




Figure 6: Graphs T_n

parameters on graphs families summarized in rows A–G of Table 4. In addition, we will use three more families, not described in this table (each of them used only for three invariants). Furthermore, the results for families in rows X, Y, Z, W will be used in proving that there is no bound between certain pairs of invariants.

Since most of the values in Table 4 can be proved as an easy exercise, we will only present here a proof for those that are a bit more involved.

Claim 5.1. *For $n \geq 3$, we have $\rho(K_n^{**}) = n$ for every $\rho \in \{\gamma_{w2}, \gamma_{\{2\}}, \gamma_{t\{2\}}, \gamma_2, \gamma_{\times 2}, \gamma_{t \times 2}\}$, and $\tilde{\gamma}_{w2}(K_n^{**}) = \gamma_R(K_n^{**}) = 2n - 2$.*

Proof. Recall that K_n^{**} is the graph obtained from the complete graph of order n by gluing two triangles along each edge, and let us denote K_n^{**} shortly by G_n . Let V_n be the set of original vertices in K_n^{**} (the vertices of the complete subgraph K_n), and V_n^{**} the set $V(G_n) \setminus V_n$ (triangle vertices). On the one hand, it is easy to see that $\gamma_{t \times 2}(G_n) \leq n$, since assigning 1 to each vertex of V_n and 0 to each other vertex yields a total double dominating function of G_n of total weight n . On the other hand, we will now show that $\tilde{\gamma}_{w2}(G_n) \geq 2n - 2$. Since the bounds from Table 2 imply that $\tilde{\gamma}_{w2}(G_n) \leq \gamma_R(G_n) \leq 2\gamma_{t \times 2}(G_n) - 2$ and $\rho(G_n) \leq \gamma_{t \times 2}(G_n)$ for all $\rho \in \{\gamma_{w2}, \gamma_{\{2\}}, \gamma_{t\{2\}}, \gamma_2, \gamma_{\times 2}\}$, the claim will then follow.

Suppose for a contradiction that there is a minimum rainbow weak 2-dominating function $f : V(G_n) \rightarrow \mathcal{P}(\{a, b\})$ of G_n of total weight at most $2n - 3$. We first argue that we may assume without loss of generality that $f(v) = \emptyset$ for all vertices $v \in V_n^{**}$. Indeed, if $f(v) \neq \emptyset$ for some vertex $v \in V_n^{**}$, then the minimality of f implies that $f(v') \neq \emptyset$, where v' denotes the unique vertex with $v' \neq v$ and $N(v) = N(v')$. Hence, assigning $\{a, b\}$ to one of the neighbors of v and

H vs. $\rho(H)$	1	2	3	4	5	6	7	8	9	10	11	12	13
	γ	γ_t	γ_{w2}	$\gamma_{\{2\}}$	$\gamma_{\{2\}}$	γ_2	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$\tilde{\gamma}_{w2}$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	γ_R
A	k	$2k$	$2k$	$2k$	$4k$	$2k$	$2k$		$2k$	$2k$	$2k$		$2k$
B	$2k$	$2k$	$2k$		$4k$	$2k$		$4k$	$2k$	$2k$	$4k$	$4k$	$3k$
C	$n-1$	$n-1$	n	n	n	n	n	n	$2n-2$				$2n-2$
D		$2k$	$4k$	$4k$	$4k$				$4k$				$4k$
E			$4k$	$4k$	$4k$	$4k$	$4k$	$4k$	$4k$	$4k$	$4k$	$4k$	$4k$
F							$n+1$	$2n+1$			$n+1$		
G			$n+1$	$2n+1$		$n+1$	$2n+1$	$3n+1$	$n+1$	$n+1$			
X	1	2	2	2	4	n	$n+1$		2	n	$n+1$		2
Y						3	5	6		$n+3$	$n+4$		
Z	2	4	4	4	8			$2n+2$	4				4
W	3	5	5	6	8	5	6	8	6	6	6	$n+9$	6

Table 4: Summary of proofs for the entries of Table 5. Only the entries that are used in the proof of Table 5 are non-empty. Each nonempty entry of the table gives the value of the domination parameter indexing the corresponding column on the graph indexing the row. The entries show that the bounds are sharp, or that a certain parameter cannot be bounded from above by a function of another one.

assigning \emptyset to each of v and v' maintains feasibility without increasing the total weight. Performing the above procedure as long as necessary eventually results in a function f such that $f(v) = \emptyset$ for all vertices $v \in V_n^{**}$. Let $V_a = \{v \in V_n \mid f(v) = \{a\}\}$; the sets V_b and V_{ab} are defined similarly. Since $2|V_{ab}| \leq f(V(G_n)) \leq 2n - 3$, we infer that $|V_{ab}| \leq n - 2$. The fact that f is a rainbow weak 2-dominating function of G_n with $f(v) = \emptyset$ for each $v \in V_n^{**}$, implies that $|V_a| \leq 1$ and $|V_b| \leq 1$. If either $V_a = \emptyset$ or $V_b = \emptyset$, then there exists a vertex $v \in V_n^{**}$ such that $|f(N(v))| \leq 1$, contrary to the fact that f is a rainbow weak 2-dominating function and $f(v) = \emptyset$. Consequently, $|V_a| = |V_b| = 1$, which implies $|V_{ab}| = n - 2$. But now, $f(V(G_n)) = 2n - 2$, a contradiction. \square

As most other values in rows A-G in Table 4 are straightforward (in particular the values for $kK_2, kC_4, kH, kK_{4,4}, F_n^3$), we continue with the class F_n^4 ; these are the graphs obtained from a set of n cycles C_4 by identifying a vertex from each of the 4-cycles to a single vertex. Let us denote by v the unique vertex of degree $2n$ in F_n^4 . First, note that $\tilde{\gamma}_{w2}(F_n^4) \leq \tilde{\gamma}_2(F_n^4) \leq n + 1$, which is proven by the inequality $\tilde{\gamma}_{w2}(F_n^4) \leq \tilde{\gamma}_2(F_n^4)$ (see Table 2) and the function $f : V(F_n^4) \rightarrow \mathcal{P}(\{a, b\})$, which assigns $f(v) = \{a\}$, $f(u) = \{b\}$ to all non-neighbors u of v , and $f(x) = \emptyset$ to the remaining vertices. The values $\gamma_{w2}(F_n^4) = \gamma_2(F_n^4) = \tilde{\gamma}_{w2}(F_n^4) = \tilde{\gamma}_2(F_n^4) = n + 1$ can be derived from the following result and the corresponding upper bounds in row 4 in Table 2.

Claim 5.2. For $n \geq 3$, $\gamma_{\{2\}}(F_n^4) = 2n + 1$.

Proof. Let S denote the set of all vertices at distance two from the central vertex v of F_n^4 . Let us denote F_n^4 shortly by G_n . Let us show that $\gamma_{\{2\}}(G_n) \geq 2n + 1$. Indeed, suppose for a contradiction that there exists a $\{2\}$ -dominating function $f : V(G_n) \rightarrow \{0, 1, 2\}$ of G_n of total weight at most $2n$. Since all vertices in S have pairwise disjoint closed neighborhoods, each of them needs weight 2 to dominate vertices in S . Since v is not contained in any closed neighborhood of a vertex from S , we deduce $f(v) = 0$; moreover, $f(N[u]) = 2$ for each vertex $u \in S$. In order to dominate the neighbors of v , we must have $f(u) = 2$ for all $u \in S$. But this implies that $f(N[v]) = 0$, a contradiction. This shows that $\gamma_{\{2\}}(G_n) \geq 2n + 1$. Since $\gamma_{\{2\}}(G) \leq 2\gamma_{w2}(G) - 1 \leq 2(n + 1) - 1$, we derive that the claim is correct. \square

Note that Claim 5.2 also implies that $\gamma_{\times 2}(F_n^4) = 2n + 1$. To see that $\gamma_{\times 2}(F_n^4) \geq 3n + 1$ one needs only to observe that for each vertex u with degree 2 a total double dominating function f of G must assign 1 to each of the neighbors of u . On the other hand, assigning 1 to all vertices yields a total double dominating function of F_n^4 , thus $\gamma_{\times 2}(F_n^4) = 3n + 1$.

Some of the (not straightforward) values in rows X, Y, Z, W will be proven in Subsection 5.3 along with the proofs of unboundedness relations.

5.2 Sharpness of the bounds

The families of graphs used to prove sharpness of the bounds in Table 2 are summarized in Table 5. Most of the required values for families in Table 5 have already been established in Table 4. In fact, in Table 5 there are only three graph families whose values have not yet been determined and are used to show the sharpness of bounds (note that the families marked in bold letters in Table 5 are used to show the non-existence of a function that would bound one parameter with another one).

We start with the family $S(K_{1,n})^-$, that appears in the entries (1,13) and (4,13) of the table. Note that the corresponding bounds are $\gamma(G) \leq \gamma_R(G) - 1$ and $\gamma_{\{2\}}(G) \leq 2\gamma_R(G) - 2$ for an arbitrary

$\rho \leq f(\rho')$	1	2	3	4	5	6	7	8	9	10	11	12	13
γ	γ_t	$\gamma_{\{2\}}$	γ_{w2}	$\gamma_{\{2\}}$	$\gamma_{t\{2\}}$	γ_2	$\gamma_{\times 2}$	$\gamma_{t \times 2}$	$\tilde{\gamma}_{w2}$	$\tilde{\gamma}_2$	$\tilde{\gamma}_{\times 2}$	$\tilde{\gamma}_{t \times 2}$	γ_R
γ	kC_4	kC_4	kC_4	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	kC_4	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	kC_4	kC_4	kK_2	kC_4	$[S(K_{1,n})^-]$
γ_t	$=$	kK_2	$S(K_{2n+1})$	kK_2	$K_n^{**}, n \geq 3$	$S(K_{2n+1})$	kK_2	$K_n^{**}, n \geq 3$	kK_2	kK_2	kK_2	kC_4	kK_2
γ_{w2}	kK_2	kH	$=$	kK_2	kH	kK_2	kK_2	$K_n^{**}, n \geq 3$	kK_2	kK_2	kK_2	$kK_{4,4}$	kK_2
$\gamma_{\{2\}}$	kK_2	kH	$[F_n^4]$	$=$	$K_n^{**}, n \geq 3$	$[F_n^4]$	kK_2	$K_n^{**}, n \geq 3$	$[F_n^4]$	$[F_n^4]$	kK_2	$kK_{4,4}$	$[S(K_{1,n})^-]$
$\gamma_{t\{2\}}$	kK_2	kK_2	kK_2	kK_2	$=$	kK_2	kK_2	$K_n^{**}, n \geq 3$	kK_2	kK_2	kK_2	kC_4	kK_2
γ_2	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$=$	kK_2	$K_n^{**}, n \geq 3$	$K_{1,n}$	kK_2	kK_2	$kK_{4,4}$	$K_{1,n}$
$\gamma_{\times 2}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	F_n^4	$=$	$K_n^{**}, n \geq 3$	$K_{1,n}$	F_n^4	kK_2	$kK_{4,4}$	$K_{1,n}$
$\gamma_{t \times 2}$	Q_n	Q_n	Q_n	Q_n	Q_n	F_n^4	F_n^3	$=$	Q_n	F_n^4	F_n^3	kC_4	Q_n
$\tilde{\gamma}_{w2}$	kK_2	kH	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	$=$	kK_2	kK_2	$kK_{4,4}$	kK_2
$\tilde{\gamma}_2$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$S(K_3^{(n)})$	$S(K_3^{(n)})$	$S(K_3^{(n)})$	$K_{1,n}$	$=$	kK_2	$kK_{4,4}$	$K_{1,n}$
$\tilde{\gamma}_{\times 2}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$K_{1,n}$	$S(K_3^{(n)})$	$S(K_3^{(n)})$	$S(K_3^{(n)})$	$K_{1,n}$	kC_4	$=$	kC_4	$K_{1,n}$
$\tilde{\gamma}_{t \times 2}$	T_n	T_n	T_n	T_n	T_n	T_n	T_n	T_n	T_n	T_n	T_n	$=$	T_n
γ_R	kK_2	kH	$S(K_n^{(2)})$	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	$S(K_n^{(2)})$	$K_n^{**}, n \geq 3$	$K_n^{**}, n \geq 3$	kC_4	kC_4	kK_2	$kK_{4,4}$	$=$

Table 5: Summary of the families of graphs that either achieve the bounds from Table 2, or demonstrate that there is no function bounding one parameter with another (the latter families are bolded). Families in a bracket [...] demonstrate the sharpness of the bounds for graphs with at least one edge.

graph G with edges. Recall that the graphs $S(K_{1,n})^-$ are obtained from the subdivision graph of the star $K_{1,n}$ by deleting a leaf. It is easy to see that $\gamma(S(K_{1,n})^-) = n$ and $\gamma_{\{2\}}(S(K_{1,n})^-) = 2n$ for $n \geq 3$, the main argument being that the closed neighborhoods of the n vertices of degree 1 are pairwise disjoint. To prove the sharpness of the two bounds it remains to prove the following.

Claim 5.3. For $n \geq 3$, $\gamma_R(S(K_{1,n})^-) = n + 1$.

Proof. Note that the function $f : V(G) \rightarrow \{0, 1, 2\}$ that assigns 2 to the unique vertex v of degree n , assigns 1 to all vertices at distance 2 from v , and assigns 0 to all the remaining vertices of $S(K_{1,n})^-$, is a Roman dominating function of the graph. $\gamma_R(S(K_{1,n})^-) \leq n + 1$. On the other hand, since $\gamma(S(K_{1,n})^-) = n$, and since $\gamma(G) \leq \gamma_R(G) - 1$ for any graph G with an edge (see Table 2), we infer the claimed result. \square

We continue with the subdivision graph $S(K_{2n+1})$ of the complete graph of odd order $2n + 1$, which appears in the entries (2,3) and (2,6). Note that the bounds from the table show that $\gamma_t(G) \leq \frac{3\gamma_{w2}(G)-1}{2} \leq \frac{3\gamma_2(G)-1}{2}$ for any graph G .

Claim 5.4. For $n \geq 2$, $\gamma_{w2}(S(K_{2n+1})) = \gamma_2(S(K_{2n+1})) = 2n + 1$ and $\gamma_t(S(K_{2n+1})) = 3n + 1$.

Proof. Note that the vertex set of $S(K_{2n+1})$ is given by $V \cup \binom{V}{2}$, where $V = V(K_{2n+1})$ and $\binom{V}{2}$ are the vertices added in the subdivision of K_{2n+1} . We denote by x_{uv} to the vertex added in the subdivision of the edge uv . Clearly $S(K_{2n+1})$ is bipartite with bipartition $\{V, \binom{V}{2}\}$. On the one hand, V is a 2-dominating set of $S(K_{2n+1})$, showing that $\gamma_{w2}(S(K_{2n+1})) \leq \gamma_2(S(K_{2n+1})) \leq 2n + 1$. On the other hand, we claim that $\gamma_t(S(K_{2n+1})) \geq 3n + 1$. Indeed, suppose to the contrary that D is a total dominating set of $S(K_{2n+1})$ with at most $3n$ vertices. Then either $|D \cap V| \leq 2n - 1$ or $|D \cap \binom{V}{2}| \leq n$. In the former case, there exists a pair $u, v \in V$ of distinct vertices not in D , and therefore $N_{S(K_{2n+1})}(x_{uv}) \cap D = \{u, v\} \cap D = \emptyset$. In the latter case, vertices of $D \cap \binom{V}{2}$ dominate at most $2n$ vertices in V , hence there exists a non-dominated vertex in V . In either case, we obtain a contradiction. Therefore,

$$3n + 1 \leq \gamma_t(S(K_{2n+1})) \leq \frac{3\gamma_{w2}(S(K_{2n+1})) - 1}{2} \leq \frac{3(2n + 1) - 1}{2} = 3n + 1$$

and equalities hold throughout. \square

Finally, the only remaining entries in Table 5 that demonstrate sharpness of the bounds and do not follow from entries in Table 4 are (13,3) and (13,6). The bounds show that $\gamma_R(G) \leq 2\gamma_{w2}(G) - 1 \leq 2\gamma_2(G) - 1$ for any graph G . The sharpness is demonstrated by the family of graphs $G_n = S(K_n^{(2)})$, which are the subdivision graphs of the multigraphs obtained from the complete graphs by duplicating each edge (see Fig. 4 for an example). On the one hand, the (weak) 2-domination number of G_n is at most n , since assigning weight 1 to each original vertex of $K_n^{(2)}$ and weight 0 to all other vertices results in a (weak) 2-dominating function of G_n of weight n . On the other hand, we will now show that the Roman domination number of G_n is at least $2n - 1$.

Claim 5.5. For $n \geq 3$, $\gamma_R(S(K_n^{(2)})) = 2n - 1$.

Proof. Among all minimum weight Roman dominating functions of G_n , choose one, say f , that minimizes the value of S_f , defined as the sum of f -weights of all subdivided vertices. First, we will show that $f(s) = 0$ for every subdivided vertex s . Indeed, suppose for a contradiction that

$f(s) > 0$ where s is a subdivided vertex of maximum f -weight. Let s' be the twin of s , that is, the vertex $s' \neq s$ such that $N_{G_n}(s') = N_{G_n}(s)$, and let t and t' be the two (common) neighbors of s and s' in G_n . We consider two cases:

- *Case 1: $f(s) = 2$.*

Then, $f(s') \leq 1$, since otherwise a Roman dominating function with smaller weight than f could be obtained, by setting $f(s') = 1$.

If both neighbors of s have f -weight 0, then we could obtain a Roman dominating function g of the same weight as f and such that $S_g < S_f$ by setting $g(s) = g(s') = 0$, $g(t) = 1$, $g(t') = 2$, and $g(u) = f(u)$ for all other vertices $u \in V(G_n)$. This contradicts the choice of f .

If both neighbors of s have positive f -weight, then a Roman dominating function with smaller weight than f could be obtained by setting $f(s) = 1$, so this case is also impossible.

Hence, we may assume that $f(t) = 0$ and $f(t') \in \{1, 2\}$. In this case, we could obtain a Roman dominating function g of the same weight as f and such that $S_g < S_f$ by setting $g(s) = 0$, $g(t) = 2$, and $g(u) = f(u)$ for all other vertices $u \in V(G_n)$.

- *Case 2: $f(s) = 1$.*

On the one hand, by the choice of s we have $f(s') \in \{0, 1\}$. On the other hand, by the minimality of the total weight of f , $f(s') = 1$, for otherwise $f(t) = 2$ or $f(t') = 2$, and so $f(s)$ could be set to 0 without violating the constraints of Roman domination. A Roman dominating function g of weight at most that of f and such that $S_g < S_f$ can be obtained by setting $g(s) = g(s') = 0$, $g(t) = 2$, and $g(u) = f(u)$ for all other vertices $u \in V(G_n)$. This contradicts the choice of f .

Since $f(s) = 0$ for every subdivided vertex s , for every pair of original vertices t and t' , either $f(t) = 2$ or $f(t') = 2$ (or both). Hence, at most one original vertex can have weight less than 2. If such a vertex exists, its weight must be 1, hence $\gamma_R(G_n) = f(V(G_n)) \geq 2n - 1$. By the above, since $\gamma_{w2}(G) \leq n$, we infer that $\gamma_R(G_n) = 2n - 1$. \square

By this it is proven that all non-bold entries of Table 5 demonstrate the sharpness of the corresponding bounds from Table 2.

5.3 Proofs of unboundedness

For the direct proofs of unboundedness of one parameter with respect to another one can use the families of graphs summarized in Table 5. We will prove in this section the correctness of these examples. As we will elaborate, some of the unboundedness proofs follow by transitivity, using the bounds in Table 2 and are summarized in Table 3.

While the values for the star $K_{1,n}$ are easy to prove, we can argue the nonexistence of corresponding functions only by focusing on two parameters, notably γ_2 and $\gamma_{t\{2\}}$.

Proposition 5.6. *There is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\gamma_2(G) \leq f(\gamma_{t\{2\}}(G))$ for every graph G admitting both parameters.*

Proof. It is easy to see that $\gamma_2(K_{1,n}) = n$ for $n \geq 2$, since the leaves cannot be dominated by a 2-dominating function f from the outside, i.e., each leaf u must be assigned $f(u) = 1$. On the other hand, assigning $f(v) = 2$ to the central vertex, and $f(u) = 2$ to one of the leaves, results in a total $\{2\}$ -dominating function of $K_{1,n}$, thus $\gamma_{t\{2\}}(K_{1,n}) \leq 4$. \square

Since the parameters in columns 1-4, 9, and 13 are bounded from above by a function of $\gamma_{t\{2\}}(G)$ (see Table 2), we derive that $K_{1,n}$ is also an example for these invariants compared with γ_2 . In addition, since the parameters $\gamma_{\times 2}, \tilde{\gamma}_2, \tilde{\gamma}_{\times 2}$ are bounded from below by γ_2 (see the diagram on Fig. 1) we infer from both observations that the entries (i, j) from the subtable $\{6, 7, 10, 11\} \times \{1, 2, 3, 4, 5, 9, 13\}$ of Table 2, are correct. That is, the family of stars $K_{1,n}$ shows that there does not exist an upper bound on a parameter ρ in terms of a function of another parameter ρ' for all the corresponding pairs (ρ, ρ') . Note that this family cannot be used to obtain similar conclusions also for row 12, that is, for the parameter $\tilde{\gamma}_{t \times 2}(G)$: this parameter is not finite on the family of stars.

Consider now the graphs Q_n , for $n \geq 3$, which can be obtained from the multigraph $K_2^{(n)}$ by replacing each edge with a path P_4 (cf. Fig. 5). These graphs will be used for the proofs of unboundedness in the row 8 of Tables 2 and 4, concerning the parameter $\gamma_{t \times 2}$.

Proposition 5.7. *There is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\gamma_{t \times 2}(G) \leq f(\gamma_{t\{2\}}(G))$ for every graph G admitting both parameters.*

Proof. It is easy to see that $\gamma_{t \times 2}(Q_n) = 2n + 2$. Indeed, if $f : V(G) \rightarrow \{0, 1\}$ is a total double dominating function, then for every vertex of degree 2 both its neighbors must receive f -value 1. This implies that all vertices of Q_n must receive value 1. To see that $\gamma_{t\{2\}}(Q_n) \leq 8$ for $n \geq 3$, consider the function f assigning 2 to both vertices of degree n , and 2 to one of the neighbors of each of these two vertices. \square

Since the parameters $\gamma, \gamma_t, \gamma_{w2}, \gamma_{\{2\}}, \tilde{\gamma}_{w2}, \gamma_R$ are all bounded from above by $\gamma_{t\{2\}}$, we infer from Proposition 5.7 that all entries (i, j) from $\{8\} \times \{1, 2, 3, 4, 5, 9, 13\}$ of Table 2 are correct. In fact, the family of graphs Q_n demonstrates the nonexistence of a function f bounding the corresponding parameters with $f(\gamma_{t \times 2}(G))$.

Recall that the graph $S(K_3^{(n)})$ is the subdivision graph of the multigraph $K_3^{(n)}$ (the multigraph obtained from K_3 by adding $n - 1$ parallel edges between each pair of vertices; cf. Fig. 5).

Proposition 5.8. *There is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\tilde{\gamma}_2(G) \leq f(\gamma_{t \times 2}(G))$ for every graph G admitting both parameters.*

Proof. The result follows from the correctness of the entries (Y,8) and (Y,10) in Table 4. To see this consider the function $f : V(S(K_3^{(n)})) \rightarrow \{0, 1\}$, which assigns 1 exactly to the three vertices of degree $2n$ and to three vertices of degree 2, one from each subdivided parallel edge. Then f is clearly a total double dominating function of the graph, with total weight 6. On the other hand, note that for $n \geq 3$, every rainbow 2-dominating function of $S(K_3^{(n)})$ that assigns the empty set to a vertex of degree $2n$ is of total weight at least $2n$. Furthermore, in every rainbow 2-dominating function of $S(K_3^{(n)})$ that assigns a non-empty set to each vertex of degree $2n$, at least two vertices of degree $2n$ receive the same value, hence the neighbors v of these two vertices must receive a non-empty value. The above arguments imply that $\tilde{\gamma}_2(S(K_3^{(n)})) \geq n + 3$. \square

Since γ_2 and $\gamma_{\times 2}$ are bounded from above by $\gamma_{t \times 2}(G)$, we derive that $S(K_3^{(n)})$ is also an example for these invariants with respect to $\tilde{\gamma}_2$. In addition, the parameter $\tilde{\gamma}_{\times 2}$ is bounded from below by $\tilde{\gamma}_2$ (see the Hasse diagram on Fig. 1), which together with the previous observation implies that the entries (i, j) from the subtable $\{10, 11\} \times \{6, 7, 8\}$ of Table 2, are correct. In fact, the family of graphs $S(K_3^{(n)})$ can be used to demonstrate that there does not exist an upper bound on a parameter ρ in terms of a function of another parameter ρ' for all the corresponding pairs (ρ, ρ') .

Proposition 5.9. *There is no function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $\tilde{\gamma}_{t \times 2}(G) \leq f(\tilde{\gamma}_{\times 2}(G))$, for every graph G admitting both parameters.*

Proof. Recall that T_n , for $n \in \mathbb{N}$, is the graph whose vertex set is $\{v_1, \dots, v_n, w_1, \dots, w_n, s_1, s_2, s_3, t_1, \dots, t_5\}$ and such that s_1, s_2, s_3 induce a triangle, t_1, \dots, t_5 induce C_5 , s_1 and s_3 are adjacent to v_i for every $1 \leq i \leq n$, t_1 and t_5 are adjacent to w_i for every $1 \leq i \leq n$, $v_i w_i \in E(T_n)$ for every $1 \leq i \leq n$, and there are no other edges (see Fig. 6).

Consider h defined as $h(s_1) = h(t_1) = h(t_4) = \{a\}$, $h(s_3) = h(t_3) = h(t_5) = \{b\}$, and $h(v) = \emptyset$ for every other $v \in V(T_n)$. It can be easily checked that h is a rainbow double domination function of T_n , as for every vertex $v \in V(T_n)$, it holds $h_{\cup}(N[v]) = \{a, b\}$. So $\tilde{\gamma}_{\times 2}(T_n) \leq 6$, and indeed it can be seen that it holds by equality, because s_2, t_3 and w_1 have disjoint closed neighborhoods.

As for the rainbow total double domination number, where $h_{\cup}(N(v)) = \{a, b\}$ is required for every vertex $v \in V(T_n)$, first note that if a vertex v has degree two in a graph then its two neighbors have to be labeled with different labels. This implies that in a rainbow total double domination function h of T_n , $h(t_1) \neq h(t_3)$ and $h(t_5) \neq h(t_3)$, hence $h(t_1) = h(t_5)$. So the vertices w_1, \dots, w_n are missing one label on their open neighborhoods, therefore vertices v_1, \dots, v_n have to have a nonempty label, implying $\tilde{\gamma}_{t \times 2}(T_n) \geq n$. Indeed, a rainbow total double domination function h of T_n can be defined as $h(s_1) = h(s_2) = h(t_1) = h(t_2) = h(t_5) = \{a\}$, $h(s_3) = h(t_3) = h(t_4) = h(w_1) = \{b\}$, $h(v_i) = \{b\}$ for every $1 \leq i \leq n$, and $h(w_i) = \emptyset$ for every $2 \leq i \leq n$. It is not hard to see that its weight is minimum possible, so $\tilde{\gamma}_{t \times 2}(T_n) = n + 9$. \square

Since all the parameters (except of course for $\tilde{\gamma}_{t \times 2}$) in graphs G are bounded by a function of $\tilde{\gamma}_2(G)$ we infer that the entries in the row 12 of Table 2 are correct. In fact, the family of graphs T_n demonstrates the nonexistence of a function f bounding any of the other parameters with $f(\tilde{\gamma}_{t \times 2}(G))$.

6 Algorithmic and complexity issues

We now discuss the algorithmic and complexity consequences of the bounds obtained in Section 3 for corresponding optimization problems. More specifically, we obtain new results regarding the existence of approximation algorithms for the studied invariants, matched with tight or almost tight inapproximability bounds, which hold even in the class of split graphs.

Recall that an algorithm \mathcal{A} for a minimization problem Π is said to be a *c-approximation algorithm* (where $c \geq 1$) if it runs in polynomial time and for every instance I of Π , we have $\mathcal{A}(I) \leq c \cdot OPT(I)$, where $\mathcal{A}(I)$ is the value of the solution produced by \mathcal{A} , given I , and $OPT(I)$ is the optimal solution value, given I . (For more details on complexity and approximation, we refer to [5, 77].) Given a graph G , let $\rho(G)$ denote the optimal value of any of the minimization parameters studied in this paper (e.g., the domination number of G , the rainbow total double domination number of G , etc.). The corresponding optimization problem is the following problem: Given a graph G , compute the value of $\rho(G)$. In the case of a *c-approximation algorithm* for the above problem, we also require that for each instance G not only an approximation to the optimal value but also a feasible solution to the problem is computed achieving value at most $c \cdot \rho(G)$. Note that in the problems relating to any of the parameters considered in this paper, a feasible solution is a function f with domain V , whose value equals the total weight $f(V)$ (see Section 2).

First we recall a simple (folklore) observation that can be useful for transferring both lower and upper bounds regarding (in)approximability of minimization problems. In order to keep the

notation as simple as possible, we keep the presentation of the result confined to the parameters defined in Section 2, however, the same result clearly applies more generally. For the sake of completeness, we include the simple proof.

Proposition 6.1. *Let ρ and ρ' be any two graph invariants defined in Section 2 and let \mathcal{G} be a class of graphs such that there exist constants $c_1, c_2 > 0$ such that for all $G \in \mathcal{G}$, we have $c_1 \cdot \rho(G) \leq \rho'(G) \leq c_2 \cdot \rho(G)$. Suppose furthermore that there exists a polynomial time algorithm that for a given graph $G \in \mathcal{G}$ and a feasible solution f to ρ , computes a feasible solution f' to ρ' with $f'(V(G)) \leq c_2 \cdot f(V(G))$. Then, for every $c \geq 1$, if there is a c -approximation algorithm for ρ on graphs in \mathcal{G} , then there is (cc_2/c_1) -approximation algorithm for ρ' on graphs in \mathcal{G} .*

Proof. Let \mathcal{A} be a c -approximation algorithm for ρ on graphs in \mathcal{G} . Consider the following algorithm for ρ' on graphs in \mathcal{G} :

1. Given a graph $G \in \mathcal{G}$, run \mathcal{A} on G and let $f_{\mathcal{A}}$ be the solution produced by \mathcal{A} .
2. Compute a feasible solution f' to ρ' with $f'(V(G)) \leq c_2 \cdot f_{\mathcal{A}}(V(G))$ using the algorithm that exists by assumption.
3. Return f' .

Since \mathcal{A} is a c -approximation algorithm for ρ on graphs in \mathcal{G} , we have $f_{\mathcal{A}}(V(G)) \leq c\rho(G)$. It follows that $f'(V(G)) \leq c_2 \cdot f_{\mathcal{A}}(V(G)) \leq c_2 \cdot c\rho(G) \leq (c_2c/c_1)\rho'(G)$, where the last inequality follows from $c_1\rho(G) \leq \rho'(G)$. As the algorithm clearly runs in polynomial time, it is a (c_2c/c_1) -approximation algorithm for ρ' for graphs in \mathcal{G} . \square

Note that all the bounds from Table 2 are of the form $\rho'(G) \leq c\rho(G) - d$ for some constants $c \geq 1$ and $d \geq 0$, hence they immediately imply bounds of the form $\rho'(G) \leq c\rho(G)$ (for some constant $c \geq 1$). Furthermore, it follows from the proofs of the bounds that all the translations between parameters involving bounds summarized in Table 2 can be efficiently constructed, in the sense that if $\rho'(G) \leq c\rho(G)$ is a bound following from bounds in Table 2, then there is a polynomial time algorithm that, given a graph $G = (V, E)$ and a feasible solution f to ρ , computes a feasible solution f' to ρ' with $f'(V) \leq c \cdot f(V)$.

6.1 Lower bounds

Several hardness and inapproximability results for variants of domination considered in the paper are already known in the literature. We list here only the strongest results known and an earliest available proof for each of them, making no attempt to survey the literature regarding hardness of the problems in various graph classes – with the single exception of the class of split graphs, which naturally appears in many of the underlying proofs. A graph $G = (V, E)$ is said to be *split* if it admits a *split partition*, that is, a pair (C, I) such that C is a clique in G , I is an independent set in G , $C \cup I = V$, and $C \cap I = \emptyset$. Split graphs were introduced by Földes and Hammer in [33], where several characterizations were also given.

Theorem 6.2 (combining results from [17, 19, 60, 69]). *For every $\rho \in \{\gamma, \gamma_t, \gamma_2, \gamma_{\times 2}, \gamma_{t \times 2}\}$ and every $\epsilon > 0$, there is no polynomial time algorithm approximating ρ for n -vertex split graphs without isolated vertices within a factor of $(1 - \epsilon) \ln n$, unless $\text{NP} \subseteq \text{DTIME}(n^{O(\log \log n)})$.*

The statement of Theorem 6.2 was proved:

- (i) for domination and total domination (γ, γ_t) by Chlebík and Chlebíková in [17],
- (ii) for 2-domination (γ_2) by Cicalese et al. [19] (in the more general context of k -domination),
- (iii) for double domination $(\gamma_{\times 2})$ by Klasing and Laforest [60] (in the more general context of k -tuple domination),
- (iv) for total double domination $(\gamma_{t \times 2})$ independently by Pradhan [69] and by Cicalese et al. [19] (in both cases in the more general context of k -tuple domination).

Only the results by Chlebík and Chlebíková were mentioned explicitly for split graphs. However, since the corresponding reductions from [19, 60, 69] are performed from either domination or total domination by simply adding a number of universal vertices to the input graph, all of the above results also hold for split graphs.

The basis of the inapproximability results from [17, 19, 60, 69] summarized in Theorem 6.2 is the analogous result due to Feige for the well-known SET COVER problem: Given a set system (S, \mathcal{F}) where S is a finite set (also called a *ground set*) and \mathcal{F} is a family (multiset) of subsets of S , find a smallest *set cover* of \mathcal{F} , that is, a sub-collection $\mathcal{F}' \subseteq \mathcal{F}$ such that $\bigcup \mathcal{F}' = S$ (that is, such that every element of S appears in some member of \mathcal{F}').

Theorem 6.3 (Feige [30]). *For every $\epsilon > 0$, there is no polynomial time algorithm approximating SET COVER within a factor of $(1 - \epsilon) \ln n$ (where n is the size of the ground set), unless $\text{NP} \subseteq \text{DTIME}(n^{O(\log \log n)})$.*

In 2014, Dinur and Steurer improved Feige’s inapproximability result by weakening the hypothesis to $\text{P} \neq \text{NP}$.

Theorem 6.4 (Dinur and Steurer [24]). *For every $\epsilon > 0$, there is no polynomial time algorithm approximating SET COVER within a factor of $(1 - \epsilon) \ln n$, unless $\text{P} = \text{NP}$.*

An essential fact in proving the bounds from Theorem 6.2 is that the instances of SET COVER arising in Feige’s construction are such that $\ln(|S| + |\mathcal{F}|) \approx \ln |S|$, that is, the ratio $\ln(|S| + |\mathcal{F}|) / \ln |S|$ can be assumed to be arbitrarily close to 1. This is also true for the instances of arising in the construction proving Theorem 6.4. Consequently, Theorem 6.2 can be improved as follows:

Theorem 6.5. *For every $\rho \in \{\gamma, \gamma_t, \gamma_2, \gamma_{\times 2}, \gamma_{t \times 2}\}$ and every $\epsilon > 0$, there is no polynomial time algorithm approximating ρ for n -vertex split graphs without isolated vertices within a factor of $(1 - \epsilon) \ln n$, unless $\text{P} = \text{NP}$.*

In particular, the above results imply that the decision variants of the corresponding optimization problems are NP-complete.

We are not aware of inapproximability results for any of the invariants $\rho \in \{\gamma_R, \gamma_{w2}, \gamma_{\{2\}}, \gamma_{t\{2\}}, \tilde{\gamma}_2, \tilde{\gamma}_{w2}, \tilde{\gamma}_{\times 2}, \tilde{\gamma}_{t \times 2}\}$. (Recall that invariants $\tilde{\gamma}_2$, $\tilde{\gamma}_{\times 2}$, and $\tilde{\gamma}_{t \times 2}$ are, to the best of our knowledge, considered for the first time in this paper.) The following NP-completeness results for some of these parameters are available in the literature:

- The NP-completeness of Roman domination (γ_R) was proved by Dreyer in [28]. (The problem was already claimed to be NP-complete in [20], referring to a private communication with A.A. McRae.)

- The weak 2-domination (γ_{w2}) and the rainbow weak 2-domination ($\tilde{\gamma}_{w2}$) problems were proved NP-complete by Brešar and Kraner Šumenjak in [10].
- The NP-completeness of $\{2\}$ -domination ($\gamma_{\{2\}}$) was proved by Gairing et al. in [36] (in the more general context of $\{k\}$ -domination).
- The NP-completeness of rainbow double domination ($\tilde{\gamma}_{\times 2}(G)$; for graphs without isolated vertices) follows from the analogous result due to Hedetniemi et al. [49] for disjoint domination (cf. Proposition 2.1).

We are not aware of any published hardness results about total $\{2\}$ -domination ($\gamma_{\{2\}}$).

In the rest of this subsection, we strengthen the above NP-completeness results by showing that all the domination parameters studied in this paper, except for the rainbow total double domination number, admit an inapproximability bound of the form $\Omega(\ln n)$ for n -vertex split graphs, unless $P = NP$. Before doing that, we show that for the rainbow total double domination number ($\tilde{\gamma}_{\times 2}$; recall that this is the topmost parameter in the diagram in Fig. 2), the situation is even worse. We say that a graph G is $\tilde{\gamma}_{\times 2}$ -feasible if $\tilde{\gamma}_{\times 2}(G)$ is finite (cf. Proposition 2.2 on p. 10).

Theorem 6.6. *There is no polynomially computable function f such that there exists an $f(n)$ -approximation algorithm for rainbow total double domination on n -vertex $\tilde{\gamma}_{\times 2}$ -feasible split graphs, unless $P = NP$.*

Proof. Suppose for a contradiction that there exists a polynomially computable function f such that there exists an $f(n)$ -approximation algorithm for rainbow total double domination on n -vertex $\tilde{\gamma}_{\times 2}$ -feasible split graphs. We will show that this implies $P = NP$, by designing a polynomial time algorithm for the NP-complete HYPERGRAPH 2-COLORABILITY problem [41], which asks whether a given hypergraph is 2-colorable. A *hypergraph* \mathcal{H} is a pair (V, \mathcal{E}) where V is a finite set and \mathcal{E} is a set of subsets of V . A hypergraph is said to be 2-colorable if its vertex set V admits a partition into two independent sets A and B , where a set $X \subseteq V$ is *independent* if it does not contain any hyperedge $e \in \mathcal{E}$. We may assume that $|A| \geq 2$ and $|B| \geq 2$ in every partition as above since otherwise the problem can be solved in polynomial time.

Given an input $\mathcal{H} = (V, \mathcal{E})$ to the HYPERGRAPH 2-COLORABILITY problem, construct the split graph $G = (V', E)$ with split partition (C, I) where $C = V$, $I = \mathcal{E}$, and there is an edge in G between $v \in C$ and $e \in I$ if and only if $v \in e$. Clearly, G can be constructed from \mathcal{H} in polynomial time.

We claim that \mathcal{H} is 2-colorable if and only if G is $\tilde{\gamma}_{\times 2}$ -feasible. First, suppose that \mathcal{H} is 2-colorable, and let $\{A, B\}$ be a partition of V into two independent sets. Then, the function $g : V(G) \rightarrow \{\emptyset, \{a\}, \{b\}\}$ defined by

$$g(v) = \begin{cases} \{a\}, & \text{if } v \in A; \\ \{b\}, & \text{if } v \in B; \\ \emptyset, & \text{otherwise.} \end{cases}$$

is a rainbow total double dominating function of G . Indeed, the assumption $|A| \geq 2$ and $|B| \geq 2$ implies that $g_{\cup}(N(v)) = \{a, b\}$ for all $v \in C$, while the fact that A and B are both independent in \mathcal{H} implies that g also dominates vertices in I . It follows that G is $\tilde{\gamma}_{\times 2}$ -feasible. Conversely, suppose that G is $\tilde{\gamma}_{\times 2}$ -feasible, with a rainbow total double dominating function $g : V(G) \rightarrow \{\{a\}, \{b\}, \emptyset\}$. Modify g if necessary by setting $g(v) = \{a\}$ for every $v \in C$ with $g(v) = \emptyset$; clearly, the so

obtained function is still a rainbow total double dominating function of G . Moreover, the sets $A = \{v \in V : g(v) = a\}$ and $B = \{v \in V : g(v) = b\}$ form a partition of V , the vertex set of \mathcal{H} . Since $g_{\cup}(N(v)) = \{a, b\}$ for all $v \in I$, each of the sets A and B is independent in \mathcal{H} , and thus \mathcal{H} is 2-colorable.

Now, let $n = |V'|$, and let \mathcal{A} be an $f(n)$ -approximation algorithm for rainbow total double domination on n -vertex $\tilde{\gamma}_{t \times 2}$ -feasible split graphs. We know that \mathcal{A} computes a rainbow total double dominating function on $\tilde{\gamma}_{t \times 2}$ -feasible split graphs, but if the input graph is not of this form, there is no guarantee about what \mathcal{A} computes or whether it even halts. By definition \mathcal{A} runs in polynomial time on n -vertex $\tilde{\gamma}_{t \times 2}$ -feasible split graphs, say its running time is bounded by a polynomial $p(n)$.

The polynomial time algorithm that decides whether \mathcal{H} is 2-colorable goes as follows.

1. Construct the split graph G as specified above.
2. Compute $n = |V(G)|$ and $f(n)$, and let \mathcal{A} be an $f(n)$ -approximation algorithm for rainbow total double domination on n -vertex $\tilde{\gamma}_{t \times 2}$ -feasible split graphs.
3. Run \mathcal{A} on G for at most $p(n)$ steps.
4. If \mathcal{A} did not compute anything, then G is not $\tilde{\gamma}_{t \times 2}$ -feasible. We conclude that \mathcal{H} is not 2-colorable.
5. If \mathcal{A} computed something, then check whether what it computed is a rainbow total double dominating function on G .

If it is, then G is $\tilde{\gamma}_{t \times 2}$ -feasible, and we conclude that \mathcal{H} is 2-colorable. (In this case we also have that the total weight of the computed function is at most $f(n)\tilde{\gamma}_{t \times 2}(G)$, but we will not need this fact.)

If it is not, then G is not $\tilde{\gamma}_{t \times 2}$ -feasible, and we conclude that \mathcal{H} is not 2-colorable.

It is clear that the algorithm runs in polynomial time. Its correctness follows from the correctness of \mathcal{A} and from the fact that \mathcal{H} is 2-colorable if and only if G is $\tilde{\gamma}_{t \times 2}$ -feasible. Thus, the above algorithm efficiently solves the NP-complete HYPERGRAPH 2-COLORABILITY problem, implying that $P = NP$. This completes the proof. \square

We now turn our attention to the remaining parameters. Proposition 6.1 and the discussion following it show that in order to prove an inapproximability bound of the form $\Omega(\ln n)$ for each of the remaining considered parameters, namely $\rho \in \{\gamma_{w2}, \gamma_{\{2\}}, \gamma_{t\{2\}}, \gamma_R, \tilde{\gamma}_2, \tilde{\gamma}_{w2}, \tilde{\gamma}_{\times 2}\}$, it suffices to show an inapproximability bound of the same type for just one parameter in each of the bottom three equivalence classes in the diagram of Fig. 2. As mentioned above, such bounds already exist, even for the class of split graphs, for any $\rho \in \{\gamma, \gamma_t, \gamma_2, \gamma_{\times 2}, \gamma_{t \times 2}\}$, which takes care of the invariants appearing in the bottom two equivalence classes in Fig. 2. We summarize this in the following theorem.

Theorem 6.7. *For every $\rho \in \{\gamma_{w2}, \gamma_{\{2\}}, \gamma_{t\{2\}}, \gamma_R, \tilde{\gamma}_{w2}, \tilde{\gamma}_{\{2\}}, \tilde{\gamma}_{t\{2\}}\}$ and every $\epsilon > 0$, there is no polynomial time algorithm approximating ρ for n -vertex split graphs without isolated vertices within a factor of $(1/2 - \epsilon) \ln n$, unless $P = NP$.*

Proof. Recall from Section 3 that for every graph G without isolated vertices, we have $\gamma(G) \leq \gamma_{w2}(G) \leq \tilde{\gamma}_{w2}(G) \leq \gamma_R(G) \leq \tilde{\gamma}_{\{2\}}(G) = 2\gamma(G)$ and $\gamma_t(G) \leq \gamma_{\{2\}}(G) \leq \gamma_{t\{2\}}(G) \leq \tilde{\gamma}_{t\{2\}}(G) = 2\gamma_t(G)$. Thus, the theorem follows from the inapproximability bound for domination (resp., total domination), see Theorem 6.5, the above inequalities, and Proposition 6.1. We prove the statement formally only for the weak 2-domination number (γ_{w2}); the proofs for the other parameters are analogous.

Let \mathcal{G} be the class of split graphs without isolated vertices and suppose that there is some $\epsilon > 0$ such that there is a polynomial time algorithm approximating the weak 2-domination number on n -vertex graphs in \mathcal{G} within a factor of $(1/2 - \epsilon) \ln n$. For every graph G , we have $\frac{1}{2} \gamma_{w2}(G) \leq \gamma(G) \leq \gamma_{w2}(G)$. Moreover, for every weak 2-dominating function f of G , the set $\{v \in V(G) : f(v) > 0\}$ is a dominating function of G of weight at most $f(V(G))$. Therefore, Proposition 6.1 applies with $c_1 = 1/2$, $c_2 = 1$, and hence there is a polynomial time algorithm approximating the domination number on n -vertex graphs in \mathcal{G} within a factor of $(1 - 2\epsilon) \ln n$. By Theorem 6.5, this is only possible if $P = NP$. \square

We also explicitly state the following consequence of Theorem 6.7 for total $\{2\}$ -domination ($\gamma_{t\{2\}}$), which does not seem to be yet available in the literature.

Corollary 6.8. *The decision version of the total $\{2\}$ -domination problem is NP-complete.*

The remaining equivalence class from Fig. 2 contains two parameters, namely rainbow 2-domination ($\tilde{\gamma}_2$) and rainbow double domination ($\tilde{\gamma}_{\times 2}$). Using a reduction from SET COVER, we now prove the inapproximability bounds for the rainbow 2-domination ($\tilde{\gamma}_2$) and the rainbow double domination ($\tilde{\gamma}_{\times 2}$) problems in split graphs. As discussed above, it would suffice to prove a bound for only one of the two parameters. We give a direct proof for both parameters, since with almost no additional work, we save a multiplicative factor of 2 in one of the two bounds compared to the bounds we would obtain using the above approach.

Theorem 6.9. *For every $\rho \in \{\tilde{\gamma}_2, \tilde{\gamma}_{\times 2}\}$ and every $\epsilon > 0$, there is no polynomial time $(1 - \epsilon) \ln n$ -approximation algorithm for computing ρ on n -vertex split graphs, unless $P = NP$.*

Proof. Fix $\rho \in \{\tilde{\gamma}_2, \tilde{\gamma}_{\times 2}\}$ and suppose for some $\epsilon > 0$, there is a polynomial time $(1 - \epsilon) \ln n$ -approximation algorithm, say \mathcal{A} , for computing ρ on n -vertex split graphs.

Let $J = (S, \mathcal{F})$ be an instance to the SET COVER problem. First, note that we may assume that

$$\ln 3 + \ln(|S| + |\mathcal{F}|) \leq (1 + \epsilon/2) \ln(|S| + |\mathcal{F}|) \leq (1 + \epsilon) \ln |S|. \quad (2)$$

Indeed, if the first inequality above is violated, then $\ln(|S| + |\mathcal{F}|)$ is bounded by $2 \ln 3/\epsilon$ and the problem can be solved in constant time. The second inequality follows from the fact that the ratio $\ln(|S| + |\mathcal{F}|)/(\ln |S|)$ can be made arbitrarily close to 1 (as remarked right after Theorem 6.4).

Consider the split graph $G_J = (V, E)$ with split partition (C, I) where $C = A \cup B$ with $A = \{a_F \mid F \in \mathcal{F}\}$, $B = \{b_F \mid F \in \mathcal{F}\}$, $I = S_1 \cup S_2 \cup S_3$ with $S_j = \{s_j : s \in S\}$ for $j \in \{1, 2, 3\}$, and there is an edge between F_i for $i \in \{a, b\}$, $F \in \mathcal{F}$ and $s_j \in S_j$ for $j \in \{1, 2, 3\}$, $s \in S$ if and only if $s \in F$.

Graph G_J has $3|S| + 2|\mathcal{F}| \leq 3(|S| + |\mathcal{F}|) \leq |S|^{1+\epsilon}$ vertices and can be computed in polynomial time from J . Let OPT denote the minimum size of a set cover for \mathcal{F} . First, we prove the following claim.

Claim: $\tilde{\gamma}_2(G_J) = \tilde{\gamma}_{\times 2}(G_J) = 2 \cdot OPT$.

Proof of claim: The inequality $\tilde{\gamma}_2(G_J) \leq \tilde{\gamma}_{\times 2}(G_J)$ always holds (see Table 2). Thus, it remains to prove $\tilde{\gamma}_{\times 2}(G_J) \leq 2 \cdot OPT$ and $2 \cdot OPT \leq \tilde{\gamma}_2(G_J)$.

We first prove that $\tilde{\gamma}_{\times 2}(G_J) \leq 2 \cdot OPT$. First, let \mathcal{F}' be a minimum set cover for \mathcal{F} . Consider the function $f : V(G_J) \rightarrow \{\emptyset, \{a\}, \{b\}\}$ defined as follows:

$$f(v) = \begin{cases} \{a\}, & \text{if } v = a_F \in A \text{ and } F \in \mathcal{F}'; \\ \{b\}, & \text{if } v = b_F \in B \text{ and } F \in \mathcal{F}'; \\ \emptyset, & \text{otherwise.} \end{cases}$$

Clearly, $f(V(G_J)) = 2|\mathcal{F}'| = 2 \cdot OPT$. Thus, to prove that $\tilde{\gamma}_{\times 2}(G_J) \leq 2 \cdot OPT$, it suffices to check that f is a rainbow double dominating function of G_J , that is, that $f_{\cup}(N[v]) = \{a, b\}$ holds for all $v \in V$. If $v \in I$, then $v = s_j$ for some $s \in S$ and some $j \in \{1, 2, 3\}$. There exists some $F \in \mathcal{F}'$ with $s \in F$. This implies that a_F and b_F are adjacent to s_j in G_J , and by construction these two vertices are labeled $\{a\}$ and $\{b\}$, respectively. If $v \in C$, then we have either $f(v) = \{a\}$ or $f(v) = \{b\}$ or $f(v) = \emptyset$. If $f(v) = \{a\}$ then since C is a clique, any vertex b_F with $F \in \mathcal{F}'$ is a neighbor of v labeled $\{b\}$. The case when $f(v) = \{b\}$ is symmetric. Finally, if $f(v) = \emptyset$, then we similarly observe that v is adjacent to both a vertex of the form a_F and a vertex of the form b_F (with $F \in \mathcal{F}'$). It follows that f is a rainbow $\{2\}$ -dominating function of G_J , which implies $\tilde{\gamma}_{\times 2}(G_J) \leq 2 \cdot OPT$.

Now, we prove that $OPT \leq \tilde{\gamma}_2(G_J)/2$. Let $f : V(G_J) \rightarrow \{\emptyset, \{a\}, \{b\}\}$ be a minimum rainbow 2-dominating function. We therefore have $f_{\cup}(N(v)) = \{a, b\}$ for all $v \in V(G_J)$ with $f(v) = \emptyset$. First, we will show that we have $f(v) = \emptyset$ for all $v \in I$. Suppose for a contradiction that $f(v) \neq \emptyset$ for some $v \in I$. By minimality of f , the function obtained by relabeling v to \emptyset is not a rainbow 2-dominating function of G_J , which implies that v does not have both labels $\{a\}$ and $\{b\}$ in its neighborhood. Assume that $a \notin f_{\cup}N(v)$ (the other case is symmetric). Let $s \in S$ and $j \in \{1, 2, 3\}$ be such that $v = s_j$. Then $a \notin f_{\cup}N(u)$ for all $u \in \{s_1, s_2, s_3\}$, which implies that $f(s_j) \neq \emptyset$ for all $j \in \{1, 2, 3\}$. Let $F \in \mathcal{F}$ such that $s \in F$, and consider the function f' obtained from f by relabeling each of s_j to \emptyset , and by setting $f'(a_F) = \{a\}$ and $f'(b_F) = \{b\}$ (and leaving all other values unchanged). It is easy to see that f' is a rainbow 2-dominating function of smaller total weight than f . This is a contradiction with the minimality of f and proves that $f(v) = \emptyset$ for all $v \in I$. This assumption implies that every $v \in I$ has both labels $\{a\}$ and $\{b\}$ in its neighborhood.

The minimality of f implies that for every $F \in \mathcal{F}$, at most one of a_F and b_F gets label $\{a\}$. (If both a_F and b_F would get label $\{a\}$, then replacing one of them with \emptyset would result in a rainbow 2-dominating function of G_J of smaller total weight than f .) Similarly, at most one of a_F and b_F gets label $\{b\}$. Also, by the symmetry of the construction, we may assume that if one of a_F and b_F gets label $\{a\}$, then $f(a_F) = \{a\}$, and that if one of a_F and b_F gets label $\{b\}$, then $f(b_F) = \{b\}$. Thus, $A' = \{v \in C : f(v) = \{a\}\}$ and $B' = \{v \in C : f(v) = \{b\}\}$ satisfy $A' \subseteq A$ and $B' \subseteq B$. Without loss of generality assume that $|A'| \leq |B'|$. We claim that $\mathcal{F}' = \{F \in \mathcal{F} : a_F \in A'\}$ is a set cover of \mathcal{F} . Indeed, if $s \in S$, then the fact that $s_1 \in I$ and every vertex in I has label $\{a\}$ in its neighborhood implies that there is a vertex $a_F \in N(s_1)$ such that $f(a_F) = \{a\}$, in other words $a_F \in A'$, which implies that $s \in F$ (since $a_F \in N(s_1)$) and $F \in \mathcal{F}'$ (since $a_F \in A'$). Since \mathcal{F}' is a set cover of \mathcal{F} , it follows that

$$OPT \leq |\mathcal{F}'| = |A'| \leq \frac{|A'| + |B'|}{2} = \frac{f(V(G_J))}{2} = \frac{\tilde{\gamma}_2(G_J)}{2}.$$

This completes the proof of the claim. \blacktriangle

Now we can complete the proof of the theorem. Recall that \mathcal{A} is a polynomial time $(1 - \epsilon) \ln n$ -approximation algorithm for computing ρ on n -vertex split graphs. Using \mathcal{A} , we can design an

approximation algorithm for SET COVER, transforming an instance $J = (S, \mathcal{F})$ to the split graph G_J , computing an approximate solution f to ρ on G_J , and returning the corresponding set cover \mathcal{F}' obtained from f as in the above proof of the claim. Letting $n = |V(G_J)|$, we can bound the size of \mathcal{F}' from above as

$$\begin{aligned}
|\mathcal{F}'| &\leq f(V(G_J))/2 && \text{(by the above proof of the claim)} \\
&\leq (1 - \epsilon)(\ln n)\rho(G_J)/2 && \text{(since } f \text{ was computed using the} \\
&&& \text{(1 - } \epsilon \text{) ln } n\text{-approximation algorithm } \mathcal{A}\text{)} \\
&\leq (1 - \epsilon) \ln(3(|S| + |\mathcal{F}|))OPT && \text{(since } n \leq 3(|S| + |\mathcal{F}|) \text{ and } \rho(G_J) = 2 \cdot OPT\text{)} \\
&\leq (1 - \epsilon)(1 + \epsilon)(\ln |S|)OPT && \text{(by (2))} \\
&\leq (1 - \epsilon^2)(\ln |S|)OPT,
\end{aligned}$$

Therefore, there exists a polynomial time algorithm that computes a $(1 - \epsilon^2) \ln |S|$ -approximation to SET COVER. By Theorem 6.4, this is only possible if $P = NP$. \square

6.2 Upper bounds

The following theorem summarizes the upper bounds on approximability of domination parameters considered in this paper available in the literature:

Theorem 6.10 (combining results from [19, 25, 60, 69]).

1. For each $\rho \in \{\gamma, \gamma_{\times 2}\}$, there is a $(\ln(\Delta(G) + 1) + 1)$ -approximation algorithm for ρ .
2. For each $\rho \in \{\gamma_t, \gamma_{t \times 2}\}$, there is a $(\ln(\Delta(G)) + 1)$ -approximation algorithm for ρ .
3. For each $\rho \in \{\gamma_2, \gamma_{w2}\}$, there is a $(\ln(\Delta(G) + 2) + 1)$ -approximation algorithm ρ .

The result for domination and total domination follows from the fact that these two problems can be easily modeled as special cases of SET COVER. It is well known that a simple greedy algorithm for SET COVER produces a solution that is always within a factor $\ln \Delta + 1$ of the optimum, where Δ is the maximum size of a set in \mathcal{F} [25]. As proved independently by Dobson [25] and by Klasing and Laforest [60], the same is true for the more general problem in which the task is to find a minimum size subcollection $\mathcal{F}' \subseteq \mathcal{F}$ such that every element s appears in at least k sets in \mathcal{F}' (Dobson's result is in fact more general: each vertex can have a different coverage requirement). In turn, this implies the above-mentioned approximation results for double domination ($\gamma_{\times 2}$) and total double domination ($\gamma_{t \times 2}$); see [19, 60, 69]. The result for 2-domination (γ_2) can be obtained with a straightforward modification of the proof of [19, Theorem 3]. That result gives an approximation algorithm for the more general problem called *vector domination* (in which one seeks a small subset S of vertices of a graph such that any vertex outside S has at least a prescribed number of neighbors in S), using a reduction to the so-called MINIMUM SUBMODULAR COVER problem and applying a result of Wolsey [78].

Without trying to optimize the obtained approximation ratios, let us simply note that Theorem 6.10 and a similar approach to that used in the proof of Theorem 6.7 implies the following result.

Theorem 6.11. For every $\rho \in \{\gamma_{\{2\}}, \gamma_{t\{2\}}, \tilde{\gamma}_{w2}, \gamma_R, \tilde{\gamma}_{\{2\}}, \tilde{\gamma}_{t\{2\}}\}$, there is a $2(\ln(\Delta(G) + 2) + 1)$ -approximation algorithm for ρ .

To the best of our knowledge, these are the first results regarding approximation algorithms for any of these parameters. Development of approximation algorithms for rainbow 2-domination and rainbow double domination remains an open question.

We conclude with the following related questions, which we leave for future research:

- Can the factors $1/2 - \epsilon$ in the inapproximability bounds from Theorem 6.7 be improved to $1 - \epsilon$? Possible approaches to this question include a development of direct reductions from SET COVER and a study of the inequalities relating the relevant parameters in the class of split graphs.
- Can the approximation ratios given by Theorem 6.11 be further improved?
- The only known inapproximability bound for the rainbow 2-domination and rainbow double domination problems are those given by Theorem 6.9, and no nontrivial approximation algorithms for these two problems are known. It would be interesting to settle the (in-)approximability status of these two problems. The case of the rainbow double domination number $\tilde{\gamma}_{\times 2}(G)$ of a graph G without isolated vertices is particularly interesting, because the parameter coincides with the previously studied disjoint domination number $\gamma\gamma(G)$ (cf. Proposition 2.1).

Acknowledgements

The work for this paper was done in the framework of a bilateral project between Argentina and Slovenia, financed by the Slovenian Research Agency (BI-AR/12–14–012, BI-AR/12–14–013) and MINCyT, Argentina (SLO/11/12 and SLO/11/13). F. Bonomo, L.N. Grippo, and M.D. Safe were partially supported by UBACyT Grant 20020130100808BA, CONICET PIP 112-201201-00450CO, and ANPCyT PICT-2012-1324 (Argentina). F. Bonomo was partially supported by ANPCyT PICT-2015-2218 (Argentina). L.N. Grippo and M.D. Safe were partially supported by PIO CONICET UNGS 144-20140100027-CO (Argentina). B. Brešar was supported in part by the Slovenian Research Agency (ARRS) under the grant P1-0297. Research of M. Milanič was supported in part by the Slovenian Research Agency (I0-0035, research program P1-0285 and research projects N1-0032, J1-5433, J1-6720, J1-6743, and J1-7051).

References

- [1] B. D. Acharya. Domination in hypergraphs II. New directions. In *Advances in discrete mathematics and applications: Mysore, 2008*, volume 13 of *Ramanujan Math. Soc. Lect. Notes Ser.*, pages 1–18. Ramanujan Math. Soc., Mysore, 2010.
- [2] M. Adabi, E. E. Targhi, N. J. Rad, and M. S. Moradi. Properties of independent Roman domination in graphs. *Australas. J. Combin.*, 52:11–18, 2012.
- [3] V. Anusuya and R. Kala. A note on disjoint dominating sets in graphs. *Int. J. Contemp. Math. Sci.*, 7(41-44):2099–2110, 2012.
- [4] H. Aram, S. M. Sheikholeslami, and L. Volkmann. On the total $\{k\}$ -domination and total $\{k\}$ -domatic number of graphs. *Bull. Malays. Math. Sci. Soc. (2)*, 36(1):39–47, 2013.

- [5] G. Ausiello, P. Crescenzi, G. Gambosi, V. Kann, A. Marchetti-Spaccamela, and M. Protasi. *Complexity and Approximation*. Springer-Verlag, Berlin, 1999.
- [6] S. C. Barman, S. Mondal, and M. Pal. Minimum 2-tuple dominating set of permutation graphs. *J. Appl. Math. Comput.*, 43(1-2):133–150, 2013.
- [7] M. Blidia, M. Chellali, and O. Favaron. Ratios of some domination parameters in graphs and claw-free graphs. In *Graph theory in Paris*, Trends Math., pages 61–72. Birkhäuser, Basel, 2007.
- [8] B. Brešar, M. A. Henning, and S. Klavžar. On integer domination in graphs and Vizing-like problems. *Taiwanese J. Math.*, 10(5):1317–1328, 2006.
- [9] B. Brešar, M. A. Henning, and D. F. Rall. Rainbow domination in graphs. *Taiwanese J. Math.*, 12(1):213–225, 2008.
- [10] B. Brešar and T. Kraner Šumenjak. On the 2-rainbow domination in graphs. *Discrete Appl. Math.*, 155(17):2394–2400, 2007.
- [11] Y. Caro and R. Pepper. Degree sequence index strategy. *Australas. J. Combin.*, 59:1–23, 2014.
- [12] G. J. Chang, B.-J. Li, and J. Wu. Rainbow domination and related problems on strongly chordal graphs. *Discrete Appl. Math.*, 161(10-11):1395–1401, 2013.
- [13] G. J. Chang, J. Wu, and X. Zhu. Rainbow domination on trees. *Discrete Appl. Math.*, 158(1):8–12, 2010.
- [14] M. Chellali, T. W. Haynes, and S. T. Hedetniemi. Bounds on weak roman and 2-rainbow domination numbers. *Discrete Appl. Math.*, 178:27–32, 2014.
- [15] M. Chellali and N. Jafari Rad. On 2-rainbow domination and Roman domination in graphs. *Australas. J. Combin.*, 56:85–93, 2013.
- [16] M. Chellali, A. Khelladi, and F. Maffray. Exact double domination in graphs. *Discuss. Math. Graph Theory*, 25(3):291–302, 2005.
- [17] M. Chlebík and J. Chlebíková. Approximation hardness of dominating set problems in bounded degree graphs. *Inform. and Comput.*, 206(11):1264–1275, 2008.
- [18] K. Choudhary, S. Margulies, and I. V. Hicks. Integer domination of Cartesian product graphs. *Discrete Math.*, 338(7):1239–1242, 2015.
- [19] F. Cicalese, M. Milanič, and U. Vaccaro. On the approximability and exact algorithms for vector domination and related problems in graphs. *Discrete Appl. Math.*, 161(6):750–767, 2013.
- [20] E. J. Cockayne, P. A. Dreyer, Jr., S. M. Hedetniemi, and S. T. Hedetniemi. Roman domination in graphs. *Discrete Math.*, 278(1-3):11–22, 2004.
- [21] E. J. Cockayne and S. T. Hedetniemi. Towards a theory of domination in graphs. *Networks*, 7(3):247–261, 1977.

- [22] E. DeLaViña, W. Goddard, M. A. Henning, R. Pepper, and E. R. Vaughan. Bounds on the k -domination number of a graph. *Appl. Math. Lett.*, 24(6):996–998, 2011.
- [23] W. J. Desormeaux, T. W. Haynes, and L. Vaughan. Double domination in complementary prisms. *Util. Math.*, 91:131–142, 2013.
- [24] I. Dinur and D. Steurer. Analytical approach to parallel repetition. In *STOC'14—Proceedings of the 2014 ACM Symposium on Theory of Computing*, pages 624–633. ACM, New York, 2014.
- [25] G. Dobson. Worst-case analysis of greedy heuristics for integer programming with nonnegative data. *Math. Oper. Res.*, 7(4):515–531, 1982.
- [26] G. S. Domke, S. T. Hedetniemi, R. C. Laskar, and G. Fricke. Relationships between integer and fractional parameters of graphs. In *Graph theory, combinatorics, and applications, Vol. 1 (Kalamazoo, MI, 1988)*, Wiley-Intersci. Publ., pages 371–387. Wiley, New York, 1991.
- [27] P. Dorbec, B. Hartnell, and M. A. Henning. Paired versus double domination in $K_{1,r}$ -free graphs. *J. Comb. Optim.*, 27(4):688–694, 2014.
- [28] P. A. Dreyer, Jr. *Applications and variations of domination in graphs*. ProQuest LLC, Ann Arbor, MI, 2000. Thesis (Ph.D.)—Rutgers The State University of New Jersey - New Brunswick.
- [29] O. Favaron, A. Hansberg, and L. Volkmann. On k -domination and minimum degree in graphs. *J. Graph Theory*, 57(1):33–40, 2008.
- [30] U. Feige. A threshold of $\ln n$ for approximating set cover. *J. ACM*, 45(4):634–652, 1998.
- [31] H. Fernau. Roman domination: a parameterized perspective. *Int. J. Comput. Math.*, 85(1):25–38, 2008.
- [32] J. F. Fink and M. S. Jacobson. n -domination in graphs. In *Graph theory with applications to algorithms and computer science (Kalamazoo, Mich., 1984)*, Wiley-Intersci. Publ., pages 283–300. Wiley, New York, 1985.
- [33] S. Földes and P. L. Hammer. Split graphs having Dilworth number two. *Canad. J. Math.*, 29(3):666–672, 1977.
- [34] S. Fujita and M. Furuya. Difference between 2-rainbow domination and Roman domination in graphs. *Discrete Appl. Math.*, 161(6):806–812, 2013.
- [35] M. Furuya. A note on total domination and 2-rainbow domination in graphs. *Discrete Appl. Math.*, 184:229–230, 2015.
- [36] M. Gairing, S. Hedetniemi, P. Kristiansen, and A. McRae. Self-stabilizing algorithms for $\{k\}$ -domination. In S.-T. Huang and T. Herman, editors, *Self-Stabilizing Systems*, volume 2704 of *Lecture Notes in Computer Science*, pages 49–60. Springer Berlin Heidelberg, 2003.
- [37] T. Gallai. Gráfokkal kapcsolatos maximum-minimum tételek (i. rész). *Magyar Tud. Akad. Mat. Fiz. Oszt. Közl.*, 7:305–338, 1957.
- [38] T. Gallai. Gráfokkal kapcsolatos maximum-minimum tételek (ii. rész). *Magyar Tud. Akad. Mat. Fiz. Oszt. Közl.*, 8:1–40, 1958.

- [39] T. Gallai. Maximum-minimum Sätze über Graphen. *Acta Math. Acad. Sci. Hungar.*, 9:395–434, 1958.
- [40] T. Gallai. Über extreme Punkt- und Kantenmengen. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.*, 2:133–138, 1959.
- [41] M. R. Garey and D. S. Johnson. *Computers and Intractability*. W. H. Freeman and Co., San Francisco, Calif., 1979. A guide to the theory of NP-completeness, A Series of Books in the Mathematical Sciences.
- [42] A. Hansberg and R. Pepper. On k -domination and j -independence in graphs. *Discrete Appl. Math.*, 161(10-11):1472–1480, 2013.
- [43] F. Harary and T. W. Haynes. Nordhaus-Gaddum inequalities for domination in graphs. *Discrete Math.*, 155(1-3):99–105, 1996. Combinatorics (Acireale, 1992).
- [44] F. Harary and T. W. Haynes. Double domination in graphs. *Ars Combin.*, 55:201–213, 2000.
- [45] B. L. Hartnell and D. F. Rall. On dominating the Cartesian product of a graph and K_2 . *Discuss. Math. Graph Theory*, 24(3):389–402, 2004.
- [46] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, editors. *Domination in Graphs. Advanced Topics*, volume 209 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1998.
- [47] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, editors. *Fundamentals of Domination in Graphs*, volume 208 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1998.
- [48] T. W. Haynes and P. J. Slater. Paired-domination in graphs. *Networks*, 32(3):199–206, 1998.
- [49] S. M. Hedetniemi, S. T. Hedetniemi, R. C. Laskar, L. Markus, and P. J. Slater. Disjoint dominating sets in graphs. In *Discrete mathematics*, volume 7 of *Ramanujan Math. Soc. Lect. Notes Ser.*, pages 87–100. Ramanujan Math. Soc., Mysore, 2008.
- [50] S. T. Hedetniemi, R. R. Rubalcaba, P. J. Slater, and M. Walsh. Few compare to the great Roman empire. *Congr. Numer.*, 217:129–136, 2013.
- [51] M. A. Henning and S. T. Hedetniemi. Defending the Roman Empire—a new strategy. *Discrete Math.*, 266(1-3):239–251, 2003.
- [52] M. A. Henning and A. P. Kazemi. k -tuple total domination in graphs. *Discrete Appl. Math.*, 158(9):1006–1011, 2010.
- [53] M. A. Henning, C. Löwenstein, and D. Rautenbach. Remarks about disjoint dominating sets. *Discrete Math.*, 309(23-24):6451–6458, 2009.
- [54] M. A. Henning and A. Yeo. Strong transversals in hypergraphs and double total domination in graphs. *SIAM J. Discrete Math.*, 24(4):1336–1355, 2010.
- [55] M. A. Henning and A. Yeo. 2-colorings in k -regular k -uniform hypergraphs. *European J. Combin.*, 34(7):1192–1202, 2013.

- [56] M. A. Henning and A. Yeo. *Total Domination in Graphs*. Springer Monographs in Mathematics. Springer, New York, 2013.
- [57] X. Hou and Y. Lu. On the $\{k\}$ -domination number of Cartesian products of graphs. *Discrete Math.*, 309(10):3413–3419, 2009.
- [58] N. John and S. Suen. Graph products and integer domination. *Discrete Math.*, 313(3):217–224, 2013.
- [59] B. K. Jose and Z. Tuza. Hypergraph domination and strong independence. *Appl. Anal. Discrete Math.*, 3(2):347–358, 2009.
- [60] R. Klasing and C. Laforest. Hardness results and approximation algorithms of k -tuple domination in graphs. *Inform. Process. Lett.*, 89(2):75–83, 2004.
- [61] M. Krzywkowski. On trees with double domination number equal to 2-domination number plus one. *Houston J. Math.*, 39(2):427–440, 2013.
- [62] V. R. Kulli. Inverse domination and inverse total domination in digraphs. *International Journal of Advanced Research in Computer Science & Technology*, 2(1):106–109, 2014.
- [63] N. Li and X. Hou. On the total $\{k\}$ -domination number of Cartesian products of graphs. *J. Comb. Optim.*, 18(2):173–178, 2009.
- [64] C.-H. Liu and G. J. Chang. Roman domination on 2-connected graphs. *SIAM J. Discrete Math.*, 26(1):193–205, 2012.
- [65] C.-H. Liu and G. J. Chang. Roman domination on strongly chordal graphs. *J. Comb. Optim.*, 26(3):608–619, 2013.
- [66] C. Löwenstein and D. Rautenbach. Pairs of disjoint dominating sets and the minimum degree of graphs. *Graphs Combin.*, 26(3):407–424, 2010.
- [67] P. Pavlič and J. Žerovnik. Roman domination number of the Cartesian products of paths and cycles. *Electron. J. Combin.*, 19(3):Paper 19, 37, 2012.
- [68] M. Pilipczuk, M. Pilipczuk, and R. Škrekovski. Some results on Vizing’s conjecture and related problems. *Discrete Appl. Math.*, 160(16-17):2484–2490, 2012.
- [69] D. Pradhan. Algorithmic aspects of k -tuple total domination in graphs. *Inform. Process. Lett.*, 112(21):816–822, 2012.
- [70] N. J. Rad. Critical concept for 2-rainbow domination in graphs. *Australas. J. Combin.*, 51:49–60, 2011.
- [71] E. Sampathkumar and L. P. Latha. Strong weak domination and domination balance in a graph. *Discrete Math.*, 161(1-3):235–242, 1996.
- [72] A. Schrijver. *Combinatorial Optimization. Polyhedra and Efficiency (3 volumes)*, volume 24 of *Algorithms and Combinatorics*. Springer–Verlag, Berlin, 2003.

- [73] W. Shang, X. Wang, and X. Hu. Roman domination and its variants in unit disk graphs. *Discrete Math. Algorithms Appl.*, 2(1):99–105, 2010.
- [74] Z. Shao, M. Liang, C. Yin, X. Xu, P. Pavlič, and J. Žerovnik. On rainbow domination numbers of graphs. *Inform. Sci.*, 254:225–234, 2014.
- [75] I. Stewart. Defend the roman empire! *Sci. Am.*, 281:136–139, 1999.
- [76] T. K. Šumenjak, D. F. Rall, and A. Tepeh. Rainbow domination in the lexicographic product of graphs. *Discrete Appl. Math.*, 161(13-14):2133–2141, 2013.
- [77] V. V. Vazirani. *Approximation Algorithms*. Springer-Verlag, Berlin, 2001.
- [78] L. A. Wolsey. An analysis of the greedy algorithm for the submodular set covering problem. *Combinatorica*, 2(4):385–393, 1982.
- [79] Y. Wu and H. Xing. Note on 2-rainbow domination and Roman domination in graphs. *Appl. Math. Lett.*, 23(6):706–709, 2010.
- [80] B. Zelinka. Total domatic number of cacti. *Math. Slovaca*, 38(3):207–214, 1988.
- [81] B. Zelinka. Domatic numbers of graphs and their variants: a survey. In *Domination in graphs*, volume 209 of *Monogr. Textbooks Pure Appl. Math.*, pages 351–377. Dekker, New York, 1998.