

NITRILE HYDROLYSIS IN A RHENIUM(I)–RUTHENIUM(III) DINUCLEAR COMPLEX

FLORENCIA FAGALDE*, MARÍA GABRIELA MELLACE,
NOEMÍ D. LIS DE KATZ and NÉSTOR E. KATZ

*Instituto de Química Física, Facultad de Bioquímica, Química y Farmacia,
Universidad Nacional de Tucumán, Ayacucho 491,
(T4000INI) San Miguel de Tucumán, Argentina*

(Received 23 January 2003; Revised 6 October 2003; In final form 27 February 2004)

Kinetic data for the hydrolysis reaction: $[(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{4+} + \text{H}_2\text{O} \rightarrow [(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-C(O)NHpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{3+} + \text{H}^+$ (bpy = 2,2'-bipyridine, 4-CNpy = 4-cyanopyridine and 4-C(O)NHpy = isonicotinamido), that occurs after oxidation with $\text{S}_2\text{O}_8^{2-}$ of the dinuclear species $[(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{II}}(\text{NH}_3)_5]^{3+}$, have been obtained in aqueous solutions by spectrophotometric techniques. The observed rate constant, $k_h = (8.6 \pm 0.5) \times 10^{-3} \text{ s}^{-1}$, at 25°C, pH = 3.0 (CF₃COOH) and $I = 0.1 \text{ M}$ (KCl), is *ca.* three times higher than the corresponding one for the mononuclear complex $[(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{3+}$, indicating that the catalytic effect of the $\text{Ru}^{\text{III}}(\text{NH}_3)_5$ moiety is enhanced by coordination of the free N of 4-CNpy to the $\text{Re}^{\text{I}}(\text{CO})_3(\text{bpy})$ moiety. The value of k_h is even higher than that of the dinuclear complex $[(\text{trpy})(\text{bpy})\text{Ru}^{\text{II}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{5+}$ (trpy = 2,2':6',2''-terpyridine), reflecting the fact that carbonyls are much stronger π -acceptors than polypyridines.

Keywords: Nitrile hydrolysis; Mixed-valent complexes; 4-cyanopyridine; Catalysis; Dinuclear complexes

INTRODUCTION

The hydrolysis of nitriles to amides catalyzed by transition metals is a subject of considerable interest [1,2]. For some coordinated nitriles, selective hydrolysis is possible, so catalytic hydrolytic systems can be designed. Since the pioneering work by Zanella and Ford [3], ruthenium-catalyzed nitrile hydrolysis reactions have been extensively investigated. Tfouni *et al.* [2] have reviewed the chemistry of Ru(II) and Ru(III) ammine complexes with coordinated nitriles.

Recently we studied [4] the preparation and characterization of a new dinuclear complex, of formula $[(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{II}}(\text{NH}_3)_5]^{3+}$, (**I**), (with bpy = 2,2'-bipyridine and 4-CNpy = 4-cyanopyridine, coordinated to Ru through the nitrile N). The asymmetric mixed-valent species, $[(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{4+}$, (**II**),

*Corresponding author. Fax: +54-3814248169. E-mail: ffagald@unt.edu.ar

was obtained “*in situ*” by oxidation of (I) with *p*-fluorobenzene-diazonium hexafluorophosphate in CH₃CN. In order to extend our knowledge of the catalytic effect of remote metallic sites, we report in this work the hydrolysis reaction that occurs after formation of the mixed-valent complex (II) by oxidizing (I) with S₂O₈²⁻ in aqueous solutions.

EXPERIMENTAL

Syntheses

The Br⁻ salt of (I) was obtained by dissolving 50 mg of the PF₆⁻ salt, prepared as in reference [4], in 2 mL of acetone and adding 1 g of Buⁿ₄NBr, previously dissolved in 2 mL of acetone. The precipitate was collected by filtration, washed with cold acetone and dried *in vacuo* over P₄O₁₀. Yield: 67%. (Anal. Calcd. for ReRuC₁₉H₂₇N₉O₃P₃F₁₈: C, 19.8; H, 2.4; N, 10.9%. Found: C, 19.7; H, 2.7; N, 9.7%).

The mixed-valent complex (II) was obtained “*in situ*” by oxidation of (I) with S₂O₈²⁻ in aqueous solutions.

The hydrolysis product, [(CO)₃(bpy)Re^I(4-C(O)NHpy)Ru^{III}(NH₃)₅]³⁺, (III), (4-C(O)NHpy = isonicotinamido), was obtained as a PF₆⁻ salt by adding ten-fold excess of K₂S₂O₈ to (I), previously dissolved in 6 mL of 10⁻³ M CF₃COOH. After 30 min of reaction, NH₄PF₆ was added to precipitate the complex. The solid was filtered off, washed with water and dried *in vacuo* over P₄O₁₀. (Anal. Calcd. for ReRuC₁₉H₂₈N₉O₄P₃F₁₈ · 9NH₄KS₂O₈: C, 6.2; N, 7.3%. Found C, 6.2; N, 7.5%).

Materials, Instrumentation and Techniques


All chemicals were reagent grade and used as received. Double distilled water was used for all kinetic determinations. pH measurements were carried out with a precision of ±0.05 pH units, using a Mehtrom 744 pHmeter. UV-visible spectra were recorded with a Shimadzu UV-160A spectrophotometer, equipped with a thermostatted cell compartment. IR spectra were measured (as KBr pellets) using a double-beam Perkin-Elmer 983G spectrophotometer. Chemical analyses were done at INQUIMAE, University of Buenos Aires, Argentina.

The hydrolysis reactions were studied under *pseudo*-first-order conditions, at temperatures between 15 and 30°C, pH = 3 (CF₃COOH) and pH = 5 (buffer acetic acid/acetate) and *I* = 0.1 M (KCl). Absorbance (*A*) vs. time (*t*) data were recorded at λ = 470 nm. Duplicate or triplicate runs were made at each temperature and pH. Rate constants were obtained from least-squares fits of ln(*A*_{*t*} - *A*_∞) vs. *t*, which were linear up to three half-lives. The errors in rate constants were estimated to be ± 2–5%.

RESULTS AND DISCUSSION

In previous work [4], we demonstrated that 4-CNpy can act as a bridging ligand in the dinuclear species [(CO)₃(bpy)Re^I(4-CNpy)Ru^{II}(NH₃)₅]³⁺ (I), where the pyridine N (of 4-CNpy) is coordinated to a Re^I(CO)₃(bpy) group and the nitrile N (of 4-CNpy) to a Ru^{II}(NH₃)₅ moiety. In aqueous solutions, S₂O₈²⁻ rapidly oxidizes the ammine ruthenium, as studied previously [5–7]. In this work we have investigated

the hydrolysis of the oxidized cyanopyridine complex $[(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{4+}$ (**I**) by conventional spectrophotometric techniques, to give $[(\text{CO})_3(\text{bpy})\text{Re}^{\text{I}}(4\text{-C(O)NHpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{3+}$, (**III**), according to equation (1):



The identity of the product is confirmed by comparing the IR spectra of the PF_6^- salts of (**I**) and (**III**). The nitrile stretching band, which appears at 2177 cm^{-1} in (**I**), disappears completely in (**III**), while the ammonia symmetric deformation frequency, $\delta_{\text{sym}}(\text{NH}_3)$, shifts from 1282 cm^{-1} in (**I**) to 1312 cm^{-1} in (**III**), a clear indication of the oxidation state for the ammine Ru(III) center [4]. A new and intense band appears at 1405 cm^{-1} , which can be assigned to stretching vibrations of an amide ($-\text{C(O)NH}-$) group.

Figure 1 shows the consecutive spectra obtained at $\text{pH}=3.0$ (CF_3COOH), $I=0.1\text{ M}$ (KCl) and 25°C upon mixing aqueous solutions of (**I**) ($C=1.1 \times 10^{-4}\text{ M}$) and $\text{S}_2\text{O}_8^{2-}$ ($C=1.1 \times 10^{-3}\text{ M}$). During a time scale of 30 min, the oxidized cyanopyridine complex (**I**), with a shoulder at λ ca. 470 nm (deconvoluted band maximum of the metal-to-metal charge transfer (MMCT) transition $\text{Re}(\text{I}) \rightarrow \text{Ru}(\text{III})$) [4], evolved exponentially into that of amido complex (**III**) (shoulder at λ ca. 440 nm); an isosbestic point at 425 nm being obtained. The spectrum of the final product is coincident with that obtained when dissolving the solid salt (**III**) in CF_3COOH (10^{-3} M).




FIGURE 1 Visible spectra of $[(\text{bpy})(\text{CO})_3\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{II}}(\text{NH}_3)_5]^{3+}$ and $\text{S}_2\text{O}_8^{2-}$ in water at $[\text{Ru}(\text{II})]=1.1 \times 10^{-4}\text{ M}$, $I=0.1\text{ M}$ and $T=25^\circ\text{C}$. Reaction times were: 0, 90, 150, 300, 600, 900 and 1500 s from the initial (**I**) to the final (**F**) spectra.

The MMCT (or intervalence) band of (**II**) (detected at a similar wavelength in CH₃CN solution [4]) is shifted to higher energy in (**III**), as expected from the difference in redox potentials [8], but is masked by the long tail of the metal-to-ligand charge transfer (MLCT) band corresponding to a $d_{\pi}(\text{Re}) \rightarrow \pi^*(\text{bpy})$ transition.

By fixing the measuring wavelength at 470 nm, k_h values have been determined under *pseudo*-first-order conditions: $[\text{complex}] = 1.3 \times 10^{-4} \text{ M}$ and $[\text{S}_2\text{O}_8^{2-}] = 1.3 \times 10^{-3} \text{ M}$. A value of $k_h = (8.6 \pm 0.5) \times 10^{-3} \text{ s}^{-1}$ was obtained at 25°C, pH = 3.0 (CF₃COOH) and $I = 0.1 \text{ M}$ (KCl), which can be compared with other metal-catalysed hydrolysis rate constants, as shown in Table I. A similar value was obtained at pH = 5, so we conclude that the nitrile hydrolysis rate constants are pH-independent.

The value obtained for the complex studied in this work is *ca.* three times *higher* than the corresponding one for the mononuclear complex $[(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{3+}$ [8], indicating that the catalytic effect of a $\text{Ru}^{\text{III}}(\text{NH}_3)_5$ moiety is enhanced by coordination of the pyridine N of 4-CNpy to a $\text{Re}^{\text{I}}(\text{CO})_3(\text{bpy})$ moiety, probably due to an inductive effect. Moreover, the value of k_h is 1.5 times *higher* than that of the complex $[(\text{trpy})(\text{bpy})\text{Ru}^{\text{II}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{5+}$ (trpy = 2,2':6',2''-terpyridine) [5], in spite of lower charge, evidence that π -backbonding of the $\text{Re}^{\text{I}}(\text{CO})_3(\text{bpy})$ group to 4-CNpy is less than that of the $\text{Ru}^{\text{II}}(\text{trpy})(\text{bpy})$ group to 4-CNpy, a fact which can be explained on the basis that carbonyls are much stronger π -acceptors than polypyridines. However, the measured value of k_h of $[(\text{bpy})(\text{CO})_3\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{4+}$ is still *lower* (*ca.* 30 times) than that of $[(\text{NH}_3)_5\text{Ru}^{\text{III}}(4\text{-NCpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{6+}$ [6], as expected from differences in charge and π -backbonding effects.

Table II shows the value of k_h at different temperatures. From Eyring's rate equation [9], values of $\Delta H^\ddagger = 39 \pm 3 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -95 \pm 14 \text{ J mol}^{-1} \text{ K}^{-1}$ have been determined, which are consistent with activation parameters obtained for similar nitrile hydrolysis reactions and indicate an associative mechanism.

We conclude that the $\text{Re}^{\text{I}}(\text{CO})_3(\text{bpy})$ group enhances the catalytic effect of $\text{Ru}(\text{NH}_3)_5^{3+}$ on the hydrolysis of coordinated 4-CNpy by a factor of three. The measured rate constant at acidic conditions is even higher than that of the $\text{Ru}^{\text{II}}(\text{trpy})(\text{bpy})$ group, due to the fact that carbonyls are much better π -acceptors than polypyridines.

TABLE I Hydrolysis rate constants for coordinated nitriles in some ruthenium complexes

Complex	$k_h, \text{ s}^{-1}$	Ref.
$[(\text{NH}_3)_5\text{Ru}(4\text{-NCpy})]^{3+}$	2.9×10^{-3}	8
$[(\text{trpy})(\text{bpy})\text{Ru}^{\text{II}}(4\text{-NCpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{5+}$	5.8×10^{-3}	5
$[(\text{bpy})(\text{CO})_3\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{4+}$	8.6×10^{-3}	This work
$[(\text{NH}_3)_5\text{Ru}^{\text{III}}(4\text{-NCpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{6+}$	25.0×10^{-2}	6

TABLE II Hydrolysis rate constants for $[(\text{bpy})(\text{CO})_3\text{Re}^{\text{I}}(4\text{-CNpy})\text{Ru}^{\text{III}}(\text{NH}_3)_5]^{4+}$ at different temperatures

$T, \text{ K}$	$k_h, \text{ s}^{-1}$
288.2	$(4.6 \pm 0.2) \times 10^{-3}$
293.2	$(6.2 \pm 0.5) \times 10^{-3}$
298.2	$(8.6 \pm 0.5) \times 10^{-3}$
303.2	$(1.1 \pm 0.3) \times 10^{-2}$

Acknowledgments

We thank Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina), Consejo de Investigaciones de la Universidad Nacional de Tucumán (CIUNT, Argentina), Agencia Nacional de Promoción Científica y Técnica (ANPCyT, Argentina) and Fundación Antorchas for financial help. M.G.M. thanks CONICET for a graduate fellowship. F.F. and N.E.K are Members of the Research Career (CONICET).

References

- [1] Y.N. Belokon', V.I. Tararov, T.F. Savel'eva, S.V. Vitt, E.A. Paskonova, S.Ch Dotdayev, Y.A. Borisov, Y.T. Struchkov, A.S. Batasanov and V.M. Belikov, *Inorg. Chem.* **27**, 4046 (1988).
- [2] Z.N. da Rocha, G. Chiericato Jr. and E. Tfouni, *Adv. Chem. Ser. ACS* **297** (1997).
- [3] A.W. Zanella and P.C. Ford, *Inorg. Chem.* **14**, 42 (1975).
- [4] M.G. Mellace, F. Fagalde and N.E. Katz, *Polyhedron* **22**, 369 (2003).
- [5] N.D. Lis de Katz, F. Fagalde and N.E. Katz, *Polyhedron* **14**, 3111 (1995).
- [6] M.H. Chou, C. Creutz and N. Sutin, *Inorg.Chem.* **31**, 2318 (1992).
- [7] F. Fagalde, N.E. Katz, V. Povse and J.A. Olabe, *Polyhedron* **18**, 25 (1998).
- [8] M.Y. Huang, W.J. Chen, C.C. Yang and A. Yeh, *Inorg. Chem.* **30**, 1862 (1991).
- [9] R.G. Wilkins, *Kinetics and Mechanism of Reactions of Transition Metal Complexes* (VCH, Weinheim, 1991) 2nd Edn., p. 88.