

Seed Science Research

Phosphorylated 11S globulins in sunflower seeds

--Manuscript Draft--

Manuscript Number:	SSR-D-13-00031R2
Full Title:	Phosphorylated 11S globulins in sunflower seeds
Article Type:	Short Communication
Corresponding Author:	Mariana Regente ARGENTINA
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	Ivana Quiroga
First Author Secondary Information:	
Order of Authors:	Ivana Quiroga
	Mariana Regente
	Luciana Pagnussat
	Ana Maldonado
	Jesús Jorrín
	Laura de la Canal
Order of Authors Secondary Information:	
Abstract:	Helianthinins are storage proteins present in <i>Helianthus annuus</i> seeds belonging to the 11S globulin family. We describe here that a fraction of the helianthinins is phosphorylated. This conclusion is supported by different criteria including identification by MALDI-TOF of major protein bands revealed with a specific dye for phosphoproteins, anti-phosphoserine antibody and binding to a phosphoprotein affinity matrix. Moreover, we show that the phosphorylation status of helianthinins changes following germination.

Manuscript number: SSR-D-13-00031R1

Title: Phosphorylated 11S globulins in sunflower seeds

Editor's decision: Minor Revision

Dear Dr. Derek Bewley,

We appreciate the reconsideration of the MS entitled *Phosphorylated 11S globulins in sunflower seeds*, by Quiroga *et al.*

Reviewers requested a few minor revisions that have been taken into account and modified in the present version. We hope that the changes introduced will fully meet the requirements of SSR.

Kind regards,

Mariana Regente

Response to the Reviewers

Comments to Author:

Thank you for making the changes as requested. There are still a few minor revisions that need to be made, particularly in the Tables and Figure legends, but then the paper is acceptable.

Answer: All the suggestions of the reviewer have been taken into account and modified in the revised MS.

1 **Phosphorylated 11S globulins in sunflower seeds**

2

3

4 **Running head title:** Phosphorylation of sunflower storage proteins.

5

6

7 Ivana Quiroga^{†a}, Mariana Regente^{†a}, Luciana Pagnussat^a, Ana Maldonado^b, Jesús Jorrín^b, Laura de
8 la Canal^{a*}.

9

10 ^a Instituto de Investigaciones Biológicas, Universidad Nacional de Mar del Plata-CONICET, Mar
11 del Plata, Argentina.

12 ^b Departamento de Bioquímica y Biología Molecular, Universidad de Córdoba, Córdoba, España.

13

14

15 † Both authors have equally contributed to this work

16

17 * **Corresponding author**

18 Mariana Regente. Instituto de Investigaciones Biológicas, Universidad Nacional de Mar del Plata -
19 CONICET, Funes 3250, 7600 Mar del Plata, Argentina. Tel 54-223-4753030. Fax 54-223-
20 4724143. E-mail address: mregente@mdp.edu.ar

21

22

23 **Keywords**

24 *Helianthus annuus*, Germination, 11S Globulins, Helianthinin, Phosphorylation, Storage proteins.

25

26

27

28

1 **Abstract**

2

3 **Phosphorylation of sunflower storage proteins.**

4

5 Helianthinins are storage proteins present in *Helianthus annuus* seeds belonging to the 11S
6 globulin family. We describe here that a fraction of the helianthinins is phosphorylated. This
7 conclusion is supported by different criteria including identification by MALDI-TOF of major
8 protein bands revealed with a specific dye for phosphoproteins, anti-phosphoserine antibody and
9 binding to a phosphoprotein affinity matrix. Moreover, we show that the phosphorylation status of
10 helianthinins changes following germination.

11

12

1 **Introduction**

2

3 Seed storage proteins accumulate in high amounts during the late stages of seed
4 development and provide nitrogen for seed germination and early seedling growth. These proteins
5 have been classified into albumins, globulins and glutelins based on their solubility in water,
6 saline, and NaOH solutions respectively (Osborne, 1924). Helianthinin, the main storage protein of
7 sunflower seeds, belongs to the salt soluble 11S globulin fraction (Raymond *et al.*, 1995). These
8 globulins are composed by α and β subunits assembled as a hexamer (Shewry *et al.*, 1995). They
9 are synthesized as a precursor in the rough endoplasmic reticulum and follow the secretory
10 pathway to be delivered into storage vacuoles. Sequence analysis revealed that the gene
11 encoding the sunflower 11 S precursor (Helianthinin G3) contains a signal peptide responsible for
12 its targeting to the secretory pathway, followed by the coding sequence for α and β chains (Vonder
13 Haar *et al.*, 1988). In fact, the proglobulin containing the α and β polypeptides is proteolytically
14 cleaved after disulfide bond formation and transported through the secretory system as
15 intermediate trimers that further assemble into the mature hexameric form (Shewry *et al.*, 1995).
16 On the other hand, helianthinins are polymorphic and their subunit composition varies among
17 different sunflower cultivars. At least six types of subunits with molecular masses ranging from 40
18 to 64 kDa have been described so far, consisting in different combinations of the large (32-40
19 kDa) acidic α polypeptide and the small (20-25 kDa) basic β polypeptide (Rahma and Narasinga,
20 1979; Schwenke *et al.*, 1979).

21 Reversible protein phosphorylation is one of the most frequent post-translational modifications
22 and its relevance is that it regulates diverse cellular processes. Protein phosphorylation in
23 eukaryotes mostly occurs by the addition of a phosphate group on certain serine and threonine
24 residues, whereas phosphorylation on tyrosine residues is less abundant. Post-translational
25 modifications of a protein can determine its activity state, localization, turnover, and interactions
26 with other proteins (Zolnierowics and Bollen, 2000). Besides its well-known role in several
27 signaling pathways activating and deactivating enzymes, phosphorylation participates in an
28 expanding catalogue of physiological functions in plants (Kline-Jonakin *et al.*, 2011). Among them,
29 it plays a key role during embryogenesis and seed germination (Fujii and Zhu, 2009; Nakashima *et*
30 *al.*, 2009). Also, Wan *et al.* (2007) have shown that the 12S globulin cruciferin is the major
31 phosphorylated protein in *Arabidopsis thaliana* seeds. A total of 20 phosphorylation sites were
32 identified on this globulin, including serine, threonine and tyrosine residues (Wan *et al.*, 2007).

1 Although a specific function was not assigned to this post-translational modification, cruciferin
2 processing and mobilization were suggested. In this respect, Ghelis *et al.* (2008) detected changes
3 in cruciferin tyrosine-phosphorylation in response to ABA treatment, suggesting this is induced by
4 ABA to prevent cruciferin proteolysis. Agrawal and Thelen (2006) identified in a proteomic study
5 of *B. napus* phosphoproteins belonging to 10 major functional categories, including storage
6 proteins during seed filling. A recent analysis of seed maturation in *Arabidopsis*, rapeseed, and
7 soybean detected novel orthologs of seed storage proteins using a phosphoproteomic approach
8 (Meyer *et al.*, 2012).

9 Taking into account these observations, the aim of this work was to elucidate whether
10 phosphorylated storage proteins are present in sunflower, a phylogenetically distant species.

11

12

1 **Materials and Methods**

2

3 **Protein sample preparation and analysis**

4 *Helianthus annuus* L. seeds (line 10347 Advanta Semillas SAIC) were sterilized in 27.5 mg/ml
5 sodium hypochlorite for 30 min, rinsed with sterile water and subjected to imbibition for the
6 indicated times at 25° C. Petri dishes containing wet Whatman filter paper under sterile conditions
7 were used. Dried seeds, germinating seeds (16 and 24 h of imbibition) and seedlings (48, 90 and
8 96 h of imbibition) were subjected to pulverization and extracted in the following buffer: 50 mM
9 Tris-HCl pH 7.5, 2 mM DTT, 0.1 M EDTA, protease and phosphatase inhibitors (Martin *et al.*,
10 2007).

11 Protein concentration was determined by the bicinchoninic acid method (Smith *et al.*, 1985) using
12 bovine serum albumin as standard. Protein fractions (50 µg) were resolved in 12 % SDS-PAGE
13 (Laemmli, 1970) and stained with Coomassie Brilliant Blue R-250 or Colloidal Coomassie Blue
14 (Neuhoff *et al.*, 1988). Western blot assays were carried out by protein transfer onto nitrocellulose
15 membranes. Rabbit anti-phosphoserine 1:500 (Invitrogen) or sunflower anti-11S globulin
16 antibodies provided by Dr. S. Petruccelli (1:1000) were used as primary antibodies and an anti-
17 rabbit IgG coupled to alkaline phosphatase (Sigma) was used as secondary antibody (1:10000).

18

19 **2D-PAGE**

20 Homogenates from imbibed seeds (100 µg protein) were subjected to chloroform-methanol
21 precipitation and the pellet was suspended in 7 M urea, 2 M thiourea, 4 % (w/v) CHAPS, 100 mM
22 DTT, 0.2 % (v/v) ampholytes pH 3-10 (BioRad), trace of bromophenol blue. Samples were loaded
23 onto pH 3–10 nonlinear, immobilized pH gradient strips (7 cm, ReadyStrips, BioRad). Isoelectric
24 focusing was performed applying the following conditions: for the rehydration step the voltage
25 was maintained for 16 h at 50 V, and then the proteins were focused at 9000 V/h at 20°C. After
26 isoelectric focusing the strips were equilibrated in 375 mM Tris-HCl pH 8.8, 6 M urea, 20 % (v/v)
27 glycerol, 2 % (w/v) SDS, 2 % (w/v) DTT, followed by 375 mM Tris-HCl pH 8.8, 6 M urea, 20 %
28 (v/v) glycerol, 2 % (w/v) SDS, 2.5 % (w/v) iodoacetamide. Second dimension was carried out
29 according to Laemmli (1970) in 12 % gels at 150 V and PeppermintStick phosphoprotein
30 molecular mass standards (Invitrogen) were loaded in the first gel lane. After SDS-PAGE, gels
31 were fixed with 50 % (v/v) methanol and 10 % (v/v) acetic acid for 30 min, washed with distilled
32 water, stained with Pro Q Diamond phosphoprotein stain (Life Technologies) according to the

1 manufacturer's instructions and scanned with a FluorImager instrument (Amersham Biosciences)
2 using 514 nm laser as excitation source and a 570 nm band-pass emission filter. After, gels were
3 stained with SYPRO Ruby (Life Technologies) and scanned using a 488 laser as excitation source
4 and a 610 band-pass emission filter.

5

6 **MALDI-TOF**

7 Protein bands and spots indicated in the text were automatically excised from 1D and 2D gels
8 employing Investigator ProPic robotic workstation (Genomic Solutions, UK). Gel pieces were
9 digested with trypsin according to standard protocols in a ProGest station (Genomic Solutions).
10 MS analyses of peptides were performed in a 4700 Proteomics Station (Applied Biosystems,
11 USA) as previously described (Pérez-Reinado *et al.*, 2007). Protein identification was assigned by
12 comparing the obtained peptide mass fingerprinting with the non-redundant plant database, using a
13 Mascot 1.9 search engine (Matrixscience, UK) (Pinedo *et al.*, 2012).

14

15 **Phosphoprotein purification**

16 Phosphoprotein enrichment was performed employing phosphoprotein affinity columns (Qiagen)
17 according to the manufacturer's instructions and the previously described modifications adapted
18 for plant extracts (Meimoun *et al.*, 2007). Seeds were decorticated and protein was extracted using
19 25 mM MES, 1 M NaCl, 0.25 % (w/v) CHAPS, protease inhibitors and Benzonase Nuclease stock
20 solution (Phosphoprotein Purification Kit, Qiagen). Protein extracts (5 mg) were loaded onto a
21 Qiagen column and after washing with loading buffer (8 ml) bound phosphoproteins were eluted
22 with 3 ml of elution buffer (Qiagen) and submitted to MALDI TOF for identification.

23

24

1 **Results and Discussion**

2
3 As a first approach to detect putative phosphorylated proteins in sunflower seeds we performed 2D
4 SDS-PAGE of 16-h-imbibed seed protein extracts and the resulting gels were sequentially stained
5 for phosphoprotein detection (Pro-Q Diamond stain) and total protein (Sypro Ruby). Pro-Q
6 Diamond phosphoprotein stain allows in-gel detection of phosphate groups attached to tyrosine,
7 serine or threonine residues (Oh *et al.*, 2009; Chitteti and Peng, 2007; Laugesen *et al.*, 2006). Fig.
8 1B shows that a limited number of spots were clearly detected upon Pro-Q Diamond staining.
9 Among them, three spots that were consistently reproducible and could be major seed proteins
10 were further analyzed by peptide mass fingerprinting to identify them. They were digested with
11 trypsin, submitted to MALDI-TOF and putatively identified using Mascot software on the
12 Genbank database. Table 1 summarizes the data obtained. The three spots matched to the 11S
13 globulin seed storage protein G3 (Helianthinin G3) from *Helianthus annuus* (Uniprot ID P19084).
14 Also, good agreement was observed between the experimentally-determined molecular mass of
15 those spots and that predicted for β subunit of Helianthinin G3 (20.98 kDa). Helianthinin G3
16 corresponds to the precursor protein of 493 amino acids which, as described for other 11S
17 globulins, is further cleaved into two chains: the acidic chain (α subunit), and the basic chain (β
18 subunit) (Rahma and Narasinga, 1979; Schwenke *et al.*, 1979; Vonder Haar *et al.*, 1988; Shewry *et*
19 *al.*, 1995). Indeed, according to proteomic criteria all the three spots detected with the
20 phosphoprotein specific stain can be assigned as 11S globulins. The phosphorylated spots between
21 35-40 kDa correlate with major spots in Fig 1A (total proteins). One major spot (~35 kDa and acid
22 isoelectric point) was clearly visible under Pro-Q Diamond staining, whose molecular mass and
23 isoelectric point are consistent with the α subunit. Even though these spots could not be identified
24 by peptide mass fingerprinting, their features suggest that they correspond to α subunits of 11S
25 globulins.

26 Supporting the suggestion that 11S globulin subunits are phosphorylated *in vivo*, a bioinformatic
27 tool (NetPhos program, <http://www.cbs.dtu.dk/services/NetPhos/>) (Blom *et al.*, 1999) predicts 16
28 putative phosphorylation sites on the Helianthinin G3 precursor sequence. These correspond to 14
29 serines, 1 threonine and 1 tyrosine (Fig. 2), covering both α and β subunits. To experimentally
30 confirm this presumption we used a strategy based on the purification of phosphoproteins by
31 affinity chromatography; the robustness of this procedure has been reported (Laugesen *et al.*, 2006
32 Meimoun *et al.*, 2007). Total seed extracts were loaded onto a Qiagen affinity matrix and eluted

1 proteins were fractionated by SDS-PAGE followed by MALDI-TOF analysis of the major
2 proteins. Among them, four bands could be identified and all matched to the 11S globulin
3 precursor G3 (Table 2). Thus several criteria indicate that 11S globulins from imbibed sunflower
4 seeds are phosphorylated: they are detected by a phosphoprotein specific stain, they are retained in
5 a specific affinity matrix and, in both cases, their identity can be confirmed by peptide mass
6 fingerprinting.

7 In *Arabidopsis* the phosphorylation of cruciferins involves key residues allowing protein
8 assembly, processing and mobilization (Wan *et al.*, 2007), and changes in phosphorylation are
9 regulated by ABA treatment (Ghelis *et al.*, 2008). In this context, it can be hypothesized that
10 globulins can vary in their phosphorylation status according to their physiological state. Thus, we
11 have investigated whether phosphorylation of helianthinins can be regulated during seed
12 germination and seedling growth. We have compared the total seed protein profile (Fig. 3A) and
13 that of phosphoproteins detected with an anti-P-Ser antibody (Fig. 3B). As observed in Fig 3A, the
14 expected protein pattern was observed in dry seeds (Raymond *et al.*, 1995; Molina *et al.*, 2004;
15 Serre *et al.*, 2001) with major bands of around 32-40 and 20-25 kDa corresponding to 11S
16 globulin α and β subunits, respectively. In addition, there were bands of around 60 kDa that could
17 correspond to an $\alpha\beta$ subunit (Molina *et al.*, 2004). The identity of these bands was also confirmed
18 by immunoblotting because they were recognized by sunflower anti-11S globulins serum (not
19 shown). The protein pattern in dry seeds does not change during the first 24 h of imbibition, but
20 modification in the protein profile occurs thereafter (Fig. 3 A). Particularly, bands around 40 kDa
21 (α subunit) consistently decrease due to their mobilization, as previously reported (Shewry *et al.*,
22 1995). The 11S subunits appear to be differently phosphorylated during seedling growth while the
23 band of around 60 kDa retains the same phosphorylation status (Fig. 3B). Even if bands of ~22
24 kDa (β subunit) are barely detected by the anti-P-Ser antibodies, subunits around 40-kDa are
25 clearly visible from 16 to 48 h of seed imbibition while they are almost undetectable in dried seeds
26 and after 90-96 h. The absence of phosphorylated α subunits after 2 days could be attributed to
27 their degradation, since they are also undetectable by Coomassie Blue staining (Fig 3A). Dry
28 seeds show a typical pattern of helianthinins but are barely detected by anti-P-Ser.

29 In conclusion, the phosphorylation of helianthinins appears to be induced following the
30 completion of germination. Even though the basis and role of this post-translational modification
31 is unknown, our results are indicative of the necessity for future investigations to analyze the

1 relationship between storage protein phosphorylation and mobilization during post-germinative
2 growth.
3
4

1 **Acknowledgements**

2 The authors thank Dr. Silvana Petruccelli for providing the sunflower anti-11S globulins antibody.

3 This work was supported by the AECI (Agencia Española de Cooperación Internacional), AUIP

4 (Asociación Universitaria Iberoamericana de Postgrado) and additional grants from CONICET and

5 the University of Mar del Plata (Argentina).

6

1 **References**

2
3 **Agrawal, G.K., Thelen, J.J.** (2006) Large-scale identification and quantitative profiling of
4 phosphoproteins expressed during seed filling in oilseed rape. *Molecular and Cellular Proteomics*
5 **5**, 2044-2059.

6
7 **Blom, N., Gammeltoft, S., Brunak, S.** (1999) Sequence- and structure-based prediction of
8 eukaryotic protein phosphorylation sites. *Journal of Molecular Biology* **294**, 1351-1362.

9 **Chitteti, B.R., Peng, Z.** (2007) Proteome and phosphoproteome differential expression under
10 salinity stress in rice (*Oryza sativa*) roots. *Journal of Proteome Research* **6**, 1718-1727.

11 **Fujii, H., Zhu, J-K.** (2009) *Arabidopsis* mutant deficient in 3 abscisic acid-activated protein
12 kinases reveals critical roles in growth, reproduction, and stress. *Proceedings of the National*
13 *Academy of Sciences, USA* **106**, 8380-8385.

14 **Ghelis, T., Bolbach, G., Clodic, G., Habricot, Y., Miginiac, E., Sotta, B., Jeannette, E.** (2008)
15 Protein tyrosine kinases and protein tyrosine phosphatases are involved in abscisic acid-dependent
16 processes in *Arabidopsis* seeds and suspension cells. *Plant Physiology* **148**, 1668-80.

17 **Kline-Jonakin, K.G., Barrett-Wilt, G.A., Sussman, M.R.** (2011) Quantitative plant
18 phosphoproteomics. *Current Opinion in Plant Biology* **14**, 507-11.

19 **Laemmli, U.K.** (1970). Cleavage of structural proteins during the assembly of the head of
20 bacteriophage T4. *Nature* **227**, 680-685.

21 **Laugesen, S., Messinese, E., Hem, S., Pichereaux, C., Grat, S., Ranjeva, R., Rossignol, M.,**
22 **Bono, J.J.** (2006) Phosphoproteins analysis in plants: a proteomic approach. *Phytochemistry* **67**,
23 2208-14.

24
25 **Martin, M., Espinosa Vidal, E., de la Canal, L.** (2007) Expression of a lipid transfer protein in
26 *Escherichia coli* and its phosphorylation by a membrane-bound calcium-dependent protein kinase.
27 *Protein and Peptide Letters* **14**, 793-799.

28

- 1 **Meimoun, P., Ambard-Bretteville, F., Colas-des Francs-Small, C., Valot, B., Vidal, J.** (2007).
2 Analysis of plant phosphoproteins. *Analytical Biochemistry* **371**, 238-246.
3
- 4 **Meyer, L.J., Gao, J., Xu, D., Thelen, J.J.** (2012) Phosphoproteomic analysis of seed maturation
5 in *Arabidopsis*, rapeseed, and soybean. *Plant Physiology* **159**, 517–528.
6
- 7 **Molina, M.I., Petruccelli, S., Añón, M.C.** (2004). Effect of pH and ionic strength modifications
8 on thermal denaturation of the 11S globulin of sunflower (*Helianthus annuus*). *Journal of*
9 *Agricultural and Food Chemistry* **54**, 6023-6029.
10
- 11 **Nakashima, K., Fujita, Y., Kanamori, N., Katagiri, T., Umezawa, T., Kidokoro, S.,**
12 **Maruyama, K., Yoshida, T., Ishiyama, K., Kobayashi, M., Shinozaki, K., Yamaguchi-**
13 **Shinozaki, K.** (2009) Three *Arabidopsis* SnRK2 protein kinases, SRK2D/SnRK2.2,
14 SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3, involved in ABA signaling are essential for the
15 control of seed development and dormancy. *Plant and Cell Physiology* **50**, 1345-1363.
16
- 17 **Neuhoff, V., Arold, N., Taube, D., Ehrhardt, W.** (1988). Improved staining of proteins in
18 polyacrylamide gels including isoelectric focusing gels with clear background at nanogram
19 sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis* **9**, 255-262.
20
- 21 **Oh, M.H., Wang, X., Kota, U., Goshe, M., Clouse, S., Huber, S.** (2009) Tyrosine
22 phosphorylation of the BRI1 receptor kinase emerges as a component of brassinosteroid signaling
23 in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **106**, 658-663.
24
- 25 **Osborne, T.B.** (1924) The vegetable proteins. London, Longmans, Green and Co.
- 26 **Pérez-Reinado, E., Ramírez-Boo, M., Garrido, J.J., Jorrín, J.V., Moreno, A.** (2007) Towards
27 a global analysis of porcine alveolar macrophages proteins through two-dimensional
28 electrophoresis and mass spectrometry. *Developmental and Comparative Immunology* **31**, 1220-32.
- 29 **Pinedo, M., Regente, M., Elizalde, M., Quiroga, I., Pagnussat, L., Jorrín-Novo, J.,**
30 **Maldonado, A., de la Canal, L.** (2012) Extracellular sunflower proteins: Evidence on non-
31 classical secretion of a jacalin-related lectin. *Protein and Peptide Letters* **19**, 270-276.

- 1 **Rahma, E.H., Narasinga Rao, M.S.** (1979) Characterization of sunflower proteins. *Journal of*
2 *Food Science* **44**, 579-582.
3
- 4 **Raymond, J., Robin, J.M., Azanza, J.L.** (1995). 11S seed storage proteins from *Helianthus*
5 species (*Compositae*): biochemical, size and charge heterogeneity. *Plant Systematics and*
6 *Evolution* **198**, 195-208.
7
- 8 **Schwenke, K.D., Pahtz, W.L.K.J., Schultz, M.** (1979) On oil seed proteins Part II. Purification,
9 chemical composition and some physico-chemical properties of the 11S globulin (Helinthinin) in
10 sunflower seed. *Nahrung* **23**, 241-254.
11
- 12 **Serre, M., Feingold, S., Salaberry, T., Leon, A., Berry, S.** (2001). The genetic map position of
13 the locus encoding the 2S albumin seed storage proteins in cultivated sunflower (*Helianthus*
14 *annuus* L.). *Euphytica* **121**, 273-278.
15
- 16 **Shewry, P.R., Napier, J.A., Tatham, A.S.** (1995). Seed storage proteins: Structure and
17 biosynthesis. *The Plant Cell* **7**, 945-956.
18
- 19 **Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K.** (1985). Measurement of protein using
20 bicinchoninic acid. *Analytical Biochemistry* **150**, 76-85.
21
- 22 **Vonder Haar, R.A., Allen, R.D., Cohen, E.A., Nessler, C.L., Thomas, T.L.** (1988)
23 Organization of the sunflower 11S storage protein gene family. *Gene* **74**, 433-43.
24
- 25 **Wan, I., Ross, A., Yang, J., Hegedus, D., Kermode, A.** (2007). Phosphorylation of the 12S
26 globulin cruciferin in wild-type and *abil-1* mutant *Arabidopsis thaliana* (thale cress) seeds.
27 *Biochemical Journal* **404**, 247-256.
28
- 29 **Zolnierowics, S., Bollen, M.** (2000). Protein phosphorylation and protein phosphatases. *EMBO*
30 *Journal* **19**, 483-488.
31

1 **Figure legends**

2

3 **Figure 1**

4 2D gel electrophoresis of protein extracts from 16-h-imbibed sunflower seeds. (A) Sypro Ruby
5 staining, (B) Pro-Q diamond staining. Location of phosphorylated molecular mass standards in
6 kDa is indicated on the left. Circles indicate the spots identified by MALDI-TOF peptide mass
7 fingerprinting.

8

9 **Figure 2**

10 Amino acid sequence of 11S globulin seed storage protein G3 from *Helianthus annuus* (UniProt
11 accession number P19084). Putative phosphorylated residues were predicted using the NetPhos
12 2.0 program and appear underlined in the sequence. (S: 14, T: 1, Y: 1). The signal peptide is
13 shown in dark gray boxes; α and β subunits are presented in gray and white boxes respectively.

14

15 **Figure 3**

16 Phosphoserine immunodetection of sunflower seed proteins at different times of germination and
17 seedling growth. (A) 50 μ g of total proteins extracted from dry seeds and after 16, 24, 48, 90 and
18 96 h of imbibition were fractionated by SDS-PAGE and stained with Coomassie Blue. (B)
19 Immunoblotting with anti-phosphoserine antibody of the same samples detailed in (A). Figure
20 shows a representative experiment of three replicates. Molecular mass in kDa is indicated on the
21 left.

22

23

24

25

26

27

28

29

30

31

32

Figure 1





Figure 2

MASKATLLLAFTLLFATCIARHQQRQQQQNQCQLQNI¹EAL²EPI³EVI⁴QAEAGVTEIWDAYD
QQFQCAWSILFDTGFNLVAFSCLPTSTPLFWPSSREGVILPGCRRTYEYSQEQQFSGEGG
RRGGGEGTFRTVIRKLENLKEGDVVAIPTGTAHWLHNDGNT⁵ELVVVFLDTQNHENQLDEN
QRRFFLAGNPQAQAQSQQQQQRQPRQQSPQRQRQRQRQGQGQ⁶NAGNIFNGFTPELIAQSE
NVDQETAQKLQGQNDQRGHI⁷VNVGQDLQIVRPPQDRRS⁸PRQQQE⁹QATS¹⁰PRQQQEQQQGRR
GGWSNGVEETICSMKFKVNI¹¹DNPSQAD¹²FVNPQAGSIANLNSFKFPILEHLRLSVERGELR
PNAIQSPHWTINAHNLLYVTEGALRVQIVDNQGNSVFDNELREGQVVVIPQNFAVIK¹³RAN
EQGSRWVSFKTNDNAMIANLAGRVSASAASPLTLWANRYQLSREEAQQLKFSQRET¹⁴VLFA
PSFSRGQIRASR

Figure 3

[Click here to download high resolution image](#)

Figure 3



1 **Table 1**

2

3 Proteins identified in extracts from 16-h-imbibed sunflower seeds by peptide mass
4 fingerprinting.

5

6

Spot	Protein identification	Species	Accession number UniProt	Molecular mass Exper/Theor (kDa)	Peptides matched	Coverage (%)	Score
1	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	19.97/20.98 (β)	3	5.9	154
2	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	19.97/20.98 (β)	15	27.6	113
3	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	19.46/20.98 (β)	13	21.3	86

7

8

9

10

11

12

13

14

15

16

17

18

19

20

1 **Table 2**

2

3 Proteins identified in the eluted fraction of phosphoprotein affinity chromatography from seed
4 extracts by peptide mass fingerprinting

5

6

7

Band	Protein identification	Species	Accession number UniProt	Molecular mass Exper/Theor (kDa)	Peptides matched	Coverage (%)	Score
1	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	29/20.98 (β)	10	18.5	34
2	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	25/20.98 (β)	5	9.1	110
3	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	24/20.98 (β)	18	26.4	290
4	11S globulin seed storage protein G3 precursor (α - β)	<i>Helianthus annuus</i>	P19084	22/20.98 (β)	8	15	202

8

9

10

11