

The “Living Fossil” Peccary, *Catagonus wagneri* (Tayassuidae), and Its Climatic Significance during the Pleistocene and Holocene

Germán M. Gasparini, Esteban Soibelzon, Eduardo P. Tonni, and Martín Ubilla

► **Keywords:** *Catagonus*, Tayassuidae, paleoenvironments, Pleistocene, South America

The Tayassuidae first expanded their range into North America from Eurasia and then extended into South America during the “Great American Biotic Interchange,” becoming one of the first North American mammalian immigrants. Three genera are recognized in South America: *Platygonus* Le Conte (middle Pliocene to early Pleistocene), *Catagonus* Ameghino (late Pliocene? early Pleistocene to Recent), and *Tayassu* Fischer (middle Pleistocene to Recent; see Gasparini et al. 2009, 2010).


Catagonus wagneri was assigned to the genus *Catagonus* Ameghino by Wetzel et al. (1975); it is commonly known as Chacoan peccary, *taguá* or *chancho quimilero*. It has the most restricted geographical distribution among extant Tayassuidae and inhabits semiarid thorny forests of Dry Chaco in western Paraguay, southeastern Bolivia and northern Argentina (Mayer and Wetzel 1986; Gasparini et al. 2006; see Figure 1). The genus *Catagonus* was known from early- to middle-Pleistocene deposits in Buenos Aires City, Argentina (the extinct species *C. metropolitanus*). The species *C. wagneri* was reported by Rusconi (1930) in archaeological sites from Santiago del Estero Province, Argentina (ca. 1000 RCYBP; see Tonni 2006); it was believed to be extinct until it was described by Wetzel et al. (1975). Recently a partial skull from late Pleistocene of Uruguay (Cuareim River, Artigas Department) was assigned to *C. wagneri* (Gasparini et al. 2009, 2010). This material comes from the Sopas Formation (>45,000 RCYBP; $43,500 \pm 3600$ and $58,300 \pm 7400$ CALYBP [TL dates]; see Ubilla 2004). This is the oldest fossil record of this species and shows that the species ranged out of its present distribution during late Pleistocene.

Chacoan peccary is more like the extinct *Platygonus* than the other living peccaries. Certain anatomical features of *C. wagneri* are linked with a cursorial life in open and arid or semiarid environments (e.g., elongated limbs, a reduction of the lateral digits, and a great development of the sinuses and nasal chambers; see Wetzel 1977).

Germán M. Gasparini, Esteban Soibelzon, and Eduardo P. Tonni, División Paleontología Vertebrados, Museo de La Plata, Facultad de Ciencias Naturales y Museo, UNLP, Paseo del Bosque s/n, B1900FWA La Plata, Argentina; emails: germanmgasparini@gmail.com esoibelzon@fcnym.unlp.edu.ar eptonni@fcnym.unlp.edu.ar

Martín Ubilla, Facultad de Ciencias, Iguá 4225, 11400, Montevideo, Uruguay; email: ubilla@fcien.edu.uy

Figure 1. Archaeological and paleontological sites where fossil remains of *C. wagneri* were found, and current range: **1**, Llajta Mañica archaeological site, 15 km northwest of Melero, Santiago del Estero Province; **2**, Tulip-Loman archaeological site, near Icano, Santiago del Estero Province, 45km south of Llajta Mañica; **3**, paleontological site río Cuareim, Artigas Department, Uruguay; **A**, Mariscal Estigarribia, Boquerón Department, Paraguay; **B**, Las Lomitas, Formosa Province, Argentina. Gray area indicates current range of *C. wagneri*.



At present, *C. wagneri* inhabits areas of rainfall between 800 mm (Mariscal Estigarribia, Paraguay) and 80 mm (Las Lomitas, Argentina)—concentrated in summer months—and high temperatures (mean annual temperature over 24°C). During the late Pleistocene (Uruguay) and the late Holocene (Argentina), *C. wagneri* is recorded in association with mammals that also indicate arid or semiarid conditions (e.g., *Lama guanicoe*, *Myrmecophaga tridactyla*) linked to chacoan vegetation.

During the arid phases of the Pleistocene and Holocene, *C. wagneri* extended its geographic range; during the mainly humid phases—similar to the present ones—it has survived in a scrub-thorn refugium.

The authors thank the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional de La Plata, ANPCyT, and CIC-PBA, for financial support.

References Cited

- Gasparini, G. M., E. Ortiz Jaureguizar, and A. A. Carlini 2006 Familia Tayassuidae. In *Los Mamíferos de Argentina: Sistemática y distribución*, edited by R. M. Bárbquez, M. M. Díaz, and R. A. Ojeda, pp. 114–15. Publicación Especial de la Sociedad Argentina para el Estudio de los Mamíferos (SAREM).
- Gasparini, G. M., M. Ubilla, and E. P. Tonni 2009 Tres especies de tayassuidos (*Catagonus wagneri*, *C. stenocephalus* y *Tayassu pecari*) en el Pleistoceno tardío del norte de Uruguay (Fm. Sopas). *Reunión Anual de Comunicaciones de la Asociación Paleontológica Argentina*:47.
- Gasparini, G.M., M. Ubilla, and E. P. Tonni 2010 The Chacoan Peccary, *Catagonus wagneri* (Mammalia, Artiodactyla) in the Late Pleistocene of Northern Uruguay (South America). *The Third International Palaeontological Congress, IPC3*:123.
- Mayer, J. J., and R. M. Wetzel 1986 *Catagonus wagneri*. *Mammalian Species* 259:1–5.
- Rusconi, C. 1930 Las especies fósiles argentinas de pecaríes y sus relaciones con las del Brasil y Norteamérica. *Anales del Museo Nacional de Historia Natural "Bernardino Rivadavia"*, 36:121–241.
- Tonni, E. P. 2006 Cambio climático en el Holoceno tardío de la Argentina. Una síntesis con énfasis en los últimos 1000 años. *Folia Histórica del Nordeste* 16:187–95.

- Ubilla, M. 2004 Mammalian Biostratigraphy of Pleistocene Fluvial Deposits in Northern Uruguay, South America. *Proceedings of the Geologists Association* 115:347–57.
- Wetzel, R. M. 1977 The Chacoan Peccary, *Catagonus wagneri* (Rusconi). *Bulletin of the Carnegie Museum of Natural History* 3:1–36.
- Wetzel, R. M., R. E. Dubois, R. L. Martin, and P. Myers 1975 *Catagonus* an “Extinct” Peccary, Alive in Paraguay. *Science* 189:379–81.