

Ligninolytic Ability and Potential Biotechnology Applications of the South American Fungus *Pleurotus laciniatocrenatus*

M.C.N. SAPARRAT^a, F. GUILLÉN^b

^aInstituto de Botánica Spegazzini, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata,
1900 La Plata, Argentina
e-mail masaparrat@yahoo.com.ar

^bMicrobiología Molecular, Centro de Investigaciones Biológicas, Consejo Superior de Investigaciones Científicas,
28040 Madrid, Spain

Received 22 June 2004

Revised version 21 December 2004

ABSTRACT. The extracellular ligninolytic enzyme system of *Pleurotus laciniatocrenatus*, grown under different culture conditions, was characterized and the ability of this strain to degrade different components of *Eucalyptus globulus* wood was determined. In shaken liquid cultures grown on a C-limited medium supplemented with yeast extract (0.1 %) and peptone (0.5 %), the fungus produced extracellular aryl-alcohol oxidase (Aao), laccase (Lac), manganese-dependent peroxidase (MnP) and manganese-independent peroxidase (MiP) activities, their maximum levels being, respectively, about 600, 50, 1360, and 920 pkat/mL. The supplementation of 1 mmol/L vanillic acid and 150 µmol/L CuSO₄ produced an increase of Lac activity levels up to 4-fold and 68.3-fold, respectively. No significant differences were found in the levels of the other ligninolytic enzyme activities when compared to the basal medium. Solid-state fermentation cultures on *E. globulus* wood chips revealed Lac and MiP activities. These cultures showed degradative activity on lignin and lipophilic wood extractives.

Biodiversity, defined as the variety and richness of life forms on Earth, includes genetic pools, species, ecosystems and the ecological processes of which they are a part (Hawsworth 2001). Even though South American biodiversity is well known, this knowledge is mainly based on the study of vascular plants and vertebrates. At the moment, there is no comprehensive study on the wood-rotting fungi from South America, which play a fundamental role in nutrient cycling and are potential tools for biotechnological applications. However, the available evidence suggests that the fungal diversity in this continent is high (Hawsworth 2001; Ryvarden 1996). Therefore, a contribution to the knowledge of fungal diversity from South America would be important for building a better theoretical background, as well as for applied uses in solving real-world problems. In this sense, several white-rot fungi and their extracellular ligninolytic enzymes are currently being studied (Abdullah and Zafar 1999; Cing *et al.* 2003; Johannes and Majcherczyk 2000; Kahraman and Yeşilada 2003; Martínez *et al.* 1994, 1999; Saparrat *et al.* 2002; Tomšovský and Homolka 2003; Wong and Yu 1999).

The genus *Pleurotus* is well known for the production of edible mushrooms and includes strongly ligninolytic species. Different *Pleurotus* species are being extensively studied due to their ability to degrade lignin and other aromatic pollutants causing environmental problems (Cohen *et al.* 2002; Eichlerová *et al.* 2003; Kahraman and Yeşilada 2003; Martínez *et al.* 1994; Novotný *et al.* 1999; Rodríguez *et al.* 2004). These fungi have developed a nonspecific oxidative system to degrade aromatic recalcitrant compounds, including extracellular oxidases and peroxidases, low-molar-mass metabolites and activated oxygen species (Sannia *et al.* 1986, 1991; Bourbonnais and Paice 1989; Guillén *et al.* 1994, 2000; Martínez *et al.* 1996; Muñoz *et al.* 1997; Giardina *et al.* 2000).

Pleurotus laciniatocrenatus (SPEG.) SPEG. (Agaricales, Basidiomycetes) is a South American fungus, largely found in warm and temperate regions of Argentina, Brazil, Paraguay and Uruguay, which has been relatively little studied compared to other *Pleurotus* species. *P. laciniatocrenatus* has shown capability to colonize different lignocellulosic substrates (Salusso and Moraña 1997) and ABTS-oxidizing activity in solid medium (Saparrat *et al.* 2000). However, no information about the enzyme system of this fungal species has been reported. The aim of our work was to provide knowledge on the enzyme composition of the ligninolytic system of *P. laciniatocrenatus* (CLPS 39 strain), grown under different culture conditions, as well as its ability to degrade different components from *Eucalyptus globulus* LABILL. wood. This approach

would contribute to explore autochthonous fungal species for biotechnological processes and increase our knowledge about the enzyme characteristics and ecophysiology of South American fungi.

MATERIALS AND METHODS

Fungal strain. *Pleurotus laciniatocrenatus* CLPS 39 (Culture Collection of the La Plata Spengazzini Institute, Argentina) was isolated from the tissue of a fruiting-body collected from the trunk of a *Taxodium* sp. living tree in an urban forest area (Parque Pereyra woodland; Buenos Aires Province, Argentina). Stock cultures were maintained on slants containing malt extract agar supplemented with 0.4 % yeast extract and *Populus nigra* L. wood chips at 4 °C.

Liquid cultures. The production of extracellular ligninolytic enzymes was carried out on a basal C-limited medium supplemented with 0.1 % yeast extract and 0.5 % peptone (Saparrat *et al.* 2002). The effect of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (150 $\mu\text{mol/L}$) and vanillic acid (1 mmol/L) on the extracellular ligninolytic enzyme system was tested by adding them on the 3rd day of incubation to cultures grown on basal medium. The strain was cultivated in 1-L Erlenmeyer flasks with 200 mL medium, under shaking (2.5 Hz) at 25 ± 1.5 °C; a mycelial suspension was used as inoculum (Saparrat *et al.* 2002). The cultures were done in three replicates. Aliquots of 5 mL were taken periodically from each replicate flask. The mycelium was separated from the culture liquid by centrifugation (10 000 g, 4 °C, 30 min). The supernatant of the liquid culture was collected to measure reducing sugars, protein, pH and enzyme activity.

Solid-state fermentation (SSF). The CLPS 39 strain was cultivated in 100-mL Erlenmeyer flasks containing 2 g (dry mass) of *E. globulus* wood chips ($1-2 \times 10-20$ mm) and 5 mL water (sterilized at 121 °C for 2 periods of 30 min) that were inoculated with two mycelium plugs (diameter 5 mm) from cultures on 2 % malt extract agar. The inoculated flasks were incubated at 28 ± 1.5 °C under stationary conditions and constant humidity. Six replicates were done in parallel. Uninoculated sterilized flasks containing wood were incubated under the same conditions as control. After 15 and 30 d, three replicate flasks were analyzed for the presence of ligninolytic enzyme activity (expressed as pkat/g wood). The extraction of enzymes from wood treated with the fungus was carried out by adding 10 mL of 50 mmol/L sodium tartrate buffer (pH 5) to SSF cultures and shaking (0.83 Hz, 2 h, 4 °C); the enzyme extracts were obtained by filtration and dialysis against the same extraction buffer. Simultaneously, the other three cultures were analyzed in order to estimate wood mass loss and the degradation of Klason lignin and acetone extractives of eucalypt wood treated with the fungus.

Analytical methods. Extracellular protein was determined according to Bradford (1976), using *Bio-Rad* protein assay and bovine serum albumin as standard. Reducing sugars were assayed by the Somogyi and Nelson method (Somogyi 1945), using glucose as standard. Dry mass of wood was determined at 60 ± 5 °C in an aerated oven for 12 h. Lipophilic wood-acetone extract compounds were obtained *via* Soxhlet extraction of sawdust (<0.4 mm), from control wood and from wood treated with the fungus, with acetone for 6 h (Martínez *et al.* 1999). The acetone extracts were evaporated to dryness for mass estimation. Klason lignin content was determined according to TAPPI (1993). The determinations were done in triplicates.


Enzyme assays. Aryl-alcohol oxidase (EC 1.1.3.7; Aao) activity was estimated through veratraldehyde formation from 5 mmol/L veratryl alcohol (*Fluka*) in 0.1 mol/L phosphate buffer, pH 6 (Guillén *et al.* 1994). Laccase (EC 1.10.3.2; Lac) activity was measured using 5 mmol/L 2,6-dimethoxyphenol (*Fluka*) in 0.1 mol/L sodium tartrate buffer, pH 3 (Martínez *et al.* 1996). Lignin peroxidase (EC 1.11.1.14; LiP) activity was determined by H_2O_2 -dependent veratraldehyde formation from 2 mmol/L veratryl alcohol in 0.1 mol/L sodium tartrate buffer, pH 3; all reactions were started by the addition of 0.4 mmol/L H_2O_2 (Saparrat *et al.* 2002). Manganese-dependent peroxidase (EC 1.11.1.13; MnP) activity was estimated by the formation of Mn^{3+} -tartrate complex during the oxidation of 0.1 mmol/L Mn^{2+} (as MnSO_4) in 0.1 mol/L sodium tartrate buffer (pH 5) in the presence of 0.1 mmol/L H_2O_2 (Martínez *et al.* 1996). Manganese-independent peroxidase (EC 1.11.1.7, MiP) activity was assayed as Lac activity in the presence of 0.1 mmol/L H_2O_2 . All the oxidation rates were determined at 25 °C, using a *Beckman* DU 640 UV-VIS spectrophotometer. The enzyme activity is expressed in pkat (picokatal; 1 pkat is the enzyme activity releasing 1 pmol of oxidized product per s).

Statistical analysis. The effect of vanillic acid and CuSO_4 on extracellular ligninolytic enzyme activity levels, compared to cultures grown on basal medium, was analyzed statistically using one-way ANOVA with Tukey's honestly significant difference contrasts.

RESULTS AND DISCUSSION

P. laciniatocrenatus grown under shaking in a C-limited medium with 0.1 % yeast extract and 0.5 % peptone was found to be in trophophase until the 6th day of incubation, exhibiting an idiophase under C-limited conditions during the rest of the incubation (Fig. 1). A slight increase in the initial pH (5.8) was observed during the trophophase, followed by an acidification at the beginning of idiophase and a further increment of pH to values close to neutrality. Extracellular proteins exhibited a pattern of net consumption during the trophophase, followed by a production peak ($121 \pm 9.6 \mu\text{g/mL}$) after 9 d. Shaken cultures of *P. laciniatocrenatus* on C-limited peptone–yeast extract medium revealed different ligninolytic enzyme activities. Peptone has been reported to contain peptides that induce the secretion of peroxidases in *Pleurotus* species and other basidiomycetes (Kaal *et al.* 1993; Martínez *et al.* 1996). Whereas MnP activity with maximum levels $\approx 1360 \text{ pkat/mL}$ was detected during the trophophase, Aao activity, which attained a peak of 600 pkat/mL , was only detected after glucose depletion. However, Lac activity, which revealed low levels (50 pkat/mL), and MiP activity, whose maximum was about 920 pkat/mL , were detected in relation to both the trophophase and under C-limited conditions. No LiP activity was detected in the supernatant of liquid cultures under our working conditions. In this sense, it is necessary to remark that a Mn^{2+} -oxidizing peroxid-

Fig. 1. Time course of enzyme activities (nkat/mL; Aao, open squares; Lac, circles; MiP, closed squares; MnP, triangles) and reducing sugars (mmol/L; closed triangles) in the extracellular fluid of shaken *Pleurotus laciniatocrenatus* cultures grown in C-limited peptone–yeast extract medium; means of three replicates \pm SD.



ase, sharing catalytic properties with LiP and MnP, was first reported for *P. eryngii* (Martínez *et al.* 1996). This enzyme, called afterwards versatile peroxidase (VP) and reported in other *Pleurotus* and *Bjerkandera* species (Camarero *et al.* 1996; Sarkar *et al.* 1997; Heinfling *et al.* 1998; Mester and Field 1998), showed a high activity on Mn^{2+} but also on phenolic and nonphenolic aromatic compounds without Mn^{2+} . The MnP and MiP activities detected in *P. laciniatocrenatus* could correspond to the VP enzyme. However, the differences in the production pattern of MnP and MiP in this fungus suggest that the MnP activity could be related to the Mn^{2+} -oxidizing enzyme reported for other *Pleurotus* species, whereas the MiP activity could correspond to a different type of peroxidase in these fungi. Such activity had been reported only in *Bjerkandera* sp. (de Jong *et al.* 1992).

The addition of CuSO_4 or vanillic acid to the basal medium did not produce significant differences in the patterns and levels of extracellular Aao and MnP activities, and in the pH, proteins and reducing sugars under our experimental conditions. However, the presence of both compounds affected the Lac and MiP activities (Table I). Lac activity was significantly increased by the addition of vanillic acid to a maximum level of 200 pkat/mL on the 5th day ($p \leq 0.05$). This value represents a 4-fold increase, compared to cultures grown on basal medium. The increase in Lac activity levels by this compound was previously reported for other white-rot basidiomycetes, including *Pleurotus* species (Hatakka 1994; Muñoz *et al.* 1997; Farnet *et al.* 1999). The Lac activity of *P. laciniatocrenatus* was also increased by the addition of CuSO_4 , showing a peak titer after 14 d of incubation (3420 pkat/mL , $p \leq 0.01$). In this case, the resulting activity level was 68 times higher than that found in the basal medium. A strong induction of Lac activity by copper had been also described in the basidiomycetes *Corioloopsis rigida*, *Phanerochaete chrysosporium*, *Pleurotus ostreatus* and *Trametes versicolor* (Collins and Dobson 1997; Dittmer *et al.* 1997; Palmieri *et al.* 2000; Saparrat *et al.* 2002). No significant differences were found in MiP activity levels in the supplemented cultures in relation to control ones, although the presence of CuSO_4 revealed the maximum activity earlier than in the basal medium, which was significantly different from the results obtained with vanillic acid ($p \leq 0.05$). Although no increase of peroxidase activity has been found in *P. laciniatocrenatus* cultures with the above

compounds, it has been reported that they are effective inducers of these enzymes in other basidiomycetes (Rogalski *et al.* 1991; Hatakka 1994; Schlosser *et al.* 1997; Levin *et al.* 2002).

Table I. Effect of 150 $\mu\text{mol/L}$ CuSO_4 (Cu) and 1 mmol/L vanillic acid (Va) on the levels of extracellular Lac and MiP activities (pkat/mL; mean \pm SD of three replicates) in liquid cultures of *P. laciniatocrenatus* grown on a C-limited peptone–yeast extract medium.

Culture time d	Lac activity			MiP activity		
	control	+ Cu	+ Va	control	+ Cu	+ Va
4	20 \pm 0	50 \pm 30	180 \pm 20	170 \pm 240	290 \pm 230	240 \pm 0
5	40 \pm 10	160 \pm 60	200 \pm 10	320 \pm 230	820 \pm 220	360 \pm 310
6	20 \pm 20	670 \pm 220	40 \pm 10	40 \pm 60	360 \pm 40	0 \pm 10
8	30 \pm 0	1300 \pm 350	30 \pm 10	60 \pm 90	300 \pm 140	10 \pm 10
10	30 \pm 10	1480 \pm 570	30 \pm 10	250 \pm 360	570 \pm 180	140 \pm 70
12	40 \pm 10	2910 \pm 470	20 \pm 10	500 \pm 330	460 \pm 230	330 \pm 120
15	50 \pm 0	3420 \pm 680	30 \pm 10	810 \pm 200	330 \pm 120	1300 \pm 70
18	50 \pm 10	3120 \pm 100	40 \pm 0	920 \pm 270	260 \pm 110	1250 \pm 70
20	30 \pm 0	3040 \pm 110	30 \pm 10	870 \pm 210	180 \pm 130	1200 \pm 160

The existence of different ligninolytic enzymes depends on the fungal species, though it has been reported that differences in the enzyme expression could be a consequence of the culture conditions (Rogalski *et al.* 1991; Hatakka 1994; Ruiz-Dueñas *et al.* 1999). The ligninolytic enzymes of *P. laciniatocrenatus* on eucalypt wood SSF cultures, as well as the degradation of lignin and lipophilic wood extracts, are shown in Table II. This wood was chosen since it is currently used as raw material for paper pulp manufacturing in certain Latin American countries, *e.g.*, Brazil, Argentina and Uruguay. In spite of the inducing effect of lignin and/or lignocellulose on the activity and expression of the ligninolytic enzyme system in different basidiomycetes (Rogalski *et al.* 1991; Muñoz *et al.* 1997; Schlosser *et al.* 1997), only extracellular Lac and MiP activities were found in *P. laciniatocrenatus* SSF enzyme extracts. Similar results were reported for other *Pleurotus* species under similar conditions (Rodríguez *et al.* 2004). In *P. laciniatocrenatus* SSF cultures the mass loss of eucalypt wood was also determined, as well as the degradation of lignin and acetone wood extractives after 15 and 30 d of incubation (Table II). The low mass loss of the wood treated with *P. laciniatocrenatus* is a prerequisite for wood pretreatment in paper pulp manufacture (to avoid yield loss in the process). The percentage of lignin and acetone wood extractive degradation was increased about twice after 30 d compared to those obtained after 15 d. The capacity of different *Pleurotus* species and other basidiomycetes to degrade lignin has been related to ligninolytic enzyme production (Hatakka 1994; Camarero *et al.* 1996). It is necessary to emphasize that basidiomycetes have been described as the most adequate fungi for pith biocontrol in eucalypt wood SSF cultures (Martínez *et al.* 1999) and the use of laccases and peroxidases seems to be also involved in the transformation of wood extractives (Buchert *et al.* 2001; del Río *et al.* 2002).

Table II. *Eucalyptus globulus* wood treatment with *Pleurotus laciniatocrenatus*

Incubation time, d	Enzyme activity ^a , pkat/g wood		Wood mass loss ^{a,b} %	Lignin degradation ^{a,b,c} %	Acetone-extractive removal ^{a,b} , %
	Lac	MiP			
15	470 \pm 120	150 \pm 60	1.8 \pm 0.5	1.9 \pm 0.6	11.7 \pm 3.6
30	480 \pm 160	160 \pm 50	2.4 \pm 1.0	3.6 \pm 1.1	21.6 \pm 4.1

^aMeans \pm SD of three replicates.

^bDegradation in relation to the content of uninoculated sterilized wood chips (control).

^cAnalyzed by the Klason method.

The genus *Pleurotus* and/or its enzymes are being studied for applications in different industries (edible protein production, applications in paper and pulp production and bioremediation studies). Our work provides preliminary information on the ability of a South American *Pleurotus* strain to produce enzymes that are currently being tested for biotechnological applications from other wood fungi, and to degrade lignin and extractives from *E. globulus* wood. Since a thorough knowledge of the biodiversity of a given area

permits the exploration of potential uses of autochthonous biological resources for the implementation of new environmentally-sound methods and technology, the relevance of studies focusing on our fungal diversity must be emphasized.

The first author is recipient of a scholarship from the *National Council of Scientific and Technological Research*. This research was supported by grants from the *National Council of Scientific and Technological Research* and the *National University of La Plata*, Argentina.

REFERENCES

- ABDULLAH N., ZAFAR S.I.: Lignocellulose biodegradation by white-rot *Basidiomycetes*: overview. *Internat.J.Mushroom Sci.* **2**, 59–78 (1999).
- BOURBONNAIS R., PAICE M.G.: Oxidative enzymes from the lignin-degrading fungus *Pleurotus sajor-caju*, pp. 472–481 in N.G. Lewis, M.G. Paice (Eds): *Plant Cell Wall Polymers: Biogenesis and Biodegradation*. ACS Symp. Ser. Vol. 399. American Chemical Society, Washington (DC) 1989.
- BRADFORD M.M.: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal.Biochem.* **72**, 248–254 (1976).
- BUCHERT J., MISTRANTA A., KONKANEN H., KARLSSON S., TENKANEN M., HOLMBOM B.: Enzymatic control of wood extractives. *Proc. 11th Internat. Symp. Wood Pulp Chemistry* **3**, 375–378 (2001).
- CAMARERO S., BÖCKLE B., MARTÍNEZ M.J., MARTÍNEZ A.T.: Manganese-mediated lignin degradation by *Pleurotus pulmonarius*. *Appl. Environ.Microbiol.* **62**, 1070–1072 (1996).
- CING S., ASMA(HAMAMCI) D., APOHAN E., YEŞİLADA O.: Decolorization of textile dyeing wastewater by *Phanerochaete chrysosporium*. *Folia Microbiol.* **47**, 639–642 (2003).
- COHEN R., PERSKY L., HADAR Y.: Biotechnological applications and potential of wood-degrading mushrooms of the genus *Pleurotus*. *Appl.Microbiol.Biotechnol.* **58**, 582–594 (2002).
- COLLINS P.J., DOBSON A.D.W.: Regulation of laccase gene transcription in *Trametes versicolor*. *Appl.Environ.Microbiol.* **63**, 3444–3450 (1997).
- DITTMER K., PATEL N., DHAWALE S.W., DHAWALE S.S.: Production of multiple laccase isoforms by *Phanerochaete chrysosporium* grown under nutrient sufficiency. *FEMS Microbiol.Lett.* **149**, 65–70 (1997).
- EICHLEROVÁ I., HOMOLKA L., NERUD F.: Decolorization of orange G by *Pleurotus ostreatus* monokaryotic isolates with different laccase activity. *Folia Microbiol.* **48**, 775–779 (2003).
- FARNET A.M., TAGGER S., LE PETIT J.: Effects of copper and aromatic inducers on the laccases of the white-rot fungus *Marasmius quercophilus*. *Acad.Sci.Paris, Sciences de la vie* **322**, 499–503 (1999).
- GIARDINA P., PALMIERI G., FONTANELLA B., RIVIECCIO V., SANNIA G.: Manganese peroxidase isoenzymes produced by *Pleurotus ostreatus* grown on wood sawdust. *Arch.Biochem.Biophys.* **376**, 171–179 (2000).
- GUILLÉN F., MARTÍNEZ A.T., MARTÍNEZ M.J., EVANS C.S.: Hydrogen peroxide-producing system of *Pleurotus eryngii* involving the extracellular enzyme aryl-alcohol oxidase. *Appl.Microbiol.Biotechnol.* **41**, 465–470 (1994).
- GUILLÉN F., GÓMEZ-TORIBIO V., MUÑOZ C., MARTÍNEZ M.J., MARTÍNEZ A.T.: Production of hydroxyl radical by the synergistic action of fungal laccase and aryl-alcohol oxidase. *Arch.Biochem.Biophys.* **382**, 142–147 (2000).
- HATAKKA A.: Lignin-modifying enzymes from selected white-rot fungi – production and role in lignin degradation. *FEMS Microbiol. Rev.* **13**, 125–135 (1994).
- HAWKSWORTH D.L.: The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycol.Res.* **105**, 1422–1432 (2001).
- HEINFLING A., RUIZ-DUEÑAS F.J., MARTÍNEZ M.J., BERGBAUER M., SZEWCZYK U., MARTÍNEZ A.T.: A study on reducing substrates of manganese-oxidizing peroxidases from *Pleurotus eryngii* and *Bjerkandera adusta*. *FEBS Lett.* **428**, 141–146 (1998).
- JOHANNES C., MAJCHERCZYK A.: Natural mediators in the oxidation of polycyclic aromatic hydrocarbons by laccase mediator systems. *Appl.Environ.Microbiol.* **66**, 524–528 (2000).
- DE JONG E., FIELD J.A., DE BONT J.A.M.: Evidence for a new extracellular peroxidase: manganese inhibited peroxidase from the white-rot fungus *Bjerkandera* sp. BOS55. *FEBS Lett.* **299**, 107–110 (1992).
- KAAL E.E.J., DE JONG E., FIELD J.A.: Stimulation of ligninolytic peroxidase activity by nitrogen nutrients in the white-rot fungus *Bjerkandera* sp. strain BOS55. *Appl.Environ.Microbiol.* **59**, 4031–4036 (1993).
- KAHRAMAN S., YEŞİLADA O.: Decolorization and bioremediation of molasses wastewater by white-rot fungi in a semi-solid-state condition. *Folia Microbiol.* **48**, 525–528 (2003).
- LEVIN L., FORCHIASSIN F., RAMOS A.M.: Copper induction of lignin-modifying enzymes in the white-rot fungus *Trametes trogii*. *Mycologia* **94**, 377–383 (2002).
- MARTÍNEZ A.T., CAMARERO S., GUILLÉN F., GUTIÉRREZ A., MUÑOZ C., VARELA E., MARTÍNEZ M.J., BARRASA J.M., RUEL K., PELAYO J.: Progress in biopulping of non-woody materials: chemical, enzymatic and ultrastructural aspects of wheat straw delignification with ligninolytic fungi from the genus *Pleurotus*. *FEMS Microbiol.Rev.* **13**, 265–274 (1994).
- MARTÍNEZ M.J., RUIZ-DUEÑAS F.J., GUILLÉN F., MARTÍNEZ A.T.: Purification and catalytic properties of two manganese-peroxidase isoenzymes from *Pleurotus eryngii*. *Eur.J.Biochem.* **237**, 424–432 (1996).
- MARTÍNEZ M.J., BARRASA J.M., GUTIÉRREZ A., DEL RÍO J.C., MARTÍNEZ A.T.: Fungal screening for biological removal of extractives from *Eucalyptus globulus* wood. *Can.J.Bot.* **77**, 1513–1522 (1999).
- MESTER T., FIELD J.A.: Characterization of a novel manganese peroxidase-lignin peroxidase hybrid isozyme produced by *Bjerkandera* species strain BOS55 in the absence of manganese. *J.Biol.Chem.* **273**, 15412–15417 (1998).
- MUÑOZ C., GUILLÉN F., MARTÍNEZ A.T., MARTÍNEZ M.J.: Induction and characterization of laccase in the ligninolytic fungus *Pleurotus eryngii*. *Curr.Microbiol.* **34**, 1–5 (1997).
- NOVOTNÝ Č., ERBANOVA P., ŠÁSEK V., KUBÁTOVÁ A., CAJTHAML T., LANG E., KRAHL J., ZADRAZIL F.: Extracellular oxidative enzyme production and PAH removal in soil by exploratory mycelium of white-rot fungi. *Biodegradation* **10**, 159–168 (1999).
- PALMIERI G., GIARDINA P., BIANCO C., FONTANELLA B., SANNIA G.: Copper induction of laccase isoenzymes in the ligninolytic fungus *Pleurotus ostreatus*. *Appl.Environ.Microbiol.* **66**, 920–924 (2000).

- DEL RÍO J.C., GUTIÉRREZ A., MARTÍNEZ M.J., MARTÍNEZ A.T.: Identification of a novel series of alkylitaconic acids in wood cultures of *Ceriporiopsis subvermispota* by gas chromatography/mass spectrometry. *Rapid Commun.Mass Spectrom.* **16**, 62–68 (2002).
- RODRÍGUEZ E., NUERO O., GUILLÉN F., MARTÍNEZ A.T., MARTÍNEZ M.J.: Degradation of phenolic and non-phenolic aromatic pollutants by four *Pleurotus* species: the role of laccase and versatile peroxidase. *Soil Biol.Biochem.* **36**, 909–916 (2004).
- ROGALSKI J., LUNDELL T., LEONOWICZ A., HATAKKA A.: Production of laccase, lignin peroxidase and manganese-dependent peroxidase by various strains of *Trametes versicolor* depending on culture conditions. *Acta Microbiol.Polon.* **40**, 221–234 (1991).
- RUIZ-DUEÑAS F.J., GUILLÉN F., CAMARERO S., PÉREZ-BOADA M., MARTÍNEZ M.J., MARTÍNEZ A.T.: Regulation of peroxidase transcript levels in liquid cultures of the ligninolytic fungus *Pleurotus eryngii*. *Appl.Environ.Microbiol.* **65**, 4458–4463 (1999).
- RYVARDEN L.: Biodiversity in polypore fungi. A comparison between tropical Africa and America. *Rev.Biol.Trop.* **44**, 125–129 (1996).
- SALUSSO M., MORAÑA L.B.: Cultivo de *Pleurotus laciniatocrenatus* en Argentina. *Rev.Iberoam.Micol.* **14**, 129–130 (1997).
- SANNIA G., GIARDINA P., LUNA M., ROSSI M., BUONOCORE F.: Laccase from *Pleurotus ostreatus*. *Biotechnol.Lett.* **8**, 797–800 (1986).
- SANNIA G., LIMONGI P., COCCA E., BUONOCORE F., NITTI G., GIARDINA P.: Purification and characterization of a veratryl-alcohol oxidase enzyme from the lignin degrading basidiomycete *Pleurotus ostreatus*. *Biochim.Biophys.Acta* **1073**, 114–119 (1991).
- SAPARRAT M.C.N., BUCSINSZKY A.M.M., TOURNIER H.A., CABELLO M.N., ARAMBARRI A.M.: Extracellular ABTS-oxidizing activity of autochthonous fungal strains from Argentina in solid medium. *Rev.Iberoam.Micol.* **17**, 64–68 (2000).
- SAPARRAT M.C.N., GUILLÉN F., ARAMBARRI A.M., MARTÍNEZ A.T., MARTÍNEZ M.J.: Induction, isolation, and characterization of two laccases from the white-rot basidiomycete *Coriolopsis rigida*. *Appl.Environ.Microbiol.* **68**, 1534–1540 (2002).
- SARKAR S., MARTÍNEZ A.T., MARTÍNEZ M.J.: Biochemical and molecular characterization of a manganese peroxidase isoenzyme from *Pleurotus ostreatus*. *Biochim.Biophys.Acta* **1339**, 23–30 (1997).
- SCHLOSSER D., GREY R., FRITSCH W.: Patterns of ligninolytic enzymes in *Trametes versicolor*: distribution of extra- and intracellular enzyme activities during cultivation on glucose, wheat straw and beech wood. *Appl.Microbiol.Biotechnol.* **47**, 412–418 (1997).
- SOMOGYI M.: A new reagent for determination of sugars. *J.Biol.Chem.* **160**, 61–73 (1945).
- TAPPI: *Test Methods 1992–1993*. Technical Association of the Pulp and Paper Industry, Atlanta (USA) 1993.
- TOMŠOVSKÝ M., HOMOLKA L.: Laccase and other ligninolytic enzyme activities of selected strains of *Trametes* spp. from different localities and substrates. *Folia Microbiol.* **48**, 413–416 (2003).
- WONG Y., YU J.: Laccase-catalyzed decolorization of synthetic dyes. *Water Res.* **33**, 3512–3520 (1999).