



Fungal Planet description sheets: 1042–1111

P.W. Crous^{1,2}, M.J. Wingfield², Y.-H. Chooi³, C.L.M. Gilchrist³, E. Lacey⁴, J.I. Pitt⁴, F. Roets⁵, W.J. Swart⁶, J.F. Cano-Lira⁷, N. Valenzuela-Lopez^{7,8}, V. Hubka^{9,10}, R.G. Shivas¹¹, A.M. Stchigel⁷, D.G. Holdom¹², Ž. Jurjević¹³, A.V. Kachalkin^{14,15}, T. Lebel¹⁶, C. Lock¹², M.P. Martín¹⁷, Y.P. Tan¹⁸, M.A. Tomashevskaya¹⁵, J.S. Vitelli¹², I.G. Baseia¹⁹, V.K. Bhatt²⁰, T.E. Brandrud²¹, J.T. De Souza²², B. Dima²³, H.J. Lacey⁴, L. Lombard¹, P.R. Johnston²⁴, A. Morte²⁵, V. Papp²⁶, A. Rodríguez²⁵, E. Rodríguez-Andrade⁷, K.C. Semwal²⁷, L. Tegart²⁸, Z.G. Abad²⁹, A. Akulov³⁰, P. Alvarado³¹, A. Alves³², J.P. Andrade^{33,34}, F. Arenas²⁵, C. Asenjo³⁵, J. Ballarà³⁶, M.D. Barrett³⁷, L.M. Berná²⁵, A. Berraf-Tebbal³⁸, M.V. Bianchinotti³⁹, K. Bransgrove¹⁸, T.I. Burgess⁴⁰, F.S. Carmo²², R. Chávez⁴¹, A. Čmoková⁹, J.D.W. Dearnaley¹¹, A.L.C.M. de A. Santiago⁴², J.F. Freitas-Neto⁴³, S. Denman⁴⁴, B. Douglas⁴⁵, F. Dovana⁴⁶, A. Eichmeier³⁸, F. Esteve-Raventós⁴⁷, A. Farid⁴⁸, A.G. Fedosova⁴⁹, G. Ferisin⁵⁰, R.J. Ferreira⁵¹, A. Ferrer⁵², C.N. Figueiredo⁵³, Y.F. Figueiredo²², C.G. Reinoso-Fuentealba³⁹, I. Garrido-Benavent⁵⁴, C.F. Cañete-Gibas⁵⁵, C. Gil-Durán⁴¹, A.M. Glushakova^{14,56}, M.F.M. Gonçalves³², M. González⁵⁷, M. Gorczak⁵⁸, C. Gorton⁴⁴, F.E. Guard⁵⁹, A.L. Guarnizo²⁵, J. Guarro⁷, M. Gutiérrez³⁵, P. Hamal⁶⁰, L.T. Hien⁶¹, A.D. Hocking⁶², J. Houbraken¹, G.C. Hunter⁶³, C.A. Inácio⁶⁴, M. Jourdan⁶⁵, V.I. Kapitonov⁶⁶, L. Kelly⁶⁷, T.N. Khanh⁶¹, K. Kisto⁵⁸, L. Kiss¹¹, A. Kiyashko⁴⁹, M. Kolařík¹⁰, J. Kruse¹¹, A. Kubátová⁹, V. Kučera⁶⁸, I. Kučerová⁹, I. Kušan⁶⁹, H.B. Lee⁷⁰, G. Levicán⁴¹, A. Lewis⁴⁴, N.V. Liem⁶¹, K. Liimatainen⁴⁵, H.J. Lim⁷⁰, M.N. Lyons⁷¹, J.G. Maciá-Vicente⁷², V. Magaña-Dueñas⁷, R. Mahiques⁷³, E.F. Malysheva⁴⁹, P.A.S. Marbach⁵³, P. Marinho⁷⁴, N. Matočec⁶⁹, A.R. McTaggart⁷⁵, A. Mešić⁶⁹, L. Morin⁶³, J.M. Muñoz-Mohedano²⁵, A. Navarro-Ródenas²⁵, C.P. Nicolli²², R.L. Oliveira⁷⁶, E. Otsing⁷⁷, C.L. Ovrebo⁷⁸, T.A. Pankratov^{14,79}, A. Paños²⁵, A. Paz-Conde⁸⁰, A. Pérez-Sierra⁴⁴, C. Phosri⁸¹, Á. Pintos⁸², A. Pošta⁶⁹, S. Prencipe⁸³, E. Rubio⁸⁴, A. Saitta⁸⁵, L.S. Sales⁵³, L. Sanhueza⁵², L.A. Shuttleworth⁴⁴, J. Smith⁸⁶, M.E. Smith⁸⁷, D. Spadaro⁸³, M. Spetik³⁸, M. Sochor⁸⁸, Z. Sochorová⁸⁹, J.O. Sousa⁴³, N. Suwannasai⁹⁰, L. Tedersoo⁷⁷, H.M. Thanh⁶¹, L.D. Thao⁶¹, Z. Tkalčec⁶⁹, N. Vaghefi¹¹, A.S. Venzhik¹⁴, A. Verbeken⁹¹, A. Vizzini⁹², S. Voyron⁴⁶, M. Wainhouse⁹³, A.J.S. Whalley⁹⁴, M. Wrzosek⁹⁵, M. Zapata⁹⁶, I. Zeil-Rolfe⁶³, J.Z. Groenewald¹

Key words

ITS nrDNA barcodes
LSU
new taxa
systematics

Abstract Novel species of fungi described in this study include those from various countries as follows: **Antarctica**, *Cladosporium arenosum* from marine sediment sand. **Argentina**, *Kosmimatomyces alatophyllus* (incl. *Kosmimatomyces* gen. nov.) from soil. **Australia**, *Aspergillus banksianus*, *Aspergillus kumbius*, *Aspergillus luteorubrus*, *Aspergillus malvicolor* and *Aspergillus nanangensis* from soil, *Erysiphe medicaginis* from leaves of *Medicago polymorpha*, *Hymenotorrendiella communis* on leaf litter of *Eucalyptus bicostata*, *Lactifluus albopici* and *Lactifluus austropiperatus* on soil, *Macalpinomyces collinsiae* on *Eriachne benthamii*, *Marasmius vagus* on soil, *Microdochium dawsoniorum* from leaves of *Sporobolus natalensis*, *Neopestalotiopsis nebuloides* from leaves of *Sporobolus elongatus*, *Pestalotiopsis etonensis* from leaves of *Sporobolus jacquemontii*, *Phytophthora personensis* from soil associated with dying *Grevillea mcutcheonii*. **Brazil**, *Aspergillus oxumiae* from soil, *Calvatia baixaverdensis* on soil, *Geastrum calycicoriaceum* on leaf litter, *Greeneria kielmeyerae* on leaf spots of *Kielmeyera coriacea*. **Chile**, *Phytophthora aysenensis* on collar rot and stem of *Aristotelia chilensis*. **Croatia**, *Mollisia gibbospora* on fallen branch of *Fagus sylvatica*. **Czech Republic**, *Neosetophoma hnaniceana* from *Buxus sempervirens*. **Ecuador**, *Exophiala frigidotolerans* from soil. **Estonia**, *Elaphomyces bucholtzii* in soil. **France**, *Venturia paralias* from leaves of *Euphorbia paralias*. **India**, *Cortinarius balteatoindicus* and *Cortinarius ulkhagarhiensis* on leaf litter. **Indonesia**, *Hymenotorrendiella indonesiana* on *Eucalyptus urophylla* leaf litter. **Italy**, *Penicillium taurinense* from indoor chestnut mill. **Malaysia**, *Hemileucoglossum kelabitense* on soil, *Satchmopsis pini* on dead needles of *Pinus tecunumanii*. **Poland**, *Lecanicillium praecognitum* on insects' frass. **Portugal**, *Neodevriesia aestuarina* from saline water. **Republic of Korea**, *Gongronella namwonensis* from freshwater. **Russia**, *Candida pellucida* from *Exomias pellucidus*, *Heterocephalacria septentrionalis* as endophyte from *Cladonia rangiferina*, *Vishniacozyma phoenicis* from dates fruit, *Volvariella paludosa* from swamp. **Slovenia**, *Mallochybe crassivelata* on soil. **South Africa**, *Beltraniella podocarpi*, *Hamatocanthoscypha podocarpi*, *Coleophoma podocarpi* and *Nothoseiridium podocarpi* (incl. *Nothoseiridium*


Abstract (cont.)

gen. nov.) from leaves of *Podocarpus latifolius*, *Gyrothrix encephalarti* from leaves of *Encephalartos* sp., *Paraphyton cutaneum* from skin of human patient, *Phacidiella alsophilae* from leaves of *Alsophila capensis*, and *Satchmopsis metrosideri* on leaf litter of *Metrosideros excelsa*. **Spain**, *Cladophialophora cabanerensis* from soil, *Cortinariopsis paezii* on soil, *Cylindrium magnoliae* from leaves of *Magnolia grandiflora*, *Trichophoma cylindrospora* (incl. *Trichophoma* gen. nov.) from plant debris, *Tuber alcaracense* in calcareous soil, *Tuber buendiae* in calcareous soil. **Thailand**, *Annulohyphoxylon spougei* on corticated wood, *Poaceascoma filiforme* from leaves of unknown *Poaceae*. **UK**, *Dendrostoma luteum* on branch lesions of *Castanea sativa*, *Ypsilina buttingtonensis* from heartwood of *Quercus* sp. **Ukraine**, *Myrmecridium phragmiticola* from leaves of *Phragmites australis*. **USA**, *Absidia pararepens* from air, *Juncomyces californiensis* (incl. *Juncomyces* gen. nov.) from leaves of *Juncus effusus*, *Montagnula cylindrospora* from a human skin sample, *Muriphila oklahomaensis* (incl. *Muriphila* gen. nov.) on outside wall of alcohol distillery, *Neofabraea eucalyptorum* from leaves of *Eucalyptus macrandra*, *Diabolocovidia claustris* (incl. *Diabolocovidia* gen. nov.) from leaves of *Serenoa repens*, *Paecilomyces penicilliformis* from air, *Pseudopezicula betulae* from leaves of leaf spots of *Populus tremuloides*. **Vietnam**, *Diaporthe durionigena* from branches of *Durio zibethinus* and *Roridomyces pseudoirritans* on rotten wood. Morphological and culture characteristics are supported by DNA barcodes.

Article info Received: 1 April 2020; Accepted: 30 May 2020; Published: 29 June 2020.

- 1 Westerdijk Fungal Biodiversity Institute, P.O. Box 85167, 3508 AD Utrecht, The Netherlands.
- 2 Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), Faculty of Natural and Agricultural Sciences, University of Pretoria, Private Bag X20, Hatfield 0028, Pretoria, South Africa.
- 3 School of Chemistry and Biochemistry, University of Western Australia, Perth, WA 6009, Australia.
- 4 Microbial Screening Technologies, 28 Percival Rd, Smithfield, NSW 2164, Australia.
- 5 Department of Conservation Ecology and Entomology, Stellenbosch University, Stellenbosch 7600, South Africa.
- 6 Department of Plant Sciences (Division of Plant Pathology), University of the Free State, P.O. Box 339, Bloemfontein 9300, South Africa.
- 7 Mycology Unit, Medical School and IISPV, Universitat Rovira i Virgili (URV), Sant Llorenç 21, 43201 Reus, Tarragona, Spain.
- 8 Unidad de Microbiología, Departamento de Tecnología Médica, Facultad de Ciencias de la Salud, Universidad de Antofagasta, Av. Universidad de Antofagasta 02800, Antofagasta, Chile.
- 9 Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 01 Prague 2, Czech Republic.
- 10 Laboratory of Fungal Genetics and Metabolism, Institute of Microbiology of the Czech Academy of Sciences, v.v.i, Vídeňská 1083, 142 20 Prague 4, Czech Republic.
- 11 Centre for Crop Health, University of Southern Queensland, Toowoomba 4350, Queensland, Australia.
- 12 Biosecurity Queensland, Department of Agriculture and Fisheries, Dutton Park 4102, Queensland, Australia.
- 13 EMSL Analytical, Inc., 200 Route 130 North, Cinnaminson, NJ 08077 USA.
- 14 Lomonosov Moscow State University, 119234, Moscow, Leninskíe Gory Str. 1/12, Russia.
- 15 All-Russian Collection of Microorganisms, G.K. Skryabin Institute of Biochemistry and Physiology of Microorganisms RAS, 142290, Pushchino, pr. Nauki 5, Russia.
- 16 Royal Botanic Gardens Victoria, Private Bag 2000, Victoria 3141, Australia.
- 17 Departamento de Micología, Real Jardín Botánico, RJB-CSIC, Plaza de Murillo 2, 28014 Madrid, Spain.
- 18 Plant Pathology Herbarium, Department of Agriculture and Fisheries, Dutton Park 4102, Queensland, Australia.
- 19 Departamento Botânica e Zoologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Campus Universitário, 59072–970 Natal, RN, Brazil.
- 20 Navdanya, 105, Rajpur Road, Dehradun, Uttarakhand, India.
- 21 Norwegian Institute for Nature Research, Gaustadalléen 21, NO-0349 Oslo, Norway.
- 22 Federal University of Lavras, Minas Gerais, Brazil.
- 23 Department of Plant Anatomy, Institute of Biology, Eötvös Loránd University, Pázmány Péter sétány 1/C, H-1117, Budapest, Hungary.
- 24 Manaaki Whenua – Landcare Research, Private Bag 92170, Auckland 1142, New Zealand.
- 25 Departamento de Biología Vegetal (Botánica), Facultad de Biología, Universidad de Murcia, 30100 Murcia, Spain.
- 26 Department of Botany, Faculty of Horticultural Science, Szent István University, P.O. Box 53, H-1518, Budapest, Hungary.
- 27 Department of Biology, College of Sciences, Eritrea Institute of Technology, Mai Nefhi, Asmara, Eritrea.
- 28 Menzies Institute for Medical Research, University of Tasmania, Tasmania, Australia.
- 29 USDA-APHIS-PPQ-Science & Technology Beltsville Laboratory, Bldg 580-East, Powder Mill Rd, Beltsville, MD 20705 USA.
- 30 Department of Mycology and Plant Resistance, V. N. Karazin Kharkiv National University, Maidan Svobody 4, 61022 Kharkiv, Ukraine.
- 31 ALVALAB, Dr. Fernando Bongera st., Severo Ochoa bldg. S1.04, 33006 Oviedo, Spain.
- 32 Departamento de Biologia, CESAM, Universidade de Aveiro, 3810-193 Aveiro, Portugal.
- 33 Universidade Estadual de Feira de Santana, Bahia, Brazil.
- 34 Faculdades Integradas de Sergipe, Sergipe, Brazil.
- 35 Servicio Agrícola y Ganadero, Laboratorio Regional Osorno, Unidad de Fitopatología, Ruta a Puerto Octay U-55-V, Osorno, Chile.
- 36 C/ Tossalet de les Forques, 44, E-08600, Berga, Catalonia, Spain.
- 37 Australian Tropical Herbarium, James Cook University, Smithfield Queensland 4878, Australia.
- 38 MENDELEUM – Institute of Genetics, Mendel University, Valtická 334, 69144, Czech Republic.
- 39 Laboratorio de Micología, Fitopatología y Control Biológico, Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS-CONICET). Camino La Carrindanga, Km 7. Dto. de Biología, Bioquímica y Farmacia. Universidad Nacional del Sur (DBBF-UNS). San Juan 670. (B8000ICN) Bahía Blanca, Argentina.
- 40 Phytothora Science and Management, Centre for Climate Impacted Terrestrial Ecosystems, Harry Butler Institute, Murdoch University, Murdoch, WA 6150, Australia.
- 41 Facultad de Química y Biología, Universidad de Santiago de Chile (USACH), Alameda 3363, Estación Central, 9170022, Santiago, Chile.
- 42 Department of Mycology, Federal University of Pernambuco, Recife, Brazil.
- 43 Universidade Federal do Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil.
- 44 Forest Research, Alice Holt Lodge, Farnham, Surrey, GU10 4LH, UK.
- 45 Jodrell Laboratory, Royal Botanic Gardens, Kew, Surrey TW9 3AB, UK.
- 46 Department of Life Sciences and Systems Biology, University of Turin, Viale P.A. Mattioli 25, 10125, Torino, Italy.
- 47 Departamento de Ciencias de la Vida, Universidad de Alcalá, Campus universitario 28805, Alcalá de Henares (Madrid), Spain.
- 48 Herbarium, Department of Cell, Microbiology & Molecular Biology, 4202 E Fowler Avenue, Tampa FL 33620, USA.
- 49 Komarov Botanical Institute of the Russian Academy of Sciences, 197376, 2 Prof Popov Str., Saint Petersburg, Russia.
- 50 Via A. Vespucci 7, 1537, 33052 Cervignano del Friuli (UD), Italy.
- 51 Departamento de Micología, Universidade Federal de Pernambuco, 50670-420 Recife, PE, Brazil.
- 52 Facultad de Estudios Interdisciplinarios, Núcleo de Química y Bioquímica, Universidad Mayor, Santiago, Chile.
- 53 Federal University of Recôncavo da Bahia, Bahia, Brazil.
- 54 Department of Biogeochemistry and Microbial Ecology, National Museum of Natural Sciences, CSIC, E-28002, Madrid, Spain.
- 55 Fungus Testing Laboratory, Department of Pathology and Laboratory Medicine, Long School of Medicine, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, Texas, USA.
- 56 Mechnikov Research Institute for Vaccines and Sera, 105064, Moscow, Maly Kazenny by-street, 5A, Russia.
- 57 Torrecerredo 11 1F, 33211 Gijón, Spain.
- 58 Institute of Evolutionary Biology, Faculty of Biology, University of Warsaw, Żwirki i Wigury 101, 02–089 Warsaw, Poland.
- 59 Maleny, Queensland, Australia.
- 60 Department of Microbiology, Faculty of Medicine and Dentistry, Palacky University, Olomouc, Czech Republic.
- 61 Division of Plant Pathology, Plant Protection Research Institute (PPRI), Duc Thang, Bac Tu Liem, Hanoi, Vietnam.
- 62 CSIRO Agriculture and Food, North Ryde, NSW 2113, Australia.

- ⁶³ CSIRO Health and Biosecurity, GPO Box 1700, Canberra, ACT 2601, Australia.
- ⁶⁴ Federal and Rural University of Rio de Janeiro, Seropedica, Rio de Janeiro, Brazil.
- ⁶⁵ CSIRO European Laboratory, Campus International de Baillarguet, Montferrier sur lez 34980, France.
- ⁶⁶ Tobolsk Complex Scientific Station of the Ural Branch of the Russian Academy of Sciences, 626152, 15 Academic Yuri Osipov Str., Tobolsk, Russia.
- ⁶⁷ Department of Agriculture and Fisheries, Queensland Government, Toowoomba 4350, Queensland, Australia.
- ⁶⁸ Plant Science and Biodiversity Centre, Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23, Bratislava, Slovakia.
- ⁶⁹ Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia.
- ⁷⁰ Environmental Microbiology Lab, Dept. of Agricultural Biological Chemistry, College of Agriculture and Life Sciences, Chonnam National University, Gwangju 61186, Korea.
- ⁷¹ Ecosystem Science, Department of Biodiversity, Conservation and Attractions, Kensington 6151, Western Australia.
- ⁷² Institute of Ecology, Evolution and Diversity, Goethe University Frankfurt, Max-von-Laue-Str. 13, 60438, Frankfurt am Main, Germany.
- ⁷³ C/ Dr. Climent, 26, E-46837, Quatretonda, València, Spain.
- ⁷⁴ Departamento de Biologia Celular e Genética, Universidade Federal do Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil.
- ⁷⁵ Queensland Alliance for Agriculture and Food Innovation, University of Queensland, Brisbane 4102, Australia.
- ⁷⁶ Centro de Biociências, Universidade Federal do Rio Grande do Norte, Av. Senador Salgado Filho, 3000, 59072-970 Natal, RN, Brazil.
- ⁷⁷ Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, 40 Lai St., 51005 Tartu, Estonia.
- ⁷⁸ Department of Biology, University Central Oklahoma Edmond, Oklahoma, 73034 USA.
- ⁷⁹ S.N. Winogradsky Institute of Microbiology, Research Center of Biotechnology of the Russian Academy of Sciences, 119071, Moscow, Russia.
- ⁸⁰ Agropacio Micologica Berguedana, carrera Vall-Ter 791, apto. correos 6, 17455, Girona, Spain.
- ⁸¹ Biology programme, Faculty of Science, Nakhon Phanom University, Nakhon Phanom, 48000, Thailand.
- ⁸² Interdisciplinary Ecology Group, Universitat de les Illes Balears, ctra. de Valldemossa Km 7.5. 07122 Illes Balears, Spain.
- ⁸³ University of Turin - Department of Agricultural, Forestry and Food Sciences, Largo Paolo Braccini 2, 10095, Grugliasco, Turin, Italy.
- ⁸⁴ José Cueto 3 5B, 33401 Avilés, Spain.
- ⁸⁵ Department of Agricultural, Food and Forest Sciences, University of Palermo, Viale delle Scienze, Palermo, 90128, Italy.
- ⁸⁶ School of Forest Resources and Conservation, University of Florida, Gainesville, FL 32611-0410 USA.
- ⁸⁷ Department of Plant Pathology & Florida Museum of Natural History, 2527 Fifield Hall, Gainesville FL 32611, USA.
- ⁸⁸ Centre of the Region Haná for Biotechnological and Agricultural Research, Crop Research Institute, Šlechtitelů 29, 78371, Olomouc, Czech Republic.
- ⁸⁹ Department of Botany, Faculty of Science, Palacký University Olomouc, Šlechtitelů 27, 78371, Olomouc, Czech Republic.
- ⁹⁰ Department of Microbiology, Faculty of Science, Srinakharinwirot University, Bangkok, 10110 Thailand.
- ⁹¹ Campus Ledeganck, Ghent University, Belgium.
- ⁹² Institute for Sustainable Plant Protection (IPSP) – CNR, Viale P.A. Mattioli 25, 10125, Torino, Italy.
- ⁹³ Organisms and Environment Research Division, School of Biosciences, Cardiff University, Cardiff, UK.
- ⁹⁴ School of Pharmacy and Biomolecular Sciences, Liverpool John Moores, Byrom Street, Liverpool, L3 3 AF, UK.
- ⁹⁵ Botanic Garden, Faculty of Biology, University of Warsaw, Aleje Ujazdowskie 4, 00-478 Warsaw, Poland.
- ⁹⁶ Servicio Agrícola y Ganadero, Laboratorio Regional Chillán, Unidad de Fitopatología, Claudio Arrau 738, Chillán, Código Postal 3800773, Chile.

Kosmimatamyces alatophylus

Fungal Planet 1085 – 29 June 2020

***Kosmimatamyces* Bianchin., Reinoso F., Rodr.-Andr., Cano & Stchigel, gen. nov.**

Etymology. From Greek *κοσμήματα*-, jewellery, and *-μύκης*, fungus, because of the microscopic look of the fungus.

Classification — *Capnodiaceae*, *Capnodiales*, *Dothideo-mycetidae*, *Dothideomycetes*.

Mycelium consisting of branched, septate, pale to dark brown, thick-walled hyphae, sometimes coarsely ornamented. *Conidiophores* solitary, macronematous or semimacronematous, erect, straight to flexuous, from hyaline to dark brown, thick- and smooth- to rough-walled, cylindrical, narrow, branched or not, branches

terminal and lateral, in angles of 45 to 90°. *Conidiogenous cells* determinate, integrated, terminal and intercalary, mono- or polyblastic, pale to dark brown, verrucose, scars truncate. *Conidia* holoblastic, 0–1-septate, brown to dark brown, thick-walled, globose, ovoid or ellipsoid, ornamented with spines and crater-like warts, with dark scars at one or both ends, arranged in branching acropetal chains.

Type species. *Kosmimatamyces alatophylus* Bianchin., Reinoso F., Rodr.-Andr., Cano & Stchigel.
Mycobank MB833527.

***Kosmimatamyces alatophylus* Bianchin., Reinoso F., Rodr.-Andr., Cano & Stchigel, sp. nov.**

Etymology. From Greek *αλατος*-, salt, and *-φιλος*, lover, because of the environment from which the fungus was recovered.

Mycelium consisting of branched, septate, thick-walled, 2.5–4.5 µm wide hyphae. *Conidiophores* solitary, macronematous or semimacronematous, erect, straight to flexuous, from hyaline to dark brown, thick- and smooth-walled to verrucose along its length, branched or unbranched, branches terminal or lateral, in an angle 45 to 90°, 13–100 × 3.5–6 µm. *Conidiogenous cells* determinate, integrated, terminal or intercalary, pale to dark brown, verrucose, mono- or polyblastic, 8–12 × 5–7.5 µm, scars truncate of 2–3.5 µm wide. *Ramiconidia* aseptate, pale to dark brown, thick- and smooth-walled to verrucose, subcylindrical, 8.5–25 × 3–6 µm. *Conidia* 0–1-septate, brown to dark brown, thick-walled, with a spinulose, digitate, pustulate to crater-like ornamentation, globose, limoniform to ovoid or ellipsoid, 6–11 × 5–10 µm, with one or more notorious scars, arranged in branching acropetal chains, of schizolytic secession.

Culture characteristics — (after 2 wk in darkness at 25 °C). Colonies on oatmeal agar (OA) up to 37 mm diam, flat, slightly dusty to floccose, greyish sepia (Rayner 1970), aerial mycelium scarce, margins entire, exudates as olivaceous brown; reverse black, diffusible pigments absent. Colonies on potato dextrose agar (PDA) up to 39 mm diam, flat, velvety, radiate and sulcate, greyish sepia at centre, greyish white to the borders, margins regular, scarce droplets of olivaceous exudates; reverse olive black to greyish sepia, diffusible pigments absent. On malt extract agar (MEA) up to 34 mm diam, velvety, zonate, radially folded and somewhat elevated, pale olivaceous grey, mostly consisting of vegetative mycelium, margins irregular; reverse greenish black, diffusible pigments absent. On potato carrot agar (PCA) up to 41 mm diam, floccose, olivaceous black, radiate, margin filamentous; reverse olivaceous black, diffusible pigments absent. On SNA up to 40 mm diam, flat, radiate, olivaceous at the centre and isabelline to the margins, margin entire; reverse olivaceous black at the centre, borders olive, diffusible pigments absent.

Colour illustrations. *Kosmimatamyces alatophylus*, Salitral de la Vidriera. Colony on OA at 2 wk; conidiophores, conidiogenous cells and conidia. Scale bar = 10 µm.

Typus. ARGENTINA, Buenos Aires province, Salitral de la Vidriera, S38 44.816 W62 33.251, from soil collected in a saltmarsh, 28 Aug. 2015, C. G. Reinoso Fuentealba & M.V. Bianchinotti (holotype CBS H-24325, culture ex-type FMR 15091; ITS and LSU sequences GenBank LR588887 and LR588888, MycoBank MB833528).

Notes — *Kosmimatamyces* is a new genus that groups in the *Capnodiaceae*, a family whose members are known as sooty molds whose dark hyphae cover the surface of living leaves and twigs of many plants (Hughes 1976, Abdollahzadeh et al. 2020). Hypersaline soil represents a new ecological niche, reinforcing the hypothesis of Crous et al. (2009) and Chomnunti et al. (2011) that plant surfaces are not the only environmental niche for this group of fungi. Based on a megablast search of NCBI's GenBank nucleotide database, the closest hit using the ITS sequence was *Microxiphium theae* CBS 202.30 (GenBank MH855113; Identities = 475/514 (92 %), 11 gaps (2 %)), *Antennariella placitae* AS01 (GenBank MG583755; Identities = 472/511 (92 %), 11 gaps (2 %)), and *Leptoxiphium kurandae* MCC1085 (GenBank KF826942; Identities = 470/510 (92 %), 9 gaps (2 %)); using the LSU sequence the closest hit were *Capnodium coartatum* MFLUCC10-0066 (GenBank JN832613; Identities = 547/555 (99 %), no gaps), *Microxiphium aciculiforme* CBS 892.73 (GenBank GU301847; Identities = 547/555 (99 %), no gaps), and *Conidioxiphium gardeniorum* CPC 14327 (GenBank GU301807; Identities = 547/555 (99 %), no gaps). The LSU phylogenetic tree corroborated the placement of our isolate close to the genus *Leptoxiphium*. The species of *Leptoxiphium* are characterised by pycnidial conidiomata with a bulbous swollen base and cylindrical neck that expands at the apex to become funnel-shaped (Hughes 1976, Chomnunti et al. 2011), whereas *Kosmimatamyces* produces single conidiophores.

Supplementary material

FP1085 Maximum likelihood tree obtained from the LSU sequence of our isolate and those retrieved from GenBank.

Cintia Reinoso-Fuentealba & M. Virginia Bianchinotti, Laboratorio de Micología, Fitopatología y Control Biológico, Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS-CONICET), Camino La Carrindanga, Km 7. Dto. de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur (DBBF-UNS), San Juan 670, (B8000ICN) Bahía Blanca, Argentina; e-mail: cin.reinoso@gmail.com & vbianchi@uns.edu.ar

Ernesto Rodríguez-Andrade, José F. Cano-Lira & Alberto M. Stchigel, Mycology Unit, Medical School and IISPV, Universitat Rovira i Virgili (URV), Sant Llorenç 21, 43201 Reus, Tarragona, Spain; e-mail: dc.ernesto.roan@outlook.com, jose.cano@urv.cat & albertomiguel.stchigel@urv.cat