



Published in final edited form as:

Hippocampus. 2019 November ; 29(11): 1038–1048. doi:10.1002/hipo.23094.

TRPC channels are not required for graded persistent activity in entorhinal cortex neurons

Alexei V. Egorov¹, Dagmar Schumacher², Rebekka Medert², Lutz Birnbaumer³, Marc Freichel^{2,*}, Andreas Draguhn^{1,*}

¹Institute of Physiology and Pathophysiology, Heidelberg University, 69120 Heidelberg, Germany

²Institute of Pharmacology, Heidelberg University, 69120 Heidelberg, Germany

³Neurobiology Laboratory, National Institute of Environmental Health Sciences, Durham, North Carolina 27709, USA and School of Medical Sciences, Catholic University of Argentina, Institute of Biomedical Research (BIOMED UCA-CONICET), Buenos Aires, C1107AAZ Argentina

Abstract

Adaptive behavior requires the transient storage of information beyond the physical presence of external stimuli. This short-lasting form of memory involves sustained (“persistent”) neuronal firing which may be generated by cell-autonomous biophysical properties of neurons or/and neural circuit dynamics. A number of studies from brain slices reports intrinsically generated persistent firing in cortical excitatory neurons following suprathreshold depolarization by intracellular current injection. In layer V (LV) neurons of the medial entorhinal cortex (mEC) persistent firing depends on the activation of cholinergic muscarinic receptors and is mediated by a calcium-activated nonselective cation current (I_{CAN}). The molecular identity of this conductance remains, however, unknown. Recently, it has been suggested that the underlying ion channels belong to the canonical transient receptor potential (TRPC) channel family and include heterotetramers of TRPC1/5, TRPC4 and/or TRPC1/4/5 channels. While this suggestion was based on pharmacological experiments and on effects of TRP-interacting peptides, an unambiguous proof based on TRPC channel-depleted animals is pending. Here, we used two different lines of TRPC channel knockout mice, either lacking TRPC1-, TRPC4-, and TRPC5-containing channels or lacking all seven members of the TRPC family. We report unchanged persistent activity in mEC LV neurons in these animals, ruling out that muscarinic-dependent persistent activity depends on TRPC channels.

Keywords

TRPC1/4/5 KO mice; hepta-TRPC KO mice; persistent firing; muscarinic

Correspondence Alexei V. Egorov, Institute of Physiology and Pathophysiology, Heidelberg University, Im Neuenheimer Feld 326, 69120 Heidelberg, Germany, alexei.egorov@urz.uni-heidelberg.de.

* shared senior authorship

Conflict of interest

The authors declare that they have no conflict of interest. The authors declare no competing financial interests.

Introduction

Working memory represents the ability of neuronal networks to maintain information for short periods of time after disappearance of the triggering stimulus. This transient storage may involve sustained (“persistent”) neuronal firing during the delay period, as suggested by different studies in non-human primates (Funahashi et al., 1989; Miller et al., 1996; Romo et al., 1999), humans (Kamitsuki et al., 2017) and rodents (MacDonald et al., 2011, Harvey et al., 2012; Pastalkova et al., 2008). Memory-associated persistent firing has been observed in multiple brain areas, including cortical networks, e.g. prefrontal, parietal, inferior temporal, auditory, somatosensory, and entorhinal cortex (EC) (Zylberberg & Strowbridge, 2017). Persistent activity may be caused by cell-autonomous (intrinsic) biophysical properties, neural circuit dynamics (i.e., synaptic reverberation in recurrent circuits) or a combination of both (Major & Tank, 2004; Zylberberg & Strowbridge, 2017). A number of studies from brain slices reports that suprathreshold depolarization by intracellular current injection can initiate firing that outlasts the stimulus even during pharmacological blockade of synaptic transmission. Such intrinsically generated persistent firing has been observed in excitatory neurons of prefrontal cortex (Lei et al., 2014), perirhinal cortex (Navaroli et al., 2012), entorhinal cortex (Egorov et al., 2002a; Tahvildari et al., 2007), amygdala (Egorov et al., 2006), postsubiculum (Yoshida & Hasselmo, 2009), and hippocampus (Larimer & Strowbridge, 2010; Jochems & Yoshida, 2013; Knauer et al., 2013). Persistent activity in the medial entorhinal cortex (mEC) is mediated by a calcium-activated nonselective cation current (I_{CAN}) (Egorov et al., 2002a; Fransén et al., 2006; Tahvildari et al., 2008). It has been suggested that the channels underlying I_{CAN} in mEC layer V (LV) neurons belongs to the transient receptor potential (TRP) channel family (Al-Yahya et al., 2003), specifically to the canonical TRP channel subfamily (TRPC) (Reboreda et al., 2011; Zhang et al., 2011). Indeed, the seven members of TRPC subfamily (TRPC1 to TRPC7) form non-selective cation channels that are activated in response to stimulation of phospholipase C-coupled receptors (Montell et al., 2002; Wu et al., 2010). These properties are apparently in line with the observed muscarinic receptor-dependent persistent activity in entorhinal cortex LV neurons (Egorov et al., 2002a), which has been suggested to be mediated heterotetramers consisting of either TRPC1 and TRPC5, TRPC1 and TRPC4 and/or TRPC1 with TRPC4 and TRPC5 channels (Zhang et al., 2011). Recently, we analyzed the assembly of TRPC1, TRPC4, and TRPC5 in the central nervous system (CNS) by means of quantitative high-resolution mass spectrometry (Bröker-Lai et al., 2017). Indeed, affinity purifications of TRPC isoforms from mouse hippocampal membranes antibodies demonstrated robust co-assembly of TRPC1, TRPC4 and TRPC5, while no other TRP channels were detected. Thus, TRPC1/4/5 channels form heteromers from three (TRPC1-TRPC4-TRPC5) or two subunits (TRPC1-TRPC4, TRPC1-TRPC5, TRPC4-TRPC5) (Bröker-Lai et al., 2017). Despite such insights about the molecular composition of TRPC subunits, the functional contribution of TRPC channels to intrinsic neuronal properties is less clear. Based on pharmacological experiments using flufenamic acid (FFA, 100 μ M), 2-APB (100 μ M) or SKF-96365 (50 μ M) and on effects of TRPC4/TRPC5-interacting peptides TRPC channels have been proposed to mediate sustained neuronal firing (Zhang et al., 2011). However, an unambiguous proof based on TRPC channel-depleted animals is pending.

Here, we used two different lines of TRPC channel knockout (KO) mice, either lacking TRPC1-, TRPC4-, and TRPC5-containing channels (Bröker-Lai et al., 2017) or lacking all seven members of the TRPC family (Birnbaumer, 2015). We report unchanged persistent activity in mEC LV neurons in these animals, ruling out that muscarinic-dependent persistent activity depends on TRPC channels.

Materials and Methods

Animals

Experiments were performed using male TRPC1/4/5-triple-knockout and Trpc1/2/3/4/5/6/7-hepta-knockout mice. All experimental procedures were approved and performed in accordance with the ethic regulations and the animal welfare committee of Heidelberg University and with the approval of the state government of Baden-Württemberg. Housing was provided in Makrolon II cages with a maximum of three animals in a temperature-controlled holding room ($23 \pm 1^\circ\text{C}$) on a 12/12-hr light/dark cycle. Animals had ad libitum access to food and water.

A triple-knockout mouse line Trpc1/4/5-/- lacking TRPC1, TRPC4 and TRPC5 was generated by intercrossing mice of the three mouse lines Trpc1-/- (Dietrich et al, 2007), Trpc4-/- (Freichel et al, 2001), and Trpc5-/- (Xue et al., 2011). Each had been backcrossed to the C57Bl6/N strain (Charles River) for at least seven generations before they were used to generate the Trpc1/4/5-/- line. For all details see (Bröker-Lai et al., 2017).

The hepta-knockout mouse line Trpc1/2/3/4/5/6/7-/- lacking all seven TRPC subtypes (TRPC1, TRPC2, TRPC3, TRPC4, TRPC5, TRPC6, TRPC7) was generated by intercrossing mice of the following mouse lines: Trpc1-/- (Dietrich et al., 2007), Trpc2-/- (Stowers et al., 2002), TRPC3-/- (Hartmann et al., 2008), Trpc4-/- (Freichel et al., 2001), Trpc5-/- (Phelan et al., 2013), Trpc6-/- (Dietrich et al., 2005) and Trpc7-/- (Perez-Leighton et al., 2011). Trpc1/2/3/4/5/6/7-/- (hepta-TRPC KO) mice are viable and fertile (Birnbaumer, 2015). First generation (F1) offspring of C57Bl6/N and 129SvJ intercrosses were used as wild type (WT) controls for Trpc1/2/3/4/5/6/7-/- mice that were also on mixed genetic background derived from C57 and 129 strains. All mice used in experiments were genotyped for deficiency in Trpc alleles (for details see Supporting Information Figure S1 and Table S1).

Preparation of mouse brain slices

Horizontal brain slices (450 μm thick) containing the hippocampus and entorhinal cortex were obtained from 3–4 month old mice using standard techniques (Roth et al., 2016). In order to minimize animal suffering we decapitated mice under deep CO_2 -induced anesthesia and used up to 6 slices per animal for experiments. After decapitation, brains were rapidly removed and placed in cold ($1-4^\circ\text{C}$) oxygenated artificial cerebrospinal fluid (ACSF) containing (in mM): 124 NaCl, 3 KCl, 1.8 MgSO_4 , 1.6 CaCl_2 , 10 glucose, 1.25 NaH_2PO_4 , 26 NaHCO_3 , saturated with carbogen (95% O_2 / 5% CO_2 , pH 7.4 at 34°C). Brain slices were cut using a vibratome slicer (Leica VT1200S, Nussloch, Germany), then transferred into a Haas-type interface chamber (Haas et al., 1979), superfused with ACSF at a rate of 1.5–2

ml/min, and maintained at $34 \pm 1^\circ\text{C}$. Prior to electrophysiological recordings, slices were allowed to recover for at least two hours.

Recording procedures

Intracellular recordings were obtained using sharp microelectrodes pulled on a Flaming/Brown puller P-1000 (Sutter Instruments, Novato, CA, USA) from 1.5 mm borosilicate glass capillaries (Science Products, Hofheim, Germany, Cat. No. GB150F-10). Electrode tips were backfilled with 1% biocytin (Sigma-Aldrich, Taufkirchen, Germany, Cat. No. B4261) dissolved in 2 M K-acetate (tip resistance of 70–110 M Ω). We used biocytin staining to confirm cell location after the experiment. Potentials were amplified using an Axoclamp-2 amplifier (Axon Instruments, Burlingame, CA, USA), filtered at 3 kHz and digitized at 20 kHz with an ADC (model MICRO 1401 mkII, CED, Cambridge, UK). Signals were stored on a computer and visualized using Spike2 software (CED, Cambridge, UK). Intracellular potentials were recorded in bridge mode and the bridge balance was monitored throughout the experiment. Durations of positive and negative current injections (range 0.1 – 0.8 nA) were controlled using a Master-8 VP stimulator (AMPI, Jerusalem, Israel). Membrane potential was manually adjusted by intracellular injection of DC current through the recording electrode and held near firing threshold (approximately –60 mV) for suprathreshold conditions. The mEC was identified with a dissecting microscope by transillumination. EC LV excitatory neurons were identified electrophysiologically on the basis of their firing characteristics as previously described (Hamam et al., 2000; Egorov et al., 2002b).

Chemicals

Carbachol (CCh, 10 – 20 μM , Cat. No. C4382) and atropine (10 μM , Cat. No. A0257, both from Sigma-Aldrich, Taufkirchen, Germany) were bath-applied by continuous perfusion. Since the muscarinic-induced phenomena did not desensitize, the neurons were directly impaled in the presence of CCh. All recordings were performed during blockade of ionotropic glutamate- and GABA-mediated neurotransmission with a drug mixture consisting of a cocktail of DL-2-amino-5-phosphonovaleric acid (APV, 30 μM , Tocris Bristol, UK, Cat. No. 0105), 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, 10 μM , Tocris Bristol, UK, Cat. No. 1045) and picrotoxin (100 μM , Sigma-Aldrich, Taufkirchen, Germany, Cat. No. P1675). Picrotoxin was applied from stock solution made in DMSO. The final concentration of DMSO in ACSF was 0.1%.

Data analysis

Electrophysiological data were analyzed using Spike 2 laboratory software. Spectral (Fourier) analysis and peristimulus histograms were made using Spike 2. Averaged data are given as mean \pm SD or as median and the first and third quartiles (P_{25} and P_{75}). Calculation of the statistical significance was performed using unpaired, two-tailed Student's t-test for normally distributed data or Mann-Whitney U-test for non-normally distributed data. Statistical analysis was performed using GraphPad (InStat, San Diego, USA) software. Regression analysis was performed using simple linear regression in GraphPad (InStat, San Diego, USA), quantified by correlation coefficient r^2 .

Intrinsic properties of neurons are summarized in Supporting Information Table S2. All values were obtained in the presence of CCh and during blockade of AMPA-, NMDA- and GABA(A)-mediated neurotransmission. Resting membrane potential (RMP) was estimated by subtraction of the tip potential following withdrawal of the pipette from the cell. Input resistance was determined at RMP by passing current pulses (-0.2 nA, 200 ms) through the recording electrode and measuring the resulting voltage deflections (at late steady-state level). For the analysis of action potential (AP) properties, we injected DC current with slowly rising amplitude until the cell started firing at low frequency. AP threshold was calculated as the first peak of the second derivative. AP amplitude was calculated as difference between the membrane voltage at AP threshold and AP peak. AP half-width was measured at the membrane voltage halfway between AP threshold and AP peak. AP maximum rise slope was calculated as maximum of the first derivative of membrane voltage between AP threshold and AP peak. AP maximum repolarization slope was calculated as maximal negative amplitude of the first derivative of membrane voltage following AP peak. Fast afterhyperpolarization (fAHP) following single AP was determined from the notch at the end of the fast repolarization phase within 5 ms after AP peak. fAHP amplitude was calculated as difference between AP threshold and the notch potential. Amplitude of medium afterhyperpolarization (mAHP) was calculated similarly from the time interval of 5 – 50 ms after AP peak. Trains of APs were induced by positive current injection of 1 s duration starting from ~ 12 mV negative to spike threshold. In this case, amplitude of the resulting mAHP was calculated as difference between minimum membrane voltage within 200 ms after current-step offset and AP threshold or recording membrane potential, as indicated. The frequency of persistent firing was calculated as average value from at least 20 s of recording. During persistent spiking, membrane potential oscillates between afterhyperpolarizations and spikes. We typically observed a short inflection (or plateau) immediately prior to the action potential. These potentials were taken as estimated membrane potential during persistent firing and were compared to baseline potential to calculate the depolarization during persistent firing.

Expression analysis by qPCR

For qPCR analysis of the expression of Trpc1-Trpc7 transcripts in the entorhinal cortex (Figure 2a) horizontal brain slices (1800 μm thick) were obtained from 3 – 4 month old mice (F1, see section animals) using standard procedures as described above. Area of interest (the entorhinal cortex together with the perirhinal and postrhinal cortices; see Figure 2a right) was dissected from individual slices under visual control (2 slices pro animal; i.e., one slice from each brain hemisphere). Tissue was placed into greiner bio-one tubes, cooled with liquid nitrogen and then stored at -80°C . For RNA isolation tissue was homogenized in Qiagen's RLT buffer first by using a Polytron (Kinematica PT 1200 E), followed by a glass homogenizer (Kimble Kontes Tenbroeck 2mL) and centrifugation in the QIAshredder column according to manufacturer's recommendations. RNA isolation was performed using the RNeasy Mini kit (Qiagen) according to manufacturer's recommendations for tissue, including on-column DNase digest. cDNA synthesis was carried out using the SensiFAST cDNA synthesis kit (Bioline) according to manufacturer's recommendations. Primers were designed with the online tool provided by Roche (https://lifescience.roche.com/en_de/brands/universal-probe-library.html) and the best primer pair for each target out of 2 – 3 was

chosen from an initial qPCR screen. Quantitative expression analysis was performed using the Universal Probe system (Roche) with the corresponding FastStart Essential DNA Probes Master (Roche) on a LightCycler 96 Instrument (Roche, Mannheim, Germany). Relative expression levels were obtained by normalising to H3F3A (H3 histone family member 3A), AIP (aryl-hydrocarbon receptor-interacting protein) and CXXC1 (CXXC finger protein 1) expression levels. Primer sequences are listed in Table 1.

For analysis of the expression of *Trpc1-Trpc7* transcripts in hepta-TRPC knockout tissue as compared to control mice (Figure 2b) we designed the primer pairs listed in Table 2. For each intron spanning assay one primer was located in the exon that was deleted in the corresponding *Trpc* genes, and the second primer was placed in the adjacent exon upstream or downstream of the deleted exon. Quantitative expression analysis was performed using RNA obtained from frontal lobes of the brain of four independent hepta-TRPC KO mice and wild type controls, respectively. The amplification reaction was performed using the iTaq™ Universal SYBR® Green Supermix (Bio Rad) on a LightCycler 96 Instrument (Roche, Mannheim, Germany). Expression levels are displayed as Cq values (limited to 35 cycles) for each of the seven *Trpc* genes in comparison to the H3F3A, AIP and CXXC1 housekeeping genes.

Results

Persistent activity in TRPC1/4/5-triple-knockout mice

Muscarinic-dependent plateau potential and persistent activity in entorhinal cortex layer V neurons are mediated by a calcium-activated nonselective cation current (I_{CAN}) (Egorov et al., 2002a; Fransen et al., 2006). Transient receptor potential (TRP) channels are permeable to both monovalent and divalent cations and are promising candidates underlying I_{CAN} (Ramsey et al., 2006). Indeed, there is pharmacological evidence that I_{CAN} is mediated by TRP channels in the mEC (Al-Yahya et al., 2003; Reboreda et al., 2011). These channels are likely composed of the canonical channel subunits TRPC1, TRPC4 and TRPC5 (Zhang et al., 2011). However, a more direct proof using genetic interventions is missing. We therefore recorded intracellular potentials from individual mEC layer V neurons of mice lacking the implicated channel subunits (*Trpc1/Trpc4/Trpc5*-triple-knockout; TRPC1/4/5 KO; Bröker-Lai et al., 2017). Sharp microelectrode recordings were performed from entorhinal cortex slices of TRPC1/4/5 KO mice in the presence of the cholinergic receptors agonist CCh (10 – 20 μ M) and during blockade of AMPA-, NMDA- and GABA(A)-mediated neurotransmission. Under these conditions the resting membrane potential (RMP) was -65.3 ± 1.2 mV ($n = 8$) and average input resistance was 66 ± 27 M Ω ($n = 8$). Intrinsic properties of neurons are summarized in Supporting Information Table S2. To our surprise, we found that all tested mEC LV neurons from TRPC1/4/5 KO mice (8 neurons from 4 mice) were able to generate persistent firing: neurons responded to a suprathreshold current-step stimulus with delayed firing at a constant frequency (4.5 – 14 Hz) for an apparently indefinite period of time (tested up to 5 min) (Figure 1). Stimulation parameters required to elicit persistent activity varied from cell to cell. In general, positive current injections of 1 – 4 s duration starting from ~ 12 mV or less negative to spike threshold and eliciting spike trains of 20 – 51 Hz were sufficient. Similar to our previous observations in rats (Egorov et

al., 2002a; Egorov et al., 2006), persistent firing displayed pronounced activity- and voltage-dependence (i.e., the probability to induce sustained firing activity increased with rising duration/intensity of stimulation and with membrane depolarization, and vice versa). Figure 1 illustrates an example recording where current pulses of different duration were applied at a given voltage level. At this voltage level stimulation for 1 or 2 s triggered only after-depolarizations or prolonged plateau potentials. Increasing stimulus duration, however, led to a stable state of sustained spiking (Figure 1, left). After membrane hyperpolarization, trigger pulses of equivalent strength were not able to induce persistent firing (Figure 1, right). In the same neurons persistent firing could be elicited by a short stimulation from more depolarized voltage levels at about 10 mV or less negative to spike threshold. Thus, a stimulation with one second long suprathreshold current-step (spike train at 44.5 ± 5 Hz) caused sustained firing at a frequency of 9.0 ± 3.6 Hz in all tested cells ($n = 8/8$). The sustained firing was accompanied by a post-stimulus persistent depolarization of 4.1 ± 1.3 mV ($n = 8$) measured as the difference between the estimated membrane potential during persistent firing and the baseline potential before its induction (see Methods). The quantitative parameters of persistent activity are summarized in Table 3. In 2 out of 4 neurons persistent firing could be induced with strong current steps of 0.2 s duration (spike train frequency during stimulation was 70 – 85 Hz). Together, this data shows the ability of entorhinal cortex layer V neurons to generate persistent activity in the absence of TRPC1-, TRPC4-, and TRPC5-containing channels.

Persistent activity in TRPC hepta-knockout mice

It cannot be excluded that the deleted channel subunits (TRPC1, TRPC4 and TRPC5) were functionally substituted by other members of the TRPC-family. Moreover, qPCR analysis using RNA isolated from the entorhinal, perirhinal and postrhinal cortices indicates robust expression of other TRPC subtypes, particularly TRPC7, but also TRPC2, TRPC3 and TRPC6 (Figure 2a), that indeed might contribute to persistent firing of these neurons. We therefore repeated the experiments in cells from mice lacking expression of all seven *Trpc* genes (termed “hepta-TRPC KO”, Supporting Information Figure S1 and Table S1, and Birnbaumer, 2015). Quantitative PCR (qPCR) analysis using RNA isolated from brain tissue showed that all TRPC subunits are absent in the hepta-TRPC KO mice (Figure 2b and Table 2). Resting membrane potential of these neurons was -73.4 ± 5.4 mV ($n = 11$) and input resistance was 83 M Ω (median, $P_{25} = 59$ M Ω and $P_{75} = 89$ M Ω) ($n = 11$; recordings in the presence of CCh, CNQX, APV and picrotoxin). Intrinsic properties of neurons are summarized in Supporting Information Table S2. We found that mEC LV neurons lacking all seven TRPC-channel subunits could indeed respond to a suprathreshold current step with persistent firing that is comparable to wild type (WT) controls (Figure 3a, left; 11 out of 12 tested cells from 7 hepta-TRPC KO mice; see below for WT control values). The persistent activity relied on the activation of muscarinic receptors since its induction was completely blocked by atropine ($n = 3$; Figure 3a right). Similar to the above-described observations in TRPC1/4/5 KO mice, the plateau-potential that sustained persistent firing in hepta-TRPC KO mice displayed pronounced activity- and voltage-dependence (Figure 3b, c). Indeed, persistent firing could be elicited by a short stimulation from voltage levels less ~ 10 mV negative to spike threshold. At this potential, a 1 s spike train (32.9 ± 5.8 Hz) elicited sustained firing at a frequency of 6.7 ± 3.7 Hz (8 out of 11 cells). These values were not

significantly different from similar experiments in WT control mice (stimulation-induced firing 37.4 ± 3.3 Hz, persistent firing 8.7 ± 2.3 Hz; $n = 6$; no difference between WT controls and hepta-TRPC KO, $p = 0.113$ for induced firing, $p = 0.253$ for persistent firing, t-test; Figure 3d). Resting membrane potential of neurons from WT control mice was more depolarized than RMP in hepta-TRPC KO mice (-64.9 ± 2.2 mV ($n = 6$) versus -73.4 ± 5.4 mV ($n=11$), $p = 0.002$, t-test, measured in presence of carbachol). In line with previous findings, frequency of persistent firing in hepta-TRPC KO mice correlated positively with frequency of spike trains during current-step injections (Figure 3e).

The marked post-stimulus membrane depolarization underlying the persistent firing was also not significantly different between WT control and hepta-TRPC KO mice (4.4 ± 0.8 mV ($n = 6$) vs. 4.4 ± 1.4 mV ($n = 8$), $p = 0.963$, t-test; Figure 3f). A 2-s spike train at 32 Hz (median) elicited a post stimulus membrane depolarization of 5.0 ± 2.2 mV and sustained firing at a frequency of 7.4 ± 4.2 Hz ($n = 9$ for all values, 9/10 tested neurons). Quantitative parameters of persistent activity were not statistically different between hepta-TRPC KO and WT control mice (Table 3). Similar to TRPC1/4/5 KO mice, persistent firing in hepta-TRPC KO mice could be elicited by a 0.2 s current step (1/3 tested neurons; Figure 4a). Analogous to our previous observations (Egorov et al., 2002a), persistent firing could be reversibly turned off by prolonged membrane hyperpolarization for 6 – 12 s to below ~ -80 mV (Figure 4b).

Graded persistent activity in TRPC hepta-knockout mice

Medial EC LV neurons generate graded persistent activity upon repetitive stimulation (Egorov et al., 2002a; Fransén et al., 2006; Reboreda et al., 2007). We tested this behaviour in cells from hepta-TRPC KO mice. As illustrated in Figure 5a, repetitive application of an activating current step gave rise to graded increases of sustained discharge rates (8 neurons from 6 mice; 3 to 7 levels following repetitive stimulation with a 1 – 4 s depolarizing step). The maximum persistent firing frequency induced in this manner in hepta-TRPC KO mice was 9 Hz (median, $n = 18$, minimal value: 6.3 Hz, maximal value: 19 Hz). In addition, repetitive application of hyperpolarizing current pulse steps led to graded decreases in firing rate ($n = 3$, Figure 5b). Stimulus durations of at least 6 – 8 s were required for this graded decrease in persistent firing rate.

In 3 out of 8 neurons in hepta-TRPC KO mice repetitive depolarizing current steps initially gave rise to graded increases of firing rate followed by a decrease in firing frequency and, finally, termination (Figure 6a). This sequence could be repeated after ~ 100 s of silence, suggesting an activity- (e.g., calcium-) dependent mechanism of inactivation. This behavior is well compatible with the inverted U-shape of the Ca^{2+} dependence of persistent firing in mEC cells from wild type rats (Zhang et al., 2011). In these experiments Zhang and colleagues tested effects of intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) on the persistent spiking activity in mEC LV neurons by clamping $[\text{Ca}^{2+}]_i$ with BAPTA in the patch pipette. The authors found that Ca^{2+} influx from the extracellular space is required to establish an optimal window of $[\text{Ca}^{2+}]_i$ (70–500 nM) for CCh-evoked persistent activity. When $[\text{Ca}^{2+}]_i$ was maintained at a low (20 nM) or high (1 mM) levels, CCh-evoked persistent firing in response to depolarizing current pulses was absent. Persistent firing relied on the activation

of muscarinic receptors, since its induction was completely blocked after adding atropine (Figure 6b).

Discussion

Persistent and graded persistent firing during a delayed response task, in which the animal is required to keep information about a sensory signal across a delay period between the stimulus and the behavioral response, may represent a cellular analog of working memory (Zylberberg & Strowbridge, 2017). In brain slices, persistent activity or slow afterdepolarization (sADP), following spike firing during activation of muscarinic acetylcholine receptors have been reported in many types of cortical neurons, including pyramidal cells of the entorhinal cortex, hippocampus, and neocortex (Egorov et al., 2002a; Knauer et al., 2013; Lei et al., 2014).

In this study, we investigated the potential contribution of TRPC channels to generation of persistent activity in mEC LV neurons. Involvement of TRPC channels in cholinergic persistent activity has been suggested by various observations (including our own). This was assumed, however, based on nonspecific pharmacological experiments as well as peptides that interfere with TRP channel functions (Al-Yahya et al., 2003; Reboreda et al., 2011; Zhang et al., 2011; Tai et al., 2011). Previous studies have proposed that the nonselective cation current underlying sustained firing in the mEC is mediated by heteromultimeric TRPC channels containing TRPC1, TRPC4 or/and TRPC5 subunits (Zhang et al., 2011). Indeed, these subunits are expressed in the EC (von Bohlen und Halbach et al., 2005). We here report recordings from individual mEC layer V neurons obtained from mice lacking any TRPC1-, TRPC4-, and TRPC5-containing channels (Bröker-Lai et al., 2017) or a mouse line lacking all seven TRPC subtypes (Birnbaumer, 2015) that might contribute to formation of alternative TRPC heteromers. Our results lead, however, to the conclusion that TRPC channels are not necessary for persistent firing in the EC. We note that some hepta-TRPC KO neurons showed only transient plateau potentials but no persistent firing (see Table 3). This is, however, possibly due to inter-cell variance or to small differences in intrinsic neuronal properties which impair induction of persistent activity. In any case, hepta-TRPC KO neurons are clearly able to express persistent activity.

The potential involvement of TRPC channels to persistent firing or sADP in neocortical neurons has remained a controversial issue, and the ability of TRP proteins to form multimers complicates testing this hypothesis. It has been reported that expression of dominant negative TRPC subunits can attenuate sADP responses in prefrontal cortex neurons (Yan et al., 2009). However, genetic deletion of several TRPC subunits, including TRPC1, TRPC5, and TRPC6 (single knockouts), or both TRPC5 and TRPC6 together (double knockout), failed to reduce the amplitude of cholinergic sADPs in layer V pyramidal neurons of the mouse medial prefrontal cortex (Dasari et al., 2013). Indeed, besides TRPC, other subfamily member of TRP channels, that are expressed in the brain (Sawamura et al., 2017), could underlie persistent activity in cortical neurons. For instance, the TRP melastatin 5 channel (TRPM5) contributes importantly to the slow afterdepolarization in the mouse prefrontal cortex (Lei et al., 2014). By examining animals in which TRPM4 and TRPM5 channels were genetically deleted either alone or in combination this study further showed

that TRPM5 but not TRPM4 channels are involved in the generation of the sADP in layer V neurons. Nevertheless, since a significant sADP was still observed in TRPM4/5 double knockout mice, the slow afterdepolarization must be mediated by more than one type of ion channel (Lei et al., 2014). Pharmacology-based approaches suggest involvement of TRPM channels (particularly Ca^{2+} activated cation channel TRPM4) (Nilius et al., 2003) in the regulation of neuronal excitability in diverse cell types. Thus, TRPM4 and TRPM5 are the ion channels underlying I_{CAN} in preBötzing complex neurons during inspiratory burst activity (Crowder et al., 2007). TRPM2/TRPM4 channels are involved in generation of bursting activity in dopamine neurons of the substantia nigra (Mrejeru et al., 2011). In the mEC TRPM (probably TRPM4) mediates I_{CAN} which is responsible for the generation of hyperexcitable bursts at proximal dendrites of LV neurons. In the same preparation, antagonists of TRPC did not affect bursting behavior (Lin et al., 2017). These data show a limited contribution of TRPC channels to the regulation of neuronal excitability in mEC LV neurons, in line with our current results. In contrast, TRPM channels may be involved in generation of graded persistent firing in the mEC.

In addition to I_{CAN} , different inward or outward currents which are active near the resting potential could support intrinsic persistent activity (Zylberberg & Strowbridge, 2017, Debanne et al., 2018). Thus, using both biophysical and pharmacological assays, Cui and Strowbridge nicely demonstrated that in neocortical neurons carbachol mediated persistent firing may result from the modulation of ether-á-Go-Go related gene (ERG) K^+ channels (Cui and Strowbridge, 2018). Accordingly, ERG channels mediate a leak potassium current which is downregulated by calcium entry induced by repetitive spiking.

There is convergent evidence showing involvement of TRPC channels in cognitive functions. Therefore, TRPC1/4/5 KO mice (i.e. animals that we used in our current study) display deficits in spatial working memory and flexible spatial relearning as demonstrated in hippocampus-specific tasks, while spatial reference memory assessed using the Morris water maze is unaltered (Bröker-Lai et al., 2017). Deletion of the TRPC1 gene alone impairs spatial working memory and fear memory (Lepannetier et al., 2018). Mice lacking TRPC4 or TRPC5 channels have been shown to exhibit less anxious behaviors (decreased innate fear) than their WT counterparts (Riccio et al., 2009, Riccio et al., 2014). In contrast, targeted knockdown of TRPC3 channel in the hippocampus enhanced contextual fear conditioning (Neuner et al., 2015). Nevertheless, the role of TRPC channels in memory-related functions and particularly the link between sustained firing activity and working memory are still not completely understood. Our present findings show that one potential mechanism of working memory, persistent neuronal firing, is not related to the presence of any of the TRPC proteins, at least in mEC neurons. Further studies will need to define the molecular identity of the calcium-activated nonselective cation channels underlying persistent firing in the EC.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (SFB 1134, A01 to A.V.E. and A.D. and SFB-TRR 152, P07, P21, SFB 1118, B02, FOR 2289, P02 to M.F.) and by the Intramural Research Program of the NIH (project Z01-ES-101684 to L.B.). We thank Prof. Klaus Groschner and Patrick Wiedner (University of Graz) for great help with transgenic animals. We also thank Yoko Oguchi and Christin Matka for the expert assistance in the qPCR analysis and Dr. Angela Wirth for her help during preparation of this manuscript.

References

- Al-Yahya E, Hamel E, Kennedy TE, Alonso AA, & Egorov AV (2003). Persistent activity in entorhinal cortex neurons induced by muscarinic and metabotropic glutamate receptor activation and its dependence on TRP channels. *Society for Neuroscience Abstract*, 3775.
- Birnbaumer L (2015). From GTP and G proteins to TRPC channels: a personal account. *Journal of Molecular Medicine (Berlin)*, 93(9), 941–553.
- Bröker-Lai J, Kollewe A, Schindeldecker B, Pohle J, Nguyen Chi V, Mathar I, Guzman R, Schwarz Y, Lai A, Weißgerber P, Schwegler H, Dietrich A, Both M, Sprengel R, Draguhn A, Köhr G, Fakler B, Flockerzi V, Bruns D, & Freichel M (2017). Heteromeric channels formed by TRPC1, TRPC4 and TRPC5 define hippocampal synaptic transmission and working memory. *EMBO Journal*, 36(18), 2770–2789. [PubMed: 28790178]
- Crowder EA, Saha MS, Pace RW, Zhang H, Prestwich GD, & Del Negro CA (2007). Phosphatidylinositol 4,5-bisphosphate regulates inspiratory burst activity in the neonatal mouse preBötzing complex. *Journal of Physiology*, 582, 1047–1058. [PubMed: 17599963]
- Cui ED, & Strowbridge BW (2018). Modulation of Ether-à-Go-Go Related Gene (ERG) Current Governs Intrinsic Persistent Activity in Rodent Neocortical Pyramidal Cells. *The Journal of Neuroscience*, 38(2), 423–440. [PubMed: 29175952]
- Dasari S, Abramowitz J, Birnbaumer L, & Gullledge AT (2013). Do canonical transient receptor potential channels mediate cholinergic excitation of cortical pyramidal neurons? *Neuroreport*, 24(10), 550–554. [PubMed: 23652155]
- Debanne D, Inglebert Y, & Russier M (2018). Plasticity of intrinsic neuronal excitability. *Current Opinion in Neurobiology*, 54, 73–82. [PubMed: 30243042]
- Dietrich A, Kalwa H, Storch U, Mederos y Schnitzler M, Salanova B, Pinkenburg O, Dubrovskaya G, Essin K, Gollasch M, Birnbaumer L, & Gudermann T (2007). Pressure-induced and store-operated cation influx in vascular smooth muscle cells is independent of TRPC1. *Pflügers Archiv*, 455(3), 465–477. [PubMed: 17647013]
- Dietrich A, Mederos Y Schnitzler M, Gollasch M, Gross V, Storch U, Dubrovskaya G, Obst M, Yildirim E, Salanova B, Kalwa H, Essin K, Pinkenburg O, Luft FC, Gudermann T, & Birnbaumer L (2005). Increased vascular smooth muscle contractility in TRPC6^{-/-} mice. *Molecular and Cellular Biology*, 25(16), 6980–6989. [PubMed: 16055711]
- Egorov AV, Hamam BN, Fransén E, Hasselmo ME, & Alonso AA (2002a). Graded persistent activity in entorhinal cortex neurons. *Nature*, 420(6912), 173–178. [PubMed: 12432392]
- Egorov AV, Heinemann U, & Müller W (2002b). Differential excitability and voltage-dependent Ca²⁺ signalling in two types of medial entorhinal cortex layer V neurons. *European Journal of Neuroscience*, 16(7), 1305–12. [PubMed: 12405991]
- Egorov AV, Unsicker K, & von Bohlen und Halbach O (2006). Muscarinic control of graded persistent activity in lateral amygdala neurons. *European Journal of Neuroscience*, 24(11), 3183–3194. [PubMed: 17156379]
- Fransén E, Tahvildari B, Egorov AV, Hasselmo ME, & Alonso AA (2006). Mechanism of graded persistent cellular activity in entorhinal cortex layer V neurons. *Neuron*, 49(5), 735–746. [PubMed: 16504948]
- Freichel M, Suh SH, Pfeifer A, Schweig U, Trost C, Weissgerber P, Biel M, Philipp S, Freise D, Droogmans G, Hofmann F, Flockerzi V, & Nilius B (2001). Lack of an endothelial store-operated Ca²⁺ current impairs agonist-dependent vasorelaxation in TRP4^{-/-} mice. *Nature Cell Biology*, 3(2), 121–127. [PubMed: 11175743]

- Funahashi S, Bruce CJ, & Goldman-Rakic PS (1989). Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *Journal of Neurophysiology*, 61(2), 331–349. [PubMed: 2918358]
- Haas HL, Schaerer B, & Vosmansky M (1979). A simple perfusion chamber for the study of nervous tissue slices in vitro. *Journal of Neuroscience Methods*, 1(4), 323–325. [PubMed: 544974]
- Hamam BN, Kennedy TE, Alonso A, & Amaral DG (2000). Morphological and electrophysiological characteristics of layer V neurons of the rat medial entorhinal cortex. *The Journal of Comparative Neurology*, 418(4), 457–472. [PubMed: 10713573]
- Hartmann J, Dragicevic E, Adelsberger H, Henning HA, Sumser M, Abramowitz J, Blum R, Dietrich A, Freichel M, Flockerzi V, Birnbaumer L, & Konnerth A (2008). TRPC3 channels are required for synaptic transmission and motor coordination. *Neuron*, 59(3), 392–398. [PubMed: 18701065]
- Harvey CD, Coen P, & Tank DW (2012). Choice-specific sequences in parietal cortex during a virtual-navigation decision task. *Nature*, 484(7392), 62–68. [PubMed: 22419153]
- Jochems A, & Yoshida M (2013). Persistent firing supported by an intrinsic cellular mechanism in hippocampal CA3 pyramidal cells. *European Journal of Neuroscience*, 38(2), 2250–2259. [PubMed: 23651161]
- Kami ski J, Sullivan S, Chung JM, Ross IB, Mamelak AN, & Rutishauser U (2017). Persistently active neurons in human medial frontal and medial temporal lobe support working memory. *Nature Neuroscience*, 20(4), 590–601. [PubMed: 28218914]
- Knauer B, Jochems A, Valero-Aracama MJ, & Yoshida M (2013). Long-lasting intrinsic persistent firing in rat CA1 pyramidal cells: a possible mechanism for active maintenance of memory. *Hippocampus*, 23(9), 820–831. [PubMed: 23609880]
- Larimer P, & Strowbridge BW (2010). Representing information in cell assemblies: persistent activity mediated by semilunar granule cells. *Nature Neuroscience*, 13(2), 213–222. [PubMed: 20037579]
- Lei YT, Thuault SJ, Launay P, Margolskee RF, Kandel ER, & Siegelbaum SA (2014). Differential contribution of TRPM4 and TRPM5 nonselective cation channels to the slow afterdepolarization in mouse prefrontal cortex neurons. *Frontiers in Cellular Neuroscience*, 8, 267. [PubMed: 25237295]
- Lepanetier S, Gualdani R, Tempesta S, Schakman O, Seghers F, Kreis A, Yerna X, Slimi A, de Clippele M, Tajeddine N, Voets T, Bon RS, Beech DJ, Tissir F, & Gailly P (2018) Activation of TRPC1 Channel by Metabotropic Glutamate Receptor mGluR5 Modulates Synaptic Plasticity and Spatial Working Memory. *Frontiers in Cellular Neuroscience*, 12, 318. [PubMed: 30271326]
- Lin EC, Combe CL, & Gasparini S (2017). Differential Contribution of Ca²⁺-Dependent Mechanisms to Hyperexcitability in Layer V Neurons of the Medial Entorhinal Cortex. *Frontiers in Cellular Neuroscience*, 11,182. [PubMed: 28713246]
- MacDonald CJ, Lepage KQ, Eden UT, & Eichenbaum H (2011). Hippocampal “time cells” bridge the gap in memory for discontinuous events. *Neuron*, 71(4), 737–749. [PubMed: 21867888]
- Major G, & Tank D 2004 Persistent neural activity: prevalence and mechanisms. *Current Opinion in Neurobiology*, 14(6), 675–684. [PubMed: 15582368]
- Miller EK, Erickson CA, & Desimone R (1996). Neural mechanisms of visual working memory in prefrontal cortex of the macaque. *Journal of Neuroscience*, 16(16), 5154–5167. [PubMed: 8756444]
- Montell C, Birnbaumer L, & Flockerzi V (2002). The TRP channels, a remarkably functional family. *Cell*, 108(5), 595–598. [PubMed: 11893331]
- Mrejeru A, Wei A, & Ramirez JM (2011). Calcium-activated non-selective cation currents are involved in generation of tonic and bursting activity in dopamine neurons of the substantia nigra pars compacta. *Journal of Physiology*, 589, 2497–2514. [PubMed: 21486760]
- Navaroli VL, Zhao Y, Boguszewski P, & Brown TH (2012). Muscarinic receptor activation enables persistent firing in pyramidal neurons from superficial layers of dorsal perirhinal cortex. *Hippocampus*, 22(6), 1392–1404. [PubMed: 21956787]
- Neuner SM, Wilmott LA, Hope KA, Hoffmann B, Chong JA, Abramowitz J, Birnbaumer L, O'Connell KM, Tryba AK, Greene AS, Savio Chan C, & Kaczorowski CC (2015). TRPC3 channels critically regulate hippocampal excitability and contextual fear memory. *Behavioural Brain Research*, 281, 69–77. [PubMed: 25513972]

- Nilius B, Prenen J, Droogmans G, Voets T, Vennekens R, Freichel M, Wissenbach U, & Flockerzi V (2003). Voltage dependence of the Ca²⁺-activated cation channel TRPM4. *Journal of Biological Chemistry*, 278(33), 30813–30820. [PubMed: 12799367]
- Pastalkova E, Itskov V, Amarasingham A, & Buzsaki G (2008). Internally generated cell assembly sequences in the rat hippocampus. *Science*, 321(5894), 1322–1327. [PubMed: 18772431]
- Perez-Leighton CE, Schmidt TM, Abramowitz J, Birnbaumer L, & Kofuji P (2011). Intrinsic phototransduction persists in melanopsin-expressing ganglion cells lacking diacylglycerol-sensitive TRPC subunits. *European Journal of Neuroscience*, 33(5), 856–867. [PubMed: 21261756]
- Phelan KD, Shwe UT, Abramowitz J, Wu H, Rhee SW, Howell MD, Gottschall PE, Freichel M, Flockerzi V, Birnbaumer L, & Zheng F (2013). Canonical transient receptor channel 5 (TRPC5) and TRPC4 contribute to seizure and excitotoxicity by distinct cellular mechanisms. *Molecular Pharmacology*, 83(2), 429–438. [PubMed: 23188715]
- Ramsey IS, Delling M, & Clapham DE (2006). An introduction to TRP channels. *Annual Review of Physiology*, 68, 619–647.
- Reboreda A, Raouf R, Alonso A, & Séguéla P (2007) Development of cholinergic modulation and graded persistent activity in layer v of medial entorhinal cortex. *Journal of Neurophysiology*, 97(6), 3937–3947. [PubMed: 17442765]
- Reboreda A, Jiménez-Díaz L, & Navarro-López JD (2011). TRP channels and neural persistent activity. *Advances in Experimental Medicine and Biology*, 704, 595–613. [PubMed: 21290318]
- Riccio A, Li Y, Moon J, Kim KS, Smith KS, Rudolph U, Gapon S, Yao GL, Tsvetkov E, Rodig SJ, Van't Veer A, Meloni EG, Carlezon WA Jr., Bolshakov VY, & Clapham DE (2009). Essential role for TRPC5 in amygdala function and fear-related behavior. *Cell*, 137(4), 761–772. [PubMed: 19450521]
- Riccio A, Li Y, Tsvetkov E, Gapon S, Yao GL, Smith KS, Engin E, Rudolph U, Bolshakov VY, & Clapham DE (2014). Decreased anxiety-like behavior and Gq/11-dependent responses in the amygdala of mice lacking TRPC4 channels. *The Journal of Neuroscience*, 34(10), 3653–3667. [PubMed: 24599464]
- Romo R, Brody CD, Hernández A, & Lemus L (1999). Neuronal correlates of parametric working memory in the prefrontal cortex. *Nature*, 399(6735), 470–473. [PubMed: 10365959]
- Roth FC, Beyer KM, Both M, Draguhn A, & Egorov AV (2016). Downstream effects of hippocampal sharp wave ripple oscillations on medial entorhinal cortex layer V neurons in vitro. *Hippocampus*, 26(12), 1493–1508. [PubMed: 27479916]
- Sawamura S, Shirakawa H, Nakagawa T, Mori Y, & Kaneko S (2017). *Neurobiology of TRP Channels*. (2nd ed.). Boca Raton (FL): CRC Press/Taylor & Francis; Chapter 16. TRP Channels in the Brain: What Are They There For?.
- Stowers L, Holy TE, Meister M, Dulac C, & Koentges G (2002). Loss of sex discrimination and male-male aggression in mice deficient for TRP2. *Science*, 295(5559), 1493–1500. [PubMed: 11823606]
- Tahvildari B, Alonso AA, & Bourque CW (2008). Ionic basis of ON and OFF persistent activity in layer III lateral entorhinal cortical principal neurons. *Journal of Neurophysiology*, 99(4), 2006–2011. [PubMed: 18256167]
- Tahvildari B, Fransén E, Alonso AA, & Hasselmo ME (2007). Switching between “On” and “Off” states of persistent activity in lateral entorhinal layer III neurons. *Hippocampus*, 17(4), 257–263. [PubMed: 17315198]
- Tai C, Hines DJ, Choi HB, & MacVicar BA (2011). Plasma membrane insertion of TRPC5 channels contributes to the cholinergic plateau potential in hippocampal CA1 pyramidal neurons. *Hippocampus*, 21(9), 958–967. [PubMed: 20865744]
- von Bohlen und Halbach O, Hinz U, Unsicker K, & Egorov AV (2005). Distribution of TRPC1 and TRPC5 in medial temporal lobe structures of mice. *Cell and Tissue Research*, 322(2), 201–206. [PubMed: 16044320]
- Wu LJ, Sweet TB, & Clapham DE (2010). International Union of Basic and Clinical Pharmacology. LXXVI. Current progress in the mammalian TRP ion channel family. *Pharmacological Reviews*, 62(3), 381–404. [PubMed: 20716668]

- Xue T, Do MT, Riccio A, Jiang Z, Hsieh J, Wang HC, Merbs SL, Welsbie DS, Yoshioka T, Weissgerber P, Stolz S, Flockerzi V, Freichel M, Simon MI, Clapham DE, & Yau KW (2011). Melanopsin signalling in mammalian iris and retina. *Nature*, 479(7371), 67–73. [PubMed: 22051675]
- Yan HD, Villalobos C, & Andrade R (2009). TRPC channels mediate a muscarinic receptor-induced afterdepolarization in cerebral cortex. *The Journal of Neuroscience*, 29(32), 10038–10046. [PubMed: 19675237]
- Yoshida M, & Hasselmo ME (2009). Persistent firing supported by an intrinsic cellular mechanism in a component of the head direction system. *Journal of Neuroscience*, 29(15), 4945–4952. [PubMed: 19369563]
- Zhang Z, Reboreda A, Alonso A, Barker PA, & Séguéla P (2011). TRPC channels underlie cholinergic plateau potentials and persistent activity in entorhinal cortex. *Hippocampus*, 21(4), 386–397. [PubMed: 20082292]
- Zylberberg J, & Strowbridge BW (2017). Mechanisms of Persistent Activity in Cortical Circuits: Possible Neural Substrates for Working Memory. *Annual Review of Neuroscience*, 40, 603–627.




Figure 1.

CCh-induced persistent firing in TRPC1/4/5-triple-knockout mice.

Responses of mEC LV neuron to depolarizing current steps of different duration (left).

Current pulse of equivalent strength was not able to elicit persistent firing after slight membrane hyperpolarization (right). Recording was obtained during blockade of neurotransmission with CNQX (10 μ M), APV (30 μ M) and picrotoxin (100 μ M).





Figure 2. qPCR analysis of the *Trpc* subunits. (a) qPCR analysis of the *Trpc* subunits made by RNA isolated from the entorhinal cortex (EC) together with the perirhinal cortex (PER) and the postrhinal cortex (POR). Quantitative expression analysis was performed using high efficiency probe-based assays. Data are given as mean \pm SD (four animals). (b) qPCR analysis using SYBR green I assays specifically designed to verify the respective *Trpc* deletion in brain tissue of hepta-TRPC KO mice. The primer efficiency varied between assays and accordingly the level of expression is displayed as Cq values. Each bar represents the value of one animal.

**Figure 3.**

Muscarinic-dependent persistent activity in hepta-TRPC knockout mice.

(a) CCh-induced persistent firing in mEC LV neuron in hepta-TRPC KO mice (left) and its complete block by the muscarinic antagonist atropine (1 μ M; right). (b) Activity-dependence of persistent firing in hepta-TRPC KO mice. Responses of mEC LV neuron to depolarizing current steps of different duration. (c) Voltage-dependence of persistent firing in hepta-TRPC KO mice. Responses to depolarizing current steps at two different membrane potentials. The arrowhead indicates d.c. shift. (d) Box plots of firing frequency during 1 s

current-step injection (initiation firing) and during persistent activity. No significant differences between WT control and hepta-TRPC KO mice were observed. (e) Frequency of persistent firing correlated positively with frequency of spike trains during current-step injections for WT control (black) and KO mice (red). Plotted data from d. Dotted lines represent the 95% confidence intervals. (f) Box plots of membrane depolarisation during persistent firing for WT control and KO mice. Box plots indicating median, 25th and 75th percentiles and individual values. Whiskers show 5th and 95th percentiles, squares indicate mean. ns, not significant, t-test. All recordings were obtained during blockade of neurotransmission with CNQX, APV and picrotoxin.




Figure 4.

Induction and termination of persistent firing in hepta-TRPC knockout mice. (a) An example of a neurons that showed voltage-dependent persistent firing in response a brief stimulus (0.2 s, +0.4 nA). The arrowhead indicates d.c. shift. (b) Persistent firing termination by prolonged hyperpolarizing current pulse (6 s, -0.6 nA). All recordings were performed in the presence of CCh, CNQX, APV and picrotoxin.




Figure 5.

Graded persistent activity in hepta-TRPC knockout mice.

(a) Repetitive stimulation with a 1 s depolarizing current step gives rise to four distinct increases of stable discharge rate in mEC LV neuron in hepta-TRPC KO mice. (b) Repetitive application of 6 s hyperpolarizing steps gives rise to discrete decreases of firing rate. Short intervals of firing are shown at an expanded time scale for each level below voltage traces. The lower diagrams correspond to the peristimulus histograms (bin width in (a) and (b): 300 ms and 700 ms respectively). All recordings were performed in the presence of CCh, CNQX, APV and picrotoxin.




Figure 6. Activity dependent modulation of graded persistent firing in hepta-TRPC knockout mice. (a) Specimen trace showing that repetitive depolarizing current step gives initially increases of stable discharge rate followed, however, by its discrete decreases and full termination. The lower diagram correspond to the peristimulus histograms (bin width 700 ms). (b) Repetitive stimulation in same neuron after adding atropine. Recording was performed in the presence of CCh, CNQX, APV and picrotoxin.

TABLE 1

Primer sequences used for qPCR analysis with FastStart Essential DNA Probes

| | | primer sequence | probe |
|-----------------|-----|---------------------------|-------|
| TRPC1 | Fw | agggtacctccaagagcag | 105 |
| | Rev | ccaatgaacgagtggaaggt | |
| TRPC2 | Fw | tcctgtcttcctcggagtc | 52 |
| | Rev | ttcacagatagggcactggac | |
| TRPC3 | Fw | aagttcatggftctctcattatgg | 72 |
| | Rev | gaataaagtatgaacatgccaatca | |
| TRPC4 | Fw | ggaatcatgggacatgtgg | 53 |
| | Rev | cggaggggaactgaagatgttt | |
| TRPC5 | Fw | gtggattcacggaatacatcc | 31 |
| | Rev | ttgccaggtagaggagttc | |
| TRPC6 | Fw | cggggagagactgtcgttag | 78 |
| | Rev | gaggggctctgctctatctg | |
| TRPC7 | Fw | gtcatcatgaagtgggtctta | 93 |
| | Rev | tectcccagatctcttgc | |
| Reference genes | | | |
| | | primer sequence | probe |
| H3F3A | Fw | gccatcttcaattgtgttcg | 19 |
| | Rev | agccatggtaaggacacctc | |
| AIP | Fw | cgctgtgtaagcagaga | 29 |
| | Rev | aagcgactttgtctect | |
| CXXC1 | Fw | tagtgccgaccgtgact | 26 |
| | Rev | ggcctcctccctaactgaat | |

TABLE 2

Primer sequences used for qPCR analysis with iTaq™ Universal SYBR® Green

| | | primer sequence |
|-----------------|-----|-------------------------------|
| TRPC1 | Fw | acctttgccctcaaagtgt |
| | Rev | gcccaaatagagctgggtg |
| TRPC2 | Fw | cgctccaagctctacctgc |
| | Rev | ggatggcaggatgttaaagg |
| TRPC3 | Fw | cctgctttaccacggftga |
| | Rev | ctgtcatcctcgatctcttgg |
| TRPC4 | Fw | ggaatcatgggacatgtgg |
| | Rev | cggagggaaactgaagatgttt |
| TRPC5 | Fw | gtggattcacgaatacatcc |
| | Rev | ttgccaggtagaggagttc |
| TRPC6 | Fw | tgacaaaagtcacactgggg |
| | Rev | ggatcagtagggteccactt |
| TRPC7 | Fw | getcatcatgaagtgggtctta |
| | Rev | tectcccagatctcttgc |
| Reference genes | | |
| | | primer sequence |
| H3F3A | Fw | gccatcttcaattgtgttcg |
| | Rev | accagtcacccaagagg |
| AIP | Fw | aggcgatggcgtcatagta |
| | Rev | aagcgagctttgtctect |
| CXXC1 | Fw | tagtgccgaccgtgact |
| | Rev | ggcctctcccctaactgaat |

TABLE 3

The quantitative parameters of persistent activity in mEC LV neurons of TRPC1/4/5 knockout mice, and hepta-TRPC knockout versus WT control mice

| mEC LV | TRPC1/4/5 KO | | hepta-TRPC KO | | vs. | | WT control | |
|-----------------------------------|--------------|----------|-------------------|----------|----------|---------------|-----------------|----------|
| | | <i>n</i> | | <i>n</i> | <i>p</i> | <i>test</i> | | <i>n</i> |
| 1 s current step | | | | | | | | |
| Persistent firing (neurons) | 8 out of 8 | | 8 out of 11 | | | | 6 out of 6 | |
| Initiation firing (Hz) | 44.5 ± 5.0 | 8 | 32.9 ± 5.8 | 8 | 0.113 | <i>t-test</i> | 37.4 ± 3.3 | 6 |
| Persistent firing (Hz) | 9.0 ± 3.6 | 8 | 6.7 ± 3.7 | 8 | 0.253 | <i>t-test</i> | 8.7 ± 2.3 | 6 |
| Post. stim. depolarization (mV) | 4.1 ± 1.3 | 8 | 4.4 ± 1.4 | 8 | 0.963 | <i>t-test</i> | 4.4 ± 0.8 | 6 |
| Recording membrane potential (mV) | -59.8 ± 3.1 | 8 | -61.2 ± 3.0 | 8 | 0.373 | <i>t-test</i> | -59.9 ± 1.9 | 6 |
| 2 s current step | | | | | | | | |
| Persistent firing (neurons) | 7 out of 7 | | 9 out of 10 | | | | 4 out of 4 | |
| Initiation firing (Hz) | 40.7 ± 5.1 | 7 | 32 (29; 34) | 9 | 0.534 | <i>U-test</i> | 33 (31; 35) | 4 |
| Persistent firing (Hz) | 8.7 ± 2.3 | 7 | 7.4 ± 4.2 | 9 | 0.825 | <i>U-test</i> | 5.2 (4.6; 6.7) | 4 |
| Post. stim. depolarization (mV) | 5.4 ± 1.3 | 7 | 5.0 ± 2.2 | 9 | 0.999 | <i>U-test</i> | 4.8 (4.0; 6.2) | 4 |
| 3 s current step | | | | | | | | |
| Persistent firing (neurons) | 5 out of 5 | | 4 out of 5 | | | | 4 out of 4 | |
| Initiation firing (Hz) | 34.9 ± 7.7 | 5 | 32.5 (29.8; 33.3) | 4 | 0.772 | <i>U-test</i> | 31 (29.5; 32.4) | 4 |
| Persistent firing (Hz) | 10.1 ± 2.7 | 5 | 6.2 (4.8; 7.8) | 4 | 0.886 | <i>U-test</i> | 6.8 (6.0; 7.1) | 4 |
| Post. stim. depolarization (mV) | 4.3 ± 1.8 | 5 | 4.9 (4.4; 6.3) | 4 | 0.885 | <i>U-test</i> | 5.9 (5.2; 6.7) | 4 |

Averaged data are given as mean ± SD or as median (P25; P75). P values for hepta-TRPC KO vs. WT control (Unpaired two-tailed t-test or Mann-Whitney U-test).