

CHANGES IN SYNAPTIC TRANSMISSION AND PROTEIN EXPRESSION IN THE BRAINS OF ADULT OFFSPRING AFTER PRENATAL INHIBITION OF THE KYNURENINE PATHWAY

C. M. FORREST,^a O. S. KHALIL,^a M. PISAR,^a
K. MCNAIR,^a E. KORNISIUK,^b M. SNITCOFSKY,^b
N. GONZALEZ,^b D. JERUSALINSKY,^{b,c}
L. G. DARLINGTON^d AND T. W. STONE^{a*}

^a *Institute of Neuroscience and Psychology, West Medical Building, University of Glasgow, Glasgow G12 8QQ, UK*

^b *Instituto de Biología Celular & Neurociencias “Prof Eduardo De Robertis”, Facultad de Medicina, University of Buenos Aires and CONICET, Buenos Aires, Argentina*

^c *CBC, Universidad de Buenos Aires, Buenos Aires, Argentina*

^d *Epsom General Hospital, Dorking Road, Epsom, Surrey KT18 7EG, UK*

Abstract—During early brain development, N-methyl-D-aspartate (NMDA) receptors are involved in cell migration, neurogenesis, axon guidance and synapse formation, but the mechanisms which regulate NMDA receptor density and function remain unclear. The kynurenine pathway of tryptophan metabolism includes an agonist (quinolinic acid) and an antagonist (kynurenic acid) at NMDA receptors and we have previously shown that inhibition of the pathway using the kynurenine-3-monooxygenase inhibitor Ro61-8048 in late gestation produces rapid changes in protein expression in the embryos and effects on synaptic transmission lasting until postnatal day 21 (P21). The present study sought to determine whether any of these effects are maintained into adulthood. After prenatal injections of Ro61-8048 the litter was allowed to develop to P60 when some offspring were euthanized and the brains removed for examination. Analysis of protein expression by Western blotting revealed significantly reduced expression of the GluN2A subunit (32%) and the morphogenetic protein sonic hedgehog (31%), with a 29% increase in the expression of doublecortin, a protein associated with neurogenesis. No changes were seen in mRNA abundance using quantitative real-time polymerase chain reaction. Neuronal excitability was normal in the CA1 region of hippocampal slices but paired-pulse stimulation revealed less inhibition at short interpulse inter-

vals. The amount of long-term potentiation was decreased by 49% in treated pups and recovery after low-frequency stimulation was delayed. The results not only strengthen the view that basal, constitutive kynurenine metabolism is involved in normal brain development, but also show that changes induced prenatally can affect the brains of adult offspring and those changes are quite different from those seen previously at weaning (P21). Those changes may be mediated by altered expression of NMDAR subunits and sonic hedgehog. © 2013 IBRO. Published by Elsevier Ltd. All rights reserved.

Key words: kynurenines, kynurenic acid, tryptophan, neurodevelopment, doublecortin, hedgehog.

INTRODUCTION

There is increasing interest in the role of epigenetic modification in the early development of the nervous system. A wide range of factors can modify brain development after fetal or early postnatal exposure, including diet, stress, therapeutically used medicines and environmental agents. There is relatively little information, however, on the mechanisms by which such external influences induce changes in brain development.

One possibility is that changes are induced in the activation of glutamate receptors for N-methyl-D-aspartate (NMDA), since these are known to be intimately involved in early phases of brain development. NMDA receptors play key roles in the initial formation and guidance of axon branches, the establishment and stabilization of synaptic contacts and the induction and maintenance of dendritic spines (Simon et al., 1992; Rajan and Cline, 1998; Ernst et al., 1998; Heng et al., 1999; Cuppini et al., 1999; Udin and Grant, 1999; Colonnese et al., 2005; Alvarez et al., 2007; Ultanir et al., 2007). These and other aspects of neuronal and synaptic development ultimately determine synaptic function and plasticity in the mature, postnatal, offspring (Iwasato et al., 2000; Myers et al., 2000; Ramoa et al., 2001; Fagiolini et al., 2003).

In addition, antagonists acting at NMDA receptors prevent many of these neuro-developmental processes and, when administered during late fetal or early postnatal life, increase the natural loss of neurons and synapses (Ikonomidou et al., 1999; Dikranian et al., 2001; Harris et al., 2003; Vincent et al., 2004).

*Corresponding author. Tel: +44-(0)141-330-4481; fax: +44-(0)141-330-5481.

E-mail address: Trevor.Stone@glasgow.ac.uk (T. W. Stone).

Abbreviations: aCSF, artificial cerebrospinal fluid; ANOVA, analysis of variance; CED, Cambridge Electronic Design; COX-2, cyclo-oxygenase-2; DISC-1, Disrupted in Schizophrenia-1; fEPSP, field excitatory postsynaptic potential; HRP, horseradish peroxidase; IDO, indoleamine-2,3-dioxygenase; KMO, kynurenine-3-monooxygenase; LTP, long-term potentiation; LTD, long-term depression; NMDA, N-methyl-D-aspartate; P, postnatal day; PCNA, Proliferating Cell Nuclear Antigen; PPF, paired-pulse facilitation; PPI, paired-pulse inhibition; PSD-95, Post-Synaptic Density molecule-95; qt-PCR, quantitative real-time polymerase chain reaction; Ro61-8048, 3,4-dimethoxy-N-[4-(3-nitrophenyl)thiazol-2-yl]-benzenesulfonamide; TBST, Tris-buffered saline containing 0.05% Tween; TH, tyrosine hydroxylase; VAMP-1, Vesicle-Associated Membrane Protein-1.

The neuronal and synaptic disruption produced by exogenously administered NMDA receptor antagonists results in profound abnormalities of neuronal development, brain structure and behavior reminiscent of those seen in schizophrenia (Harris et al., 2003; du Bois and Huang, 2007). However, previous studies do not address the question of whether manipulating physiological, endogenous factors affecting NMDA receptor function could also result in disordered neuronal development.

One potential method for modulating the activation of NMDA receptors by endogenous ligands is to interfere with the kynurenine pathway. This pathway is the major route for the metabolism of tryptophan and generates quinolinic acid, a selective agonist at NMDA receptors (Stone and Perkins, 1981; Stone and Darlington, 2002) as well as kynurenic acid, which is an antagonist at all ionotropic glutamate receptors, though with greatest potency at NMDA receptors (Perkins and Stone, 1982; Stone et al., 2013). Kynurenic acid may also block nicotinic receptors in the CNS (Hilmas et al., 2001) although this has been disputed (Mok et al., 2009; Dobelis et al., 2012). The ratio between the endogenous levels of quinolinic acid and kynurenic acid will, therefore, influence neuronal excitability and viability.

In this study we have used an inhibitor of kynurenine-3-monoxygenase (KMO) to alter the relative concentrations of endogenous quinolinic acid and kynurenic acid. Previous work has shown that the major effect of inhibition by 3,4-dimethoxy-N-[4-(3-nitrophenyl)thiazol-2-yl]-benzenesulfonamide (Ro61-8048; Röver et al., 1997) is to raise the levels of kynurenic acid in the blood and brain (Röver et al., 1997; Cozzi et al., 1999; Clark et al., 2005; Forrest et al., 2013). The compound was administered to gestating female rats and the offspring were allowed to develop normally until day 60 (P60). The effects of the exposure to Ro61-8048 were then examined on hippocampal synaptic transmission and plasticity in the brains of those P60 animals. It is around this age that many studies have shown behavioral changes resulting from prenatal infection or mimetic agents (Iwasato et al., 2000; Fatemi et al., 2005; Zuckerman and Weiner, 2005) and we therefore tested behavior in the open-field test of exploration and a step-down inhibitory avoidance task of learning. Since many of the neurodevelopmental functions of NMDA receptors are probably mediated via their interactions with cell proliferation, axonal guidance and cytoskeletal molecules which are crucial to the development and maintenance of synaptic contacts (Ozaki et al., 2000; Hoffman et al., 2001; Ramoa et al., 2001; Fagiolini et al., 2003), we have also included an examination of the expression of representative proteins as well as the mRNA for key proteins by quantitative real-time polymerase chain reaction (qRT-PCR). The target molecules examined are all known to play key roles in neuronal development, neurite outgrowth, synapse formation and neurotransmitter release. They include components of the synaptic vesicle and neurotransmitter release machinery such as

synaptophysin, Vesicle-Associated Membrane Protein-1 (VAMP-1; synaptobrevin) and the vesicular release calcium sensor synaptotagmin. As markers of axon guidance and synapse formation, the proteins EphA4, Unc5H1 and Unc5H3 were studied, in addition to the cytoskeletal modulators RhoA and RhoB, the glutamate complex component Post-Synaptic Density molecule-95 (PSD-95) and tyrosine hydroxylase (TH).

We have also examined the expression of two proteins that might be important indicators of the basic mechanisms underlying the observed changes: the interneuronal maturation molecule doublecortin and the morphogenetic protein sonic hedgehog.

Finally, whereas the juvenile (P21) study was performed on whole cerebral hemispheres, the present, more detailed analysis of adults at P60 was focussed on the hippocampus.

EXPERIMENTAL PROCEDURES

This study was carried out according to the regulations of the Animals (Scientific Procedures) act 1986 of the UK, administered and monitored by the Home Office. Male and female Wistar rats were housed together for mating and inspected daily for the occurrence of a vaginal plug. Thereafter, the pregnant females were housed alone with free access to food and water.

To inhibit tryptophan oxidation along the kynurenine pathway we used Ro61-8048 (Röver et al., 1997). This compound is an inhibitor of KMO, a key enzyme in the pathway which shifts the balance of tryptophan metabolism away from the generation of the NMDA receptor agonist, quinolinic acid, toward the antagonist, kynurenic acid (Cozzi et al., 1999; Clark et al., 2005; Forrest et al., 2013). From those earlier studies we identified the dose of 100 mg/kg (i.p.) as one which can be administered repeatedly to the same animal (Clark et al., 2005; Rodgers et al., 2009). In order to maximize the period of development during which the activity of the kynurenine pathway is affected, we administered this compound to the pregnant dam at days E14, E16 and E 18 of gestation. Groups of control animals were injected with the saline vehicle. In most experiments, gestation was allowed to proceed normally, with neonates being removed from the home cage for euthanasia followed by removal of the brain when adult at postnatal day (P60). In an earlier study of postnatal animals at the time of weaning (P21) we confirmed that prenatal exposure to Ro61-8048 had the predicted effects on levels of kynurenine and kynurenic acid, concentrations of both being increased between 10 and 100-fold in the maternal blood and embryo brains (Forrest et al., 2013).

Electrophysiology

Electrophysiological studies were performed on male and female animals which were allowed to wean and grow under normal conditions to 60 days of age (P60 animals) with food and water available *ad libitum*. Animals were killed by administration of an overdose of

urethane (2.5 g/kg rat delivered as an i.p. injection of a 25% solution in water) followed by cervical dislocation. The brain was removed into ice-cold artificial cerebrospinal fluid (aCSF) of composition: (in mM) NaCl 115; KH₂PO₄ 2.2; KCl 2; MgSO₄ 1.2; NaHCO₃ 25; CaCl₂ 2.5; glucose 10, gassed with 5% CO₂ in oxygen. The hippocampi were rapidly removed and chopped into 450- μ m transverse slices using a Mcllwain tissue chopper. The slices were preincubated at room temperature for at least 1 h in a water-saturated atmosphere of 5% CO₂ in O₂ before individual slices were transferred to a 1-ml capacity superfusion chamber for recording. Slices were superfused at 28–30 °C using aCSF at a flow rate of 3–4 ml/min. A concentric bipolar electrode was used for stimulation of the Schaffer collateral and commissural fibers in stratum radiatum, using stimuli delivered at 0.1 Hz or 0.05 Hz with a pulse width of 50–300 μ s, adjusted to evoke a response amplitude of approximately 70% of maximum to allow increases or decreases in size to be detected. Extracellular recordings were made via glass microelectrodes containing 1 M NaCl (tip diameter approximately 2 μ m, DC resistances 2–5 M) with the tip positioned under microscopic visualization in the stratum radiatum of the CA1 region to evoke field excitatory postsynaptic potentials (fEPSPs). Potentials were amplified, digitized and stored on a computer via a CED (Cambridge Electronic Design, Cambridge, UK) micro1401 interface. The fEPSPs were routinely quantified by measurement of the early positive slope of the potential, using Signal software (CED, Cambridge, UK). The axonal volley was monitored wherever it was possible to distinguish it clearly from the fEPSP in order to ensure that no change occurred during the experiments.

Once placed into the recording chamber, the recording of fEPSPs was allowed to stabilize and a minimum period of 10 min obtained at a stable baseline. The degree of long-term potentiation (LTP) and long-term depression (LTD) was quantified by measuring the size of the evoked potentials once a post-stimulation plateau had been obtained and comparing with the size of the potential before stimulation.

Paired-pulse interactions were assessed in slices not used for the examination of LTP or LTD, using pairs of stimuli S1 and S2 with inter-stimulus intervals of 10–100 ms. In the case of fEPSPs the 10-ms interval resulted in a substantial overlap between successive potentials and an electronic subtraction was performed in which a single evoked potential at time S1 was subtracted from a subsequent paired-pulse response to reveal the true magnitude of the response to S2.

qRT-PCR

Total RNA was isolated from snap frozen, whole hippocampi removed from P60 rats (as previously described, $n = 3$ /litter, $n = 3$ litters/treatment group, $n = 9$ control and $n = 9$ treated) using the Qiagen RNeasy lipid mini kit with DNaseI treatment, according to the manufacturer's instructions. The RNA

concentration of each sample was determined using a NanoDrop (Thermo Scientific) and the integrity of each mRNA sample was assessed using the Agilent 2100 Bioanalyser (Glasgow Polyomics Facility, University of Glasgow). Only samples with RNA integrity number (RIN) values greater than 8 were used (values ranged from 8.70 to 9.30).

First strand cDNAs were synthesized from total RNA using random hexamers and Superscript III Reverse transcriptase in the VILO cDNA synthesis kit (Life Technologies) following the manufacturer's instructions.

qRT-PCR was carried out in an ABI cyclor (SDS7000) using the TaqMan Gene expression assay system for the following assays: doublecortin (*Dcx*; Rn00670392_m1), sonic hedgehog (*Shh*; Rn00568129_m1), NMDA receptor subunit 2A (*Grin2a*; Rn00561341_m1) and NMDA receptor subunit 2B (*Grin2b*; Rn00680474_m1), (Applied Biosystems). In addition, four reference genes were used: beta actin (*Actb*; Rn00667869_m1), glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*; Rn01775763_g1), beta-2 microglobulin (*B2m*; Rn00560865_m1) and 18S ribosomal RNA (*Rn18s*; Mm03928990_g1) (Applied Biosystems).

Immunoblotting

Brain sample homogenates were prepared in RIPA buffer (50 mM Tris, 150 mM NaCl, 0.1% SDS, 0.5% Triton X-100, 1% IGEPAL, and a Roche complete protease inhibitor tablet) and centrifuged at 18,000g for 5 min at 4 °C. Supernatants were collected for protein concentration determination using the Bio-Rad Coomassie Blue protein assay (Bio-Rad, Hemel Hempstead, UK). Samples were then normalized to 10 μ g and prepared as; 65% protein sample, 25% sample buffer and 10% reducing agent (Life Technologies, Paisley, UK) and heated at 70 °C for 10 min. The protein samples were loaded onto NuPAGE Novex 4–12% Bis-Tris (1.0 mm) 15 lane gels (Life Technologies, Paisley, UK) and run at 150 V for 80 min to separate proteins according to their molecular weight. SeeBlue pre-stained standard (Life Technologies, Paisley, UK) was included on each gel as a molecular weight marker. The separated proteins were then blotted onto Invitrolon poly(vinylidene difluoride) membranes (Life Technologies, Paisley, UK) at 30 V for 60 min. The membranes were blocked for 1 h in 5% non-fat dried milk solution in Tris-buffered saline containing 0.05% Tween (TBST) before overnight incubation at 4 °C with the appropriate primary antibody (diluted in 5% milk-TBST). Membranes were then washed three times for 15 min with TBST and incubated with the appropriate horseradish peroxidase (HRP) conjugated secondary antibody (prepared in 5% milk-TBST) for 1 h at room temperature. Following secondary antibody incubation, blots were washed 3 times for 15 min with TBST then visualized using Enhanced Chemiluminescence Plus detection kit (GE Healthcare, Chalfont St Giles, UK).

Western blot analysis was carried out using the following primary antibodies:

From Millipore, Watford, UK: GluN1 (mouse monoclonal, 05-432, 1:1000 dilution); TH (mouse monoclonal, MAB5280, 1:10000 dilution), synaptophysin (mouse monoclonal, MAB368, 1:40000 dilution);

From R&D Systems, Abingdon, UK: GluN2A (rabbit polyclonal, PPS012, 1:5000 dilution); GluN2B (rabbit polyclonal, PPS013, 1:5000); VAMP-1/synaptobrevin (goat polyclonal, AF4828, 1: 10,000 dilution), synaptotagmin (MAB 43641, 1:5000 dilution);

From Cell Signalling, New England Biolabs, Hitchin, Herts, UK: PSD-95 (rabbit monoclonal, #3450, 1:10,000 dilution);

From Santa Cruz, Insight Biotechnology, Wembley, UK: doublecortin (goat polyclonal, sc-8066, 1:1000 dilution); α -synuclein (mouse monoclonal, sc-65500, 1:1000 dilution); actin (goat polyclonal, sc-1615, 1:10,000 dilution); Disrupted in Schizophrenia-1 (DISC-1) (goat polyclonal, sc-47990, 1:1000 dilution); Unc5H1 (goat polyclonal, sc-67902, 1:1000 dilution), Unc5H3 (goat polyclonal, sc-54442, 1:1000 dilution), SHH (goat polyclonal, sc-1194, 1:1000 dilution); NF κ B (rabbit polyclonal, sc-372, 1:5000 dilution); cyclo-oxygenase-2 (COX-2) (goat polyclonal, sc-1745, 1:1000 dilution); RhoA (mouse monoclonal, sc-418, 1:1,000 dilution); RhoB (mouse monoclonal, sc-8048, 1:1000 dilution); EphA4 (rabbit polyclonal, sc-921, 1:5000); Proliferating Cell Nuclear Antigen (PCNA) (mouse monoclonal, sc-56, 1:1000 dilution); 5HT-2c (mouse monoclonal, sc-17797, 1:1000 dilution).

The following secondary HRP-conjugated antibodies were used at 1:5000 dilution: goat anti-rabbit HRP (12-348) (Millipore, Watford, UK); donkey anti-goat HRP (sc-2020) and goat anti-mouse (sc-2005) (Santa Cruz, Insight Biotechnology, Wembley, UK).

Behavior

All experiments were performed in accordance with the Review Committee of the Veterinary School (CICUAL), University of Buenos Aires and the International Brain Research Organization (IBRO), and are in compliance with the US National Institutes of Health Guide for Care and Use of Laboratory Animals (Publication No. 85-23, revised 1985) and the European Communities Council Directive of 24 November 1986 (86/609/EEC).

Wistar rats were treated as previously described (see Methods above); pregnant Wistar rats were injected i.p. either with Ro61-8048 or the saline vehicle. At P21 female and male rats (35–40 g) bred in-house were separated and maintained in groups of 5–6 per cage, under a 12-h light/dark inverted cycle with water and food available *ad libitum* until P60 (180–220 g).

Open field test of locomotor and exploratory activities. To evaluate possible changes of locomotor activity and/or exploratory behavior, P60 female and male rats – born from mothers treated with Ro61-8048 or saline – were initially allowed to freely explore an open square field (measuring 60.0 cm long \times 60.0 cm wide \times 50.0 cm high) for 5 min (OF1). The floor of this arena was divided into 16 sectors of 15.0 cm \times 15.0 cm

each, and the number of rearings, groomings and crossings from one sector to another per minute was counted in the initial 5-min session (OF1) (carried out at about 3 h before avoidance training); the second 5-min session (OF2) was performed in the following day, about 2 h after the avoidance test, between 27 and 28 h from the first OF session.

Step-down inhibitory avoidance. Effects on learning and memory were tested using a step-down inhibitory avoidance model of learning in rats as described previously (Harvey et al., 2012).

Rats were trained in the step-down inhibitory avoidance task at about 3 h after the first open field session. A rat was placed on an elevated isolated platform (25 \times 7 \times 2.5 cm high) at the left side of an acrylic box (50 \times 25 \times 25 cm), with the floor made of parallel bronze bars (0.5 cm caliber, 0.5 cm apart). The latency was measured for the animal to step-down from the platform, placing all four paws on an electrifiable grid (training latency), at which point the animal was given a 2.0 s, 0.75 mA scrambled foot-shock, causing it to return to the platform or remain on the grid from which it was immediately removed and returned to its home-box.

On the next day (about 20 h after the training session), a retention test was performed in which the step-down latency (test latency) was recorded up to a maximum of 300 s, but no shock was delivered. The difference between test and training latencies was taken as an indication of retention: a better memory should result in a higher test latency and a greater difference from the training latency

Data analysis and statistics

Immunoblotting. All Western blots were quantified using the Image J software (<http://rsb.info.nih.gov/ij/>) and comparisons were made statistically between groups of pups born to mothers treated with Ro61-8048 and groups born to mothers injected with saline vehicle. This protocol allowed the use of a *t*-test to examine differences between the two groups. To control for variations in the total amount of protein loaded onto gels all samples were examined after staining with the Ponceau S stain. In addition, actin levels were examined in each series of blots and the ratio taken of the intensity of target protein to the intensity of actin. A probability value of 0.05 was adopted as the criterion for significance. Actual *p*-values are indicated when > 0.0001 .

Electrophysiology. Data from hippocampal slices are presented as mean \pm standard error. Baseline values were obtained from a stable 10-min period of evoked potential size prior to the addition of any drugs, with the first of those potentials being defined as 100%. This allowed the 10-min pre-drug period to provide an indication of baseline variance. Repeated measures ANOVA (analysis of variance) were used for statistical

comparisons followed by the Bonferroni *post hoc* test for individual comparisons.

qRT-PCR. PCR Ct values and reaction efficiencies were obtained from the raw fluorescent data using LinReg PCR (Ramakers et al., 2003). TaqMan assay reaction efficiencies were *Dcx* 1.802, *Shh* 1.876, *Grin2a* 1.858, *Grin2b* 1.866, *Actb* 1.807, *Gapdh* 1.729, *B2m* 1.866 and *Rn18s* 1.781. Samples were run in triplicate and Ct outliers were determined using the Grubb's test (<http://graphpad.com/quickcalcs/Grubbs>). Significant Ct outliers ($P > 0.05$) were removed before statistical analysis. The mean Ct values for each sample were analyzed using REST 2009 v2.0.13 (Pfaffl et al., 2002). REST calculates P values on the basis of 2000 permutations of a pair-wise fixed reallocation randomisation test in which Ct values for the reference and target gene are reallocated to sample and control groups and calculates the resulting expression ratios on the basis of the mean values.

Behavior. For the behavioral tasks, non-parametric statistics were used because an upper time limit (300 s) was specified for stepping-down from the platform in the avoidance task. Results are presented as medians with interquartile ranges (Q_{25}/Q_{75}). Statistical differences between test latencies were evaluated using a Kruskal–Wallis ANOVA. When significant differences were found, training and test latencies in each group were compared by the Wilcoxon test. Latencies for both training and test sessions, and differences between them (test minus training latencies) were compared between groups (Ro61-8048 vs. saline) using the Mann–Whitney 'U' test, to evaluate the effects of drug treatment.

Data from the open-field exploration task are expressed as the medians of the number of crossings or rearings performed during a 5-min session (OF1 or OF2) by animals exposed prenatally to vehicle (OF1 control and OF2 control) or Ro61-8048 (OF1 treated and OF2 treated). Groups were compared using Kruskal–Wallis ANOVA. When significant differences were found, the numbers of crossings or rearings in each group was compared by the Wilcoxon test. Further comparison between groups (Ro61-8048 vs. vehicle) was made using the Mann–Whitney 'U' test.

RESULTS

Electrophysiology

Initial experiments on each hippocampal slice established the absolute levels of excitability by measuring the size of the fEPSP in stratum radiatum as the stimulus was increased from the threshold potential that elicited a discernible response (Fig. 1A). The slopes of the potentials were compared up to the stimulus level at which the first sign of a population spike indentation appeared in the fEPSP. There were no significant differences between any of the data points in the stimulus–response relationship between slices prepared from control animals and those from animals born to dams treated with Ro61-8048 (Fig. 1A).

A similar analysis was performed for population spike amplitude (Fig. 1B), but no significant difference was detected between the stimulus response curves for control and Ro61-8048-exposed animals.

In order to obtain more detailed information about the pre- and post-synaptic function of synapses in these animals, pairs of stimuli were used, at interpulse intervals ranging from 100 ms (10 Hz) to 10 ms (100 Hz), to examine the occurrence and magnitude of paired-pulse facilitation (PPF) and paired-pulse inhibition (PPI). At the 10-ms interval there was too much overlap between the evoked fEPSPs to allow a direct comparison of the first (fEPSP1) and second (fEPSP2) potentials, and the Signal software was used to perform a digital subtraction of fEPSP1 from the paired-pulse potentials (Fig. 1C, a–e). While the overall profile of the potential ratios at varying interpulse intervals are similar between slices taken from control and Ro61-8048-treated animals, there was a significant difference between the fEPSP responses of the slices at the shortest interpulse interval of 10 ms (Fig. 1C).

Digital subtraction was not necessary for the analysis of population spikes, for which the relatively rapid time course of the spike, together with the much greater degree of facilitation or inhibition, made the contribution of any overlap negligible. A similar relationship was found to that observed with fEPSPs, although control slices showed a significant degree of PPI at the 10-ms interpulse interval as exemplified by the potentials illustrated in Fig. 1D, a–d). In contrast, slices from Ro61-8048-exposed animals showed less facilitation at the 10-ms interval compared with longer interpulse intervals, but no overall PPI, leading to a highly significant difference compared with the controls animals (Fig. 1D), and an exaggeration of the difference noted above with fEPSPs.

However, it is recognized that, primarily at smaller interpulse intervals, the nature of the interaction between potential pairs of population spikes is highly dependent on the frequency with which the pulse pairs are elicited. In view of the difference between the test and control slices at the 10-ms interpulse interval, therefore, this relationship was examined further. It was found that, at increasing frequencies of paired-pulse presentation, control slices showed little change in the degree of PPI seen at 10 ms (Fig. 1C, E), whereas in slices from Ro61-8048-treated animals the initial PPF was lost over a series of 5–10 stimulus pairs, leaving PPI at higher presentation frequencies as illustrated in the potentials of Fig. 1E, a,b. The switch from facilitation to inhibition was sufficiently pronounced that a very significant difference was demonstrated at a paired-pulse frequency of 1 Hz (Fig. 1E) as well as at the lower frequencies of 0.1 and 0.05 Hz.

LTP. Two aspects of synaptic plasticity were examined in the slices – LTP and LTD. LTP was tested using a burst of theta stimulation (5 bursts per second, for 2 s, of four stimuli delivered at 100 Hz; Larson et al., 1986). In control slices this level of stimulation induced an increase of fEPSP amplitude which reached a




Fig. 1. Hippocampal function at P60 after prenatal exposure to Ro61-8048. The increase in size of (A) field excitatory postsynaptic potentials (EPSPs) and (B) population spikes (PS) as a function of stimulus current in the hippocampal CA1 region of adult (P60) rats exposed prenatally to Ro61-8048. The symbols indicate mean \pm s.e.m. ($n = 12$). There were no significant differences between any of the pairs of data points. Paired-pulse data are shown for the ratio of fEPSPs recorded in the stratum radiatum (C). The sample recordings alongside the graph illustrate the digital subtraction performed for the recordings made at an interpulse interval of 10 ms, with (a) being a normal single fEPSP, (b) two responses at the 10-ms interval, and (c) the second response alone generated by the subtraction of (a) from (b). Record (d) shows the directly superimposed traces (a) and (c) while (d) illustrates the superimposed traces adjusted to the same stimulus point. Paired pulse data for the ratio of PS recorded in stratum pyramidale are shown in (D). The sample traces alongside illustrate (a) a single PS while other traces illustrate pairs of PS at interpulse intervals of (b) 50 ms (c) 20 ms and (d) 10 ms. The change in PS ratio at different rates of paired-pulse presentation from 0.05 to 1 Hz is shown in (E). The sample traces show individual records of paired PS at presentation rates of (a) 0.05 Hz and (b) 1 Hz. Symbols in the graphs indicate mean \pm s.e.m. ($n = 12$). ** $P < 0.01$; *** $P < 0.001$ using the Bonferroni test for multiple comparisons following ANOVA.

plateau after approximately 20 min and was stable until the end of the recording period at 50 min (Fig. 2A, B). In order to compare potential size at the plateau with their pre-stimulation amplitude an analysis of variance was performed using the fEPSP slopes recorded over the 10-min period before stimulation and between 40 to 50 min after theta stimulation. In control animals this increase was statistically significant ($F_{[19,80]} = 17.462$; $p < 0.0001$, $n = 5$) with an increase of 34.8% between the mean fEPSP slope at 0 min ($98.1 \pm 3.5\%$) and that at 50 min ($133.0 \pm 5.8\%$; $p = 0.0009$, $n = 5$).

For slices prepared from animals subjected to Ro61-8048 administration *in utero*, a very similar temporal profile of LTP was obtained, but of a lower magnitude than in control slices (Fig. 2A, C). Analysis of variance confirmed that the fEPSP slopes were still significantly greater than the corresponding pre-stimulation level ($F_{[19,80]} = 2.248$; $p = 0.0066$, $n = 5$) with an increase of 17.7% between the mean fEPSP

slope at 0 min ($97.3 \pm 2.5\%$) and that at 50 min ($115.0 \pm 7.4\%$, $p = 0.05$, $n = 5$). A comparison of the fEPSP slopes of control and Ro61-8048-exposed slices between 45 and 50 min after stimulation indicated that the LTP was very significantly smaller in treated animals compared with controls ($F_{[19,80]} = 4.98$, $p < 0.0001$, $n = 5$) (Fig. 2A).

LTD. Previous studies of LTD in this laboratory have indicated that a clear depression of fEPSP slope can be obtained using a modification of the protocol published by Kemp and Bashir (1999). Stimulus trains of three pulses at 200 Hz were delivered every second for 5 min (stim1, Fig. 2D), and the same protocol repeated 20 min later (stim2). In slices from control animals, a clear LTD was obtained in which the fEPSP slope between 30–40 min after stim2 was very significantly lower than in the 10-min baseline period preceding stim1 ($F_{[19,80]} = 6.94$, $p < 0.0001$, $n = 5$ slices), with a




Fig. 2. Adult hippocampal plasticity after prenatal exposure to Ro61-8048. Panel (A) illustrates long-term potentiation in the two groups of offspring following theta-burst stimulation at the arrow. The inset records illustrate a control fEPSP immediately before stimulation and the potentiated response 40 min later in a slice from a control animal (B) or an animal exposed *in utero* to Ro61-8048 (C). The symbols indicate mean \pm s.e.m. ($n = 6$). Calibrations 0.5 mV and 10 ms. *** $P < 0.001$ for the final 5 min of recording (ANOVA). Panel (D) illustrates long-term depression in the two groups of offspring induced by two periods of low-frequency stimulation indicated by the bars on the abscissa at stim1 and stim2 (see text). The inset records illustrate a control fEPSP immediately before stim1 and the depressed responses 20 min after stim1 and 20 min after stim2 in a slice from a control animal (E) or an animal exposed *in utero* to Ro61-8048 (F). The symbols indicate mean \pm s.e.m. ($n = 6$). Calibrations 1 mV, 10 ms for the control records and 0.5 mV, 10 ms for drug-exposed slices. There were no significant differences between levels of LTD in the control and treated slices over the last 5 min of recording.

reduction of 32.0% between the fEPSP at 0 min immediately preceding stim1 ($102.7 \pm 6.1\%$) and the fEPSP at 40 min following stim2 ($69.7 \pm 5.6\%$; $p = 0.004$, $n = 5$) (Fig. 2D, E). In comparison, the slices taken from animals exposed to Ro61-8048 showed a similar degree of LTD such that the potential size between 30–40 min after stim2 was reduced very significantly ($F_{[19,80]} = 16.25$, $p < 0.0001$, $n = 5$ slices) with a reduction of fEPSP slope by 42.3% between the potentials at 0 min preceding stim1 ($105.2 \pm 3.1\%$) and the mean potential recorded 40 min after stim2 ($63.0 \pm 2.1\%$; $p < 0.0001$, $n = 5$) (Fig. 2D, F). The level of LTD between 30–40 min after stim2 was not significantly different between slices from control animals and those exposed to Ro61-8048 (ANOVA, $F_{[19,80]} = 0.106$, $p > 0.99$).

There was, however, a difference in the time course of LTD development in the two groups (Fig. 2D). Slices from Ro61-8048-treated animals recovered more slowly from stim2 stimulation, and this was borne out by a comparison of the individual fEPSP slopes between 30 and 50 min (0–20 min after stim2). All points in this

period between 31 and 37 min were significantly different at $p < 0.001$ (two-tailed t test, $n = 5$), while points between 38 and 40 min were different at $p < 0.01$.

Adenosine sensitivity. While LTP and LTD are primarily dependent on postsynaptic phenomena, adenosine acts on A1 receptors located almost exclusively on presynaptic terminals, thus providing a relatively selective test of presynaptic function (Goncalves and Queiroz, 2008; Stone et al., 2009). In order to determine whether Ro61-8048 treatment had produced any change specifically in presynaptic function, therefore, slices were exposed to 10 and 30 μ M concentrations of adenosine perfused for 10 min. This procedure reduced the slope of the fEPSP to a plateau level approximately 25% and 75% respectively below the baseline potential size in control animals (Fig. 3A–C). Using slices from animals exposed to Ro61-8048, a similar degree of reduction was obtained, the magnitude of the responses to adenosine being not significantly different from controls at either concentration tested (Fig. 3D).




Fig. 3. Sensitivity to adenosine of adult hippocampal slices after prenatal exposure to Ro61-8048. (A) Graph summarizing changes in fEPSP amplitude in response to the application of adenosine at concentrations of 10 and 30 μ M. Adenosine was superfused for 10 min indicated by the solid bar. Symbols represent responses recorded in control slices superfused with 10 μ M (filled squares, ■), or 30 μ M (filled circles, ●), and slices from animals exposed prenatally to Ro61-8048 superfused with 10 μ M (open diamonds, ◇) or 30 μ M (open triangles, △) adenosine. (B and C) Show individual sample records of fEPSPs from control (B) or Ro61-8048 exposed animals (C), recorded before and after recovery from the application of adenosine at 30 μ M, and at the peak of the response to adenosine, illustrating complete recovery after the adenosine application. (D) Summarizes the pooled data on the degree of fEPSP depression produced by adenosine at 10 and 30 μ M, illustrating the absence of any difference between slices from control and Ro61-8048 treated animals. Calibrations 1 mV, 10 ms.

qRT-PCR

REST 2009 was used to analyze relative gene expression. B2 m expression was significantly reduced in the Ro61-treated rats ($p = 0.009$). This reference gene was therefore excluded from further analysis. The reference genes used to analyze relative gene expression were Actb, Gapdh and Rn18s as these were not significantly different between control and treated rats. mRNA expression of Dcx, Shh, Grin2a and Grin2b transcripts did not change relative to Actb, Gapdh and Rn18s (Fig. 4).

Protein expression

The range of proteins examined in this study was similar to that targeted in our previous study of animals at

21 days of age (Forrest et al., 2013). They included proteins known to be involved in neuronal migration, early axon guidance, dendritogenesis and the formation of spines and synaptic contacts. Significant differences were noted in the expression of three of these proteins, in marked contrast to the variety of changes seen at P21, where nine of 20 proteins examined showed significant changes of expression.

In several cases proteins which had shown no change at P21 remained unchanged at P60. These included the presynaptic vesicular release protein synaptophysin ($p = 0.57$), the vesicle release calcium sensor synaptotagmin ($p = 0.14$) and VAMP-1/synaptobrevin ($p = 0.41$). There was also no evidence of change for the dependence receptors Unc5H1 ($p = 0.24$) and Unc5H3 ($p = 0.29$), which are dependent on the secreted paracrine protein family of netrins and which play crucial roles in establishing the balance between attraction and repulsion between growing axons, dendrites and postsynaptic targets. Similarly, the pro-inflammatory transcription factor NF κ B as well as COX-2 – an important enzyme in generating the oxidative stress that accompanies tissue inflammation – were not different between test and control animals ($p = 0.25$ and $p = 0.43$ respectively).

In view of the role of NMDAR receptors for both synaptic development and function, particular importance was attached to the expression of the major receptor subunits especially since, at P21, there had been no change in GluN1 but very significant changes in the expression of both GluN2A and GluN2B subunits. In the present study, expression of the GluN2A subunit was significantly lower in P60 offspring exposed *in utero* to Ro61-8048 compared with controls (Fig. 5A) and this was a decrease in expression whereas levels were increased at P21. No difference was detected in the expression of GluN2B or GluN1 subunits (Fig. 5B, C).

The levels of several proteins with strong links to NMDA receptor function were examined but neither the GluN1 subunit ($p = 0.20$, Fig. 5C) nor PSD-95 ($p = 0.14$, Fig. 5D) showed a clear change even though the latter had been significantly increased at the P21 age point. The two small GTPase enzymes RhoA and RhoB, which are involved in some of the plastic phenomena triggered by NMDA receptors (O’Kane et al., 2003) were similarly unchanged ($p = 0.95$, Fig. 5E; $p = 0.31$, not shown, respectively) even though there had been a very significant increase in RhoB at P21.

In other cases, such as the axon guidance and synapse formation receptor EphA4 ($p = 0.70$, Fig. 5F) and PCNA ($p = 0.19$, not shown) changes seen at P21 had been resolved or compensated so that no significant difference was apparent by the age of P60.

There were, however, marked and highly significant differences in the expression of two of the proteins concerned with cell proliferation, differentiation and maturation, and which had shown altered expression in animals at the time of weaning (P21). Doublecortin, which is associated with the undifferentiated state of cells and declines with differentiation, exhibited a clearly significant alteration in the pups exposed to Ro61-8048,




Fig. 4. Analysis of mRNA by qRT-PCR. Effect of prenatal administration of Ro61-8048 on mRNA expression in the hippocampus of offspring at P60. qRT-PCR analysis of the expression of transcripts (*Grin2a*, *Grin2b*, *Dcx*, *Shh*) in relation to three reference genes *Actb*, *Gapdh* and *Rn18s*. *Grin2a* (A, E, J), *Grin2b* (B, F, K), *Dcx* (C, G, L) and *Shh* (D, H, M). Results are shown as mean \pm s.e.m. of $2^{-\Delta Ct}$ ($n = 9$ animals per group).

with a 30% increase in expression compared with saline-treated animals ($p = 0.03$; Fig. 5G). This was very similar to the 30% increase seen in animals at P21.

The sonic hedgehog protein (Shh), was clearly detected in the adult hippocampus. Expression of this molecule was decreased significantly in the Ro61-8048-treated animals compared with controls ($p = 0.05$; Fig. 5H), a result again very close to that seen at P21.

Lastly, four proteins with a major relevance to dopaminergic neuronal systems were examined in this study: TH ($p = 0.58$), α -synuclein ($p = 0.52$), DISC-1 ($p = 0.89$) and the 5HT_{2C} receptor ($p = 0.58$), but none of these showed any change between treated and control animals in the hippocampus.

Behavior

Open field exploration. Tests were performed on P60 rats after prenatal maternal administration of Ro61-8048. Animals were exposed twice to an open field arena on two consecutive days. As indicated in

Fig. 6, there were no significant differences between the performance of control ($n = 38$) and Ro61-8048-treated ($n = 53$) offspring on their first exposure (OF1) to the arena, assessed either as the total number of line crossings (Fig. 6A) or the number of rearings in the 5-min session (Fig. 6B) (OF1; $p > 0.05$, Mann–Whitney test). When tested in the arena about 20 h later (OF2), both groups of animals showed a significant degree of habituation learning with fewer line crossings and rearings (Fig. 6A, B) ($p < 0.05$, Wilcoxon signed rank test after Kruskal–Wallis ANOVA for non-parametric samples). Furthermore, the variables were recorded and compared every minute in the open field to assess short term recognition of the environment. In all groups the exploratory parameters decreased during the session confirming that short term habituation to the arena was also preserved (Kruskal–Wallis followed by Dunn’s multiple comparison test, not shown).

Step-down avoidance. In the inhibitory step-down avoidance task, animals in the control ($n = 34$) and treated ($n = 50$) groups showed a significant difference




Fig. 5. Expression of proteins in the hippocampus of control and Ro61-8048-treated animals. The panels summarize the expression of several of the proteins examined in this study using Western blots. Data are shown for the optical density of blots relative to actin for (A) GluN2A, (B) GluN2B and (C) GluN1 subunits, (D) PSD-95, (E) RhoA, (F) EphA4, (G) doublecortin (DCX) and (H) sonic hedgehog (Shh). * $p < 0.05$; ** $p < 0.01$ (t test, $n = 6$ for Ro61-8048, $n = 4$ for controls).

between latencies in training (TR) and test (TEST) sessions, the test latencies being significantly lower (Fig. 7A; Wilcoxon test), indicating that both groups learned the avoidance task and remembered it in the testing session 24 h later. There was no significant

difference between the control and test groups in their response latencies in either the training or test sessions, indicating that Ro61-8048 had not affected the initial acquisition or the subsequent memory consolidation and further recall of the task.




Fig. 6. Open-field exploration task. Animals were exposed twice to an open field arena on two consecutive days. The bar diagrams summarize (A) the number of times the animals cross into new sectors of the open field arena (OF) and (B) the number of rearings for control and treated rats, during the 5-min periods they were exposed to the arena. Results are shown from the first session (OF1) and from the second session (OF2) performed 27–28 h later. The bars represent median with interquartile ranges. *Significant difference between control and treated animal ($p < 0.05$, Wilcoxon signed rank test after Kruskal–Wallis ANOVA for non-parametric samples). All animals: control $n = 38$; treated $n = 53$. There were no significant differences between control and treated groups in crossings and rearings during the first (OF1, $P > 0.05$) or second open field exposure (OF2; $p > 0.05$, Mann–Whitney test).

In order to exclude any confounding effect of gender factors on these results, a separate analysis was performed of male (Fig. 7B; control $n = 20$, treated $n = 19$) and female (Fig. 7C: control $n = 14$, treated $n = 31$) animals. Although female rats showed greater variability in their response latencies, there remained no significant differences between control and treated groups.

DISCUSSION

We have reported previously that the administration of Ro61-8048 to pregnant rats rapidly produces a substantial increase of between 10 and 100-fold in the levels of kynurenine and kynurenic acid in the maternal blood and in the embryo brains within 5 h (Forrest et al., 2013). After 24 h the maternal levels have normalized whereas they remain significantly elevated in the




Fig. 7. Step-down inhibitory avoidance in control and treated animals. The bar diagrams show the latencies to step-down from an isolated platform to the electrified grid, in an inhibitory avoidance paradigm. Control and treated rats were trained (TR) and tested (TEST) after 20 h. The bars represent median with interquartile ranges. TR: training session; TEST: test session; DIFF: differences between test and training sessions. *Significant difference between TR and TEST latencies ($p < 0.05$, Wilcoxon signed rank test after Kruskal–Wallis ANOVA for non-parametric samples). (A) All animals were 60 day old adults treated prenatally with Ro61–8048: control $n = 34$; treated $n = 50$. Analysis was also performed for males and females separately to exclude any gender effects: (B) male rats: control $n = 20$; treated $n = 19$; (C) female rats: control, $n = 14$; treated, $n = 31$. There were no significant differences between groups in latency differences ($p > 0.05$, Mann–Whitney test).

embryonic brain. It is likely, therefore, that the changes described previously at P21 and reported here for animals at P60 following the same prenatal injection protocol result from the increased levels of these tryptophan metabolites. It is unlikely that changes were

produced in the concentrations of quinolinic acid since previous work did not reveal any change in resting levels of this compound after Ro61-8048 (Chiarugi and Moroni, 1999; Clark et al., 2005).

It was also reported that the use of Ro61-8048 was not accompanied by any changes in the levels of pro-inflammatory cytokines such as interleukin-1 α , IL-1B, IL-6, tumor necrosis factor- α , monocyte chemoattractant protein-1, macrophage inflammatory protein-1 α , Regulated on Activation, Normal T cell expressed and secreted (RANTES), or interferon receptor (Forrest et al., 2013), confirming that the activation of immune function was unlikely to contribute significantly to the observed results.

Electrophysiology

While the various biochemical changes generated in the adult brain by exposure to Ro61-8048 *in utero* indicate that abnormalities of neurochemistry can be produced by changing the kynurenine environment during early brain development, those changes may not be sufficient to cause gross functional change. In order to assess the existence and degree of functional change we have analyzed several basic aspects of hippocampal electrophysiology.

The results indicate that there are no apparent alterations to basic aspects of neuronal excitability and synaptic transmission since there were no differences between control and treated animals in the stimulus–response relationships for either fEPSPs or population spikes. These data also imply that there are no changes in the relationship between presynaptic transmitter release and postsynaptic excitability (E–S coupling).

There are, however, clearly demonstrable changes in some aspects of synaptic plasticity. While slices from control and treated animals exhibit closely similar PPF across the range of interpulse intervals from 20 to 100 ms, marked differences are apparent at the 10-ms interval. The importance of this observation lies in the different mechanisms underlying PPF and PPI. The latter is believed to result from the short-lived depletion of synaptic vesicles following the first stimulus (Rosenmund and Stevens, 1996; Wang and Kaczmarek, 1998), whereas PPF is believed to result from the accumulation of residual calcium within synaptic terminals, leading to increased transmitter release and an increased size of the second potential of the pair (Zucker et al., 1991). A reduced, or absent, PPI at the smallest interpulse interval would, therefore, suggest less vesicular depletion, perhaps implying either an increased density of vesicles available for release, or a facilitated movement of vesicles to the active zone for release. The overall result would be an increased probability of vesicular release. It is also possible that the accumulation of intra-terminal calcium begins to occur earlier in synaptic terminals from treated animals so that the balance of mechanisms causing PPI and PPF is altered at this earliest time interval.

Another possibly relevant factor stems from the observation by Al-Hayani and Davies (2002) that the occurrence of secondary spikes predisposes slices to generate relatively weak PPI. However, care was taken in this study to examine only those slices which did not exhibit a secondary spike, making this factor an unlikely complication.

The occurrence of altered synaptic behavior was strongly supported by extending this analysis to an examination of paired-pulse interactions at different frequencies of paired-pulse delivery. Increasing the frequency of stimulation from 0.05 to 1 Hz has been shown previously to induce a shift from PPF to PPI in the hippocampus (Saviane et al., 2002). A similar change from PPF to PPI was reported following exposure to 4-aminopyridine, a potassium channel blocker which facilitates transmitter release (Pena et al., 2002) while increased PPI was reported on exposure to bicuculline (Higgins and Stone, 1993) or following seizure activity (Stringer and Lothman, 1989). It has also been reported that a similar shift from PPF to PPI can be induced by raising the stimulation strength (Huang et al., 2007).

In the present study, increasing stimulation frequency had little effect on the PPI seen in control slices, whereas the early PPF seen in slices from experimental animals was converted to PPI which, at a frequency of 1 Hz, significantly exceeded the magnitude of PPI in control slices. Together, these various paired-pulse results suggest a facilitated release of transmitter in treated animals at small interpulse intervals, with either an increased depletion of vesicles at higher pair frequencies or a reduction in the opposing accumulation of calcium, which normally maintains transmitter release approximately constant over the frequency range examined.

There are, however, other factors that need to be examined in further detail in future work, since it has been suggested that presynaptic inhibitory receptors (Forsythe and Clements, 1990) including those for adenosine (Higgins and Stone, 1996), receptor desensitization (Otis et al., 1996; Neher and Sakaba, 2001) or some form of refractory state (Waldeck et al., 2000) may contribute to paired-pulse interactions. GABA receptors may also be involved (Nathan et al., 1990) although this is likely to affect multiple spiking rather than primary potentials (Leung and Fu, 1994) and care was always taken in this study to limit investigation to slices showing no secondary spikes following single stimuli. While there is relatively little evidence that these processes contribute significantly to paired-pulse phenomena under normal conditions, it is of course possible that they may assume greater importance in the animals exposed to Ro61-8048.

Finally, changes were sought in the amplitude of LTP and LTD, the former induced by theta-burst stimulation which reproduces natural patterns of hippocampal activity and produces an optimal level of LTP (Larson et al., 1986). Exposure to Ro61-8048 *in utero* led to a defect in theta-burst LTP which, while still occurring to a

significant extent, was significantly smaller than in slices from control animals. The specificity of the phenomenon was emphasized by the absence of any significant difference in the amplitude of LTD, although the significant differences described in the time course of recovery from the second period of stimulation might reflect a degree of functional synaptic change. Indeed, both the reduced LTP amplitude and the slowed recovery toward the plateau LTD could be accommodated broadly within a hypothesis that explained the increased ratio of inhibition to facilitation at the higher frequencies of paired-pulse delivery. It is interesting to note a recent report that the expression of GluN2A, but not of GluN2B, was increased in the rat hippocampus 1 h after LTP induction (Baez et al., 2013). Hence, the decrease in LTP reported here may be, at least in part, a consequence of the reduced expression of GluN2A in the offspring from Ro61-8048-treated rats.

Mechanisms underlying altered development

It is well established that neuronal receptors for glutamate and acetylcholine are heavily involved in brain development. Glutamate receptors, especially those sensitive to NMDA affect neuronal migration, synapse formation (Udin and Grant, 1999; Dikranian et al., 2001), neurite growth (Udin and Grant, 1999), spine formation (Ultanir et al., 2007) and neuronal plasticity (Iwasato et al., 2000; Drian et al., 2001; Fagiolini et al., 2003; du Bois and Huang, 2007). The expression of neural cell adhesion molecules is partly controlled by NMDAR activation. The pharmacological blockade of NMDARs in neonatal rats causes a loss and disruption of synapses with profound abnormalities of brain structure and behavior in adulthood (Dikranian et al., 2001). NMDAR blockade also produces a substantial loss of neurons partly resulting from the regulation by NMDARs of the p53 tumor suppressor gene. A similar loss of NMDAR function during development could contribute to the programmed elimination of neurons which shapes the early nervous system.

The major route for the metabolism of tryptophan is the oxidative pathway via kynurenine (Stone and Darlington, 2002). In most other tissues, including blood monocytes, macrophages and their descendant microglia, the first enzyme is indoleamine-2,3-dioxygenase (IDO), which is potently induced by inflammatory mediators such as IFN- γ , IL-1 β and TNF- α which mediate the effects of lipopolysaccharides and poly(I:C). Since the pathway includes quinolinic acid as an agonist at NMDARs (Stone and Perkins, 1981) and kynurenic acid as an antagonist at all ionotropic glutamate receptors, though with greatest potency at NMDARs (Perkins and Stone, 1982), it represents a major link between activation of the immune system and the modulation of glutamate (especially NMDA) receptors. In addition, the kynurenine pathway is present partly in neurons and partly in glia (Guidetti et al., 2007; Schwarcz and Pellicciari, 2002; Guillemin et al., 2001, 2005). Since glial cells present in the early stages of CNS development regulate the appearance and function of NMDARs, the location of the

pathway is consistent with the possibility that the activation of NMDARs is determined partly by quinolinic acid balanced by the glutamate receptor antagonist kynurenic acid. Indeed, it has been proposed that tissue formation and polarity may be determined largely by a gradient for a signal antagonist, rather than an agonist (Gurdon and Bourillot, 2001), which would fit well with the concept of a major developmental role for kynurenic acid. With its selectivity for NMDARs, kynurenic acid would also alter the balance of activation of the various subpopulations of glutamate receptors.

The kynurenine pathway is present during embryonic development (Beal et al., 1992; Saito et al., 1993; Walker et al., 1999), as reflected in the expression of one or more enzymes or the presence of key components such as kynurenine, kynurenic acid or quinolinic acid. The ability of prenatal inhibition of the kynurenine pathway to produce significant changes in synaptic properties in the adult offspring, as shown in this report, strongly suggests that the pathway normally plays a significant role in cerebral development. There are several ways in which this could occur, the most likely being changes in cerebral concentration of kynurenic acid which may vary during pregnancy in response to external factors such as infection and stress. Indeed any immune challenge to the mother or neonate which results, directly or indirectly, in the activation of central glia or peripheral macrophages, or changes in the levels of cytokines or kynurenines in the fetal or neonatal CNS would alter the balance of quinolinic acid and kynurenic acid concentrations and could significantly affect neural development and plasticity. This proposal is consistent with evidence that genetic abnormalities of the kynurenine pathway are linked to disorders such as schizophrenia (Miller et al., 2006; Holtze et al., 2012).

Protein and mRNA expression

Analysis of mRNA levels using qRT-PCR failed to reveal any changes in gene expression. This strongly suggests that the various changes in synaptic function and protein expression have developed either from alterations in the translation of unchanged RNA into protein or from short-term changes in gene expression that may have been limited to the time during which Ro61-8048 was active in raising the levels of kynurenic acid. It should also be noted that temporal and quantitative differences are frequently encountered in comparisons of mRNA and protein expression, resulting in poor correlations between these parameters (English et al., 2011)

The investigation of protein expression was performed to identify changes that might result from kynurenine pathway inhibition. Many of the changes that had been recorded at P21 (Forrest et al., 2013) were no longer demonstrable in this study at P60, suggesting that those aspects of neuronal and synaptic development and function which were abnormal compared with control animals at P21 had largely returned to normal by P60. However, the existence of changes at P21 implies an effect of kynurenine pathway inhibition on both the extent of expression at this time and on the rate of

appearance or disappearance of those proteins in the course of brain development. Clearly, either quantitative or temporal disturbances of protein expression are likely to lead to a degree of abnormal expression and synaptic function at P60. Since the P21 stage of postnatal development in the rat brain is estimated to be equivalent to the early stage of brain development seen near the end of the second trimester of pregnancy in humans (Dobbing and Sands, 1979) it is interesting to speculate that any factors affecting brain development at these times are likely to result in some degree of abnormal brain function.

Of the nine target proteins whose expression was modified at P21, three of them are molecules that remain significantly changed at P60: the GluN2A subunit, doublecortin and sonic hedgehog. Of note is the finding that the difference in GluN2B expression seen at P21 was not apparent at P60 and that expression of GluN2A was significantly different but in the opposite direction to that at P21 – a decrease in expression compared with the earlier increase. Since the ratio between GluN2A and GluN2B subunits is known to change during the course of early brain development and to have a major influence on behavioral and electrophysiological outcomes by adulthood (Traynelis et al., 2010), the shifts we have reported in these two subunits may be among the more important factors in generating the functional synaptic changes described above. As noted earlier, the reduced LTP and GluN2A expression seen here is entirely compatible with the recently reported increase in expression of GluN2A induced by LTP (Baez et al., 2013).

Doublecortin

The increased expression of doublecortin at P60 – a change similar in magnitude and direction to that recorded at P21 – may provide an explanation for the electrophysiological changes reported here. Although the name ‘doublecortin’ was derived from the role of the protein in disorders of human neuronal migration in the neocortex which resulted in mental retardation and hyperexcitability (Gleeson et al., 1999; Nacher et al., 2001), the protein has also been linked with cellular lamination in the hippocampus (Corbo et al., 2002). It is a microtubule-associated protein which is expressed primarily in newly formed neurons (Couillard-Despres et al., 2005) and its deletion results in reduced neurogenesis and poor recovery after stroke injury (Jin et al., 2010). Doublecortin is certainly active in neurogenesis during the earliest phases of brain development, with a progressive decline in its levels in the mature brain although it continues to be associated with new cell generation in the adult brain. The hippocampus, especially the subgranular zone of the dentate gyrus, exhibits high levels of doublecortin in the adult, probably related to the high levels of neurogenesis here. The significant increase in doublecortin levels, therefore, would imply an increased rate of new neurogenesis in the hippocampus following exposure to Ro61-8048 *in utero*.

While there were no overall changes in LTD, the more prolonged recovery from low-frequency stimulation may be related to the association that has been reported between doublecortin expression and GABAergic function. The neuronal hyperexcitability which results from mutations or deletions of doublecortin results from a reduction of synaptic inhibition (Kerjan et al., 2009) which in turn is consistent with the co-localization of doublecortin with GABA, its synthetic enzyme glutamate decarboxylase, or parvalbumin, a marker for GABA-releasing neurons (Cai et al., 2009). Indeed, neurons expressing doublecortin may be primarily destined to become GABAergic interneurons (Xiong et al., 2008) and it has been proposed that doublecortin is required for the early migration of inhibitory interneurons in the cortex (Kappeler et al., 2006; Friocourt et al., 2007).

Sonic hedgehog

The only other protein to remain significantly changed at P60 was sonic hedgehog (Shh). This protein is involved in the early development of tissue polarization and the generation of morphogenetic and orientational gradients (Palma et al., 2005; Traiffort et al., 2010). Despite its importance in the earliest stages of CNS formation, it continues to exist in the adult brain, notably in regions such as the cerebellum (Vaillant and Monard, 2009) and hippocampus (Traiffort et al., 1998; Charytoniuk et al., 2002; Ahn and Joyner, 2005; Dellovade et al., 2006) where neurogenesis continues in the adult. Shh may also modulate cell proliferation and the migration of progenitor cells (Traiffort et al., 2001; Charytoniuk et al., 2002; Palma et al., 2005; Sims et al., 2009; Dave et al., 2011).

In addition, Shh is thought to modulate the mutual attraction or repulsion between neuritic branches and potential sites of synaptic contact (Angot et al., 2008; Hor and Tang, 2010) and the reduction in Shh may interfere with appropriate contact formation during development. This may contribute to the functional differences observed in hippocampal transmission. In view of the evidence that Shh can promote the functional development of inhibitory neurons such as cerebellar granule cells (Spassky et al., 2008; Prajerova et al., 2010) the combination of these factors could be partly responsible for the observed decreased in PPI.

Since doublecortin has been linked with neurogenesis, it is perhaps surprising that the levels of Shh protein decrease, while those of doublecortin increase, given the role of the former in neuronal orientation and contact formation. However, the roles attributed to doublecortin temporally precede those ascribed to Shh, leading to the possibility that there is increased production of new neurons which then mature and differentiate more slowly than in control animals.

Although several of the proteins examined were altered at P21 after exposure to Ro61-8048 *in utero*, the fact that no changes persisted into young adulthood at P60 suggests that the brain is able to adapt well to this particular interference. However, the protein and functional changes still persisting at P60 indicate that, despite any adaptations that have occurred between

drug administration *in utero* and P21 or P60, the effects of kynurenine pathway inhibition *in utero* can clearly generate a sufficient disturbance of brain development that changes in three proteins with fundamental roles in the formation and maintenance of neuronal numbers, growth, connectivity and plasticity persist until early adulthood. This conclusion raises the intriguing possibility that physiological or environmental influences on the kynurenine pathway *in utero* may contribute to brain function and behavior in the postnatal years and adulthood. Since Shh plays a particularly prominent role in the development of dopaminergic neuron projections (Wang et al., 1995; Hynes et al., 1995; Parish et al., 2008; Tang et al., 2010; Wu et al., 2012), its loss may be particularly relevant to the emergence of parkinsonian symptoms in adulthood.

Behavior

The changes in neurodevelopmental proteins and synaptic physiology might be expected to lead to alterations in behavior. The administration of kynurenine to adult or perinatal rats (Erhardt et al., 2004; Nilsson et al., 2006; Pocivavsek et al., 2012) or the depletion of kynurenic acid levels by deleting kynurenine aminotransferase (Potter et al., 2010) has demonstrated effects on learning in complex tasks such as pre-pulse inhibition (Erhardt et al., 2004; Nilsson et al., 2006), a water maze (Chess et al., 2007), stimulus discrimination (Alexander et al., 2012; Pocivavsek et al., 2012) or attentional set shifting (Pocivavsek et al., 2012; Alexander et al., 2013). In the present study behavioral testing was limited to a classical step-down avoidance task which can simplify the interpretation of drug effects on registration, consolidation and recall, but no significant differences were noted in the offspring of animals treated with Ro61-8048 compared with saline controls.

Differences between the results from these various studies may be caused by differences in the neurochemical effects of Ro61-8048 and kynurenine administration, the timing of prenatal treatment (three injections of Ro61-8048 during pregnancy in this study compared with kynurenine administration to adults or continual intake in the diet pre- and post-natally), or the nature of the behavioral tests. The simple step-down avoidance procedure may not be appropriate to detect cognitive changes that are only revealed in more complex tasks (Chess et al., 2007; Pocivavsek et al., 2012; Alexander et al., 2012, 2013), or which require testing of aspects of behavior more subtle than a simple learning task. It has also been noted previously that some, but not all aspects of cognitive function are influenced by elevated levels of kynurenic acid (Chess and Bucci, 2006; Chess et al., 2009). In addition, it has been reported that simpler learning tasks such as step-down avoidance task can be achieved by neural pathways involving or excluding the hippocampus, so that even major hippocampal damage can be circumvented by alternative cerebral pathways (Martel et al., 2010). This would certainly account for the failure

of the subtle hippocampal changes described here to alter step-down avoidance.

Additional consequences of KMO inhibition

In addition to the major increase in kynurenic acid concentrations, the inhibition of KMO may also have effects on neuronal excitability and function by reducing the levels of other compounds normally generated as secondary products of kynurenine oxidation (Stone and Darlington, 2002, 2013; Schwarcz et al., 2012; Stone et al., 2013). The immediate product of KMO, 3-hydroxykynurenine (3HK) is an effective neurotoxin (Okuda et al., 1998) and is converted by kynureninase to 3-hydroxy-anthranilic acid (3HAA) which, in addition to direct actions on T cell viability (Munn, 2011), can auto-oxidize to the dimeric cinnabarinic acid (Dykens et al., 1987), a compound which is at least 10-fold more active as an inducer of T cell death (Hiramatsu et al., 2008). This compound also activates the metabotropic glutamate receptor mGluR4 (Fazio et al., 2012) which, as a member of the Group III metabotropic receptor family, has a range of actions in the CNS including modulation of anxiety, depression (Wierońska et al., 2010) and basal ganglia function (Niswender et al., 2008; Broadstock et al., 2012). Another potentially important product is xanthurenic acid which has several characteristics of a neurotransmitter candidate (Gobaille et al., 2008), activates group II metabotropic glutamate receptors and can modify subcortical neurotransmission (Copeland et al., 2013). It may also activate cationic channels that enhance neuronal excitability (Taleb et al., 2012) and may modulate excitability indirectly via inhibition of vesicular glutamate transporters (Neale et al., 2013).

SUMMARY

Overall, the evidence from this study suggests that there is a basal, constitutive level of activity in the kynurenine pathway which is normally active in the embryonic, developing CNS. The pathway can influence the composition and function of NMDA receptors and the expression of proteins critical to normal brain development and function, some of which can persist into adulthood. These changes result in abnormal neurotransmission and plasticity which could cause disease or predispose individuals to the development of CNS disorders. The changes seen at P60 are remarkable in several respects. Firstly, many of the changes in protein expression seen at P21 are no longer present at P60, suggesting that the influence of modifying kynurenine function prenatally produced temporary, not permanent, alterations. The early changes, however, may well have caused or contributed to the changes noted here.

Secondly, the functional changes in synaptic interactions and plasticity are different at P60 from those seen at P21, indicating that the processes of network development and possible remodeling have been significantly modified by the earlier effects of Ro61-8048.

Thirdly, since the kynurenine pathway (mainly IDO and KMO) is activated by interferons and pro-inflammatory cytokines (Alberati-Giani et al., 1996), the ability of the pathway to regulate brain development carries the implication that it could be responsible for some of the recognized influence of infection during pregnancy on brain development of the embryo (Brown, 2006, 2011; Meyer and Feldon, 2010; Hornig et al., 1999). There are also some compounds in the diet, such as the brassinins found in cruciferous vegetables (cabbage, broccoli, etc.), which inhibit the kynurenine pathway (Banerjee et al., 2008), mimicking the effects of Ro61-8048. According to our results the changes in the levels of kynurenic acid and other kynurenine metabolites which they could produce might affect brain development. There are also simple metabolites such as oxoglutarate and pyruvate (Shin et al., 1982) as well as vitamins such as pyridoxal (Breton et al., 2000) which can activate or inhibit KMO directly depending on concentration and which might have pathological relevance.

Finally, two of the proteins that are fundamental players in the early generation, maturation and organization of cells remain significantly altered at P60 in the same direction as found at P21 – doublecortin levels are increased and sonic hedgehog levels are decreased at both time points. Together with the persisting increase in GluN2A expression it seems probable that these proteins might represent key links between the kynurenine pathway and brain development and future work should be directed at exploring the sites and mechanisms of this relationship.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest in relation to the publication of this work.

Acknowledgments—OSK is in receipt of a BBSRC Scholarship in Organismal Biology (Biotechnology and Biological Sciences Research Council). MP is supported by funding from the Malaysian Government. DAJ thanks F Hoffman-La Roche AG, Pharmaceuticals Division, Basel, Switzerland for the gift of Ro61-8048 and we thank Dr. Catherine Winchester for help and advice in qRT-PCR methodology. We are grateful to Epsom Medical Research, the Peacock Trust, the Haddon Family Trust, the National Council of Scientific and Technological Investigations (CONICET) and the University of Buenos Aires, Argentina for financial assistance.

REFERENCES

- Ahn S, Joyner AL (2005) In vivo analysis of quiescent adult neural stem cells responding to sonic hedgehog. *Nature* 437:894–897.
- Alberati-Giani D, Ricciardi-Castiglioni P, Kohler C, Cesura AM (1996) Regulation of the kynurenine metabolic pathway by interferon-gamma in murine cloned macrophages and microglial cells. *J Neurochem* 66:996–1004.
- Alexander KS, Pociavsek A, Wu H-Q, Pershing ML, Schwarcz R, Bruno JP (2013) Early developmental elevations of brain kynurenic acid impair cognitive flexibility in adults: reversal with galantamine. *Neuroscience* 234:19–28.
- Alexander KS, Wu H-Q, Schwarcz R, Bruno JP (2012) Acute elevations of brain kynurenic acid impair cognitive flexibility: normalization by the alpha7 positive modulator galantamine. *Psychopharmacology* 220:627–637.
- Al-Hayani A, Davies SN (2002) Effect of cannabinoids on synaptic transmission in the rat hippocampal slice is temperature-dependent. *Eur J Pharmacol* 442:47–54.
- Alvarez VA, Ridenour DA, Sabatini BL (2007) Distinct structural and ionotropic roles of NMDA receptors in controlling spine and synapse stability. *J Neurosci* 27:7365–7376.
- Angot E, Loulier K, Nguyen-Ba-Charvet KT, Gadeau A-P, Ruat M, Traiffort E (2008) Chemoattractive activity of sonic hedgehog in the adult subventricular zone modulates the number of neural precursors reaching the olfactory bulb. *Stem Cells* 26:2311–2320.
- Baez MV, Oberholzer MV, Cercato MC, Snitcofsky M, Aguirre AI, Jerusalinsky DA (2013) NMDA receptor subunits in the adult rat hippocampus undergo similar changes after 5 minutes in an open field and after LTP induction. *PLOS One* 8:e55244.
- Banerjee T, Duhadaway JB, Gaspari P, Sutanto-Ward E, Munn DH, Mellor AL, Malachowski WP, Prendergast GC, Muller AJ (2008) A key in vivo antitumor mechanism of action of natural product-based brassinins is inhibition of indoleamine 2,3-dioxygenase. *Oncogene* 27:2851–2857.
- Beal MF, Swartz KJ, Isacson O (1992) Developmental changes in brain kynurenic acid concentrations. *Dev Brain Res* 68:136–139.
- Breton J, Avanzini S, Covini N, Magistrelli G, Cozzi L, Isacchi A (2000) Functional characterisation and mechanism of action of recombinant human kynurenine-3-hydroxylase. *Eur J Biochem* 267:1092–1099.
- Broadstock M, Austin PJ, Betts MJ, Duty S (2012) Antiparkinsonian potential of targeting group III metabotropic glutamate receptor subtypes in the rodent substantia nigra pars reticulata. *Br J Pharmacol* 165:1034–1045.
- Brown AS (2006) Prenatal infection as a risk factor in schizophrenia. *Schizophren Bull* 32:200–202.
- Brown AS (2011) The environment and susceptibility to schizophrenia. *Prog Neurobiol* 93:23–58.
- Cai Y, Xiong K, Chu Y, Luo D-W, Luo X-G, Yuan X-Y, Struble RG, Clough RW, Spencer DD, Williamson A, Kordower JH, Patrylo PR, Yan XX (2009) Doublecortin expression in adult cat and primate cerebral cortex relates to immature neurons that develop into GABAergic subgroups. *Exp Neurol* 216:342–356.
- Charytoniuk D, Traiffort E, Hantraye P, Hermel JM, Galdes A, Ruat M (2002) Intrastratial sonic hedgehog injection increases patched transcript levels in the adult rat subventricular zone. *Eur J Pharmacol* 16:2351–2357.
- Chess AC, Simoni MK, Alling TE, Bucci DJ (2007) Elevations of endogenous kynurenic acid produce spatial working memory deficits. *Schizophren Bull* 33:797–804.
- Chess AC, Bucci DJ (2006) Increased concentration of cerebral kynurenic acid alters stimulus processing and conditioned responding. *Behav Brain Res* 170:326–332.
- Chess AC, Landers AM, Bucci DJ (2009) L-Kynurenine treatment alters contextual fear conditioning and context discrimination but not cue-specific fear conditioning. *Behav Brain Res* 201:325–331.
- Chiarugi A, Moroni F (1999) Quinolinic acid formation in immune-activated mice: studies with (m-nitrobenzoyl)-alanine (mNBA) and 3,4-dimethoxy-[N-4-(-3-nitrophenyl)thiazol-2yl]-benzenesulfonamide (Ro 61-8048), two potent and selective inhibitors of kynurenine hydroxylase. *Neuropharmacology* 38:1225–1233.
- Clark CJ, Mackay GM, Smythe GA, Bustamante S, Stone TW, Phillips SR (2005) Prolonged survival of a murine model of cerebral malaria by kynurenine synthesis inhibition. *Infect Immun* 73:5249–5251.
- Colonnese MT, Zhao JP, Constantine-Paton M (2005) NMDA receptor currents suppress synapse formation on sprouting axons in vivo. *J Neurosci* 25:1291–1303.
- Copeland CS, Neale SA, Salt TE (2013) Actions of xanthurenic acid, a putative endogenous group II metabotropic glutamate receptor agonist, on sensory transmission in the thalamus. *Neuropharmacology* 66:133–142.

- Corbo JC, Deuel TA, Long JM, LaPorte P, Tsai E, Wynshaw-Boris A, Walsh CA (2002) Doublecortin is required in mice for lamination of the hippocampus but not the neocortex. *J Neurosci* 22:7548–7557.
- Couillard-Despres S, Winner B, Schauback S, Aigner R, Vroemen M, Weidner N, Bogdahn U, Winkler J, Kuhn HG, Aigner L (2005) Doublecortin expression levels in adult brain reflect neurogenesis. *Eur J Neurosci* 21:1–14.
- Cozzi A, Carpenedo R, Moroni F (1999) Kynurenine hydroxylase inhibitors reduce ischemic brain damage: studies with (m-nitrobenzoyl)-alanine (mNBA) and 3,4-dimethoxy-[N-4-(nitrophenyl)thiazol-2YL]-benzenesulfonamide (Ro61-8048) in models of focal or global brain ischaemia. *J Cereb Blood Flow Metab* 19:771–777.
- Cuppini R, Sartini S, Ambrogini P, Falcieri E, Maltarello MC, Gallo G (1999) Control of neuron outgrowth by NMDA receptors. *J Submicrosc Cytol Pathol* 31:31–40.
- Dave RK, Ellis T, Toumpas MC, Robson JP, Julian E, Adolphe C, Bartlett PF, Cooper HM, Reynolds BA, Wainwright BJ (2011) Sonic hedgehog and notch signaling can cooperate to regulate neurogenic divisions of neocortical progenitors. *PLOS One* 6(2). Article e14680.
- Dellovade T, Romer JT, Curran T, Rubin LL (2006) The hedgehog pathway and neurological disorders. *Ann Rev Neurosci* 29:539–563.
- Dikranian K, Ishimaru MJ, Tenkova T, Labruyere J, Qin YQ, Ikonomidou C, Olney JW (2001) Apoptosis in the in vivo mammalian forebrain. *Neurobiol Dis* 8:359–379.
- Dobbing J, Sands J (1979) Comparative aspects of the brain growth spurt. *Early Hum Dev* 3:79–83.
- Dobelis P, Staley KJ, Cooper DC (2012) Lack of modulation of nicotinic acetylcholine alpha-7 receptor currents by kynurenic acid in adult hippocampal interneurons. *PLoS One* 7. Article e41108.
- Drian MJ, Bardoul M, König N (2001) Blockade of AMPA/kainate receptors can either decrease or increase the survival of cultured neocortical cells depending on the stage of maturation. *Neurochem Int* 38:509–517.
- du Bois TM, Huang XF (2007) Early brain development disruption from NMDA receptor hypofunction: relevance to schizophrenia. *Brain Res Rev* 53:260–270.
- Dykens JA, Sullivan SG, Stern A (1987) Oxidative reactivity of the tryptophan metabolites 3-hydroxyanthranilate, cinnabarinic acid, quinolinate and picolinate. *Biochem Pharmacol* 36:211–217.
- English JA, Pennington K, Dunn MJ, Cotter DR (2011) The neuroproteomics of schizophrenia. *Biol Psychiatry* 69:163–172.
- Erhardt S, Schwieler L, Emanuelsson C, Geyer M (2004) Endogenous kynurenic acid disrupts prepulse inhibition. *Biol Psychiatry* 56:255–260.
- Ernst AF, Jurney WM, McLoon SC (1998) Mechanisms involved in development of retinotectal connections: roles of Eph receptor tyrosine kinases NMDA receptors and nitric oxide. *Prog Brain Res* 118:115–131.
- Fagiolini M, Katagiri H, Miyamoto H, Mori H, Grant SGN, Mishina M, Hensch TK (2003) Separable features of visual cortical plasticity revealed by N-methyl-D-aspartate receptor 2A signaling. *Proc Natl Acad Sci USA* 100:2854–2859.
- Fatemi SH, Pearce DA, Brooks AI, Sidwell RW (2005) Prenatal viral infection in mouse causes differential expression of genes in brains of mouse progeny: a potential animal model for schizophrenia and autism. *Synapse* 57:91–99.
- Fazio F, Lionetto L, Molinaro G, Bertrand HO, Acher F, Ngomba RT, Notartomaso S, Curini M, Rosati O, Scarselli P, Di Marco R, Battaglia G, Bruno V, Simmaco M, Pin JP, Nicoletti F, Goudet C (2012) Cinnabarinic acid, an endogenous metabolite of the kynurenine pathway, activates type 4 metabotropic glutamate receptors. *Mol Pharmacol* 81:643–656.
- Forrest CM, Khalil OS, Pisar M, Darlington LG, Stone TW (2013) Prenatal inhibition of the tryptophan–kynurenine pathway alters synaptic plasticity and protein expression in the rat hippocampus. *Brain Res* 1504:1–15.
- Forsythe ID, Clements JD (1990) Presynaptic glutamate receptors depress excitatory monosynaptic transmission between mouse hippocampal-neurons. *J Physiol (Lond)* 429:1–16.
- Friocourt G, Liu JS, Antypa M, Rakic S, Walsh CA, Parnavelas JG (2007) Both doublecortin and doublecortin-like kinase play a role in cortical interneuron migration. *J Neurosci* 27:3875–3883.
- Gleeson JG, Minnerath SR, Fox JW, Allen KM, Luo RF, Hong SE, Berg MJ, Kuzniecky R, Reitnauer PJ, Borgatti R, Mira AP, Guerrini R, Holmes GL, Rooney CM, Berkovic S, Scheffer I, Cooper EC, Ricci S, Cusmai R, Crawford TO, Leroy R, Andermann E, Wheless JW, Dobyns WB, Ross ME, Walsh CA (1999) Characterization of mutations in the gene doublecortin in patients with double cortex syndrome. *Ann Neurol* 45:146–153.
- Gobaille S, Kemmel V, Brumar D, Dugave C, Aunis D, Maitre M, Gobaille S (2008) Xanthurenic acid distribution, transport, accumulation and release in the rat brain. *J Neurochem* 105:982–993.
- Goncalves J, Queiroz G (2008) Presynaptic adenosine and P2Y receptors. *Handb Exp Pharmacol* 184:339–372.
- Guidetti P, Hoffman GE, Melendez-Ferro M, Albuquerque EX, Schwarcz R (2007) Astrocytic localization of kynurenine aminotransferase II in the rat brain visualized by immunocytochemistry. *Glia* 55:78–92.
- Guillemin GJ, Smythe G, Takikawa O, Brew BJ (2005) Expression of indoleamine 2,3-dioxygenase and production of quinolinic acid by human microglia, astrocytes, and neurons. *Glia* 49:15–23.
- Guillemin GJ, Kerr SJ, Smythe GA, Smith DG, Kapoor V, Armati PJ, Croitoru J, Brew BJ (2001) Kynurenine pathway metabolism in human astrocytes: a paradox for neuronal protection. *J Neurochem* 78:842–853.
- Gurdon JB, Bourillot PY (2001) Morphogen gradient interpretation. *Nature* 413:797–803.
- Harris LW, Sharp T, Gartlon J, Jones DNC, Harrison PJ (2003) Long-term behavioural molecular and morphological effects of neonatal NMDA receptor antagonism. *Eur J Neurosci* 18:1706–1710.
- Harvey AL, Young LC, Kornisiuk E, Snitkofsky M, Blanco C, Coletti N, Jerusalinsky D, Jamieson AG, Hartley RC, Stone TW (2012) The novel dihydro-pyrazolo(3,4-d)-(1,2,4)triazolo(1,5-a)pyrimidin-4-one (AJ23) is an antagonist at adenosine A₁ receptors and enhances consolidation of step-down avoidance. *Behav Brain Res* 234:184–191.
- Heng JE, Zurakowski D, Vorwerk CK, Grosskreutz CL, Dreyer EB (1999) Cation channel control of neurite morphology. *Dev Brain Res* 113:67–73.
- Higgins MJ, Stone TW (1993) Bicuculline-resistant paired-pulse inhibition in the rat hippocampal slice. *Eur J Pharmacol* 317:215–223.
- Higgins MJ, Stone TW (1996) The contribution of adenosine to paired-pulse inhibition in the normal and disinhibited hippocampal slice. *Br J Pharmacol* 109:1164–1168.
- Hilmas C, Pereira EFR, Alkondon M, Rassoulpour A, Schwarcz R, Albuquerque EX (2001) The brain metabolite kynurenic acid inhibits alpha 7 nicotinic receptor activity and increases non-alpha 7 nicotinic receptor expression: physiopathological implications. *J Neurosci* 21:7463–7473.
- Hiramatsu R, Hara T, Akimoto H, Takikawa O, Kawabe T, Isobe K, Nagase F (2008) Cinnabarinic acid generated from 3-hydroxyanthranilic acid strongly induces apoptosis in thymocytes through the generation of reactive oxygen species and the induction of caspase. *J Cell Biochem* 103:42–53.
- Hoffman KB, Murray BA, Lynch G, Munirathinam S, Bahr BA (2001) Delayed and isoform-specific effect of NMDA exposure on neural cell adhesion molecules in hippocampus. *Neurosci Res* 39:167–173.
- Holtze M, Saetre P, Engberg G, Schwieler L, Werge T, Andreassen OA, Hall H, Terenius L, Agartz I, Jonsson EG, Schalling M, Erhardt S (2012) Kynurenine 3-monooxygenase polymorphisms: relevance for kynurenic acid synthesis in patients with schizophrenia and healthy controls. *J Psychiatry Neurosci* 37:53–57.

- Hor CHH, Tang BL (2010) Sonic hedgehog as a chemoattractant for adult NPCs. *Cell Adhes Migr* 4:1–3.
- Hornig M, Weissenbock H, Horscroft N, Lipkin WI (1999) An infection-based model of neurodevelopmental damage. *Proc Natl Acad Sci U S A* 96:12102–12107.
- Huang F-S, Meng K, Tang J-S (2007) Properties of paired-pulse firing thresholds and the relationship with paired-pulse plasticity in hippocampal CA3–CA1 synapses. *Eur J Neurosci* 25:3253–3263.
- Hynes M, Porter JA, Chiang C, Chang D, Tessierlavigne M, Beachy PA, Rosenthal A (1995) Induction of midbrain dopaminergic neurons by sonic hedgehog. *Neuron* 15:35–44.
- Ikonomidou C, Bosch F, Miksa M, Bittigau P, Vockler J, Dikranian K, Tenkova TI, Stefovskva V, Turski L, Olney JW (1999) Blockade of NMDA receptors and apoptotic neurodegeneration in the developing brain. *Science* 283:70–74.
- Iwasato T, Datwani A, Wolf AM, Nishiyama H, Taguchi Y, Tonegawa S, Knopfel T, Erzurumlu RS, Itoharu S (2000) Cortex-restricted disruption of NMDAR1 impairs neuronal patterns in the barrel cortex. *Nature* 406:726–731.
- Jin K, Wang X, Xie L, Mao XO, Greenberg DA (2010) Transgenic ablation of doublecortin-expressing cells suppresses adult neurogenesis and worsens stroke outcome in mice. *Proc Natl Acad Sci U S A* 107:7993–7998.
- Kappeler C, Saillour Y, Baudoin JP, Tuy FPD, Alvarez C, Houbon C, Gaspar P, Hamard G, Chelly J, Metin C, Francis F (2006) Branching and nucleokinesis defects in migrating interneurons derived from doublecortin knockout mice. *Hum Mol Genet* 15:1387–1400.
- Kemp N, Bashir ZI (1999) Induction of LTD in the adult hippocampus by the synaptic activation of AMPA/kainate and metabotropic glutamate receptors. *Neuropharmacology* 38:495–504.
- Kerjan G, Koizumi H, Han EB, Dube CM, Djakovic SN, Patrick GN, Baram TZ, Heinemann SF, Gleeson JG (2009) Mice lacking doublecortin and doublecortin-like kinase 2 display altered hippocampal neuronal maturation and spontaneous seizures. *Proc Natl Acad Sci U S A* 106:6766–6771.
- Larson J, Wong DL, Lynch G (1986) Patterned stimulation at the theta frequency is optimal for the induction of hippocampal long-term potentiation. *Brain Res* 368:347–350.
- Leung LS, Fu X-W (1994) Factors affecting paired-pulse facilitation in hippocampal CA1 neurons *in vitro*. *Brain Res* 650:75–84.
- Martel G, Jaffard R, Guillou J-L (2010) Identification of hippocampus-dependent and hippocampus independent memory components in step-down inhibitory avoidance tasks. *Behav Brain Res* 207:138–143.
- Meyer U, Feldon J (2010) Epidemiology-driven neurodevelopmental animal models of schizophrenia. *Prog Neurobiol* 90:285–326.
- Miller CL, Llenos IC, Dulay JR, Weis S (2006) Upregulation of the initiating step of the kynurenine pathway in postmortem anterior cingulate cortex from individuals with schizophrenia and bipolar disorder. *Brain Res* 1073:25–37.
- Mok MHS, Fricker AC, Weil A, Kew JNC (2009) Electrophysiological characterisation of the actions of kynurenine acid at ligand-gated ion channels. *Neuropharmacology* 57:242–249.
- Munn DH (2011) Indoleamine 2,3-dioxygenase, Tregs and cancer. *Curr Med Chem* 18:2240–2246.
- Myers WA, Churchill JD, Muja N, Garraghty PE (2000) Role of NMDA receptors in adult primate cortical somatosensory plasticity. *J Comp Neurol* 418:373–382.
- Nacher J, Crespo C, McEwen BS (2001) Doublecortin expression in the adult rat telencephalon. *Eur J Neurosci* 14:629–644.
- Nathan T, Jensen MS, Lambert JDC (1990) GABA-B receptors play a major role in paired-pulse facilitation in area CA1 of the rat hippocampus. *Brain Res* 531:55–65.
- Neale SA, Copeland CS, Uebele VN, Thomson FJ, Salt TE (2013) Modulation of hippocampal synaptic transmission by the kynurenine pathway member xanthurenic acid and other VGLUT inhibitors. *Neuropsychopharmacology* 38:1060–1067.
- Neher E, Sakaba T (2001) Estimating transmitter release rates from postsynaptic current fluctuations. *J Neurosci* 21:9638–9654.
- Nilsson LK, Linderholm KR, Erhardt S (2006) Subchronic treatment with kynurenine and probenecid: effects on prepulse inhibition and firing of midbrain dopamine neurons. *J Neural Transm* 113:557–571.
- Niswender CM, Johnson KA, Weaver CD, Jones CK, Xiang Z, Luo QW, Rodriguez AL, Marlo JE, de Paulis T, Thompson A, Days EL, Nalywajko T, Austin CA, Williams MB, Ayala JE, Williams R, Lindsley CW, Conn PJ (2008) Discovery, characterization and antiparkinsonian effect of novel positive allosteric modulators of metabotropic glutamate receptor 4. *Mol Pharmacol* 74:1345–1358.
- O’Kane EM, Stone TW, Morris BJ (2003) Activation of Rho GTPases by synaptic transmission in the hippocampus. *J Neurochem* 87:1309–1312.
- Okuda S, Nishiyama N, Saito H, Katsuki H (1998) 3-Hydroxykynurenine, an endogenous oxidative stress generator, causes neuronal cell death with apoptotic features and region selectivity. *J Neurochem* 70:299–307.
- Otis TS, Wu YC, Trussell LO (1996) Delayed clearance of transmitter and the role of glutamate transporters at synapses with multiple release sites. *J Neurosci* 16:1634–1644.
- Ozaki M, Tohyama K, Kishida H, Buonanno A, Yano R, Hashikawa T (2000) Roles of neuregulin in synaptogenesis between mossy fibers and cerebellar granule cells. *J Neurosci Res* 59:612–623.
- Palma V, Lim DA, Dahmane N, Sanchez P, Brionne TC, Herzberg CD, Gitton Y, Carleton A, Alvarez-Buylla A, Altaba ARI (2005) Sonic hedgehog controls stem cell behavior in the postnatal and adult brain. *Development* 132:335–344.
- Parish CL, Castelo-Branco G, Rawal N, Tonnesen J, Sorensen AT, Salto C, Kokaia M, Lindvall O, Arenas E (2008) Wnt5a-treated midbrain neural stem cells improve dopamine cell replacement therapy in parkinsonian mice. *J Clin Invest* 118:149–160.
- Pena F, Vargas J, Tapia R (2002) Paired-pulse facilitation is turned into paired-pulse depression in hippocampal slices after epilepsy induced by 4-aminopyridine *in vivo*. *Neuropharmacology* 42:807–812.
- Perkins MN, Stone TW (1982) An iontophoretic investigation of the action of convulsant kynurenines and their interaction with the endogenous excitant quinolinic acid. *Brain Res* 247:184–187.
- Pfaffl MW, Horan GW, Dempfle L (2002) Relative expression software tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res* 30:e36.
- Pocivavsek A, Wu H-Q, Elmer GI, Bruno JP, Schwarcz R (2012) Pre- and postnatal exposure to kynurenine causes cognitive deficits in adulthood. *Eur J Neurosci* 35:1605–1612.
- Potter MC, Elmer GI, Bergeron R, Albuquerque EX, Guidetti P, Wu H-Q, Schwarcz R (2010) Reduction of endogenous kynurenine acid formation enhances extracellular glutamate, hippocampal plasticity and cognitive behaviour. *Neuropsychopharmacology* 35:1734–1742.
- Prajerova I, Honsa P, Chvatal A, Anderova M (2010) Distinct effects of sonic hedgehog and Wnt-7a on differentiation of neonatal neural stem/progenitor cells *in vitro*. *Neuroscience* 171:693–711.
- Rajan I, Cline HT (1998) Glutamate receptor activity is required for normal development of tectal cells dendrites *in vivo*. *J Neurosci* 18:7836–7846.
- Ramakers C, Ruijter JM, Deprez RHL, Moorman AFM (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci Lett* 339:62–66.
- Ramoas AS, Mower AF, Liao D, Safri SIA (2001) Suppression of cortical NMDA receptor function prevents development of orientation selectivity in the primary visual cortex. *J Neurosci* 21:4299–4309.
- Rodgers J, Stone TW, Barratt MP, Bradley B, Kennedy PG (2009) Kynurenine pathway inhibition reduces CNS inflammation in a model of human African trypanosomiasis. *Brain* 132:1259–1267.
- Rosenmund C, Stevens CF (1996) Definition of the readily releasable pool of vesicles at hippocampal synapses. *Neuron* 16:1197–1207.

- Röver S, Cesura AM, Huguenin P, Kettler R, Szente A (1997) Synthesis and biochemical evaluation of N-(4-phenylthiazol-2-yl)benzenesulfonamides as high-affinity inhibitors of kynurenine 3-hydroxylase. *J Med Chem* 40:4378–4385.
- Saito K, Nowak TS, Suyama K, Quearry BJ, Saito M, Crowley JS, Markey SP, Heyes MP (1993) Kynurenine pathway enzymes in brain – responses to ischemic brain injury versus systemic immune activation. *J Neurochem* 61:2061–2070.
- Saviane C, Savtchenko LP, Raffaelli G, Voronin LL, Cherubini E (2002) Frequency-dependent shift from paired-pulse facilitation to paired-pulse depression at unitary CA3–CA3 synapses in the rat hippocampus. *J Physiol (Lond)* 544:469–476.
- Schwarcz R, Pellicciari R (2002) Manipulation of brain kynurenines: glial targets neuronal effects and clinical opportunities. *J Pharmacol Exp Therap* 303:1–10.
- Schwarcz R, Bruno JP, Muchowski PJ, Wu HQ (2012) Kynurenines in the mammalian brain: when physiology meets pathology. *Nat Rev Neurosci* 13:465–477.
- Shin M, Yamada O, Sano Km Umezawa C (1982) Effect of branched-chain α -keto acids on kynurenine-3-hydroxylase from rat liver. *J Nutr Sci Vitaminol* 28:203–207.
- Simon DK, Prusky GT, O'Leary DDM, Constantine-Paton M (1992) NMDA receptor antagonists disrupt the formation of a mammalian neural map. *Proc Natl Acad Sci U S A* 89:10593–10597.
- Sims JR, Lee SW, Topalkara K, Qiu J, Xu J, Zhou Z, Moskowitz MA (2009) Sonic hedgehog regulates ischemia/hypoxia-induced neural progenitor proliferation. *Stroke* 40:3618–3626.
- Spassky N, Han Y-G, Aguilar A, Strehl L, Besse L, Laclef C, Ros MR, Garcia-Verdugo JM, Alvarez-Buylla A (2008) Primary cilia are required for cerebellar development and Shh-dependent expansion of progenitor pool. *Dev Biol* 317:246–259.
- Stone TW, Darlington LG (2002) Endogenous kynurenines as targets for drug discovery and development. *Nat Rev Drug Discov* 1:609–620.
- Stone TW, Darlington LG (2013) The kynurenine pathway as a therapeutic target in cognitive and neurodegenerative disorders. *Br J Pharmacol* 169:1211–1277.
- Stone TW, Perkins MN (1981) Quinolinic acid: a potent endogenous excitant at amino acid receptors in CNS. *Eur J Pharmacol* 72:411–412.
- Stone TW, Ceruti S, Abbracchio M (2009) Adenosine receptors and neurological disease: neuroprotection and neurodegeneration. *Handb Exp Pharmacol* 193:535–587.
- Stone TW, Stoy N, Darlington LG (2013) An expanding range of targets for kynurenine metabolites of tryptophan. *Trends Pharmacol Sci* 34:136–143.
- Stringer JL, Lothman EW (1989) Repetitive seizures cause an increase in paired-pulse inhibition in the dentate gyrus. *Neurosci Lett* 105:91–95.
- Taleb O, Maammar M, Brumar D, Bourguignon JJ, Schmitt M, Klein C, Kemmel V, Maitre M, Mensah-Nyagan AG (2012) Xanthurenic acid binds to neuronal G-protein-coupled receptors that secondarily activate cationic channels in the cell line NCB-20. *PLoS One* 7:e48553.
- Tang M, Villaescusa JC, Luo SX, Guitarte C, Lei S, Miyamoto Y, Taketo MM, Arenas E, Huang EJ (2010) Interactions of wnt/beta-catenin signaling and sonic hedgehog regulate the neurogenesis of ventral midbrain dopamine neurons. *J Neurosci* 30:9280–9291.
- Traiffort E, Charytoniuk DA, Faure H, Ruat M (1998) Regional distribution of sonic hedgehog patched and smoothened mRNA in the adult rat brain. *J Neurochem* 70:1327–1330.
- Traiffort E, Moya KL, Faure H, Hassig R, Ruat M (2001) High expression and anterograde axonal transport of aminoterminal sonic hedgehog in the adult hamster brain. *Eur J Neurosci* 14:839–850.
- Traiffort E, Angot E, Ruat M (2010) Sonic hedgehog in the mammalian brain. *J Neurochem* 113:576–590.
- Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, Hansen KB, Yuan H, Myers SJ, Dingledine R (2010) Glutamate receptor ion channels: structure regulation and function. *Pharmacol Rev* 62:405–496.
- Udin SB, Grant S (1999) Plasticity in the tectum of *Xenopus laevis*: binocular maps. *Prog Neurobiol* 59:81–106.
- Ultanir SK, Kim JE, Hall BJ, Deerinck T, Ellisman M, Ghosh A (2007) Regulation of spine morphology and spine density by NMDA receptor signaling in vivo. *Proc Natl Acad Sci U S A* 104:19553–19558.
- Vaillant C, Monard D (2009) SHH pathway and cerebellar development. *Cerebellum* 8:291–301.
- Vincent A, Kessler JP, Baude A, Dipasquale E, Tell F (2004) N-methyl-D-aspartate receptor activation exerts a dual control on postnatal development of nucleus tractus solitarius neurons in vivo. *Neuroscience* 126:185–194.
- Waldeck RF, Pereda A, Faber DS (2000) Properties and plasticity of paired-pulse depression at a central synapse. *J Neurosci* 20:5312–5320.
- Walker DW, Curtis B, Lacey B, Nitsos I (1999) Kynurenic acid in brain and cerebrospinal fluid of fetal newborn and adult sheep and effects of placental embolization. *Pediatr Res* 45:820–826.
- Wang MZ, Jin P, Bumcrot DA, Marigo V, McMahon AP, Wang EA, Woolf T, Pang K (1995) Induction of dopaminergic neuron phenotype in the midbrain by sonic hedgehog protein. *Nat Med* 1:1184–1188.
- Wang LY, Kaczmarek LK (1998) High-frequency firing helps replenish the readily releasable pool of synaptic vesicles. *Nature* 394:384–388.
- Wieroińska JM, Stachowicz K, Pałucha-Poniewiera A, Acher F, Brański P, Pilc A (2010) Metabotropic glutamate receptor 4 novel agonist LSP1-2111 with anxiolytic, but not antidepressant-like activity, mediated by serotonergic and GABAergic systems. *Neuropharmacology* 59:627–634.
- Wu SM, Tan KS, Chen H, Beh TT, Yeo HC, Ng SK-L, Wei S, Lee D-Y, Choo AB-H, Chan KK-K (2012) Enhanced production of neuroprogenitors dopaminergic neurons and identification of target genes by overexpression of sonic hedgehog in human embryonic stem cells. *Stem Cells Dev* 21:729–741.
- Xiong K, Luo D-W, Patrylo PR, Luo X-G, Struble RG, Clough RW, Yan X-X (2008) Doublecortin-expressing cells are present in layer II across the adult guinea pig cerebral cortex: partial colocalization with mature interneuron markers. *Exp Neurol* 211:271–282.
- Zucker RS, Delaney KR, Mulkey R, Tank DW (1991) Presynaptic calcium in transmitter release and posttetanic potentiation. *Ann N Y Acad Sci* 635:191–207.
- Zuckerman L, Weiner I (2005) Maternal immune activation leads to behavioral and pharmacological changes in the adult offspring. *J Psychiatry Res* 39:311–323.