




# REVISTA ARGENTINA DE MICROBIOLOGÍA

[www.elsevier.es/ram](http://www.elsevier.es/ram)



## ORIGINAL ARTICLE

# Tolerance of dark septate endophytic fungi (DSE) to agrochemicals *in vitro*

Federico N. Spagnoletti, Viviana M. Chiocchio\*

Cátedra de Microbiología Agrícola, Facultad de Agronomía-UBA, Instituto de Investigaciones en Biociencias Agrícolas y Ambientales-CONICET, Av. San Martín 4453, (DSE 1427) CABA, Argentina

Received 27 June 2018; accepted 21 February 2019

### KEYWORDS

Dark septate  
endophytic fungus  
(DSE);  
Wheat;  
Agrochemicals

**Abstract** Dark septate endophytes (DSE) are a heterogeneous group of fungi, mostly belonging to the Phylum Ascomycota, that are involved in a mutualistic symbiosis with plant roots. The aim of this study is to evaluate the behavior of two strains of DSE isolated from wheat roots of two cropping areas in the province of Buenos Aires, Argentina, against some agrochemicals. Of all the isolates obtained, two strains were identified as *Alternaria alternata* and *Cochliobolus* sp. These DSE were found to be tolerant to glyphosate, carbendazim and cypermethrin when evaluated at the recommended agronomic dose (AD), 2 AD and, in some cases, 10 AD. This work contributes to the study of the biology of this group of fungi and their tolerance in the presence of xenobiotics widely used in agriculture.

© 2019 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### PALABRAS CLAVE

Hongos endófitos  
septados oscuros  
(DSE);  
Trigo;  
Agroquímicos

**Tolerancia de hongos endofíticos septados oscuros a agroquímicos *in vitro***

**Resumen** Los endófitos septados oscuros (DSE) son un grupo heterogéneo de hongos que participan de una simbiosis mutualista con raíces de plantas, perteneciendo principalmente al Phylum Ascomycota. El objetivo de este estudio fue aislar DSE de raíces de trigo proveniente de dos áreas de cultivo de la provincia de Buenos Aires y evaluar el comportamiento de dos cepas de DSE aisladas de raíces de trigo frente a algunos agroquímicos en dos áreas de cultivo de la provincia de Buenos Aires. De todos los aislamientos obtenidos se seleccionaron dos cepas que se identificaron como *Alternaria alternata* y *Cochliobolus* sp. Se encontró que estos DSE

\* Corresponding author.

E-mail address: [chiocchi@agro.uba.ar](mailto:chiocchi@agro.uba.ar) (V.M. Chiocchio).

<https://doi.org/10.1016/j.ram.2019.02.003>

0325-7541/© 2019 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

son tolerantes al glifosato, el carbendazim y la cipermetrina, evaluados a las dosis agronómicas recomendadas (AD), a 2x AD y, en algunos casos, a 10x AD. Este trabajo contribuye al conocimiento de la biología de este grupo de hongos y su tolerancia a xenobióticos ampliamente utilizados en la agricultura.

© 2019 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Agricultural practices include the application of different pesticides to maintain crop performance by controlling insects, diseases and weeds. In nature, pesticide residues are subjected to physical, chemical and biochemical degradation processes, although they persist in the environment due to their high stability and water solubility<sup>22</sup>. The inadequate use of pesticides can affect microorganisms and their activity, causing alterations in the biological processes that impact on soil fertility and, consequently, on crop productivity<sup>8</sup>. Pesticides can reduce the production of microbial biomass; this change may cause alterations in the mineralization of organic matter, redox reactions, denitrification, nitrification, ammonification, and methanogenesis<sup>13</sup>. One of the most widely used pesticide is glyphosate, a potent, broad spectrum, non-selective, post-emergent herbicide that is capable of controlling 97% of the weeds. Glyphosate is generally used along with cypermethrin, an insecticide belonging to the family of pyrethroids and widely used in no-tillage systems. Among the fungicides, carbendazym is a systemic fungicide used for controlling a broad range of fungi affecting crops.

In the soil environment there are different types of microorganisms, such as fungi that participate actively in all the processes of transformation of organic matter. Fungal endophytes are organisms that colonize plant tissues during a specific period of their life cycle but cause no symptoms of tissue damage to their hosts<sup>29</sup>. Among this highly diverse group of endophytic fungi, dark septate endophytes (DSE) have received much attention in the recent years<sup>27,31</sup>. The genera *Alternaria* sp. and *Curvularia* sp. are widely known as pathogenic fungi in several cereals of commercial value<sup>1,40</sup>; however, species of the genera *Alternaria* and *Curvularia* (teleomorph: *Cochliobolus*) have been found to colonize grasses, playing a role as DSE fungi and promoting plant growth<sup>14,24</sup>. These DSE fungi are not thought to be pathogenic, as they are observed on healthy plants<sup>3</sup>.

Colonization patterns by DSE seem to be different according to the plant host and the cultivation system<sup>21</sup>. These fungi often inhabit oligotrophic soils that are associated with the roots of hundreds of plant species in all climate regions and major biome types<sup>33</sup> as well as promoters of plant growth under greenhouse conditions<sup>27</sup>. Dark septate endophytes are able to hydrolyze major C, N and P polymers into usable subunits and therefore could provide N and P to the host plant<sup>5</sup>. Newsham<sup>21</sup> performed a meta-analysis of independent studies to assess the inoculation

of DSE in different crops, concluding that the contents of N and P in inoculated plant biomass were higher compared to non-inoculated plants. Conversely, another meta-analysis suggested negative to neutral effects of DSE inoculations<sup>18</sup>.

Wheat is one of the winter cereals of major importance in Latin America. Agricultural production is one of the fundamental economic activities of Argentina, which is among the main world producers of crops such as soybeans, wheat, sunflower and corn. In Argentina, the production area destined for the wheat crop was estimated in 5.95 million hectares, producing 18 million tons of the cereal<sup>19</sup>. Most studies have evaluated mycorrhizal and DSE root colonization in different hosts and Argentine regions<sup>17,28</sup>. Nevertheless, little information is available about the biology of DSE isolated from cereals<sup>27,31</sup>. The widespread use of pesticides has raised concerns about their potential impact on microbial communities and in this sense there are no studies evaluating the effects of agrochemicals on DSE. Therefore, the aim of this study is to isolate DSE from wheat roots and to assess the tolerance of isolated strains to different chemicals commonly used in agricultural practices.

## Materials and methods

Thirty wheat plants were randomly collected from two sites located in Arenales and Ferré in July 2011 (Buenos Aires Province, Argentina). The plants were at phenological stage Z5. Samples were transported to the laboratory and placed in a cold chamber at 4 °C until their use. The roots were washed under tap water and sterilized superficially by immersing them in 70% ethanol for 2 min followed by 1% sodium hypochlorite (NaOCl) for 3 min and thoroughly rinsed five times with sterile distilled water. Subsequently, twenty five root pieces (5 mm) of each plant were placed into drops of Gel-Gro (ICN Biochemicals, USA) with streptomycin and tetracycline hydrochloride 1% (Sigma-Aldrich, USA) in sterile Petri Plates 30 Plates<sup>30</sup>. The root pieces were incubated in the dark at 25 °C and were periodically observed under a stereoscopic microscope (Nikon H550S, Japan). The segments of roots in which fungal growth was observed were placed on plates containing Malt Extract Agar (MEA) and cultured on such medium at 25 °C, thereby obtaining several isolates. The endophytic nature of the thirty isolates was corroborated following the test of resynthesis (Koch's postulates) under greenhouse conditions<sup>31</sup>. Five isolates were identified following classical<sup>10,36</sup> and molecular methodologies<sup>31</sup>. The strains were deposited at the Fungi Bank of the Facultad de Agronomía, Universidad de Buenos




Figure 1 Morphology of DSE's colony grown in MEA plate. (A) *Alternaria alternata*. (B) *Cochliobolus* sp.

Aires<sup>27</sup>. Two isolates were selected for this assay due to their extracellular enzymatic production<sup>35</sup>, their phosphorus solubilization<sup>31</sup> and their tolerance to different sodium salts<sup>32</sup>.

A culture of each strain grown on MEA for 7 days at 25 °C was used as inoculum to evaluate their tolerance to pesticides. Subsequently, two successive picks were conducted on basal salt medium (BSM)<sup>15</sup> with the purpose of causing them a nutritional stress (starvation). The BSM containing the herbicide (glyphosate), the fungicide (carbendazim) or the insecticide (cypermethrin) was used in an agronomic common dose (AD), 2 AD and 10 AD. Doses per Petri dish were 0.20 ppm, 0.40 ppm and 2 ppm for the herbicide; 18.12 ppm, 36.25 ppm and 181.25 ppm for the fungicide; and 2.02 ppm, 4.04 ppm and 20.2 ppm for the insecticide respectively. The control treatment was BSM medium plus sucrose (2%). The inoculum was standardized using 5 mm-diameter disks obtained from the zone with active growth of a colony with 7 days of growth for all strains and all tested media. Each agrochemical dose and its respective control were tested in quintuplicate. Fungal growth was assessed periodically from the third day by measuring the colony diameter at 12 days. Fungal growth rates were analyzed using the following formula<sup>26</sup>: Growth rate =  $(Fd - Id) / (Ft - It)$ , where: Fd is the final diameter (cm), Id is the initial diameter (cm), Ft is the final time (days) and It is the initial time (days).

Moreover, the inhibition percentage was calculated for each fungi and each agrochemical at 6 and 12 days using the following formula: Inhibition (%) =  $(Cd - Td) / Cd \times 100$ , where: Cd is the mean colony diameter in the control treatment (cm) and Td is the mean colony diameter in the agrochemical treatment (cm).

## Statistical analysis

A two-way analysis of variance (ANOVA) with DSE strains (*A. alternata* and *Cochliobolus* sp.) and agrochemical type (4 levels: 0; AD; 2 AD; 10 AD) as factors was performed from the data obtained on growth parameters and effective concentration (EC 50) using the statistical software Infostat Professional version 2.0. Normality and homogeneity of variance were tested. Comparisons were made using the Tukey's test with  $p < 0.05$ .

## Results

The physicochemical properties of the soil (Typic Argiudoll) were: organic carbon 1.7%, available phosphorous 15.0 mg/kg (Kurtz and Bray No. 1 method), pH 6.2, electrical conductivity 0.5 d/Sm clay 27.5%, silt 50.5% and sand 22%. Figure 1 shows the morphology of the DSE colony of *A. alternata* and *Cochliobolus* sp.

The strain *Cochliobolus* sp. reached a diameter of 75 mm of the colony after 12 days of growth while *A. alternata* completed the culture dish (90 mm) after 9 days (data not shown). There was significant change in the kinetics of growth of the strains by adding the different kinds of agrochemicals to the medium. The results from the two-way analysis of variance showed that there were significant interactions between the DSE strains and the agrochemicals in growth rate ( $p < 0.0001$ ). Therefore, this interaction was further analyzed. Table 1 shows that the growth of *A. alternata* was significantly faster than that of *Cochliobolus* sp. in the control treatment and was less affected than *Cochliobolus* sp. when the agrochemicals were added.

The presence of glyphosate reduced the growth rate of both DSE strains. *A. alternata* showed significant differences in AD and 2 AD regarding *Cochliobolus* sp.; however, when the dose was increased to 10 AD, no differences were found between both strains.

The fungicide carbendazim was the most tolerated agrochemical by the DSE strains. The growth rate of *A. alternata* was reduced by the addition of carbendazim as its dose increased in the solid media. No differences were found in 2 AD between both fungi (Table 1). On the other hand, the insecticide (cypermethrin) was the most toxic agrochemical on growth rate. *A. alternata* reduced its growth rate as the dose of cypermethrin increased in the media, while a dose equal to or greater than 2 AD inhibited the growth of *Cochliobolus* sp. (Table 1).

The percentages of growth inhibition are shown in Table 2A-C. The herbicide (Table 2A) increased this parameter ( $p < 0.05$ ) in all the agrochemicals and doses tested for both fungi and on both days evaluated (day 6 and day 12). The fungicide increased the growth inhibition percentage of *Cochliobolus* sp.; however, an induction of the growth of *A. alternata* was observed when the fungicide was added at AD and 2 AD evaluated at 6 days. In these doses, at the end of

**Table 1** Growth rate of *Cochliobolus* sp. and *Alternaria alternata* (DSE strains). Comparison between DSE strains *A. alternata* and *Cochliobolus* sp. and agrochemical type.

Agrochemicals	Treatments	DSE strains	
		<i>Cochliobolus</i> sp	<i>Alternaria alternata</i>
Glyphosate	Control	0.53 ± 0.06 b	0.58 ± 0.01 a
	AD	0.25 ± 0.01 e	0.42 ± 0.02 c
	2 AD	0.22 ± 0.01 f	0.37 ± 0.03 d
	10 AD	0.05 ± 0.01 g	0.03 ± 0.00 g
Carbendazim	Control	0.53 ± 0.06 b	0.58 ± 0.01 a
	AD	0.42 ± 0.00 c	0.51 ± 0.00 b
	2 AD	0.46 ± 0.05 c	0.46 ± 0.02 c
	10 AD	0.21 ± 0.05 f	0.33 ± 0.02 d
Cypermethrin	Control	0.53 ± 0.06 b	0.58 ± 0.01 a
	AD	0.19 ± 0.04 f	0.54 ± 0.02 b
	2 AD	0.00 ± 0.00 g	0.20 ± 0.00 f
	10 AD	0.00 ± 0.00 g	0.03 ± 0.00 g

4 levels of agrochemicals: 0; AD (agronomic dose); 2 AD; 10 AD. Means ± SD ( $n=5$ ). Different letters differ significantly ( $p < 0.05$ ).

**Table 2** Inhibition of growth rate of *Cochliobolus* sp. and *Alternaria* sp. (DSE strains) growing at different concentrations of: (A) glyphosate, (B) carbendazim and (C) cypermethrin.

(A) Strains	Day	Glyphosate			
		0	AD	2 AD	10 AD
<i>Cochliobolus</i> sp.	6	0.00 ± 0.00 a	36.70 ± 2.14 b	41.53 ± 0.91 c	78.68 ± 1.25 d
	12	0.00 ± 0.00 a	44.28 ± 1.90 b	50.56 ± 1.23 c	84.29 ± 2.25 d
<i>A. alternata</i>	6	0.00 ± 0.00 a	35.86 ± 3.25 b	46.89 ± 1.98 c	78.96 ± 1.30 d
	12	0.00 ± 0.00 a	24.95 ± 3.68 b	36.00 ± 3.14 c	81.87 ± 3.95 d
(B) Strains	Day	Carbendazim			
		0	AD	2 AD	10 AD
<i>Cochliobolus</i> sp.	6	0.00 ± 0.00 a	6.37 ± 3.42 b	10.76 ± 3.59 b	35.16 ± 4.32 c
	12	0.00 ± 0.00 a	11.70 ± 1.21 b	12.37 ± 4.28 b	49.67 ± 2.06 c
<i>A. alternata</i>	6	0.00 ± 0.00 b	-4.31 ± 1.05 a	-2.82 ± 1.96 a	36.03 ± 2.14 c
	12	0.00 ± 0.00 a	0.00 ± 0.00 a	0.00 ± 0.00 a	47.62 ± 2.18 b
(C) Strains	Day	Cypermethrin			
		0	AD	2 AD	10 AD
<i>Cochliobolus</i> sp.	6	0.00 ± 0.00 a	59.12 ± 3.24 b	84.61 ± 0.00 c	84.61 ± 0.00 c
	12	0.00 ± 0.00 a	59.87 ± 4.48 b	90.60 ± 0.00 c	90.60 ± 0.00 c
<i>A. alternata</i>	6	0.00 ± 0.00 a	26.03 ± 0.94 b	64.24 ± 2.79 c	79.82 ± 0.77 d
	12	0.00 ± 0.00 a	6.50 ± 3.73 b	62.12 ± 2.74 c	83.12 ± 0.00 d


Each value represents the mean value ± SD ( $n=5$ ). Different letters differ significantly ( $p < 0.05$ ).

the growth (day 12) no significant differences were detected with respect to the control treatment. For both evaluation days, the addition of carbendazim at 10 AD generated a significant increase ( $p < 0.05$ ) in the inhibition percentage (Table 2B).

Table 2C shows the percentages of growth inhibition generated by the insecticide. The DSE *Cochliobolus* sp. exhibited significant inhibition when the AD was placed in the culture medium ( $p < 0.05$ ); however, no changes were detected between 2 AD and 10 AD. On the contrary, *A.*

*alternata* recorded increases in inhibition in all the doses studied ( $p < 0.05$ ).

The concentrations of the agrochemicals that caused a 50% reduction in fungal growth compared to the control (effective concentration: EC 50) varied depending on the agrochemical type (Fig. 2). In all the agrochemicals analyzed, the fungus *A. alternata* was the most tolerant, presenting higher EC 50. The insecticide was the agrochemical that caused the greatest effect on DSE growth, followed by the herbicide. The fungicide proved to be the least toxic



**Figure 2** Effective concentration (EC 50), the concentrations of the herbicide, fungicide and insecticide that caused a 50% reduction in fungal growth compared to the control. Means  $\pm$  SD ( $n=5$ ). Different letters differ significantly ( $p < 0.05$ ).

agrochemical for these fungi (Fig. 2). *Cochliobolus* sp. did not differ significantly from *A. alternata* in the EC 50 against carbendazim.

## Discussion

This study evaluates the tolerance of two DSE fungi to different agrochemicals. The tested DSE fungi were isolates of *A. alternata* and *Cochliobolus* sp. It is known that these fungi are grass pathogens; however, several studies have shown that they can also be found in the roots as DSE as non-pathogenic fungi<sup>3,37</sup>. Therefore, the fungal species classified as non-pathogenic cause disease symptoms in certain host plants but are neutral or beneficial in other hosts.

Moreover, although pesticides are designed to be as specific as possible, they have effects on non-target fungal organisms. In accordance with our results, glyphosate (using equivalent herbicide concentrations) had an *in vitro* toxic effect on culturable mycobiota (*Aspergillus* section Flavi) strains from agricultural soils<sup>6</sup>. Likewise, other authors<sup>12</sup> found a reduction of microbial biomass at a higher glyphosate concentration and a temporary inhibitory effect at recommended field doses. Furthermore, effects of glyphosate on soil microbial communities and its mineralization in soils after various applications of this herbicide have been found<sup>16</sup>. A recent study<sup>11</sup> also demonstrated behavioral differences between glyphosate and its major metabolite, AMPA, related to the physical properties of saturated hydraulic conductivity (Ks) and soil moisture. Other authors found that glyphosate affects arbuscular mycorrhizal fungi due to a reduction in spore viability, root colonization and arbuscule percentage<sup>9</sup>. However, DSEs are more tolerant to different types of stress than other soil fungi<sup>23</sup>. This could be due to the fact that the chemical compounds act effectively on the mycelium, while spores and chlamydospores are more resistant to these compounds due to their low activity given their water content and the presence of a large amount of unsaturated fatty acids<sup>4</sup>.

There is much less research on the effect of carbendazim on fungi. However, this fungicide has detrimental effects to fungal and bacterial biocontrol agents such as *Trichoderma harzianum*, *Trichoderma virens*, *Bacillus subtilis* and *Pseudomonas fluorescens*<sup>20</sup>. In this research, we found that DSE strains have the potential to tolerate high concentrations (181.25 ppm, 10 AD) of fungicide; these

results are in accordance with those of others authors<sup>2</sup>. One study showed that the type of fungicide applied to the soil must be seriously considered as the authors detected negative effects on *Rhizophagus fasciculatus* (arbuscular mycorrhizal fungi) with benomyl (systemic benzimidazole); however, they found no effects when captan (non-systemic fungicide)<sup>7</sup> was applied. Nevertheless, our results showed high tolerance of DSE fungi against carbendazim, which is a derivative of benomyl. *Bipolaris tetramera* and *A. alternata* have potential to tolerate and degrade 100 ppm of carbendazim *in vitro*<sup>38</sup>. In accordance with that article, we found negative inhibition values in agronomic doses, 2 AD and no differences compared with control. These results would indicate an *Alternaria* sp. promotion growth when grown with the addition of carbendazim.

In the case of cypermethrin, there are studies that demonstrate that this insecticide caused a negative effect on soil enzymatic activities and microbial diversity<sup>34</sup>. Our DSE strains showed low cypermethrin tolerance, high inhibition percentage and low EC50. Nevertheless, a study revealed that when the pesticide is applied on pepper plants there is an increase in bacterial abundance and a change in the composition of the community of the Firmicutes phylum to Bacteroidetes and  $\gamma$ -Proteobacteria phyla<sup>39</sup>. Other authors showed the potential of *Aspergillus niger*, *Aspergillus terreus*, *Monilochætes* and *Fusarium* strains in the biodegradation of 50–150 ppm of this pyrethroid<sup>25</sup>.

This research has shown that agrochemicals commonly used in agriculture have detrimental effects on the growth of DSE fungi, even when these agrochemicals are used at recommended doses; however, these negative effects vary according to the agrochemical considered. Moreover, this paper makes evident that DSE response to this type of toxin depends on the studied species of DSE fungus. This research is a contribution to a better understanding of the behavior of DSE against xenobiotics, evaluating the tolerance and/or their use as a source of nutrients.

## Conclusion

The DSE *A. alternata* tolerates all three agrochemicals at the tested doses while *Cochliobolus* sp. is glyphosate and carbendazim-tolerant in all three doses but shows no tolerance to cypermethrin when the agronomic dose is twofold. In general, there is a decrease in the growth of the strains when increasing the dose of the agrochemicals studied.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

The present research was financially supported by the Agencia Nacional de Promoción Científica y Tecnológica, Project PICT 2016:1136 and the Universidad de Buenos Aires, UBA-CYT 068/2011, Facultad de Agronomía, Argentina. We thank Ing. Raúl Silvio Lavado the critical review of this manuscript.

## References

1. Andersen B, Nielsen KF, Pinto VF, Patriarca A. Characterization of *Alternaria* strains from Argentinean blueberry, tomato, walnut and wheat. *Int J Food Microbiol.* 2015;196:1–10.
2. Bending G, Rodrigues-Cruz MS, Lincoln SD. Fungicide impacts on microbial communities in soils with contrasting management histories. *Chemosphere.* 2007;69:8–82.
3. Bonfim JA, Vasconcellos RLF, Baldesin LF, Sieber TN, Cardoso EJBN. Dark septate endophytic fungi of native plants along an altitudinal gradient in the Brazilian Atlantic forest. *Fungal Ecol.* 2016;20:10–202.
4. Bundgard-Nielsen K, Nielsen PV. Fungicidal effect of disinfectants against fungal contaminants commonly found in bread and cheese manufacturing. *J Food Protect.* 1995;59:75–268.
5. Caldwell BA, Jumpponen A, Trappe JM. Utilization of major detrital substrates by dark-septate, root endophytes. *Mycologia.* 2000;2:2–230.
6. Carranza CS, Barberis CL, Chiacchiera SM, Dalcero AM, Magnoli CE. Isolation of culturable mycobiota from agricultural soils and determination of tolerance to glyphosate of nontoxicogenic *Aspergillus* section *Flavi* strains. *J Environ Sci Health B.* 2016;51:35–43.
7. Channabasava A, Lakshman HC, Jorquera MA. Effect of fungicides on association of arbuscular mycorrhiza fungus *Rhizophagus fasciculatus* and growth of Proso millet (*Panicum miliaceum* L.). *J Soil Sci Plant Nutr.* 2015;15:35–45.
8. Cycon M, Piotrowska-Seget Z, Kozdrój J. Responses of indigenous microorganisms to a fungicidal mixture of mancozeb and dimethomorph added to sandy soil. *Intern Biodeter Biodegrad.* 2010;64:316–23.
9. Drulle M, Omacini M, Golluscio RA, Cabello MN. Arbuscular mycorrhizal fungi are directly and indirectly affected by glyphosate application. *Appl Soil Ecol.* 2013;72:143–9.
10. Ellis MB. Dematiaceous hyphomycetes. UK: Commonwealth Mycological Institute; 1971. p. pp608.
11. Erban T, Stehlík M, Sopko B, Markovic M, Seifrtova M, Halesova T, Kovaricek P. The different behaviors of glyphosate and AMPA in compost-amended soil. *Chemosphere.* 2018;207:78–83.
12. Gomez E, Ferreras L, Lovotti L, Fernandez E. Impact of glyphosate application on microbial biomass and metabolic activity in a Vertic Argiudoll from Argentina. *Eur J Soil Biol.* 2009;45:7–163.
13. Hussain S, Siddique T, Saleem M, Arshad M, Khalid A. Impact of pesticides on soil microbial diversity, enzymes, and biochemical reactions. *Adv Agron.* 2009;102:159–200.
14. Kageyama SA, Mandyam KG, Jumpponen A. Diversity, function and potential applications of the root-associated endophytes. In: Varma A, editor. *Mycorrhiza: genetics and molecular biology, eco-function, biotechnology, ecophysiology structure and systematic.* Springer-Verlag Berlin Heidelberg; 2008. p. 29–57.
15. Kilbane JJ, Chatterjee DK, Karns JS, Kellogg ST, Chakrabarty AM. Biodegradation of 2,4,5-trichlorophenoxyacetic acid by a pure culture of *Pseudomonas cepacia*. *Appl Environ Microbiol.* 1982;44:8–72.
16. Lancaster SH, Hollister EB, Senseman SA, Gentry TJ. Effects of repeated glyphosate applications on soil microbial community composition and the mineralization of glyphosate. *Pest Manag Sci.* 2010;66:59–64.
17. Lugo MA, Menoyo E, Allione LR, Negrito MA, Henning JA, Anton AM. Arbuscular mycorrhizas and dark septate endophytes associated with grasses from the Argentine Puna. *Mycologia.* 2018;110:65–54.
18. Mayerhofer MS, Kernaghan G, Harper KA. The effects of fungal root endophytes on plant growth: a meta-analysis. *Mycorrhiza.* 2013;23:28–119.
19. Ministerio de Agroindustria. Datos Abiertos Agroindustria. Producción de trigo en Argentina; 2017. URL <https://datos.madry.gob.ar/reportes.php?reporte=Estimaciones> [accessed 10.10.18].
20. Mohiddin FA, Khan MR. Tolerance of fungal and bacterial biocontrol agents to six pesticides commonly used in the control of soil borne plant pathogens. *Afr J Agric Res.* 2013;8:5–5272.
21. Newsham KK. A meta-analysis of plant responses to dark septate root endophytes. *New Phytol.* 2011;190:93–783.
22. Odukkathil G, Vasudevan N. Toxicity and bioremediation of pesticides in agricultural soil. *Rev Environ Sci Biotechnol.* 2013;12:44–21.
23. Postma JWM, Olsson PA, Falkengren-Gerup U. Root colonization by arbuscular mycorrhizal, fine endophytic and dark septate fungi across a pH gradient in acid beech forest. *Soil Biol Biochem.* 2007;39:8–400.
24. Priyadharsini P, Muthukumar T. The root endophytic fungus *Curvularia geniculata* from *Parthenium hysterophorus* roots improves plant growth through phosphate solubilization and phytohormone production. *Fungal Ecol.* 2017;27:69–77.
25. Qin K, Zhu LS, Wang JH. Screening and degradation characteristics of cypermethrin degrading fungi. *Chin J Environ Eng.* 2010;4:4–950.
26. Rojas Barreto JA, Hormaza Anaguano A. Evaluation of growth and compatibility of white rot fungi. *Cienc Desarroll.* 2014;5:197–205.
27. Rothen C, Miranda V, Aranda-Rickert A, Fracchia S, Rodríguez MA. Characterization of dark septate endophyte fungi associated with cultivated soybean at two growth stages. *Appl Soil Ecol.* 2017;120:9–62.
28. Russo ML, Pelizza SA, Cabello MN, Stenglein SA, Vianna MF, Scorsetti AC. Endophytic fungi from selected varieties of soybean (*Glycine max* L. Merr.) and corn (*Zea mays* L.) grown in an agricultural area of Argentina. *Rev Argent Microbiol.* 2016;48:60–154.
29. Schulz B, Boyle C. The endophytic continuum. *Mycol Res.* 2005;109:661–86.
30. Silvani VA, Fracchia S, Fernández L, Pérgola M, Godeas A. A simple method to obtain endophytic microorganisms from field-collected roots. *Soil Biol Biochem.* 2008;40:63–1259.
31. Spagnoletti FN, Tobar NE, Di Pardo AF, Chiocchio VM, Lavado RS. Dark septate endophytes present different potential to solubilize calcium, iron and aluminum phosphates. *Appl Soil Ecol.* 2017;111:25–32.
32. Spagnoletti FN, Tobar NE, Ureta Suelgaray FJ, Lavado RS, Chiocchio VM. Tolerancia de los hongos endófitos septados oscuros (DSE) a diferentes concentraciones de sales de sodio. V Reunión de la Red Argentina de Salinidad. San Luis: Villa Mercedes; 2017.
33. Su ZZ, Mao LJ, Li N, Feng XX, Yuan ZL, Wang LW, Lin FC, Zhang CL. Evidence for biotrophic lifestyle and biocontrol potential of dark septate endophyte *Harpophora oryzae* to rice blast disease. *PLoS ONE.* 2013;8:e61332.
34. Tejada M, García C, Hernández T, Gómez I. Response of soil microbial activity and biodiversity in soils polluted with different concentrations of cypermethrin insecticide. *Arch Environ Contam Toxicol.* 2015;69:8–19.
35. Tobar Gomez N, Spagnoletti F, Chiocchio V, Lavado RS. Capacidad enzimática de hongos endófiticos septados oscuros (DSE) y liberación de nutrientes. In: Diaz Zorita M, Correa OS, Fernandez Caniggia MV, Lavado RS, editors. *Aportes de la microbiología a la producción de cultivos.* 2013. p. 113–8.
36. Woudenberg JHC, Groenewald JZ, Binder M, Crous PW. *Alternaria* redefined. *Stud Mycol.* 2013;75:171–212.
37. Xu R, Li T, Cui H, Wang J, Yu X, Ding Y, Wang C, Yang Z, Zhao Z. Diversity and characterization of Cd-tolerant dark septate endophytes (DSEs) associated with the roots of Nepal alder (*Alnus*

Tolerance of dark septate endophytic fungi (DSE) to agrochemicals *in vitro*

7

- nepalensis*) in a metal mine tailing of southwest China. *Appl Soil Ecol.* 2015;93:8–11.
38. Yarden O, Katan J, Aharonson N, Ben-Yephet Y. Delayed and enhanced degradation of benomyl and carbendazim in disinfected and fungicide-treated soils. *Phytopathology.* 1984;75:7–763.
39. Zhang B, Bai Z, Hoefel D, Tang L, Wang L, Li B, Li Z, Zhuang G. The impact of cypermethrin pesticide application on the non-target microbial community of the pepper plant phyllosphere. *Sci Total Environ.* 2009;407:22–1915.
40. Zhong S, Ali S, Leng Y, Wang R, Garvin DF. *Brachypodium distachyon-Cochliobolus sativus* pathosystem is a new model for studying plant-fungal interactions in cereal crops. *Phytopathology.* 2015;105:9–482.