

1 Title: **Role of calcium in the defense response induced by brassinosteroids in**
2 **strawberry plants**

3 Running title: **Importance of calcium in brassinosteroid-signaling**

4 Ramiro N. Furio¹, Gustavo M. Martínez-Zamora¹, Sergio M. Salazar^{2,3}, Yamilet Coll⁴, Silvia
5 Marisa Perato¹, Gustavo G. Martos¹, Juan C. Díaz Ricci^{1*}

6 ¹Instituto Superior de Investigaciones Biológicas (INSIBIO), CONICET-UNT, and Instituto
7 de Química Biológica “Dr. Bernabé Bloj”, Facultad de Bioquímica, Química y Farmacia,
8 UNT. Chacabuco 461, T4000ILI – San Miguel de Tucumán, Argentina.

9 ²Instituto Nacional de Tecnología Agropecuaria. EEA Famaillá, Tucumán, Argentina.

10 ³Facultad de Agronomía y Zootecnia. UNT. Argentina.

11 ⁴Laboratorio de Bioproductos, Centro de Estudio de Productos Naturales, Universidad La
12 Habana, Cuba.

13 *Corresponding author e-mail: juan@fbqf.unt.edu.ar

14 **Role of calcium in the defense response induced by brassinosteroids in strawberry plants**

15 **Highlights**

- 16 • Calcium plays a key role in the defense induced by brassinosteroids in plants.
- 17 • Calcium-sensor calmodulin-like (CML) proteins were identified in *Fragaria ananassa*.
- 18 • Brassinosteroids induce overexpression of *CMLs* and other defense related genes.

19 **Abstract**

20 Brassinosteroids (BRs) are steroidal hormones essential for the normal growth and
21 development of plants. The aim of this work was to study the role of calcium, in the defense
22 response induced by two BRs e.g. BB16 and EP24, and to investigate the expression levels of
23 defense-related genes in strawberry plants treated with these BRs. An increase in the calcium
24 influx in response to treatment with BB16 or EP24 was observed. By comparing *Fragaria*
25 *vesca* and *Fragaria ananassa* genome sequences 28 nucleotide sequences coding for calcium-
26 sensing proteins called CMLs were predicted. Bioinformatic analyses of the sequences
27 revealed that all of them contained the characteristic EF-hand motif of Ca²⁺ binding, and
28 regulatory *cis*-elements related to stress. qPCR studies showed that both BB16 and EP24 cause
29 the upregulation of selected *CMLs* genes e.g. *FaCML1a*, *FaCML36*, *FaCML42* and
30 *FaCML45*, and defense marker genes, such as *PR1*, *ERF1* and *GLS5*. Results also showed that
31 whereas BB16 induced the upregulation of *MYB30*, EP24 induced the upregulation of *PRX27*.
32 These results provide valuable information of the response mechanisms induced by BRs and
33 demonstrate the role of calcium in the signaling pathways induced by BRs in strawberry
34 plants.

35

36 **Keywords:** Brassinosteroids; Calcium; Calmodulin-like; Defense; *Fragaria ananassa*.

37 **1. Introduction**

38 To defend themselves against pathogens, plants have developed sophisticated mechanisms to
39 perceive the attacker and convert that perception into signals that trigger an effective defense
40 response [1].

41 It has been demonstrated that Ca^{2+} mediated signaling plays an important role in the response
42 of plants to stress [2]. It has been shown that a transient increase in Ca^{2+} concentration is
43 detected by several Ca^{2+} sensors or Ca^{2+} binding proteins, regulating downstream targets as
44 part of a coordinated cellular response to specific stimuli. It has also been observed that these
45 Ca^{2+} binding proteins usually have helix-loop-helix structures called EF-hand motifs. Three
46 main classes of Ca^{2+} EF-hand containing sensors have been characterized in plants so far,
47 namely: calmodulin (CAM), Ca^{2+} -dependent protein kinase (CDPK) and calcineurin B-like
48 protein (CBL) [3]. Calmodulin (CaM) is considered to be one of the most conserved Ca^{2+}
49 binding proteins in all eukaryotes. The binding of Ca^{2+} to CaM induces a conformational
50 change, which allows the interaction of the CaM/ Ca^{2+} complex with target proteins.

51 In addition to several CAMs, plants also possess another large family of Ca^{2+} binding proteins
52 called calmodulin-like (CML) proteins. These proteins extend the range of potential calcium
53 sensors present in plant cells. The structure of CMLs diverges from the structure of typical
54 CaM proteins, suggesting that these proteins must have particular functions. Gene expression
55 profiles exhibit a wide diversity of expression patterns of CAM and CML genes through the
56 plant development program, or in response to external stimuli [4].

57 Xu et al. [5], studied the expression pattern of the gene *OsMSR2* (*Oryza sativa* L. Multi- Gen-
58 Stress Responsive 2), a new calmodulin-like protein, whose gene was isolated from rice
59 (*Oryza sativa* L.). They observed that *OsMSR2* was strongly upregulated by a wide spectrum
60 of stresses, including cold, drought and heat in different tissues and at different stages of rice
61 development. It was also shown that the expression of *OsMSR2* in Arabidopsis modulates salt
62 tolerance and drought through ABA-mediated pathways, and these transgenic plants exhibited
63 hypersensitivity to ABA during germination of seeds and the stages after germination.

64 Chiasson et al. [6] characterized a tomato gene (*APR134*) encoding a calmodulin-like protein
65 that is induced in leaves of plants resistant to *Pseudomonas syringae* pv. *tomato*. They
66 demonstrated that the suppression of *APR134* expression in tomato (*Solanum lycopersicum*),
67 using virus induced gene silencing (VIGS), compromises the immune response of the plant.
68 By investigating the presence of the *APR134* gene in Arabidopsis they could identify two

69 orthologous genes called *CML42* and *CML43*. They observed that *CML43* is rapidly induced
70 in *Arabidopsis* leaves resistant to diseases after inoculation with *Pseudomonas syringae* pv.
71 *tomato*. These data support a role for *CML43* and *APR134* as important mediators of Ca^{2+} -
72 dependent signals during the immune response of plants to bacterial pathogens.

73 In a previous work, we have shown that the treatment with BB16 induces the stomatal closure
74 in *Fragaria ananassa* [7]. Since the relationship between calcium influx and stomatal closure
75 was clearly demonstrated by Gilroy et al.[8], and Ward and Schroeder [9], in this paper we
76 decided to investigate the participation of Ca^{2+} on the effect observed of the brassinosteroids
77 BB16 (a formulation based on a brassinosteroid spirotanic analogue DI-31) and EP24 (24-
78 epibrasinolide) in strawberry plants.

79 The defense response is usually associated with the expression of several biochemical and
80 molecular factors. Furio et al. [7] have analyzed the biochemical markers associated with the
81 defense response induced by BRs in strawberry, showing that they induced an oxidative burst,
82 increase in intracellular nitric oxide (NO), stomatal closure, cell wall reinforcement by lignin
83 and callose deposition, and accumulation of calcium oxalate crystals. However, molecular
84 analysis of the genes involved, and calcium participation at the onset of the defense response
85 mediated by BRs were not performed.

86 The objective of this work was to evaluate the participation of calcium in the activation of the
87 defense response activated by BB16 and EP24, studying the ion influx and the role of CMLs.
88 Since there was scarce information about CMLs proteins in strawberry, a bioinformatics
89 approach was carried out to identify and predict CML sequences of *Fragariaananassa* that
90 have not been characterized yet. The expression of defense-related genes in strawberry plants
91 treated with both BRs was also studied.

92 **2. Materials and methods**

93 **2.1. Plant material**

94 Strawberry plants (*Fragaria ananassa*) of the cv. Pájaro were obtained from the strawberry
95 BGA (Strawberry Active Germplasm Bank at Universidad Nacional de Tucumán). Healthy
96 plantlets were obtained from *in vitro* cultures in MS medium (Sigma), rooted in pots with
97 sterilized substrate (humus and perlite, 2:1), and maintained at 28°C, 70% relative humidity
98 (RH), with a light cycle of 16h (white fluorescent, 350 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$).

99 **2.2 Plant treatment**

100 Three months old plants containing three or four fully expanded leaves were used in
101 experiments. Batches of 3 plants constituted the experimental unit, were sprayed to run-off
102 with BB16 (0.1 mg l^{-1}), EP24 (0.1 mg l^{-1}) or water (control), and maintained as indicated above
103 until picking leaf samples. Young totally expanded leaves of treated plants were harvested at
104 different times for RNA extraction and qPCR assays as indicated below. For intracellular
105 calcium determinations leaf discs were obtained from not-treated plants, and treated afterward
106 with BB16 (0.1 mg l^{-1}), EP24 (0.1 mg l^{-1}) or water (control) separately as indicated below.

107 **2.3. Calcium determination**

108 Accumulation of intracellular calcium was measured in leaf discs using the fluorescent probe
109 Fura2-AM (Invitrogen) according to Martos et al. [10]. 5 mm diameter discs were obtained
110 from freshly expanded young strawberry leaves, and 10 leaf discs from 5 plants treated with
111 BB16 or EP24 were used. The discs were placed in test tubes with 1000 μl of W5 buffer and
112 15 μM Fura2-AM, 5 pulses of vacuum were applied and they were allowed to stand in the
113 dark for 3 hours. The discs were then rinsed twice with distilled water and placed in a multi-
114 well plate with 100 μl of the W5 buffer per well. Immediately before each measurement,
115 BB16 (0.1 mg l^{-1}), EP24 (0.1 mg l^{-1}) or water (control) was added. The fluorescence excitation
116 spectra from 350 nm to 420 nm were obtained at $\lambda_{\text{em}} = 532$ nm using a photon counting
117 spectrofluorometer (ISSAPC1, Owingen, Germany). The calcium level was expressed as the
118 ratio between the relative fluorescence intensities measured at 362 nm and 402 nm. The
119 spectra were obtained every 30 s for 10 min. The spectra did not change in the analyzed time
120 range. Presented values correspond to the first measurement after 30 s.

121 **2.4. Bioinformatic analysis: CML homologous searching and gene prediction**

122 To find *CML* homologous, the Blast algorithm was used, searching by comparison
123 (alignments) of the nucleotide sequences with the sequences deposited in the GenBank NR
124 database. To find *CML* homologous, a search on the genome of *Fragaria ananassa* was
125 carried out using the BlastN homologous search program
126 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>), and taking as reference *CML* sequences already
127 predicted of *Fragaria vesca* and annotated in the GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>). The sequence analysis let us find “contigs” that contained sequences with
128 highly similar to the CMLs of *Fragaria vesca* used as a query. Amino acid sequences of
129 FaCML messengers found were *in silico* predicted using the FGenesh gene prediction
130 software [11]. Sequence assemblies were made with the DNAMAN program [12]. The
131 presence of structural motifs and/or conserved domains was determined by the CDSearch
132 (from NCBI) [13] and SMART [14] programs. Multiple alignments of amino acid sequences
133 were performed by Clustal X [15]. Molecular phylogeny analysis was performed from amino
134 acid alignments with the MEGA 4 software [16], using the nearest neighbor method (Neighbor
135 Joining, NJ) [17]. The reliability of tree branches was evaluated using the Bootstrap method
136 [18].

138 Bioinformatic analysis of the 5'UTR of the predicted sequences was carried out to identify
139 conserved promoter motifs related to stress responses in plants by using the PLACE (Plant cis-
140 acting regulatory DNA elements) algorithm [19] and Softberry program [20].

141 **2.5. RNA extraction and qPCR analysis**

142 Total RNA was obtained from the youngest totally expanded leaflet of three BRs-treated or
143 control not treated plants (n=3). Leaflets were harvested at 2, 4, 6, 12, 24, 48, and 72 hours post
144 treatments (hpt), pooled, weighted, immediately frozen in liquid nitrogen, pulverized and kept
145 at -80°C , until further use. 80 mg of leaf powder was used for RNA extraction with the
146 RNAqueous-4PCR kit (Ambion, USA). RNA concentration and purity was assessed
147 spectrophotometrically at 230, 260 and 280 nm (BioSpecmini, Shimadzu). Retrotranscription
148 was carried out using the SuperScript II Reverse Transcriptase (Invitrogen) according to
149 manufacturer instructions. qPCR reactions were performed using iQ SYBR Green Supermix
150 (Bio-Rad) in a 7500 Real-Time PCR System (Applied Biosystems). Values of the gene
151 expression level were normalized using the Elongation Factor-1a (*FaEF1a*) gene (GenBank

152 accession number: XM_004307362) according to [21,22]. PCR primers efficiency and Ct
153 values were calculated using the LinRegPCR software as recommended by Ramakers et al.
154 [23], and profiles were estimated taking the gene *FaEF-1 α* as reference and using the
155 fgStatistics software [24], and the algorithms suggested by Pfaffl [25]. Two biological
156 replicates with three technical replicates were used for each treatment. Gene expression was
157 expressed as the ratio between BRs and control expression levels.

158 **2.6. Primers Design and genes analyzed**

159 Primers sequences were obtained from the *F. ananassa* nucleotide sequences freely available
160 at the NCBI GenBank website (<https://www.ncbi.nlm.nih.gov/genbank/>), and optimized using
161 the Primer Express software (Applied Biosystems, USA) (see Table 1). The primers were used
162 to evaluate the expression level of genes associated to the defense response, and selected
163 *FaCMLs* in strawberry plants treated or not with BRs. The defense genes studied were: i)
164 *FaPRI*, encoding the Protein Related to Pathogenesis 1; ii) *FaMYB30*, HR marker gene; iii)
165 *ERF1*, transcription factor of the ET signaling pathway iv) *ETR2*, ET receptor, v)
166 *FaPRX27* encoding a plant peroxidase class III, and vi) *FaGSL5* involved in the synthesis of
167 callose. Four *CMLs* genes were selected to study the effect of BRs on their level of expression,
168 one containing 2 EF-hand motif (*FaCML45*), two with 3 EF-hand motifs (*FaCML42*,
169 *FaCML36*) and one exhibiting 4 EF-hand motifs (*FaCML1a*).

170 **2.7. Accession Numbers**

171 The following sequences: *FaCML*, *FaCML1a*, *FaCML1b*, *FaCML2*, *FaCML3*, *FaCML7a*,
172 *FaCML7b*, *FaCML8*, *FaCML11*, *FaCML13*, *FaCML15*, *FaCML18*, *FaCML19*, *FaCML21*,
173 *FaCML23*, *FaCML24*, *FaCML25*, *FaCML27*, *FaCML29*, *FaCML36*, *FaCML38*, *FaCML41*,
174 *FaCML42*, *FaCML44*, *FaCML45*, *FaCML47*, *FaCML48* and *FaCML49* are available in the
175 Third Party Annotation Section of the GenBank database under the accession numbers:
176 MK965998-MK966025, respectively.

177 **3. Results**

178 **3.1. Calcium influx**

179 Evaluation of the intracellular calcium content in strawberry leaves treated with
180 brasinosteroids revealed a higher level of accumulation in the strawberry plants treated with

181 BB16 or EP24 as compared to the control water treated plants (Fig. 1). Nevertheless, the
182 influx of Ca^{2+} was greater when EP24 was used compared to BB16. The intracellular calcium
183 content in plants treated with EP24 or BB16 was 118% and 40% higher to plants treated with
184 water, respectively.

185 **3.2. Analysis of the amino acid sequences of the FaCMLs.**

186 Bioinformatic investigation let us to identify 28 CMLs sequences in *F. ananassa* genome.
187 Both programs used for the identification of structural motifs (e.g. CDSearch and SMART)
188 coincided in the identification of EF-hand motifs for calcium binding in all the FaCML
189 sequences analyzed. The only difference observed among them consisted in the number of EF-
190 hand motifs contained in each protein; FaCMLs containing 2, 3 or 4 EF-hand motifs were
191 found. In figure 2 we present the alignment of the EF-hand motif domains of three FaCML
192 sequences showing 2, 3, or 4 EF-hand motifs, selected at random just to show the structural
193 differences observed. The sequences obtained from *F. ananassa* were then compared with *F.*
194 *vesca* paralogs, and results showed an identity higher than 95%. The difference observed
195 consisted in few short gaps (up to 22 aa), but the number of FvCMLs and EF-hand motifs
196 contained were conserved (data not shown).

197 **3.3. Molecular phylogenetic analysis**

198 With the amino acid sequences of the CMLs predicted in *F. ananassa*, a molecular
199 phylogenetic analysis was performed. In the phylogenetic tree obtained we observed that the
200 FaCMLs proteins are grouped into three separate clades according to the number of EF-hand
201 motifs, all of which were supported by high bootstrap values (Fig. 3). Sequences that have two
202 EF-hand motifs (red frame) clustered in the phylogenetic clade I. Within this clade, FaCML47
203 sequence prediction resulted incomplete because the FvCML47 of *F. vesca* permitted only the
204 identification of very small contigs in *F. ananassa* (Fig. 3). Nevertheless, we assumed that it
205 would be a protein with two EF-hand motifs because the *F. vesca* orthologous has two EF-
206 hand motifs. In the case of FaCML21, although it presents four EF-hands it was included in
207 the clade I. The later may be possible due to the statistic weight attributed by the NJ method to
208 the rest of the sequences out of the EF-hand domains.

209 Sequences that contained mainly three EF-hand motifs (blue frame) were grouped in an
210 heterogeneous phylogenetic clade II. In this clade, although CMLs with only three EF-hands
211 were clearly detected (blue solid-line frame), there were some CMLs that may also present a

212 possible fourth EF-hand motif (blue dotted-line frame), but with a high E-values suggesting
213 that the amino acid sequences differ from the consensus motif; hence we considered that these
214 proteins contained only 3 EF-hand motifs (Fig. 3). In Figure 3 we can also see that some
215 FaCMLs (*e.g.* FaCML8, FaCML11, FaCML15 and FaCML8) were included in this clade
216 although they exhibit four RF-hands (green frame). In these cases we also assumed that it is
217 due to the same reason mentioned above, *e.g.* the statistic weight attributed by the NJ method
218 to the rest of the sequences out of the EF-hand domains. The clade III contains only four EF-
219 hand motif CMLs (green frame, Fig. 3). In Table S2 we present the nucleotide (bp) and amino
220 acid (aa) sizes of the FaCMLs predicted according to number of EF-hand exhibited. The
221 position of the EF-hand motifs on the amino acid sequences are also indicated (Table S1).

222 **3.4. Regulatory cis-elements associated to stress response**

223 When analyzing the 5'UTR of all the sequences corresponding to the predicted *FaCMLs*, we
224 found some regulatory *cis*-elements directly associated to the response to biotic and abiotic
225 stresses (Table 2). In the supplementary figure (S2), the position of each regulatory element is
226 detailed with respect to the initiation codon of each of the *FaCMLs* genes. Since the 5'UTR
227 sequences suggest that they may be regulated by multiple factors, we decided to investigate
228 whether the BRs studied would also affect the transcription levels of *FaCMLs*. In Table S2 we
229 present the location of each regulatory *cis*-element mentioned in Table 2, found at the 5'UTR
230 *FaCMLs* sequences. Only 22 *FaCML* were included, excluding the remaining 6 genes
231 (*FaCML*, *FaCML21*, *FaCML23*, *FaCML24*, *FaCML29*, *FaCML49*), since the “contigs” of
232 *Fragaria ananassa* that included them were not large enough, to analyze the 5'UTR region
233 and find regulatory elements.

234 **3.5. Expression of FaCMLs in response to treatment with BRs**

235 To evaluate the effect of the BRs on the expression of *CMLs*, only four of them were selected
236 to continue the study because their sequences allowed us to obtain better primers, which in
237 turns increased the reliability of the assay. For the selection of *FaCMLs* we also took into
238 account genes encoding proteins that have different number of EF-hand motifs, namely:
239 FaCML1a, FaCML42, FaCML45, and FaCML36. FaCML1a corresponds to a protein with
240 four EF-hand motifs, FaCML42 with three EF-hand motifs, FaCML45 with two EF-hand
241 motifs, and FaCML36 with three EF-hand motifs plus an altered fourth one (Fig. 3). Results
242 obtained show that the level of expression of each *FaCML* depends on the particular BRs and

243 the time elapsed after the treatment (Fig. 4). Whereas both BRs induced the upregulation of
244 *FaCML42* between 2 and 6 hpt, BB16 induced the upregulation of *FaCML36* and *FaCML1a*
245 at 4 hpt, and EP24 caused the upregulation of *FaCML45* at 6 hpt (Fig. 4).

246 **3.6. Molecular analysis of defense marker genes in strawberry plants treated with BRs**

247 Taking into account the results obtained in this and previous papers [7,26], and with the aim to
248 assess the effect of BRs on the activation of a defense response, the expression of the genes
249 *PR1*, *MYB30*, *ERF1*, *ETR1*, *FaPRX27* and *FaGSL5* were analyzed at different times post-
250 treatment (hpt). Results showed that the expression level of the genes studied depends on the
251 particular BRs used and the time elapsed after the treatment (Fig. 5). In that figure we can also
252 see that strawberry plants treated with both BRs induce the expression of the *PR1* gene at 72
253 hpt, *ERF1* at 12 hpt, and *GSL5* at 24 hpt. In addition, BB16 induced an increase of the
254 expression level of the *MYB30* gene at 12 hpt, and only plants treated with EP24 exhibited the
255 upregulation of *PRX27* at 48 hpt, and *ERF1* at 72 hpt (Fig. 5).

256 **4. Discussion**

257 Previously, we have shown that strawberry plants treated with BB16 or EP24 activated a
258 cascade of biochemical and physiological events, such as oxidative burst, production of nitric
259 oxide (NO), stomatal closure, cell wall reinforcement, and accumulation of calcium oxalate
260 crystals [7]. However, calcium involvement and genes associated to a defense response were
261 not evaluated.

262 It is well known that the change of intracellular calcium concentration is a key signal for many
263 cellular processes, including the activation of plant defenses against biotic or abiotic stresses,
264 and other processes, such as the cell wall strengthening reported by Figueroa et al. [27] which
265 prevents the cell wall degradation of strawberry fruits.

266 Changes of Ca^{2+} concentration are perceived by different proteins being calmodulins (CaM)
267 one of the most well characterized ones. In plants there is also a family of proteins called
268 calmodulins-like (CMLs), but their physiological functions and regulation are not completely
269 established. It was reported that the protein AtCML24 of Arabidopsis plays an important role
270 in the response to photoperiod, ion-homeostasis, and ABA-mediated inhibition of seed
271 germination and growth [28]. It was also shown that the Arabidopsis genes *AtCML37*,
272 *AtCML38* and *AtCML39* were upregulated in response to various biotic and abiotic stimuli
273 [29], and that the constitutive expression of the soybean *SCaM4*, and *SCaM5* genes in tobacco,

274 caused the increase of the resistance against a broad spectrum of virulent and non-virulent
275 pathogens [30]. This information stimulated us to investigate whether plants treated with EP24
276 or BB16 provoked a change of the intracellular concentration of Ca^{2+} .
277 Results obtained in this paper showed that both BRs induce the increase of the intracellular
278 concentration of Ca^{2+} after the treatment. Thus, we were interested to investigate whether the
279 transcriptional level of the *FaCMLs* would also be affected. However, since *CMLs* genes were
280 not reported in *F. ananassa* a bioinformatic investigation was performed to identify the
281 orthologous genes using as reference sequences of *F. vesca*. Outcomes of this analysis let us to
282 identify 28 genes in *F. ananassa*, and with the amino acid sequence to predict the
283 corresponding *FaCMLs*. Further structural analysis showed that all *FaCMLs* predicted
284 presented the characteristic Ca^{2+} binding EF-hand amino acid motif, varying only in the
285 number of EF-hand motifs present in each protein. Each motif consists in a 12-residue loop
286 that undergoes a conformational change when binding Ca^{2+} leading to the activation (or
287 inactivation) of target proteins [31].
288 With the *FaCMLs* amino acid sequences we performed a phylogenetic analysis and results
289 showed that they can be grouped in three clear clusters according to their EF-hand motifs.
290 However, few of them, namely: *FaCML8*, *FaCML11*, *FaCML15*, *FaCML18*, and *FaCML21*
291 did not fit into the groups. One possible hypothesis, already mentioned above, is that the NJ
292 method used attributed more statistic weight to the rest of the sequences out of the EF-hands.
293 Another plausible hypothesis would be that some of these *CMLs* may suffer post-translational
294 modifications (i.e. formation of disulfide bonds) that may affect the formation of an EF-hand,
295 as reported by Khan et al. [32]. This analysis further revealed that although the EF-hand motif
296 is an important structural feature conserved in the molecular evolution of these proteins, the
297 rest of the amino acid sequence is also important and may be implied in the interaction with
298 other molecules or cellular structures participating in its regulatory feature. Some evidence
299 supporting the latter hypotheses can be found by analyzing the information presented in Table
300 S2. In that table we can observe that the positions of the EF-hands motifs on the *FaCMLs*
301 amino acid sequences are not regular, suggesting that although the EF-hands motifs can be
302 clearly identified, the rest of the protein is not comparable in many cases.
303 The identification of the *FaCMLs* sequences on the *F. ananassa* genome also allowed us to
304 analyze the 5'UTR sequences that may correspond to putative regulatory *cis*-elements. In
305 plants, there are more than 1500 transcriptional factors (TFs) reported, controlling the

306 expression of genes in complex signaling networks [33]. From the *CMLs* nucleotide sequences
307 predicted in *Fragaria ananassa*, we could identify regulatory elements that participate in the
308 defense response against biotic and abiotic stresses (Table 2). Since these regulatory elements
309 suggested that FaCMLs are regulated by plant defense signals, the effect of BRs on the
310 expression levels of selected *FaCMLs* and defense associated genes were investigated. Results
311 confirmed that EP24 and BB16 are able to regulate the expression of *FaCML1a*, *FaCML36*,
312 *FaCML42* and *FaCML45* in a complex manner depending of the particular BR used and the
313 time elapsed after the treatment (Fig. 4).

314 It has been reported that the CMLs have numerous functions in the perception of stress and in
315 the development of plants [34]. Vanderbeld and Snedden [29] showed that the Arabidopsis
316 genes *AtCML37*, *AtCML38* and *AtCML39* are regulated by biotic and abiotic stresses, as well
317 as by hormonal and chemical treatments. They observed a strong and early upregulation
318 (usually within the first 24 hours) of these *CMLs* in response to wounds, osmotic stress,
319 drought and infections with bacterial pathogens. Similarly, Braam [35] analyzed the
320 expression of three *TCH* genes, encoding for CMLs, and observed a rapid upregulation in
321 response to stimuli such as touch, wind, rain, wounds and darkness. Likewise, Tomas-Grau et
322 al. [36] also reported that a soft mechanical stimulation (SMS) induced in strawberry plants
323 the upregulation of *TCH* genes encoding for CML proteins, demonstrating the importance of
324 these CMLs in signal transduction.

325 By analyzing the information presented in Table 2 and Table S2, we would notice that from
326 the nine regulatory elements detected at the *FaCMLs* 5'UTR sequence, three associated to
327 biotic (pathogen) responses (e.g. WRKY71OS, BIHDIOS, WBOXATNPR1) and three
328 associated to abiotic response (e.g. EBOXBNNAPA, MYBIAT, GTICONSENSUS) were
329 present in most of the FaCMLs analyzed, whereas the other regulatory *cis*-elements (e.g.
330 BOXLCOREDPCAL, ASFIMOTIFCAMV, ABRERATCAL) were found in lower frequency.
331 (Table S2). However, all FaCMLs analyzed exhibited regulatory *cis*-elements of both types.

332 This information let us conclude that the expression of FaCMLs is tightly controlled by many
333 regulatory *cis*-elements and there is no apparent correlation between the number of EF-hand
334 motifs and the regulatory elements found.

335 When evaluating the expression of some defense associated genes in plants treated with EP24
336 or BB16 different expression patterns were observed (Fig. 5).The results show that strawberry

337 plants treated with any of the BRs tested induce the expression of the *FaPR1* gene at 72 hpt
338 (Fig. 5a), suggesting the participation of SA signaling pathway [37,38,39,1].
339 The upregulation of *FaMYB30* at 12 hpt (Fig. 5b) in response to treatment with both BRs,
340 suggests its participation during the hypersensitivity response (HR) [40,41]. The relationship
341 between ROS and HR was established many years ago, when Doke [42] reported the
342 production of superoxide previous to the HR caused by *Phytophthora infestans* and the
343 tobacco mosaic virus, in potatoes and tobacco, respectively.
344 Polymerization of lignin is regulated by class III peroxidases, which are induced by different
345 stimuli, and biotic stress [43,44]. Specifically, the peroxidase PRX27 is a functional enzyme
346 that is required for the polymerization of phenylpropanoids during the formation of lignin in
347 the maturation of strawberry fruits [45]. In this study, however, we observed that the
348 *FaPRX27* gene was induced at 48 hpt, in strawberry leaves after EP24 treatment (Fig. 5e).
349 These data suggest that this peroxidase participates in the defense response induced by EP24
350 catalyzing the cross-linking of monolignoles in the cytosol, contributing to the final formation
351 and accumulation of lignin in leaves. However, we did not observe this effect in plants treated
352 with BB16. Hence, we may speculate that the induction of this gene occurs at times that were
353 not studied, or that the observed production of lignin in response to BB16 occurs through the
354 action of other peroxidases, or laccases, which were not analyzed in this study [46]. When
355 studying the level of expression of the *FaGSL5* gene, we observed that the callose synthase is
356 upregulated in response to both BRs (Fig. 5f), confirming that the deposition of callose is part
357 of the defense response induced by these compounds.
358 The role of ET as a signal molecule involved in the activation of defense mechanisms has been
359 confirmed by numerous studies [47]. When studying genes related to the ET signaling
360 pathway, a clear upregulation of *FaERF1* [48] was observed 12 hpt in response to both BRs,
361 and 72 hpt in response to EP24 (Fig.5c). It is well known that the ERF1 factor initiates a
362 transcriptional cascade that results in the activation or repression of many genes involved in
363 the regulated the ET signaling pathway [49]. Additionally, ERF1 is also a key integrator of the
364 signaling pathways of JA involved in the regulation of defense response genes such as *b-CHI*
365 and *PDF1.2* [50]. The upregulation of *FaERF1* may indicate, on one side, a strong activation
366 of genes involved in the synthesis enzymes ACS and ACO, and on the other side, the possible
367 activation of the JA signaling pathway in plants treated with BRs. The downregulation of

368 *ETR2* observed after 48 hpt (Fig. 5d) may be explained by the fact that the ET receptor
369 proteins are a negative regulators of the pathway [51,52].

370 The upregulation of genes related to ET signaling pathway, such as *ERF1*, and genes related to
371 the SA pathway, such as *PR1*, suggest that the BRs may activate both plant defense signaling
372 response but in a time dependent manner confirming that there is a complex cross-talk among
373 hormones modulating plant immunity [53].

374 This work unveils the importance of Ca^{2+} in the defense response induced by EP24 and BB16
375 in strawberry plants opening new ways to investigate the mechanisms by mean of which BRs
376 activates the plant innate immunity.

377 **Notes**

378 **Authors' contributions**

379 RNF, SMP and GGM are CONICET fellowship, and MGMZ and JCDR are members of
380 CONICET. RNF performed the experiments, analyzed and interpreted the data. MGMZ
381 collaborated in the bioinformatics analysis, SMP participated in real time assays and GGM
382 collaborated with calcium influx assay. RNF wrote the manuscript, and JCDR and MGMZ
383 critically reviewed the article. All authors approved the final version of the manuscript.

384 **Acknowledgements**

385 This paper was partially supported with grants of the Universidad Nacional de Tucumán
386 (PIUNT 26/D642), Agencia Nacional de Promoción Científica y Tecnológica (PICT 2017-
387 0653), CONICET (PUE-2016-0104) and INTA (PNHFA 1106073 and TUSGO 1231101).
388 Authors are grateful to Strawberry Active Germplasm Bank (BGA) from Universidad
389 Nacional de Tucumán (UNT) and Ing. Cecilia Lemme for providing strawberry plants. RNF,
390 SMP and GGM are CONICET fellowship, and MGMZ and JCDR are members of CONICET.

391 **Compliance with ethical standards**

392 **Ethical approval:** This article does not contain any studies with human participants or
393 animals performed by any of the authors.

394 **Conflict of interest:** The manuscript has not been published and is not under consideration for
395 publication elsewhere. All authors have approved the manuscript and agreed to submit the
396 paper to Scientia Horticulturae. The research was conducted in the absence of any commercial
397 relationships that could be considered as a potential conflict of interest.

398 **References**

- 399 [1] J.D.G. Jones, J.L. Dangl, The plant immune system, *Nature* 444 (2006) 323-329.
- 400 [2] A.M. Hetherington, C. Brownlee, The generation of Ca²⁺ signals in plants, *Annual Review of Plant*
401 *Biology* 55 (2004) 401-427.
- 402 [3] E. McCormack, J. Braam, Calmodulins and related potential calcium sensors of Arabidopsis, *New*
403 *Phytologist* 159(3) (2003) 585-598.
- 404 [4] E. McCormack, Y.C Tsai, J. Braam, Handling calcium signaling: Arabidopsis CaMs and CML,
405 *Trends in plant science* 10(8) (2005) 383-389.
- 406 [5] G. Xu, Y. Cui, M. Li, M. Wang, Y. Yu, B. Zhang, L. Huang, X. Xia, *OsMSR2*, a novel rice
407 calmodulin-like gene, confers enhanced salt tolerance in rice (*Oryza sativa* L.), *Australian Journal of*
408 *Crop Science* 7(3) (2013) 368.
- 409 [6] D. Chiasson, S.K. Ekengren, G.B. Martin, S.L. Dobney, W.A. Snedden, Calmodulin-like proteins
410 from Arabidopsis and tomato are involved in host defense against *Pseudomonas syringae*pv. Tomato,
411 *Plant molecular biology* 58(6) (2005) 887-897.
- 412 [7] R.N. Furio, P.L. Albornoz, Y. Coll, G.M. Zamora, S.M. Salazar, G.G. Martos, J.C. Díaz Ricci,
413 Effect of natural and synthetic Brassinosteroids on strawberry immune response against
414 *Colletotrichum acutatum*, *European Journal of Plant Pathology* 153(1) (2019a) 227-241.
- 415 [8] S. Gilroy, N. Read, A.J. Trewavas, Elevation of cytoplasmic calcium by caged calcium or caged
416 inositol trisphosphate initiates stomatal closure, *Nature* 346(6286) (1990) 769.
- 417 [9] J.M. Ward, J.I. Schroeder, Calcium-activated K⁺ channels and calcium-induced calcium release by
418 slow vacuolar ion channels in guard cell vacuoles implicated in the control of stomatal closure, *The*
419 *Plant Cell* 6(5) (1994) 669-683.
- 420 [10] G.G. Martos, A. Mamaní, M.P. Filippone, A.P. Castagnaro, J.C.D. Ricci, The ellagitanninHeT
421 induces electrolyte leakage, calcium influx and the accumulation of nitric oxide and hydrogen peroxide
422 in strawberry, *Plant physiology and biochemistry* 123 (2018) 400-405.
- 423 [11] V.V. Solovyev, Statistical approaches in Eukaryotic gene prediction, In *Handbook of Statistical*
424 *genetics* (eds. Balding D., Cannings C., Bishop M.), Wiley-Interscience; 3d edition, 2007, pp. 1616.

- 425 [12] C. Woffelman, DNAMAN for Windows. LydonBiosoft, Institute of Molecular Plant Sciences,
426 Leiden University, 2004.
- 427 [13] A. Marchler-Bauer, S.H. Bryant, CD-Search: protein domain annotations on the fly. Nucleic acids
428 research 32(suppl_2) (2004) 327-331.
- 429 [14] I. Letunic, R.R. Copley, B. Pils, S. Pinkert, J. Schultz, P. Bork, SMART 5: domains in the context
430 of genomes and networks, Nucleic acids research34(suppl_1) (2006) 257-260.
- 431 [15] J.D. Thompson, T.J. Gibson, F. Plewniak, F. Jeanmougin, D.G. Higgins, The CLUSTAL_X
432 windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools,
433 Nucleic acids research 25(24) (1997) 4876-4882.
- 434 [16] K. Tamura, J. Dudley, M. Nei, S. Kumar, MEGA4: molecular evolutionary genetics analysis
435 (MEGA) software version 4.0, Molecular biology and evolution 24(8) (2007) 1596-1599.
- 436 [17] N. Saitou, M. Nei, The neighbor-joining method: a new method for reconstructing phylogenetic
437 trees, Molecular biology and evolution 4(4) (1987) 406-425.
- 438 [18] J. Felsenstein, Confidence limits on phylogenesis: an approach using the bootstrap, Evolution 39
439 (1985) 783-791.
- 440 [19] K. Higo, Y. Ugawa, M. Iwamoto, T. Korenaga, Plant cis-acting regulatory DNA elements
441 (PLACE) database: 1999, Nucleic acids research 27(1) (1999) 297-300.
- 442 [20] I.A. Shahmuradov, A.J. Gammerman, J.M. Hancock, P.M. Bramley, V.V. Solovyev, PlantProm: a
443 database of plant promoter sequences, Nucleic Acids Research 31(1) (2003) 114-117.
- 444 [21] M. Guidarelli, F. Carbone, F. Mourgues, G. Perrotta, C. Rosati, P. Bertolinia, E. Baraldia,
445 *Colletotrichum acutatum* interactions with unripe and ripe strawberry fruits and differential responses
446 at histological and transcriptional levels, Plant Pathology 60 (2011) 685-697.
- 447 [22] F. Amil-Ruiz, J. Garrido-Gala, R. Blanco-Portales, K.M. Folta, J. Muñoz-Blanco, J.L. Caballero,
448 Identification and validation of reference genes for transcript normalization in strawberry (*Fragaria x*
449 *ananassa*) defense responses, PloS one 8(8) (2013) e70603.
- 450 [23] C. Ramakers, J.M. Ruijter, R.H.L. Deprez, A.F. Moorman, Assumption-free analysis of
451 quantitative real-time polymerase chain reaction (PCR) data, Neuroscience letters 339(1) (2003) 62-66.

- 452 [24] J.A. Di Rienzo, fgStatistics. Statistical software for the analysis of experiments of functional
453 genomics, 2011.
- 454 [25] M.W. Pfaffl, A new mathematical model for relative quantification in real time RT-PCR, *Nucleic
455 Acids Res* 29 (2001) 2002-2007.
- 456 [26] R.N. Furio, S.M. Salazar, G.M. Martínez-Zamora, Y. Coll, V. Hael-Conrad, J.C. Díaz-Ricci,
457 Brassinosteroids promote growth, fruit quality and protection against Botrytis on *Fragaria x*
458 *ananassa*, *European Journal of Plant Pathology* (2019b) 1-10.
- 459 [27] C.R. Figueroa, M.C. Opazo, P. Vera, O. Arriagada, M. Díaz, M.A. Moya-León, Effect of
460 postharvest treatment of calcium and auxin on cell wall composition and expression of cell wall-
461 modifying genes in the Chilean strawberry (*Fragaria chilensis*) fruit, *Food Chemistry* 132(4) (2012)
462 2014-2022.
- 463 [28] N.A. Delk, K.A. Johnson, N.I. Chowdhury, J. Braam, *CML24*, regulated in expression by diverse
464 stimuli, encodes a potential Ca^{2+} sensor that functions in responses to abscisic acid, day length, and ion
465 stress, *Plant physiology* 139(1) (2005) 240-253.
- 466 [29] B. Vanderbeld, W.A. Snedden, Developmental and stimulus-induced expression patterns of
467 Arabidopsis calmodulin-like genes *CML37*, *CML38* and *CML39*, *Plant molecular biology* 64(6) (2007)
468 683-697.
- 469 [30] W. Do Heo, S.H. Lee, M.C. Kim, J.C. Kim, W.S. Chung, H.J. Chun, K.J. Lee, C.Y. Park, H.C.
470 Park, J.Y. Choi, M.J. Cho, Involvement of specific calmodulin isoforms in salicylic acid-independent
471 activation of plant disease resistance responses, *Proceedings of the National Academy of Sciences*
472 96(2) (1999) 766-771.
- 473 [31] J.L. Gifford, M.P. Walsh, H.J. Vogel, Structures and metal-ion-binding properties of the Ca^{2+} -
474 binding helix-loop-helix EF-hand motifs, *Biochemical Journal* 405(2) (2007) 199-221.
- 475 [32] A.R. Khan, K.A. Johnson, J. Braam, M.N.G. James, Comparative modeling of the three-
476 dimensional structure of the calmodulin-related TCH2 protein from Arabidopsis, *Proteins: Structure,
477 Function, and Bioinformatics* 27(1) (1997) 144-153.
- 478 [33] H.D. Priest, S.A. Filichkin, T.C. Mockler, Cis-regulatory elements in plant cell signaling, *Current
479 opinion in plant biology* 12(5) (2009) 643-649.


- 480 [34] K.W. Bender, W.A. Snedden, Calmodulin-related proteins step out from the shadow of their
481 namesake, *Plant physiology* 163(2) (2013) 486-495.
- 482 [35] J. Braam, Regulation of expression of calmodulin and calmodulin-related genes by environmental
483 stimuli in plants, *Cell calcium* 13(6-7) (1992) 457-463.
- 484 [36] R.H. Tomas-Grau, F.J. Requena-Serra, V. Hael-Conrad, M.G. Martínez-Zamora, M.F. Guerrero-
485 Molina, J.C. Díaz-Ricci, Soft mechanical stimulation induces a defense response against *Botrytis*
486 *cinerea* in strawberry, *Plant cell reports* 37(2) (2018) 239-250.
- 487 [37] V. Hael-Conrad, S.M. Perato, M.E. Arias, M.G. Martínez-Zamora, P.D.L.Á. Di Peto, G.G.
488 Martos, A.P. Castagnaro, J.C. Díaz-Ricci, N.R. Chalfoun, The Elicitor Protein AsES Induces a
489 Systemic Acquired Resistance Response Accompanied by Systemic Microbursts and Micro-
490 Hypersensitive Responses in *Fragariaananassa*, *Molecular Plant-Microbe Interactions* 31(1) (2017)
491 46-60.
- 492 [38] L.C. Van Loon, E.A. Van Strien, The families of pathogenesis related proteins, their activities,
493 and comparative analysis of PR-1 type proteins, *Physiological and Molecular Plant Pathology* 55
494 (1999) 85-97.
- 495 [39] M.Y. Graham, J. Weidner, K. Wheeler, M.J. Pelow, T.L. Graham, Induced expression of
496 pathogenesis related protein genes in soybean by wounding and the *Phytophthora sojae* cell wall
497 glucan elicitor, *Physiological and Molecular Plant Pathology* 63 (2003) 141-149.
- 498 [40] X. Daniel, C. Lacomme, J.B. Morel, D. Roby, A novel myb oncogene homologue in *Arabidopsis*
499 *thaliana* related to hypersensitive cell death, *The Plant Journal* 20 (1999) 57-66.
- 500 [41] F. Vailliau, X. Daniel, M. Tronchet, J.L. Montillet, C. Triantaphylidés, D. Roby, A R2R3-MYB
501 gene, AtMYB30, acts as a positive regulator of the hypersensitive cell death program in plants in
502 response to pathogen attack, *Proceedings of the National Academy of Sciences of the United States of*
503 *America* 99 (2002) 10179-10184.
- 504 [42] N. Doke, Involvement of superoxide anion generation in the hypersensitive response of potato
505 tuber tissues to infection with an incompatible race of *Phytophthora infestans* and to the hyphal wall
506 components, *Physiological Plant Pathology* 23(3) (1983) 345-357.
- 507 [43] R. Zhong, Z.H. Ye, Transcriptional regulation of lignin biosynthesis, *Plant Signaling & Behavior*
508 4 (2009) 1028-1034.

- 509 [44] C. Mathé, A. Barre, C. Jourda, C. Dunand, Evolution and expression of class III peroxidases,
510 Archives of Biochemistry and Biophysics 500 (2010) 58-65.
- 511 [45] L. Ring, S.Y. Yeh, S. Hücherig, T. Hoffmann, R. Blanco-Portales, M. Fouche, C. Villatoro, B.
512 Denoyes, A. Monfort, J.L. Caballero, J. Muñoz-Blanco, J. Gershenson, W. Schwab, Metabolic
513 interaction between anthocyanin and lignin biosynthesis is associated with peroxidase FaPRX27 in
514 strawberry fruit, Plant Physiology 163 (2013) 43-60.
- 515 [46] A. Kärkönen, S. Koutaniemi, Lignin biosynthesis studies in plant tissue cultures, Journal of
516 integrative plant biology 52(2) (2010) 176-185.
- 517 [47] F.B. Abeles, P.W. Morgan, M.E. Saltveit, 1992. Ethylene in Plant Biology Academic press.
- 518 [48] Y. Zhang, X. Yin, Y. Xiao, Z. Zhang, S. Li, X. Liu, B. Zhang, X. Yang, D. Grierson, G. Jiang,
519 H.J. Klee, K. Chen, An ETHYLENE RESPONSE FACTOR-MYB transcription complex regulates
520 furaneol biosynthesis by activating QUINONE OXIDOREDUCTASE expression in strawberry, Plant
521 physiology 178(1) (2018) 189-201.
- 522 [49] H. Sakai, J. Hua, Q.G. Chen, C. Chang, L.J. Medrano, A.B. Bleecker, E.M. Meyerowitz, ETR2 is
523 an ETR1-like gene involved in ethylene signaling in Arabidopsis, Proceedings of the National
524 Academy of Sciences of the United States of America 95 (1998) 5812-5817.
- 525 [50] O. Lorenzo, R. Piqueras, J.J. Sánchez-Serrano, R. Solano, ETHYLENE RESPONSE FACTOR1
526 integrates signals from ethylene and jasmonate pathways in plant defense, Plant Cell 15 (2003) 165-78.
- 527 [51] N.E. Gapper, R.P. McQuinn, J.J. Giovannoni, Molecular and genetic regulation of fruit ripening,
528 Plant Molecular Biology 82 (2013) 575-591.
- 529 [52] F. Wang, X. Cui, Y. Sun, C.H. Dong, Ethylene signaling and regulation in plant growth and stress
530 responses, Plant Cell Reports 32(7) (2013) 1099-1109.
- 531 [53] A.M. Shigenaga, C.T. Argueso, No hormone to rule them all: interactions of plant hormones
532 during the responses of plants to pathogens, Seminars in Cell & Developmental Biology 56 (2016)
533 174-189.
- 534 [54] X. Liu, X. Bai, X. Wang, C. Chu, OsWRKY71, a rice transcription factor, is involved in rice
535 defense response. Journal of plant physiology 164(8) (2007) 969-979.

- 536 [55] H. Luo, F. Song, R.M. Goodman, Z. Zheng, Z. Up-regulation of *OsBIHD1*, a rice gene encoding
537 BELL homeodomain transcriptional factor, in disease resistance responses, *Plant Biology* 7(05) (2005)
538 459-468.
- 539 [56] T. Eulgem, P.J. Rushton, E. Schmelzer, K. Hahlbrock, I.E. Somssich, Early nuclear events in plant
540 defencesignalling: rapid gene activation by WRKY transcription factors, *The EMBO journal* 18(17)
541 (1999) 4689-4699.
- 542 [57] J. Dong, C. Chen, Z. Chen, Expression profiles of the Arabidopsis WRKY gene superfamily
543 during plant defense response, *Plant molecular biology* 51(1) (2003) 21-37.
- 544 [58] K. Maeda, S. Kimura, T. Demura, J. Takeda, Y. Ozeki, DcMYB1 acts as a transcriptional
545 activator of the carrot phenylalanine ammonia-lyase gene (DcPAL1) in response to elicitor treatment,
546 UV-B irradiation and the dilution effect. *Plant molecular biology* 59(5) (2005) 739-752.
- 547 [59] K. Stålberg, M. Ellerstöm, I. Ezcurra, S. Ablov, L. Rask, Disruption of an overlapping E-
548 box/ABRE motif abolished high transcription of the *napA* storage-protein promoter in transgenic
549 *Brassica napus* seeds, *Planta* 199(4) (1996) 515-519.
- 550 [60] H. Abe, K. Yamaguchi-Shinozaki, T. Urao, T. Iwasaki, D. Hosokawa, K. Shinozaki, Role of
551 Arabidopsis MYC and MYB homologs in drought-and abscisic acid-regulated gene expression, *The*
552 *Plant Cell* 9(10) (1997) 1859-1868.
- 553 [61] J. Redman, J. Whitcraft, C. Johnson, J. Arias, Abiotic and biotic stress differentially stimulate as-1
554 element activity in Arabidopsis, *Plant Cell Reports* 21 (2002) 180-185.
- 555 [62] A.S. Buchel, F.T. Brederode, J.F. Bol, H.J. Linthorst, Mutation of GT-1 binding sites in the *Pr-IA*
556 promoter influences the level of inducible gene expression *in vivo*, *Plant molecular biology* 40(3)
557 (1999) 387-396.
- 558 [63] D. Bonetta, P. McCourt, Genetic analysis of ABA signal transduction pathways, *Trends in plant*
559 *science* 3(6) (1998) 231-235.

Table 1: Primers sequences used in real-time expression assays.


Genes	Names	Primers sequences	Accession no.
<i>FaEF-1a</i>	Elongation factor Tu-1	Fw 5'-GAACTTCCCAGGCTGATT-3' Rv 5'-CTGACCATCCTTCGAGATAC-3'	BK009992
<i>FaPR1</i>	Pathogenesis-related protein 1	Fw 5'-CAAAGAGCTCCGGCGACTT-3' Rv 5'-TCTCCCACCCACAGGTTTAC-3'	AB462752
<i>FaMYB30</i>	HR-marker gene	Fw 5'-TGCCGGCTCAGATGGACTA-3' Rv 5'-TCGGTGAAGTTACCGGTTTT-3'	BK009993
<i>FaERF1</i>	Ethylene response factor 1	Fw 5'-CTCGATGCTGTGGCTAAA-3' Rv 5'-GTGACCTCCTCTTCCTTTAC-3'	MH332903
<i>FaETR2</i>	Ethylene response 2	Fw 5'-GGAGGGTTGGTCGTATTT-3' Rv 5'-ACTATCAGATGAGTTCTGTCTC-3'	KC852067
<i>FaPRX27</i>	Peroxidase27	Fw 5'-CACCAGCTTGCCACAGACAA-3' Rv 5'-CTGGCTTTTGGCCATCTTCT-3'	AFQ36036
<i>FaGSL5</i>	Glucansynthase-like 5	Fw 5'-GGCACTTAAAATGCGGAATCTG-3' Rv 5'-GATCGTAGGCTTCCGGATACC-3'	BK009994
<i>FaCML45</i>	Calmodulin-like 45	Fw 5'-GGAGCAGAGGAGGTTTCAAGAG-3' Rv 5'-TTCCTTCACCTCCTCCAAGCT-3'	MK966022
<i>FaCML42</i>	Calmodulin-like 42	Fw 5'-GGGTTTCATATCGGCAAAGGA-3' Rv 5'-CCTTCCGGGAACCCTAGCT-3'	MK966020
<i>FaCML36</i>	Calmodulin-like 36	Fw 5'-CGGATCACGACGGGAAGAT-3' Rv 5'-CCCCTATCGCCGTGAACAC-3'	MK966017
<i>FaCML1a</i>	Calmodulin-like 1a	Fw 5'-TTGGAAAGGGAAGCATGGTT-3' Rv 5'-CGTCCAAATCAACAAGCTTGAA-3'	MK965999



561 **Fig. 1.** Intracellular calcium content in leaf discs pre-incubated with Fura2-AM and subsequently
 562 treated with: water (control), BB16 (0.1 mg l⁻¹) and EP24 (0.1 mg l⁻¹) All measurements were taken 30
 563 seconds after the treatment. The calcium concentration is expressed as the ratio of the band intensities
 564 362 nm/402 nm. The bars represent the standard deviation (mean values ± SE). The comparison
 565 between two groups was performed using the Tukey test ($p < 0.05$). Different letters denote significant
 566 difference.

FaCML42	1	----MEAAAAAPSSAGTSSERTLSKRPSASSSFRLRCPSLNSLRLLRIFDIFDKNRDGMV
FaCML23	1	-MSCFRICLTALRRCTFGGCKSFRKKRRRSSNLDAPATAASSFAAMEVSNQFKQ----VF
FaCML45	1	MEDILAAAQPGSLYVLGLFQPLLKLWTKTHNYLNYSDPHVLQPTLDQPLQCNK-----
FaCML42	57	SVDEINQALRLLGLDVEVSELEATIRTFIQPRNEGLTYDDFVGLHQSLYDTFFFDGDDIE
FaCML23	56	EVM DANGDGKIS -- PVE LSEVLCCLG----YKNKSAAAKEAEGMVR-----EM DCNG
FaCML45	55	-----GQDEPLLG-KEEVKMVMGRLG-----LCYQD-----EG
FaCML42	117	NAVEEAAALVVAARDEEEAKLLQEESDLREAFKVF DEDGDGFISAKE LQTVLGLKGFPE
FaCML23	102	DGFIDLDEF MTAVSANDTKRSGNDGGDDLMDAFLIF DTDKNKGKISAKE LRRVLSLGCER
FaCML45	82	DDVITNEDMVGAEVSRVFSEAEPSLEEVKEAFDVF DENRDGFIGAAD LQVLCNLGLKG
FaCML42	177	GNEIDRVERMITSV DRNHDGLVDLFE FKDMMRSTVLPSSCS
FaCML23	162	SS-LKECKQMIKGV DRNGDGVVDFEE FRLMMTRNV-----
FaCML45	142	KFGLEECRGMKAV DMNQDGLVDFEE FVELMDN-----


568 **Fig. 2.** Sequences alignment of three FaCMLs obtained with the Clustal program and edited with
569 BOXSHADE. The consensus sequences corresponding to the FaCML42, FaCML23 and FaCML45
570 EF-hand motifs are presented. The number of repetition of EF-hand motif varies in the different
571 FaCMLs studied. FaCML42 presents three EF-hand motifs (blue boxes), FaCML23 presents four EF-
572 hand motifs (yellow boxes), and FaCML45 two EF-hand motifs (red boxes).




573 **Fig. 3.** NJ phylogenetic tree, created with MEGA 4, from the amino acid sequence of the FaCMLs
 574 proteins predicted in *Fragaria ananassa*. The numerals at the branches indicate bootstrap values.
 575 FaCMLs framed in red indicate CMLs containing two EF-Hand motifs, in blue (solid or dotted)
 576 indicate CMLs containing three EF-Hand motifs, and in green indicate CMLs containing four EF-Hand
 577 motifs.

578 **Table 2:** Regulatory *cis* elements associated biotic and abiotic stress responses found at the 5'UTR
 579 *CML* sequences predicted in *Fragaria ananassa*. The ambiguity code characters correspond to: W =
 580 A/T; Y = C/T; N = A/T/G/C; R = A/G; B = G/T/C; M = A/C.

Response element	cis-regulatory element	Sequence	Functions	References
Inductor/ pathogen response elements	WRKY71OS	TGAC	Defense response to infections or wounds. Induced by SA and JA	[54]
	BIHD1OS	TGTCA	Related to disease responses	[55]
	WBOXATNPR1	TTGAC	Binding site of WRKY transcription factors, involved in the expression of stress-induced genes	[56,57]
	BOXLCOREDPCAL	ACCWWCC	Recognition sites for DcMYB1 (transcriptional activator of the phenylalanine ammonia-lyase gene). Important role in SA-mediated defense responses.	[58]
Abiotic stress response elements	EBOXBNNAPA	CANNTG	Response to salt stress, water deficit. Induced by ABA	[59]
	MYBIAT	WAACCA	Response to abiotic stresses e.g. low temperatures and water stress	[60]
	ASF1MOTIFCAMV	TGACG	ASF-1 binding site (transcription factor of the bZIP family). Involved in activation of number of genes induced by SA and auxins	[61]
	GT1CONSENSUS	GRWAAW	Binding site of the transcription factor GT-1. Regulates the expression of <i>PR</i> genes inducible by SA	[62]
	ABRERATCAL	MACGYGB	Response to water deficit, mediated by ABA	[63]



581 **Fig. 4.** Quantitative analysis by qRT-PCR of genes that code for different *FaCMLs* in strawberry
 582 plants treated with BB16 or EP24. All values were normalized with respect to the expression of the
 583 internal reference gene *FaEF1 α* and are expressed in relation to the values of the control plants
 584 treated with water. Dotted line represents the normalized expression value of control (water treated) plants.
 585 Bars represent average values \pm standard error of three biological replicas. The asterisks indicate a
 586 statistically significant difference calculated by the fgStatistics software ($P \leq 0.05$).



587 **Fig. 5.** Quantitative analysis by qRT-PCR of genes related to the defense response in strawberry plants
588 treated with BRs at different times. All values were normalized with respect to the expression of the
589 internal reference gene *FaEF1 α* and are expressed in relation to the values of the control plants treated
590 with water. Dotted line represents the normalized expression value of control (water treated) plants.
591 Bars represent average values \pm standard error of three biological replicas. The asterisks indicate a
592 statistically significant difference calculated by the fgStatistics software ($P \leq 0.05$).

Table S1: Sizes and position EF-hand motifs on the amino acid sequences of the FaCMLs analyzed

EF-Hand motifs	FaCMLs	Gene length	Number of amino acids	EF-hand motif position in amino acid sequence			
				1°	2°	3°	4°
2 EF-Hand motifs	FaCML7a	477 bp	158 aa	37-65	109-137		
	FaCML7b	366 bp	122 aa	56-84	92-120		
	FaCML45	525 bp	174 aa	109-137	147-174		
	FaCML47	378 bp	126 aa	52-80	88-116		
	FaCML48	840 bp	279 aa	108-136	177-205		
	FaCML49	564 bp	187 aa	85-113	126-154		
3 EF-Hand motifs	FaCML	342 bp	113 aa	12-40	49-77	85-113	
	FaCML2	477 bp	158 aa	21-49	58-86	91-119	
	FaCML13	444 bp	147 aa	12-40	83-111	119-147	
	FaCML19	576 bp	191 aa	55-83	91-119	165-191	
	FaCML36	633 bp	210 aa	70-98	144-172	181-209	
	FaCML38	423 bp	140 aa	6-34	78-106	114-140	
	FaCML41	576 bp	191 aa	60-88	96-124	165-191	
	FaCML42	657 bp	218 aa	40-68	144-172	182-210	
	FaCML44	480 bp	159 aa	8-36	90-118	128-156	
4 EF-Hand motifs	FaCML1a	558 bp	185 aa	46-74	82-110	117-145	153-181
	FaCML1b	576 bp	191 aa	50-78	86-114	122-150	158-186
	FaCML3	660 bp	219 aa	65-93	101-129	147-175	185-213
	FaCML8	444 bp	147 aa	12-40	48-76	84-112	120-146
	FaCML11	453 bp	150 aa	12-40	48-76	85-113	121-149
	FaCML15	483 bp	160 aa	11-39	47-75	86-114	122-150
	FaCML18	513 bp	170 aa	20-48	56-84	99-127	135-163
	FaCML21	687 bp	228 aa	58-86	94-122	149-177	187-215
	FaCML23	588 bp	195 aa	50-78	89-117	129-157	166-194
	FaCML24	411 bp	136 aa	3-31	39-67	72-100	108-136
	FaCML25	486 bp	161 aa	14-42	50-78	87-115	123-151
	FaCML27	504 bp	167 aa	21-49	57-85	93-121	129-157
FaCML29	474 bp	157 aa	19-47	55-83	92-120	129-157	

Table S2: Location of regulatory *cis*-elements found on the 5'UTR of the FaCMLs analyzed

	Inductor/pathogen response elements				Abiotic stress response elements					
	1	2	3	4	5	6	7	8	9	
2 EF-Hand motifs	FaCML7a	607 (-) 329 (+)	235(+)		961 (+) 961 (-)	1016 (-)		60 (+, 533 (+)	980 (+)	
	FaCML7b	24 (-)	24 (-)			152 (+)			329 (-)	
	FaCML45	112 (+) 247 (-)		112 (+) 1314 (-)		421 (+) 421 (-),	421 (+) 440 (-)			
	FaCML47	33 (-) 303 (+)		32 (-), 130 (+)		173 (+) 173 (-)	201 (-)	16 (-) 159 (-)		
	FaCML48	51 (+) 191 (+)		592 (-)		389 (+) 389 (-)	349 (-)	593 (-)	207 (+) 366 (+)	89 (+), 766 (+)
3 EF-Hand motifs	FaCML2	254 (+) 335 (+)	199 (-) 364 (+)	335 (+) 363 (-)	298 (+)	341 (+) 341 (-)			397 (-)	212 (+)
	FaCML13	557 (+) 613 (-)	556 (-) 586 (-)			589 (+) 589 (-)	198 (+)		152 (+) 279 (-)	
	FaCML19	215 (-) 412 (-)	412 (+) 1379 (-)	411 (-) 1239 (-)	172 (+) 381 (+)	330 (+) 330 (-)	19 (+) 782 (-)	647 (+) 1277 (+)	1103 (-) 1498 (-)	1763 (+)
	FaCML36			687 (+) 882 (+)		937 (+) 937 (-)	436 (+) 580 (+)		487 (+) 583 (+)	
	FaCML38						6 (+)		27 (-)	
	FaCML41	267 (-) 623 (+)	622 (+)	415 (-)		42 (+) 42 (-)	295 (-)	225 (+)	648 (-) 659 (-)	
	FaCML42	328 (-) 555 (+)	554 (-) 1804 (-)	327 (-) 1805 (+)	2582 (+)	196 (+) 196 (-)	219 (-) 1872 (+)	2451 (-) 2674 (+)	304 (-) 336 (-)	117 (+), 557 (-),
	FaCML44	524 (+) 757 (-)	628 (-)	847 (+) 1722 (-)	1680 (-)	527 (+) 527 (-)	27 (+) 48 (+)	1560 (+)	156 (-) 694 (+)	
	FaCML1a	265 (-) 317 (-)	265 (+) 317 (+)	395 (+)			581 (+)		54 (-), 108 (-)	
FaCML1b	356 (-) 567 (-)	274 (-) 573 (+)			771 (+) 771 (-)	687 (+)		496 (-), 780 (-)		
4 EF-Hand motifs	FaCML3	18 (-)		17 (-)		170 (+) 170 (-)			10 (+), 253 (-),	
	FaCML8	104 (-) 291 (-)	137 (-) 312 (-)	103 (-) 341 (+)		174 (+) 174 (-)		104 (-)	36 (-), 76 (+)	
	FaCML11					341 (+) 341 (-)	121 (-)		34 (+)	
	FaCML15	25 (-) 80 (-)	25 (+) 278 (-)			285 (+) 285 (-)	162 (+) 258 (+)		266 (+) 402 (-)	
	FaCML18					11 (+) 11 (-)				
	FaCML25	474 (-)		640 (+)	682 (+)	352 (+) 352 (-)	92 (+) 318 (+)		548 (-), 742 (-)	
	FaCML27	278 (+) 411 (+)		411 (+) 818 (-)		642 (+) 642 (-)	18 (+)		163 (-), 219 (-)	112 (+), 759 (+)

597 Numerals correspond to the following regulatory *cis*-elements: (1), WRKY71O; (2), BIHD1OS; (3),
598 WBOXATNPR1; (4), BOXLCOREDCPAL; (5), EBOXBNNAPA; (6), MYB1AT; (7),
599 ASF1MOTIFCAMV; (8), GT1CONSENSUS; (9), ABRERATCAL. (+) and (-) indicate the location of
600 the first nucleotide of the *cis*-elements found on the positive or negative strands, respectively.