


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Inorganica Chimica Acta

journal homepage: www.elsevier.com/locate/ica

Note

Spectroscopic evidence on the interaction of prephenate, a shikimate pathway intermediate, with oxidovanadium(IV) species

Luis Lezama^a, Teófilo Rojo^a, Evelina G. Ferrer^{b,*}^aDepartamento de Química Inorgánica, Facultad de Ciencia y Tecnología, Universidad del País Vasco, Apdo 644, 48080 Bilbao, Spain^bCEQUINOR (Centro de Química Inorgánica), CONICET-UNLP-Facultad de Ciencias Exactas, Universidad Nacional de La Plata, 47 y 115 (1900) La Plata, Argentina

ARTICLE INFO

Article history:

Received 18 September 2009

Received in revised form 28 December 2009

Accepted 9 January 2010

Available online 15 January 2010

Keywords:

Prephenate

Oxidovanadium(IV)

Interactions

Electronic spectra

Electronic paramagnetic resonance

ABSTRACT

The interaction of the oxidovanadium(IV) cation with the sodium salt of prephenic acid was investigated by electron absorption and electronic paramagnetic resonance spectroscopies in aqueous solution at different pH values. The study allows to demonstrate once more the effectiveness of oxidovanadium(IV) to interact with carboxylate groups. The most probable binding modes of the solution complexes were determined by EPR method. Two main coordination environments around the metal center were established: one includes four-carboxylate moieties and the other the set probably involves alkoxide and hydroxide groups.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Commonly known by its anionic form prephenate, prephenic acid (1-(2-carboxy-2-oxido-ethyl)-4-hydroxy-cyclohexa-2,5-diene-1-carboxylic acid) is an intermediate in the biosynthesis of aromatic aminoacids in plants, fungi, and bacteria [1–3]. This reaction involves the intramolecular rearrangement of chorismate(I) to prephenate(II). Chorismate mutase catalyzes the transformation in the shikimate pathway (Scheme 1) that leads to the biosynthesis of phenylalanine and tyrosine [4].

Many hydroxyacids are important in plants metabolism and nutrition. They have been identified as exudated from living leaves and roots in agricultural weeds [5–10]. These phenolic compounds, which have several functionals available for binding of biometals, are widely recognized as antioxidants and they are involved in metal transport processes. There is a great interest on the study of the bioavailability of metals in plants, and on the inhibitory effects that the non-flavonoids compounds produce on deleterious oxidative processes (cancer and atherosclerosis) [11,12].

Vanadium is a biologically ultramicrotrace transition metal and its physiologically relevant oxidation states include mainly +4 (vanadyl cation VO²⁺) and +5 (vanadate species). These species can be classified as hard acids and can strongly interact with biochemical donor groups specially containing oxygen and nitrogen atoms. Vanadium is present in rocks (5–250 ppm) and soils

(20–500 ppm) thus being more accessible to plants and animals [13,14], and it is proved by the presence of vanadium in plants, fungi and bacteria [15]. The uptake by higher plants has been confirmed by EPR measurements and it has also been shown that plant enzymes and root growth are vanadium-dependent [15]. In fungi, vanadium is accumulated in *Amanita muscaria* and it was demonstrated that amavadin [15] involves carboxylate groups in the coordination sphere of vanadium(IV) complex. In addition, some vanadium complexes containing hydroxy-carboxylate ligands have been reported to have potential pharmacological activities [16].


Taking into account that no previous reports exist about this system, the aim of this work is to study the interaction between vanadium and prephenate anion as ligand using EPR and UV-Vis spectroscopies.

2. Experimental

The barium salt of prephenic acid (Sigma) and VOSO₄·5H₂O (Merck) were used as reagents. The experiments were performed under nitrogen atmosphere to avoid oxidation phenomena. The electronic absorption spectra were measured with a Hewlett-Packard 8452 diode-array spectrophotometer using 10-mm quartz cells. Base line corrections, normalization, curve-fitting and calculations were carried out by means of Grams/32 (Galactic Industries Corporation, USA) software. The EPR spectra were obtained using a Bruker ESP300 spectrometer operating at X-band with standard Oxford low temperature devices (ESR900/ITC4). Anisotropic X-band EPR spectra of frozen solutions were recorded at 140 K, after

* Corresponding author. Tel./fax: +54 0221 4259485.

E-mail address: evelina@quimica.unlp.edu.ar (E.G. Ferrer).



Scheme 1. Conversion of chorismate (I) to prephenate (II).

addition of 5% of DMSO to ensure good glass formation. All computer manipulations (simulation and difference) of the EPR spectra were performed using the program SimFonia (WINEPR SimFonia v1.25, Bruker Analytische Messtechnik GmbH, 1996).

3. Results and discussion

EPR spectra of the aqueous solution (1:1 and 2:1 L/M ratio) at 140 K were recorded at different pH values in order to assist in the identification of different vanadium specie and the corresponding geometries, particularly equatorial ligand donor set and the overall charges.

For the 2:1:prephenate/ VO^{2+} system at pH 2.5 (Fig. 1), the EPR signal shows the characteristics line-wide trends for free $[\text{VO}(\text{H}_2\text{O})_5]^{2+}$ ion denoting that no interaction occurs in high acidic medium. By adding NaOH 1 M aqueous solution, the increase of the pH value to 3.4 and 4.2 induces important variations on EPR features (Fig. 1) and a new resonance line appears together with the signal belonging to free vanadyl aquoion. In order to identify the specie, difference spectra were performed. The experimental and the simulated spectra at pH 4.2 are shown in Fig. 2. The simulation predicts that the new observed signal is originated by a V-chromophore with the spin Hamiltonian parameters of $g_z = 1.943$, $g_x = 1.977$, $g_y = 1.980$ and the hyperfine coupling constant of $A_z = 171 \times 10^{-4} \text{ cm}^{-1}$, $A_x = 60 \times 10^{-4} \text{ cm}^{-1}$ and $A_y = 65 \times 10^{-4} \text{ cm}^{-1}$ ($g_{\text{iso}} = 1.966$, $A_{\text{iso}} = 98.67 \times 10^{-4} \text{ cm}^{-1}$). Based on the calculated parameters, the experimental spectrum at pH 4.2 has been simulated and found fitting in good concordance with the experimental one (Fig. 3A).




Fig. 1. X-band anisotropic EPR spectra ($A_{||}$ region, Low Field) of the 2:1 prephenate-oxidovanadium(IV) system in frozen solutions (140 K) at different pH values. Total concentration VO^{2+} : 2.5 mM, prephenate: 5 mM.




Fig. 2. X-band anisotropic EPR spectra. Solid line represents difference spectrum for the 2:1 prephenate-oxidovanadium(IV) frozen solutions (140 K) at pH 4.2 minus the $\text{VO}(\text{H}_2\text{O})_5^{2+}$ contribution. Dash line, simulation. Total concentration VO^{2+} : 2.5 mM, prephenate: 5 mM.

Significant changes on the spectra are detected when the pH of the solution is modified. The highest possible value for pH is 5.5 because precipitation (probably $\text{VO}(\text{OH})_2$) takes place. In this condition (Fig. 3B) two distinct specie are possible to be observed once more: the specie detected at pH 4.2 and a new one. For the new signal, the calculated EPR parameters are $g_z = 1.951$, $g_x = 1.977$, $g_y = 1.982$ and the hyperfine coupling constant of $A_z = 156 \times 10^{-4} \text{ cm}^{-1}$, $A_x = 60 \times 10^{-4} \text{ cm}^{-1}$ and $A_y = 65 \times 10^{-4} \text{ cm}^{-1}$ ($g_{\text{iso}} = 1.970$, $A_{\text{iso}} = 93.7 \times 10^{-4} \text{ cm}^{-1}$) suggesting both the formation of a new complex with some modification of the equatorial coordinated groups. Calculated EPR spectrum at pH 5.5 is in good concordance with the experimental one (Fig. 3B). Upon increasing the pH of the solution, the relative intensity of the signals decreases (not shown), precipitation takes place and probably oligonuclear specie start to be predominant, leading to extinction of the signal [17].

For the 1:1 prephenate/ VO^{2+} system the same species have been detected. It is known that parallel component of the hyperfine coupling constant is sensitive to the type of donor atoms on the equatorial positions of the coordination sphere. The following empirical relationship is frequently used in order to determine the identity of the equatorial ligands in V(IV) complexes: $A_z = \sum n_i A_{z,i}$ (n_i : number of equatorial ligands of type I, $A_{z,i}$: contribution to the parallel hyperfine coupling from each of them) [18–20]. Considering the contributions to the parallel hyperfine coupling constant of the different coordination modes of the ligand ($\text{COO}^- = 42.72$ [17–19], $\text{ArO}^- = 38.6$ and $\text{OH}^- = 38.7 \times 10^{-4} \text{ cm}^{-1}$ [20], respectively) and using the additivity rule for the detected species at pH 4.2, the empirical parameter was calculated. Very close to the $A_z = 171 \text{ cm}^{-1}$ obtained from the simulated EPR spectrum, the obtained value of $A_z = 170.9 \text{ cm}^{-1}$ is consistent with the species which could be interpreted assuming an equatorial ligand interaction via four COO^- moieties with O atoms in the planar donor set as schematically shown in Fig. 4 [21]. Comparable values are predicted considering other complexes with similar donor sets, and the calculated parameters fit well in the corresponding $g_{||}$ versus $A_{||}$ diagram [22]. The reduction of A_z with increasing pH ($A_z \text{ calc} = 155 \times 10^{-4} \text{ cm}^{-1}$) is attributed to changes in the coordination sphere of the vanadyl moiety, probably including the dissociation and coordination of the alkoxide group of the ligand and the incorporation of OH^- to the coordination sphere [18,23,24]. At this pH value, species containing one OH^- around the oxidovanadium(IV) cation were usually identified [25]. The calculated A_z value matched well with two possibilities ($A_z (2 \text{ ArO}^- + 2 \text{ OH}^-) = 154.6$ and $A_z (1 \text{ ArO}^- + 3 \text{ OH}^-) = 154.7$) which were consistent with the hypothesis of the presence of the hydrolytic complex. Considering that undetectable precipitation began, the oxidovanadium(IV) concentration




Fig. 3. X-band anisotropic EPR spectra. (A) Solid line represents experimental spectrum for the 2:1 prephenate–oxidovanadium(IV) frozen solutions (140 K) at pH 4.2. Dash line, simulation. (B) Solid line represents experimental spectrum for the 2:1 prephenate–oxidovanadium(IV) frozen solutions (140 K) at pH 5.5. Dash line, simulation. Total concentration VO^{2+} : 2.5 mM, prephenate: 5 mM.




Fig. 4. Proposed structure for the complex at pH 4.2.

in solution could be reduced. This possibly increases the probability of the mayor formation of hydrolytic oxidovanadium(IV) species. The results suggested that species with more than one hydroxide in the coordination sphere would probably be present in this conditions (pH 5.5), but being the EPR data the only source of the information there was no complete certainty about this. The formation of the ligand-free hydrolytic species could not be accurately established (e.g. $\text{VO}(\text{OH})_3(\text{H}_2\text{O})^-$ [26]). Its occurrence is usually detected at higher pH values [26]. Besides, the calculated hyperfine coupling constant is not correlated with its presence or perhaps its occurrence in very low concentration was not possible to detect in our experimental conditions.

Supplementary information was obtained from the electronic spectrum. The best fit was obtained using the experimental data at pH 3.4. The UV–Vis spectrum of prephenate–oxidovanadium(IV) system shows a characteristic feature with two absorption bands: a broad and strong one at 778 nm and a second band at 619 nm (Fig 5). At this pH (Fig. 1) EPR spectrum displays two sets of resonances, demonstrating the existence of two, distinguishable specie in solution. In order to assign the experimental spectrum, the fit of the spectrum and its components were provided. As can be seen in




Fig. 5. Original UV–Vis absorbance spectrum (solid line) and UV–Vis absorbance spectrum after curve-fitting (dash line: $\text{VO}(\text{H}_2\text{O})_5^{2+}$ and dotted line: prephenate–oxidovanadium(IV) specie detected at pH 3.4). Total concentration VO^{2+} : 2.5 mM, prephenate: 5 mM.

Fig. 5, the fit shows bands at 850, 759, 605 and 582 nm. In order to assign them, comparison with previously reported data was taken into account. For oxidovanadium(IV) complexes, two bands are predicted [27]: band I- $3d_{xy} \rightarrow 3d_{xz}$, band II- $3d_{xy} \rightarrow 3d_{x^2-y^2}$ d–d transitions. The UV–Vis spectrum of $\text{VO}(\text{H}_2\text{O})_5^{2+}$ species shows the typical bands at 763 nm (band I) and 625 nm (band II) [22,28], then the fitted bands at 760 nm and 606 nm to this species were assigned by comparison. The bands with maxima at 850 nm and 582 nm belong to the prephenate–carboxylate–oxidovanadium(IV) complex detected at pH 3.4. This prediction is in accordance with earlier experiments, in which carboxylate moiety were involved in the coordination sphere of the metallic center [16,27]. Coordination through carboxylate moiety leads to a bathochromic effect from 763 nm to 850 nm for band I and hypsochromic effect from 606 nm to 582 nm for the band II as been stated for several carboxylate– VO^{2+} interactions [16,27–29].

4. Conclusions

The present results describe clearly the prephenate–vanadyl interactions. The EPR spectra allow the identification of both new chemical specie in solution. The binding modes of the species at pH 4.2 was determined. At higher pH the detected species contains mainly ArO^- and OH^- in the coordination sphere of the metal center. They demonstrate again the ability of vanadium to interact with oxo–functional groups and this fact would explain the possibility that this type of interactions would be involved in the uptake, transport and storage of vanadium(IV) in plants and fungi.

Acknowledgements

Authors thanks the supported by UNLP, CONICET (PIP-1125) and Eusko Jaularitza/Gobierno Vasco MV-2004-3-39. EGF is member of the Carrera del Investigador, CONICET.

References

- [1] J.V. Gray, J.R. Knowles, *Biochemistry* 33 (1994) 9953.
- [2] H. Goerisch, *Biochemistry* 17 (1978) 3700.
- [3] L.O. Zamir, E. Jung, R.A. Jensen, *J. Biol. Chem.* 258 (1983) 6492.
- [4] E. Jung, L.O. Zamir, R.A. Jensen, *Proc. Natl. Acad. Sci. USA* 83 (1986) 7231.
- [5] W. Fiddler, W.E. Parker, A.E. Wasserman, R.C. Doerr, *J. Agric. Food Chem.* 15 (1967) 757 (and references therein).
- [6] H. Kindl, *Eur. J. Biochem.* 7 (1969) 340.
- [7] E.J. Baran, *J. Inorg. Biochem.* 80 (2000) 1.
- [8] P.W. Linder, A. Voyé, *Polyhedron* 6 (1987) 53.
- [9] R.A. Olsen, J.C. Brown, J.H. Bennet, D. Blume, *J. Plant Nutr.* 5 (1982) 433.
- [10] V. Römheld, H. Marschner, *J. Plant Nutr.* 3 (1981) 551.
- [11] Y. Allegretti, E.G. Ferrer, A.C. González Baró, P.A.M. Williams, *Polyhedron* 19 (2000) 2613.

- [12] S.E. Castillo-Blum, N. Barbra-Behrens, *Coord. Chem. Rev.* 3 (2000) 196.
- [13] J.O. Nriagu (Ed.), *Vanadium in the Environment*, Adv. Environ. Sci. Technol., vol. 30, John Wiley & Sons Inc., New York, 1998 (part 1, Chemistry and Biochemistry).
- [14] J.O. Nriagu (Ed.), *Vanadium in the Environment*, Adv. Environ. Sci. Technol., vol. 31, John Wiley & Sons Inc., New York, 1998 (part 2, Health Effects).
- [15] E.J. Baran, *Adv. Plant Phys.* 10 (2008) 357.
- [16] E.G. Ferrer, M.V. Salinas, M.J. Correa, F. Vrdoljak, P.A.M. Williams, *Z. Naturforsch.* 60b (2005) 305.
- [17] A. Dörnyei, E. Garribba, T. Jakusch, P. Forgó, G. Micera, T. Kiss, *Dalton Trans.* (2004) 1882.
- [18] E.G. Ferrer, M.V. Salinas, M.J. Correa, L. Naso, D.A. Barrio, S.B. Etcheverry, L. Lezama, T. Rojo, P.A.M. Williams, *J. Biol. Inorg. Chem.* 11 (2006) 791.
- [19] T.S. Smith II, R. LoBrutto, V.L. Pecoraro, *Coord. Chem. Rev.* 228 (2002) 1.
- [20] J. Costa Pessoa, I. Cavaco, I. Correia, I. Tomaz, T. Duarte, P.M. Matias, *J. Inorg. Biochem.* 80 (2000) 35.
- [21] J. Costa Pessoa, I. Tomaz, T. Kiss, P. Buglyó, *J. Inorg. Biochem.* 84 (2001) 259.
- [22] J.T.C. Delgado, A.I. Tomaz, I. Correia, J. Costa Pessoa, J.G. Jones, C.F.G.C. Geraldes, M. Margarida, C.A. Castro, *J. Inorg. Biochem.* 99 (2005) 2328.
- [23] T. Jakusch, W. Jin, L. Yang, T. Kiss, D.C. Crans, *J. Inorg. Biochem.* 95 (2003) 1.
- [24] M. Guiotoku, F. Regina Mena Barreto Silva, J.C. Azzolini, A. Lucia Ramalho Merce, A.S. Mangrich, L.F. Sala, B. Polyhedron 26 (2007) 1269.
- [25] H. Esbak, E.A. Enyedy, T. Kiss, Y. Yoshihawa, H. Sakurai, E. Garribba, D. Rehder, *J. Inorg. Biochem.* 103 (2009) 590.
- [26] M.M. Iannuzzi, P.H. Rieger, *Inorg. Chem.* 14 (1975) 2895.
- [27] S. Berto, P.G. Daniele, C. Foti, E. Prenesti, S. Sammartano, *J. Mol. Liq.* 142 (2008) 57.
- [28] A.B.P. Lever, *Inorganic Electronic Spectroscopy*, second ed., Elsevier, The Netherlands, 1984.
- [29] P. Buglyó, E. Kiss, I. Fábán, T. Kiss, D. Sanna, E. Garribba, G. Micera, *Inorg. Chim. Acta* 306 (2000) 174.