



SPECIAL ARTICLE

Toxigenic fungal species and natural occurrence of mycotoxins in crops harvested in Argentina



María Laura Chiotta^{a,c,*}, María Verónica Fumero^{a,b}, Eugenia Cendoya^{a,b}, Juan Manuel Palazzini^{a,c}, María Silvina Alaniz-Zanon^{a,c}, María Laura Ramirez^{a,c}, Sofía Noemí Chulze^{a,c}

^a Research Institute on Mycology and Micotoxicology (IMICO) – National Scientific and Technical Research Council – Argentina (CONICET) – National University of Río Cuarto (UNRC). Road 36 Km 601 (5800), Río Cuarto - Córdoba, Argentina

^b Fellow from CONICET

^c Members of the Research Career of CONICET

Received 11 October 2019; accepted 1 June 2020

Available online 24 July 2020

KEYWORDS

Toxigenic fungi;
Mycotoxins;
Cereals;
Oilseeds;
Fruits

Abstract Mycotoxins are secondary metabolites produced by fungal species that mainly belong to *Aspergillus*, *Fusarium*, *Penicillium* and *Alternaria*, which can grow in a variety of crops including cereals, oilseeds and fruits. Consequently, their prevalence in foods and by-products not only affects human and animal health but also causes important losses in both domestic and international markets. This review provides data about toxigenic fungal species and mycotoxin occurrence in different crops commonly grown in Argentina. This information will be relevant to establish adequate management strategies to reduce the impact of mycotoxins on human food and animal feed chains and to implement future legislation on the maximum permitted levels of these fungal metabolites.

© 2020 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Hongos toxigénicos;
Micotoxinas;
Cereales;
Oleaginosas

Especies fúngicas toxigénicas y ocurrencia natural de micotoxinas en cultivos y productos cosechados en Argentina

Resumen Las micotoxinas son metabolitos secundarios producidos por diferentes especies fúngicas pertenecientes, principalmente, a los géneros *Aspergillus*, *Fusarium*, *Penicillium* y *Alternaria*. Dichos microorganismos pueden crecer en una gran variedad de cultivos, entre los que se incluyen cereales, oleaginosas y frutas. La presencia de micotoxinas en alimentos y subproductos no sólo afecta la salud humana y animal, sino que también causa pérdidas importantes en los mercados nacionales e internacionales. Esta revisión proporciona datos sobre la

* Corresponding author.

E-mail address: mchiotta@exa.unrc.edu.ar (M.L. Chiotta).

prevalencia de especies fúngicas toxigénicas y de micotoxinas en diferentes cultivos y productos cosechados en Argentina. Dicha información será relevante para establecer estrategias de manejo adecuadas para reducir la entrada de las micotoxinas en las cadenas alimentarias del hombre y de los animales, así como para establecer futuras legislaciones sobre los niveles máximos permitidos de dichos metabolitos.

© 2020 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Argentina is located in South America and due to its vast size and range of altitudes, has a large variety of agroclimatic areas. It is one of the most important food-producing countries worldwide, with almost 30 million hectares of cultivable soil. Agriculture, livestock and agroindustry are the main economic activities, which have traditionally supplied the country with 70–95% of its export earnings. The economy of Argentina is mainly focused on the import and export of cereals, oil grains, fruits, wine, tea, tobacco and cotton.^{50,66,67} Annually, 25–50% of worldwide harvested crops are contaminated with mycotoxins, which are secondary metabolites produced by several fungal species not only during crop development, but also in storage conditions. The fungal genera most commonly isolated from foods and by-products are *Aspergillus*, *Penicillium*, *Fusarium* and *Alternaria*. Different species belonging to these genera can contaminate several substrates and produce mycotoxins causing adverse health effects in consumers, in addition to economic losses in food-producing and food-exporting countries.^{1,40,95} Most mycotoxins showed immunosuppressive, hepatotoxic, nephrotoxic and neurotoxic effects and some of them are carcinogens.^{16,55} Data on the occurrence of toxigenic species and mycotoxins are relevant to determine the contamination risk both for humans and animals in different regions. The present review provides data about the occurrence of toxigenic fungal species and the most frequent mycotoxins detected in cereals, oilseeds, fruits and by-products from Argentina.

Classification of mycotoxins

The most important mycotoxins based on their occurrence and toxicity are aflatoxins (AFs), fumonisins (FBs), zearalenone (ZEA), trichothecenes (TRCs), ochratoxin A (OTA) and patulin (PAT).^{55,93} Other toxicologically important mycotoxins, but less studied, are ergot alkaloids, *Alternaria* toxins, citrinin, cyclopiazonic acid, roquefortine C, mycophenolic acid, penitrem, verruculogen, griseofulvin, citreoviridin, among others.^{62,81}

Aflatoxins and OTA are produced by a species belonging to the genus *Aspergillus*. Aflatoxins are mainly associated with *Aspergillus flavus* and *Aspergillus parasiticus* growth on a variety of food products, including cereals (e.g. maize, wheat, sorghum), peanuts, edible nuts, legumes and by-products.³² For OTA contamination in food, the most

relevant producing *Aspergillus* species belong to section *Nigri* and *Circundati*.³ Within the section *Nigri*, *Aspergillus carbonarius* and *Aspergillus niger* have been identified as one of the main sources of OTA contamination and accumulation in grapes, raisins and wines.²⁷ These species have also contributed to OTA contamination in coffee, cocoa, peanut and cereals.^{30,59,60,99}

Fumonisins, ZEA and TRC are produced by several species of the genus *Fusarium* and wheat and maize are the most frequently contaminated crops. Fumonisins are mainly produced by several members of the *Fusarium fujikuroi* species complex (FFSC), including *Fusarium verticillioides*, *Fusarium proliferatum*, *Fusarium nygamai*, and *Fusarium fujikuroi* as well as by *Fusarium oxysporum*, which is not included in the FFSC.¹⁸ Zearalenone is a mycotoxin produced particularly by *Fusarium graminearum* sensu stricto but also by *Fusarium culmorum*, *Fusarium cerealis*, *Fusarium equiseti*, *F. verticillioides*, and *Fusarium incarnatum*.⁶² TRCs are divided into four types from A to D, type A and B being the most prevalent toxins found as natural contaminants in the food and feed chains. The most important type A-trichothecenes (A-TRC) are T-2 and HT-2 toxins. Among type B-trichothecenes (B-TRC), the most frequently occurring mycotoxin is deoxynivalenol (DON), along with its biosynthetic precursors 3-acetyldeoxynivalenol (3-ADON) and 15-acetyldeoxynivalenol (15-ADON). Type A-TRCs are produced mainly by *Fusarium sporotrichioides* and *Fusarium langsethiae* whereas DON by *F. graminearum* ss and *F. culmorum*. Nivalenol (NIV) co-occurs with Fusarenon-X (FUX) and DON, and is produced mainly by the *F. graminearum* species complex (FGSC), *Fusarium crookwellense* and *Fusarium nivale*.^{35,68}

The most relevant *Alternaria* toxins are alternariol (AOH), alternariol monomethyl ether (AME), tenuazonic acid (TA), tentoxin and altenuene.³⁷ Species within this genus are commonly isolated from wheat, sorghum and barley. In oilseeds such as sunflower and rapeseed, tomato, apples, citrus fruits and vegetables, the occurrence of *Alternaria* species has also been reported.⁷³

Patulin is produced by different fungal species such as *Penicillium expansum* (*Penicillium leucopus*), *Penicillium crustosum*, *Penicillium patulum* (*Penicillium urtiae* and *Penicillium griseofulvum*) and *Aspergillus clavatus*.⁹² Patulin-producing fungi have been isolated from various fruits and vegetables mainly from apple and apple-based commodities, and occasionally from other fruits such as pears, oranges, grapes and their

sub-products.^{51–53,57,70,97,100,105} Patulin is highly soluble in water and highly stable in aqueous acid media; thus, it commonly contaminates fruit juices.⁵⁰

“Emerging mycotoxins” are defined as “mycotoxins, which are neither routinely determined, nor legislatively regulated; however, the evidence of their incidence is rapidly increasing”. The main *Fusarium* mycotoxins included in this group are enniatins, beauvericin, moniliformin, fusaproliferin, fusaric acid and culmorin. Sterigmatocystin and emodin are produced by several *Aspergillus* species and mycophenolic acid is produced by fungi of the *Penicillium* species.⁸ Recently, some researchers have paid attention to their presence in foods and feeds, especially to enniatins and beauvericin. However, data on their toxicity and incidence are still limited, and further research on these metabolites is necessary to determine the toxicological risk due to their occurrence.

Toxigenic fungal species and natural occurrence of mycotoxins on different crops from Argentina

Several fungal species are commonly isolated from a wide variety of crops commonly grown in Argentina. Their occurrence cause spoilage and reduction in food quality and safety during different stages of the chain production such as pre-harvest, post-harvest processing and transport. Thus, their presence in foods and by-products is considered a potential risk due to their ability to produce mycotoxins.

Wheat flour, bran and grains

Alternaria, *Fusarium*, *Aspergillus* and *Penicillium* species can be found as frequent fungal contaminants in wheat.^{11,89} *Alternaria* was the predominant genus isolated from wheat in different agroecological regions from Argentina such as Entre Ríos and the southeast of Buenos Aires province, where *Alternaria alternata* is the most prevalent species followed by *Alternaria infectoria*.^{11,85} In contrast, Perello et al.⁸² observed a high incidence of *A. infectoria* in wheat grains collected from Buenos Aires area, which could be explained by changes in cropping systems in the last few years. The authors associated this species as the etiological agent of black point disease in wheat grains.

The main pathogen associated with *Fusarium* head blight (FHB) or scab worldwide is *F. graminearum* ss and, it is also the principal pathogen associated with FHB in wheat in Argentina.^{56,80} However, other *Fusarium* species have been isolated from bread (*Triticum aestivum* L.) and durum (*Triticum turgidum* L. var. durum) wheat grains during FHB non-outbreak years. *Fusarium proliferatum*, *Fusarium subglutinans* and *F. verticillioides* species were recovered from bread wheat.⁸⁶ Later, Palacios et al.⁷⁷ reported the presence of *F. proliferatum*, *F. oxysporum*, *F. subglutinans* and *F. equiseti* in durum wheat.

TRC natural occurrence on bread and durum wheat grains during FHB outbreak years was evaluated across the main wheat growing regions, including Santa Fe, Buenos Aires and Córdoba provinces (Figure 1). Type A-TRC, T-2 and HT-2 toxins were detected in 2 to 30% of the samples analyzed with

levels ranging from 10 to 70 ng/g and from 10 to 30 ng/g for T-2 and HT-2, respectively. Among type B-TRC, DON was the predominant toxin detected in frequencies ranging from 50 to 85% of the samples, the maximum levels detected being higher than 10,000 ng/g.^{7,49,58,79,89} Palacios et al.⁷⁸ detected not only DON but also the glycosylated form deoxynivalenol-3-glucoside (D3G) in 100 and 94% of durum wheat samples collected from three different geographic locations in the South of Buenos Aires province. The DON and D3G levels observed ranged from 50 to 9,400 ng/g and 50 to 850 ng/g, respectively. Moreover, the DON acetylated derivatives were detected but at lower frequency (49%). Recently, Cirio et al.²⁹ reported natural DON contamination in wheat flour processed in different mills located in Buenos Aires, Santa Fe, Entre Ríos, Córdoba, and Tucumán provinces. Ninety-one percent of the samples were contaminated with DON and the mean level detected was 243,000 ng/g. Basilico et al.⁷ analyzed NIV occurrence in bread wheat harvested in Santa Fe province and detected NIV levels ranging from 100 to 1,000 ng/g in only 5% of the samples (Table 1).

Fumonisin contamination of durum wheat grain samples collected during two consecutive harvest seasons (2007 and 2008) in different commercial fields located in the South of Buenos Aires province was reported for the first time by Palacios et al.⁷⁷ Data from this survey demonstrated that 97% of the samples collected during the 2007 harvest season showed total FB levels ranging from 10 to 1300 ng/g, while very low levels were detected in samples collected during the 2008 harvest season. The authors argued that these results could be explained by differences in rainfall during both evaluated harvest seasons. Later, Cendoya et al.¹⁷ reported FB occurrence not only in durum wheat grain samples, but also in bread wheat grain samples harvested in Buenos Aires province. From the samples, 93% were positive for FBs with levels ranging from 0.10 to 700 ng/g and from 0.10 to 1400 ng/g in bread and durum wheat, respectively. Recently FB contamination in flour wheat samples (68%) collected from different retail stores in Rio Cuarto city was observed, with levels ranging from 0.05 to 18.9 ng/g.¹⁸

Natural occurrence of AOH, AME and TA in 65 samples of whole, bran, and flour wheat obtained from a mill located in Gualeguaychú (Entre Ríos) was observed by Romero Bernal et al.⁹⁰ The wheat grain samples showed low AOH and AME contamination in comparison with TA levels. AOH and AME were not detected in bran and flour samples. The mean levels found for TA in whole, bran and flour wheat were 19,200, 16,800 and 7,300 ng/g, respectively.

Maize flour and grains

The genus *Fusarium* is commonly isolated from all maize-growing regions of Argentina. Studies on *Fusarium* species incidence carried out in both the main and the secondary (Northwest) maize production regions showed that *F. verticillioides* was the predominant species followed by *F. proliferatum* and *F. graminearum* ss.^{10,46,48,69,75,89} *Fusarium subglutinans* and *Fusarium temperatum* occurrence was high (45 and 60%) in the secondary production region, where lower annual temperatures than in the other areas are recorded.⁴¹ *Penicillium* incidence on maize grains ranged from 40 to 60% and the most prevalent species



Figure 1 Main crops grown in Argentina. Scale: less than 100,000 tn, between 100,000 tn and 500,000 tn, between 500,000 tn and 1,500,000 tn, between 1,500,000 tn and 4,000,000 tn, between 4,000,000 tn and 6,000,000 tn.

isolated were *Penicillium funiculosum*, *Penicillium citrinum*, *Penicillium decumbens*, *Penicillium oxalicum*, *Penicillium brevicompactum* and *Penicillium variable*.^{46,48,60,75} The genus *Aspergillus* was also isolated from maize grains in levels that varied between 15 and 30%, *A. flavus* and *A. parasiticus* being the common observed species.^{13,36,60} Among other toxicogenic species found in maize, *A. alternata* was observed in 5 to 10% of the samples.^{10,36,48,75,89} Other non-toxicogenic fungal species belonging to *Cladosporium*, *Curvularia*, *Diplodia*, *Epicoccum*, *Mucor*, *Nigrospora*, *Phoma*, *Rhizopus* and *Trichoderma* genera were also isolated from this crop.^{10,36,48,89}

Fumonisin B₁ (FB₁) is the main mycotoxin associated with maize. It was detected in high percentages of maize flour and grains (90 to 100%) collected in all maize growing regions and in different genotypes, with levels ranging from 1,000 to 10,000 ng/g.^{10,31,44,47,63,69,75,87,89,96,98} Deoxynivalenol and ZEA

have been recovered from maize at lower levels than FB₁; in 30% and < 10% of the samples with levels ranging from 0.1 to 140 ng/g and from 0.1 to 85 ng/g, respectively.^{28,44,47,75,87,96} With regard to AFs, several studies demonstrated low incidence in maize (5 to 10%), with levels ranging from 30 to 300 ng/g. These studies showed that AF occurrence in maize is influenced by climate conditions.^{28,36,44,47,69,75,81,87,89,96}

Although at present the natural occurrence of "emerging mycotoxins" in maize has not been evaluated in Argentina yet, the presence of *F. subglutinans* and *F. temperatum* have been reported, and both species have high potential to produce beauvericin, moniliformin and fusaproliferin.^{41,88} Possible synergic effects of these metabolites with other main mycotoxins such as DON and FBs have been observed.^{38,65} There is lack of knowledge concerning most of the emerging mycotoxins; thus, further studies will be nec-

Table 1 Fungal species and principal mycotoxins in the main crops cultivated in Argentina.

Crops	Species	Fungal incidence (%)	Mycotoxins	% positive samples	Levels (ng/g)
Wheat	<i>F. graminearum</i>	20 - 30	T2	20	10 - 70
			HT2	30	10 - 30
			DON	50 - 85	1000 - 10000
			NIV	5	100 - 1000
Maize	<i>F. proliferatum</i>	10 - 70	FBs	90 - 100	10 - 1400
	<i>F. verticillioides</i>	80 - 100	FBs	90 - 100	1000 - 10000
	<i>F. proliferatum</i>	10 - 20			
	<i>F. graminearum</i>	60 - 75	DON	30	0.1 - 140
Soybean	<i>A. flavus</i>	10 - 70	ZEA	< 10	0.1 - 85
	<i>A. parasiticus</i>	10 - 30	AFs	5 - 10	30 - 300
	<i>A. alternata</i>	ND	AOH	10 - 20	20 - 200
			AME	10 - 20	60 - 1200
Grapes	<i>F. graminearum</i>	ND	DON	< 5	500 - 5000
	<i>A. niger</i>	80	OTA	40 - 50	0.1 - 5
	<i>A. carbonarius</i>	10			

ND: No data.

essary to obtain information about their natural occurrence and evaluate possible synergism with the main mycotoxins.

Soybean seeds, pods and flowers

Alternaria and *Fusarium* are the most frequent genera isolated from soybean in Argentina followed by *Sclerotinia*, *Phomopsis*, *Rhizoctonia*, *Cladosporium*, *Aspergillus*, *Chaetomium* and *Penicillium*.^{4,9,11,43,45,91,103,106} With regard to *Alternaria*, *A. alternata* was the most common species found in soybean seed samples.^{4,11} Among *Fusarium* species, *F. equiseti*, *Fusarium semitectum* and FGSC were frequently isolated from soybean seeds and pods.^{6,24} A phylogenetic species recognition of the FGSC isolated from soybean seeds, pods and flowers samples was carried out in order to identify the species responsible for TRC production. The results showed evidence of the presence of four species within the *F. graminearum* species complex: *F. graminearum* ss, *Fusarium cortaderiae*, *Fusarium meridionale* and *Fusarium boothii*; *F. graminearum* ss being the most frequently identified fungal species.²⁵

The natural occurrence of AOH and AME in 50 soybean seed samples harvested in Córdoba province was evaluated by Oviedo et al.⁷⁴ Both mycotoxins were present in 60% of the samples whereas 16% and 14% were only contaminated with AOH and AME, respectively. Fifteen of the positive samples showed co-occurrence of both mycotoxins analyzed. Alternariol was detected at levels ranging from 25 to 211 ng/g, whereas AME was detected at concentrations ranging from 62 to 1,153 ng/g. Other studies on TRC natural contamination on soybean grains were conducted in samples also harvested in Cordoba province during the 2007/08 to 2013/14 harvest seasons.^{4,6,24,26} Data showed low DON levels in the sampled area, except during the 2013/14 season where a higher incidence and contamination with levels ranging from 0.7 to 4.3 ng/g were observed. These levels of contamination could be related to higher humidity periods during soybean grain filling and ripening during the 2013/14

harvest season. Although, none of the other mycotoxins such as AFs, ZEA, FBs and OTA were detected in soybean grains, the frequent presence of toxigenic species could indicate a risk of multiple mycotoxin contamination.^{45,106}

Grapes and wines

Alternaria and *Aspergillus* species were frequently isolated in Merlot, Malbec and Cabernet Sauvignon grape varieties collected from Mendoza wine grape growing region. *Penicillium*, *Cladosporium*, *Fusarium*, *Trichoderma* and *Ulocladium* species were isolated in lower frequency and were occasionally observed.^{20,61} In San Rafael wine grape growing region, the presence of four relevant genera in Malbec grape variety was observed. Among them, *Alternaria* was predominant (81%), followed by *Cladosporium* (7%) and alternatively by *Aspergillus* (4%) or *Penicillium* (3%). Moreover, *Drechslera* and other genera that belong to Mucoromycotina subphylum, were isolated in low levels (<5%). In this study, all *Alternaria* strains were identified as *A. alternata*.⁸⁴ In another study, the incidence of *Alternaria arborescens* and *Alternaria tenuisima* was also observed in Malbec grapes.¹⁰¹

Aspergillus section *Nigri* species were frequently isolated from different vineyards in Argentina.^{21-23,61,83} Studies carried out on red grape collected from Mendoza vineyards showed that species included within the *A. niger* aggregate were dominant and the main OTA producers.^{61,83} Subsequent studies on grapes harvested in different grape growing regions such as La Rioja - Chilecito, San Juan - Tulum Valley, Mendoza - Uco Valley, Mendoza North-East, Mendoza ZARM (high area near Mendoza River), Mendoza South and Neuquén - Río Negro), confirmed that *A. niger* aggregate species were dominant, followed by *A. carbonarius* and *Aspergillus* "uniseriate". *Aspergillus carbonarius* showed the highest percentages of OTA-producer strains and also the highest toxin level produced, being mainly isolated from the warmer regions, such as La Rioja and San Juan.¹⁹ Using molecular tools, *Aspergillus* section *Nigri* species were grouped

into four main clusters: *A. carbonarius*, *Aspergillus tubingensis*, *A. niger* "aggregate," and *A. uniseriate*. The *A. tubingensis* cluster was the most prevalent group and was clearly separated from the *A. niger* "aggregate".²¹ In another study, phylogenetic data corroborated the biodiversity of *Aspergillus* section *Nigri* populations. The sequencing data showed that the strains were grouped as *A. carbonarius*, *A. tubingensis*, *A. niger*, and *Aspergillus japonicus*. Moreover, *Aspergillus homomorphus* was identified and its isolation was relevant because it was the first time that it has been found in vineyards and in Argentina.²²

Wine samples collected at manufacturers' stock and retail markets from different regions of Argentina were analyzed to detect OTA occurrence, the results showing that none of the wine samples were contaminated.⁷⁶ In another study, OTA contamination from grapes and wine samples collected in different grape-growing regions was evaluated.¹⁹ Although ochratoxigenic species were isolated from samples obtained from the regions analyzed, the OTA levels detected in grape and wine samples were low in most of the evaluated samples, ranging from 0.10 to 5.40 ng/g and from 0.01 to 4.82 ng/ml, respectively. Another study showed that imported wines, mainly from Argentina and France, were the most contaminated with OTA, with levels ranging from 0.44 to 2.77 ng/ml.³³

Other foodstuffs

Although peanuts (*Arachis hypogaea* L.), yerba mate (*Ilex paraguariensis* St. Hil.) and fruits are economically important crops in Argentina, few studies on toxicogenic fungi and mycotoxin natural occurrence incidence have been conducted.

Studies on AF-producer species in peanuts harvested in the peanut-growing area of Córdoba province showed that *A. flavus* and *A. parasiticus* were frequently isolated.^{5,102} Magnoli et al.⁵⁹ evaluated the occurrence of *Aspergillus* section *Nigri* in stored peanut grains, *A. niger* var. *niger* and *A. niger* var. *awamori*, followed by *A. carbonarius* and *A. japonicus* being the predominant isolated species. In another study, Magnoli et al.⁵⁹ analyzed stored peanut grains but during a 3-month period, in which, although *A. carbonarius* was isolated in a low percentage, the percentage of OTA-producer strains was high. Aflatoxin, cyclopiazonic acid and OTA contamination in peanuts was detected in levels ranging from 0.043 to 435 ng/g, from 493 to 4300 ng/g, from 0.5 to 170 ng/g, respectively.^{5,39,44,102}

Patulin contamination in foods have been evaluated in different countries.^{2,34,54,64,71,104} In Argentina, Funes and Resnik⁴² analyzed PAT occurrence in solid and semisolid fruit by-products such as apple marmalade, jam, sweet and puree, and pear marmalade. The authors reported that 21.6% of the samples were contaminated with PAT (range 17–221 ng/g) and the highest contamination level was detected in apple puree with 50% of positive samples. Later, Oteiza et al.⁷² evaluated PAT contamination in fruit commodities (apple, apricot, grape, orange, peach, pear and pineapple) and different types of fruit by-products (concentrated cloudy juice, single strength cloudy juice, concentrated juice, concentrated pulp, single strength pulp and sulphited juice) collected between 2005 and 2013. Data

showed that PAT was detected in 33.5% of the samples with levels ranging from 30,000 to 19,622,000 ng/ml. With regard to the effect of fruit commodity, the highest percentage of positive samples (40%) contaminated with PAT and the maximum concentration were detected in apple. The type of products was also an important factor that affected PAT contamination; the higher levels were detected in single strength cloudy juice and single strength pulp (67.4% and 54.1%, respectively). In grape apple and orange juice, the natural occurrence of *Alternaria* toxins has also been observed.^{12,94}

In a study conducted by Castrillo et al.¹⁵, *Aspergillus* section *Nigri* incidence in different commercial brands of yerba mate was evaluated. Uniseriate species were more predominant than biseriate species. The percentage of isolated species was higher in milled yerba mate (without an aging period) than in elaborated yerba mate. In a previous study, the same authors performed a phylogenetic analysis of *Aspergillus* section *Nigri* isolated from yerba mate. In contrast, the results showed that all isolated strains were included in *A. niger* and *A. carbonarius* clades.¹⁴ No studies about natural mycotoxin occurrence in yerba mate have been conducted yet.

Conclusions

Different toxicogenic fungal species belonging to genera *Fusarium*, *Aspergillus* and *Alternaria* are isolated from the crops of greatest economic importance grown in Argentina. The isolation of these species was influenced by the type of crop and agro-ecological conditions. Mycotoxin occurrence was detected in all crops at variable levels, DON, TRC, FBs and OTA being the most frequently studied.

In Argentina the production of commodities increases annually and fungal contamination and mycotoxin occurrence in the food and feed chains represent a high risk to human and animal health, as well as considerable economic losses due to restrictions to the domestic and international markets. Studies on the occurrence of mycotoxins in different crops and by-products are important because they are reliable approaches to evaluate the potential exposure risk of the populations to these contaminants.

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The present work received funding from the European Union's Horizon 2020 Research and innovation program under Grant Agreement No. 678781 (MycoKey).

References

1. Adeyeye SA. Fungal mycotoxins in foods: A review. *Cogent Food Agric.* 2016;2:1–11.
2. Amézqueta S, González-Peña E, Murillo-Arbizu M, López de Cerain A, Ochratoxin A. decontamination: A review. *Food Control.* 2009;20:326–33.

3. Amézqueta S, Schorr-Galindo S, Murillo-Arbizu M, González-Peña E, López de Cerain A, Guiraud JP. OTA-producing fungi in foodstuffs: a review. *Food Control.* 2012;26:259–68.
4. Barros G, Oviedo MS, Ramirez ML, Chulze S. Safety aspect in soybean food and feed chains: fungal and mycotoxin contamination. In: Ng TB, editor. *Soybean. Biochemistry, Chemistry and Physiology.* Rijeka. Croatia: Intech Open Access Publisher; 2011. p. 7–20.
5. Barros G, Torres A, Palacio G, Chulze S. *Aspergillus* species from section *Flavi* isolated from soil at planting and harvest time in peanut-growing regions of Argentina. *J Sci Food Agric.* 2003;83:1303–7.
6. Barros G, Zanon MA, Abod A, Oviedo MS, Ramirez ML, Reynoso MM, Torres A, Chulze S. Natural deoxynivalenol occurrence and genotype and chemotype determination of a field population of the *Fusarium graminearum* complex associated with soybean in Argentina. *Food Addit Contam.* 2012;29:293–303.
7. Basílico MLZ, Pose G, Ludemann V, Pinto VEF, Aringoli EE, Ritieni A, Basílico JC. Fungal diversity and natural occurrence of fusaproliferin, beauvericin, deoxynivalenol and nivalenol in wheat cultivated in Santa Fe Province, Argentina. *Mycotoxin Res.* 2010;26:85–91.
8. Berthiller F, Crews C, Dall'Asta C, Saeger SD, Haesaert G, Karlovsky P, Oswald IP, Seefelder W, Speijer G, Stroka J. Masked mycotoxins: A review. *Mol Nutr Food Res.* 2013;57:165–86.
9. Boca RT, Pacin AM, Gonzalez HHL, Resnik SL, Souza JC. Soja y micotoxinas: Flora fúngica- Variedades - Prácticas agronómicas. *Aceites y Grasas.* 2003;4:10–515.
10. Broggi LE, Resnik SL, Pacin AM, González HHL, Cano G, Taglieri D. Distribution of fumonisins in dry-milled corn fractions in Argentina. *Food Addit Contam.* 2002;19:465–9.
11. Broggi L, González HHL, Resnik S, Pacin A. *Alternaria alternata* prevalence in cereal grains and soybean seeds from Entre Ríos. Argentina. *Rev Iberoam Micol.* 2007;24:47–51.
12. Broggi L, Reynoso C, Resnik S, Martinez F, Drunday V, Bernal ÁR. Occurrence of alternariol and alternariol monomethyl ether in beverages from the Entre Ríos Province market. Argentina. *Mycotoxin Res.* 2013;29:17–22.
13. Camiletti BX, Torrico AK, Maurino MF, Cristos D, Magnoli C, Lucini EI, Pecci MDLPG. Fungal screening and aflatoxin production by *Aspergillus* section *Flavi* isolated from pre-harvest maize ears grown in two Argentine regions. *Crop Prot.* 2017;92:41–8.
14. Castrillo ML, Fonseca MI, Bich GA, Jerke G, Horianski MA, Zapata PD. Taxonomy and phylogenetic analysis of *Aspergillus* section *Nigri* isolated from yerba mate in Misiones (Argentina). *BAG J Basic Appl Genet.* 2012;23, 0-0.
15. Castrillo ML, Horianski MA, Jerke G. Isolation of *Aspergillus* section *Nigri* strains in yerba mate in Posadas (Misiones Argentina) and evaluation of their ochratoxigenic potential. *Rev Argent Microbiol.* 2013;45:110–3.
16. Council for Agricultural Science and Technology (CAST). Mycotoxins: risks in plant, animal, and human systems; Task force report No139, 2003. Ames, Iowa, USA.
17. Cendoya E, Monge MP, Palacios SA, Chiacchiera S, Torres AM, Farnochi MC, Ramirez ML. Fumonisin occurrence in naturally contaminated wheat grain harvested in Argentina. *Food Control.* 2014;37:56–61.
18. Cendoya E, Nichea MJ, Monge MP, Sulyok M, Chiacchiera SM, Ramirez ML. Fumonisin occurrence in wheat-based products from Argentina. *Food Addit Contam: Part B.* 2019;12:31–7.
19. Chiotta ML, Ponsone ML, Combina M, Torres AM, Chulze SN. *Aspergillus* section *Nigri* species isolated from different wine grape growing regions in Argentina. *Int J Food Microbiol.* 2009;136:137–41.
20. Chiotta ML, Ponsone ML, Torres ML, Combina M, Chulze SN. Influence of *Planococcus ficus* on *Aspergillus* section *Nigri* and ochratoxin A incidence in vineyards from Argentina. *Lett Appl Microbiol.* 2010;51:212–8.
21. Chiotta ML, Reynoso MM, Combina M, Torres AM, Chulze SN. Molecular characterization and toxigenic profile of *Aspergillus* section *Nigri* populations isolated from the main grape-growing regions in Argentina. *J Appl Microbiol.* 2011;110:445–54.
22. Chiotta ML, Susca A, Stea G, Mulè G, Perrone G, Logrieco A, Chulze SN. Phylogenetic characterization and ochratoxin A - fumonisin profile of black *Aspergillus* isolated from grapes in Argentina. *Int J Food Microbiol.* 2011;149:171–6.
23. Chiotta ML, Ponsone ML, Sosa DM, Combina M, Chulze SN. Biodiversity of *Aspergillus* section *Nigri* populations in Argentinian vineyards and ochratoxin A contamination. *Food microbiol.* 2013;36:182–90.
24. Chiotta ML, Chulze S, Barros G. Fuentes de inóculo de especies de *Fusarium* potenciales productoras de micotoxinas en el agroecosistema soja. *Rev Fac Cienc Agrar.* 2015;47:171–84.
25. Chiotta ML, Zanon MA, Gajaj-Merlera G, Tessmann D, Barros GG, Chulze SN. Phylogenetic analyses of the *Fusarium graminearum* species complex isolated from soybean in Argentina and Brazil. *Australas Plant Dis Notes.* 2015;10:32.
26. Chiotta ML, Palazzini JM, Alberione E, Barros GG, Chulze SN. Dynamic of *F. graminearum* species complex populations in soybean wheat rotation. *XI Congreso Argentino de Microbiología General SAMIGE.* 2015:495–7. Resumen, Córdoba Argentina.
27. Chiotta ML, Ponsone ML, Combina M, Chulze SN. *Aspergillus* and ochratoxin A in Latin America. In: Grarg N, Abdel-Aziz SM, Aeron A, editors. *Microbes in Food and Health.* Springer International Publishing; 2016. p. 265–87.
28. Chulze S, Bertinetti C, Dalcerio A, Etcheverry M, Farnochi C, Torres A, Rizzo I, Varsavsky E. Incidence of aflatoxin, zearalenone, and deoxynivalenol on corn in Argentina. *Mycotoxin Res.* 1989;5:9–12.
29. Cirio M, Villarreal M, Seal TML, Simón ME, Smersu CSS, Kneeteman EA. Incidence of Deoxynivalenol in Wheat Flour in Argentina and GC-ECD Method Validation. *J AOAC Int.* 2019;102:1721–4.
30. Copetti MV, Pereira JL, Iamanaka BT, Pitt JI, Taniwaki MH. Ochratoxigenic fungi and ochratoxin A in cocoa during farm processing. *Int J Food Microbiol.* 2010;143:67–70.
31. Coronel MB, Vicente S, Resnik SL, Alzamora SM, Pacin A. Fumonisins in maize and gluten meal analysed in Argentinean wet milling industrial plants by ELISA compared with HPLC-FLD method. *Food Control.* 2016;67:285–91.
32. da Rocha MEB, Freire FDCO, Maia FEF, Guedes MIF, Rondina D. Mycotoxins and their effects on human and animal health. *Food Control.* 2014;36:159–65.
33. de Abreu PS, Terra MF, Prado G, Santiago WD, das Graças Cardoso M, Valeriano C, Batista LR, Ochratoxin A. in wines and evaluation of consumer exposure. *Food Public Health.* 2016;6:107–14.
34. Delage N, d'Harlingue A, Colonna Ceccaldi B, Bompeix G. Occurrence of mycotoxins in fruit juices and wine. *Food Control.* 2003;14:225–7.
35. Desjardins AE, editor. 2006. *Fusarium* mycotoxins: chemistry, genetics, and biology. APS Press. [On-line] <https://www.cabdirect.org/cabdirect/abstract/20063036927>. Consultado el 4 de febrero de 2020.
36. Etcheverry M, Nesci A, Barros G, Torres A, Chulze S. Occurrence of *Aspergillus* section *Flavi* and aflatoxin B 1 in corn genotypes and corn meal in Argentina. *Mycopathologia.* 1999;147:37–41.
37. European Food Safety Authority (EFSA). Scientific opinion on the risks for public health related to the presence of zearalenone in food. *EFSA J.* 2011; 9:2197.
38. Fairchild AS, Grimes JL, Porter JK, Croom WJ Jr, Daniel LR, Hagler WM Jr. Effects of diacetoxyscirpenol and fusaric acid

- on poulets: Individual and combined effects of dietary diacetoxyscirpenol and fusaric acid on turkey poult performance. *Int J Poult Sci.* 2005;4:350–5.
39. Fernandez Pinto V, Patriarca A, Locani O, Vaamonde G. Natural co-occurrence of aflatoxin and cyclopiazonic acid in peanuts grown in Argentina. *Food Addit Contam.* 2001;18:1017–20.
40. Food and Agriculture Organization for the United Nations (FAO). WFP, The State of Food Insecurity in the World 2013. The multiple dimensions of food security. 2013.
41. Fumero MV, Reynoso MM, Chulze SN. *Fusarium temperatum* and *Fusarium subglutinans* isolated from maize in Argentina. *Int J Food Microbiol.* 2015;199:86–92.
42. Funes GJ, Resnik SL. Determination of patulin in solid and semisolid apple and pear products marketed in Argentina. *Food Control.* 2009;20:277–80.
43. Gally T, Gonzalez BA, Pastuso F. Efecto conjunto de *Fusarium* sp. y *Phomopsis* sp., patógenos transmitidos por las semillas en plántulas de soja [Glycine max (L.) Merrill]. *Rev Mex Fitopatol.* 2006;24:156–8.
44. Garrido CE, Pezzani CH, Pacin A. Mycotoxins occurrence in Argentina's maize (*Zea mays L.*), from 1999 to 2010. *Food Control.* 2012;25:660–5.
45. Garrido CE, González HH, Salas MP, Resnik SL, Pacin AM, Resnik S, Pacin AM. Mycoflora and mycotoxins contamination of soybean RR harvested in the Pampean Region. Argentina. *Mycotoxin Res.* 2013;29:147–57.
46. González HHL, Resnik SL, Boca RT, Marasas WFO. Mycoflora of Argentinian corn harvested in the main production area in 1990. *Mycopathologia.* 1995;130:29–36.
47. Gonzalez HH, Martinez EJ, Pacin AM, Resnik SL, Sydenham EW. Natural co-occurrence of fumonisins, deoxynivalenol, zearalenone and aflatoxins in field trial corn in Argentina. *Food Addit Contam.* 1999;16:565–9.
48. González HHL, Resnik SL, Pacin AM. Mycoflora of freshly harvested flint corn from Northwestern Provinces in Argentina. *Mycopathologia.* 2003;155:207–11.
49. González HHL, Moltó GA, Pacin A, Resnik SL, Zelaya MJ, Masana M, Martínez EJ. Trichothecenes and mycoflora in wheat harvested in nine locations in Buenos Aires province Argentina. *Mycopathologia.* 2008;165:105–14.
50. Gokmen V, Acar J. Long-term survey of patulin in apple juice concentrates produced in Turkey. *Food Addit Contam.* 2000;17:933–6.
51. Guo Y, Zhou Z, Yuan Y, Yue T. Survey of patulin in apple juice concentrates in Shaanxi (China) and its dietary intake. *Food Control.* 2012;34:570–3.
52. Harris KL, Bobe G, Bourquin LD. Patulin surveillance in apple cider and juice marketed in Michigan. *J Food Prot.* 2009;72:1255–61.
53. Iha MH, Sabino M. Incidence of patulin in Brazilian apple-based drinks. *Food Control.* 2008;19:417–22.
54. Iha MH, Barbosa CB, Heck AR, Trucksess MW. Aflatoxin M1 and ochratoxin A in human milk in Ribeirao Preto-SP. Brazil. *Food Control.* 2014;40:310–3.
55. International Agency for Research on Cancer. IARC working group reports N°9. Mycotoxin Control in low- and middle-income countries. Wild C, Miller JD, Groopman D, editors. Lyon, France, 2015, p. 54.
56. Kikot GE, Moschini R, Consolo VF, Rojo R, Salerno G, Hours RA, Gasón L, Aramburu AM, Alconada TM. Occurrence of different species of *Fusarium* from wheat in relation to disease levels predicted by a weather-based model in Argentina pampas region. *Mycopathologia.* 2011;171:139–49.
57. Lee TP, Sakai R, Manaf NA, Rodhi AM, Saad B. High performance liquid chromatography method for the determination of patulin and 5-hydroxymethylfurfural in fruit juices marketed in Malaysia. *Food Control.* 2014;38:142–9.
58. Lori GA, Sistera MN, Haidukowski M, Rizzo I. *Fusarium graminearum* and deoxynivalenol contamination in the durum wheat area of Argentina. *Microbiol Res.* 2003;158:29–35.
59. Magnoli C, Astoreca A, Ponsone L, Chiacchiera S, Dalcerio AM. Ochratoxin A and the occurrence of ochratoxin A-producing black aspergilli in stored peanut seeds from Córdoba. Argentina. *J Sci Food Agric.* 2006;86:2369–73.
60. Magnoli C, Hallak C, Astoreca A, Ponsone L, Chiacchiera S, Dalcerio AM. Occurrence of ochratoxin A-producing fungi in commercial corn kernels in Argentina. *Mycopathologia.* 2006;161:53–8.
61. Magnoli C, Violante M, Combina M, Palacio G, Dalcerio A. Mycoflora and ochratoxin-producing strains of *Aspergillus* section *Nigri* in wine grapes in Argentina. *Lett Appl Microbiol.* 2003;37:179–84.
62. Marin S, Ramos AJ, Cano-Sancho G, Sanchis V. Mycotoxins: occurrence, toxicology, and exposure assessment. *Food Chem Toxicol.* 2013;60:218–37.
63. Martínez M, Moschini R, Barreto D, Bodega J, Comerio R, Forjan H, Piatti F, Presello D, Valentín O. Factores ambientales que afectan el contenido de fumonisina en granos de maíz. *Trop Plant Pathol.* 2010;35:277–84.
64. Makun HA, Adeniran AL, Mailafiya SC, Ayanda IS, Mudashiru AT, Ojukwu UJ. Natural occurrence of ochratoxin A in some marketed Nigerian foods. *Food Control.* 2013;31:566–71.
65. May HD, Wu Q, Blake CK. Effects of the *Fusarium* spp. mycotoxins fusaric acid and deoxynivalenol on the growth of *Ruminococcus albus* and *Methanobrevibacter ruminantium*. *Can J Microbiol.* 2000;46:692–9.
66. Ministerio de agricultura, ganadería y pesca [Internet]. Argentina: MAGyP; 2018 [updated Sep 2018; cited Sep 2018]. Available from: <https://datos.magyp.gob.ar/reportes.php?reporte=Estimaciones>.
67. Ministerio de Agroindustria [Internet]. Argentina: MINAGRI; 2018 [updated Sep 2018; cited Sep 2018]. Available from: <http://www.agroindustria.gob.ar/sitio/areas/estimaciones/index.php>.
68. Miller JD. Mycotoxins in small grains and maize: old problems, new challenges. *Food Addit Contam.* 2008;25:219–30.
69. Monge M, Magnoli CE, Chiacchiera SM. Survey of *Aspergillus* and *Fusarium* species and their mycotoxins in raw materials and poultry feeds from Córdoba. Argentina. *Mycotoxin Res.* 2012;28:111–22.
70. Morales H, Marín S, Rovira A, Ramos AJ, Sanchis V. Patulin accumulation in apples by *Penicillium expansum* during postharvest stages. *Lett Appl Microbiol.* 2007;44:30–5.
71. Nguyen KTN, Ryu D. Concentration of ochratoxin A in breakfast cereals and snacks consumed in the United States. *Food Control.* 2014;40:140–4.
72. Oteiza JM, Khaneghah AM, Campagnollo FB, Granato D, Mahmoudi MR, Sant'Ana AS, Gianuzzi L. Influence of production on the presence of patulin and ochratoxin A in fruit juices and wines of Argentina. *LWT - Food Sci Technol.* 2017;80:200–7.
73. Ostry V. Alternaria mycotoxins: an overview of chemical characterization, producers, toxicity, analysis and occurrence in foodstuffs. *World Mycotoxin J.* 2008;1:175–88.
74. Oviedo MS, Barros GG, Chulze SN, Ramirez ML. Natural occurrence of alternariol and alternariol monomethyl ether in soya beans. *Mycotoxin Res.* 2012;28:169–74.
75. Pacin AM, Brogi LE, Resnik SL, González HHL. Mycoflora and mycotoxins natural occurrence in corn from Entre Ríos province. Argentina. *Mycotoxin Res.* 2001;17:39–44.
76. Pacin A, Resnik S, Vega M, Saelzer R, Ciancio B, Emilia RG. Occurrence of ochratoxin A in wines in the Argentinean and Chilean markets. *ARKIVOC.* 2005:214–23.
77. Palacios SA, Ramirez ML, Cabrera Zalazar M, Farnochi MC, Zapacosta D, Chiacchiera SM, Reynoso MM, Chulze SN, Torres AM.

- Occurrence of *Fusarium* spp. and fumonisin in durum wheat grains. *J Agric Food Chem.* 2011;59:12264–9.
78. Palacios SA, Erazo JG, Ciasca B, Lattanzio VM, Reynoso MM, Farnochi MC, Torres AM. Occurrence of deoxynivalenol and deoxynivalenol-3-glucoside in durum wheat from Argentina. *Food Chem.* 2017;230:728–34.
79. Palazzini J, Fumero V, Yerkovich N, Barros G, Cuniberti M, Chulze S. Correlation between *Fusarium graminearum* and deoxynivalenol during the 2012/13 wheat *Fusarium* head blight outbreak in Argentina. *Cereal Res Commun.* 2015;43:627–37.
80. Palazzini JM, Alberione E, Torres A, Donat C, Köhl J, Chulze S. Biological control of *Fusarium graminearum* sensu stricto, causal agent of *Fusarium* head blight of wheat, using formulated antagonists under field conditions in Argentina. *Biol Control.* 2016;94:56–61.
81. Pereira VL, Fernandes JO, Cunha SC. Mycotoxins in cereals and related foodstuffs: A review on occurrence and recent methods of analysis. *Trends Food Sci Technol.* 2014;36:96–136.
82. Perello A, Moreno M, Sisterna M. *Alternaria infectoria* species-group associated with Black point of wheat in Argentina. *Plant Pathol.* 2008;57:379.
83. Ponsone ML, Combina M, Dalcero A, Chulze S. Ochratoxin A and ochratoxigenic *Aspergillus* species in Argentinian wine grapes cultivated under organic and nonorganic systems. *Int J Food Microbiol.* 2007;114:131–5.
84. Prendes LP, Merín MG, Andreoni MA, Ramirez ML, de Ambrosini VIM. Mycobiota and toxicogenic *Alternaria* spp. strains in Malbec wine grapes from DOC San Rafael, Mendoza Argentina. *Food Control.* 2015;57:122–8.
85. Ramirez ML, Sturm ME, Oviedo MS, Chulze SN. *Alternaria* species isolated from wheat in Argentina. Proceeding of Myco-Globe: Reducing Impact of Mycotoxins in Tropical Agriculture p.35, Acra Ghana. 2005.
86. Ramirez ML, Oviedo MS, Farnochi MC, Chulze SN. Mycological and mycotoxin survey on wheat during a non epidemic FHB year in Argentina. Proceeding of 9th European Fusarium Seminar. 2006:34. Wageningen The Netherlands.
87. Resnik S, Costarrica ML, Pacin A. Mycotoxins in Latin America and the Caribbean. *Food Control.* 1995;6:19–28.
88. Reynoso MM, Torres AM, Chulze SN. Fusaproliferin, beauvericin and fumonisin production by different mating populations among the *Gibberella fujikuroi* complex isolated from maize. *Mycol Res.* 2004;108:154–60.
89. Roigé MB, Aranguren SM, Riccio MB, Pereyra S, Soraci AL, Tapia MO. Mycobiota and mycotoxins in fermented feed, wheat grains and corn grains in Southeastern Buenos Aires Province, Argentina. *Rev Iberoam Micol.* 2009;26:233–7.
90. Romero Bernal AR, Reynoso CM, García Londoño VA, Broggi LE, Resnik SL. *Alternaria* toxins in Argentinean wheat, bran, and flour. *Food Addit Contam: Part B.* 2019;12:24–30.
91. Roy KW, Baird RE, Abney TS. A review of soybean (*Glycine max*) seed, pod and flower mycoflora in North America, with methods and a key for identification of selected fungi. *Mycopathologia.* 2000;150:15–27.
92. Samson RA, Houbraken J, Varga J, Frisvad JC. Polyphasic taxonomy of the heat resistant ascomycete genus *Byssochlamys* and its *Paecilomyces* anamorphs. *Persoonia.* 2009;22:14–27.
93. Schatzmayr G, Streit E. Global occurrence of mycotoxins in the food and feed chain: facts and figures. *World Mycotoxin J.* 2013;6:213–22.
94. Scott PM, Lawrence GA, Lau BPY. Analysis of wines, grape juices and cranberry juices for *Alternaria* toxins. *Mycotoxin Res.* 2006;22:142–7.
95. Sforza S, Dall'Asta C, Marchelli R. Recent advances in mycotoxin determination in food and feed by hyphenated chromatographic techniques/mass spectrometry. *Mass Spectrom Rev.* 2006;25:54–76.
96. Solovej MMS, Somoza C, Cano G, Pacin A, Resnik S. A survey of fumonisins, deoxynivalenol, zearalenone and aflatoxins contamination in corn-based food products in Argentina. *Food Addit Contam.* 1999;16:325–9.
97. Spadaro D, Ciavarella A, Frati S, Garibaldi A, Gullino ML. Incidence and level of patulin contamination in pure and mixed apple juices marketed in Italy. *Food Control.* 2007;18:1098–102.
98. Sydenham EW, Shephard GS, Thiel PG, Marasas WF, Rheeder JP, Peralta Sanhueza CE, Gonzales HHL, Resnik SL. Fumonisins in Argentinian field-trial corn. *J Agric Food Chem.* 1993;41:891–5.
99. Taniwaki MH, Pitt JI, Teixeira AA, Iamanaka BT. The source of ochratoxin A in Brazilian coffee and its formation in relation to processing methods. *Int J Food Microbiol.* 2003;82:173–9.
100. Torović L, Dimitrov N, Assunção R, Alvito P. Risk assessment of patulin intake through apple-based food by infants and preschool children in Serbia. *Food Addit Contam Part A Chem Anal Control Expo Risk Asses.* 2017;34:2023–32.
101. Trinidad AV, Ganoza FQ, Pinto VF, Patriarca A. Determination of mycotoxin profiles characteristic of *Alternaria* strains isolated from Malbec grapes. 38th World Congress of Vine and Wine. BIO Web of Conferences. 2015:02004.
102. Vaamonde G, Patriarca A, Pinto VF, Comerio R, Degrossi C. Variability of aflatoxin and cyclopiazonic acid production by *Aspergillus* section *Flavi* from different substrates in Argentina. *Int J Food Microbiol.* 2003;88:79–84.
103. Villarroel DA, Baird RE, Trevathan LE, Watson CE, Scruggs ML. Pod and seed mycoflora on transgenic and conventional soybean [*Glycine max* (L.) Merrill] cultivars in Mississippi. *Mycopathologia.* 2004;157:207–15.
104. Wu J, Tan Y, Wang Y, Xu R. Occurrence of ochratoxin A in wine and beer samples from China. *Food Addit Contam Part B Surveill.* 2001;4:52–6.
105. Zaied C, Abid S, Hlel W, Bacha H. Occurrence of patulin in apple-based-foods largely consumed in Tunisia. *Food Control.* 2013;31:263–7.
106. Zelaya MJ, González HH, Resnik SL, Pacin AM, Salas MP, Martínez MJ. Mycobiota and potential mycotoxin contamination of soybean RR in different production areas in Argentina. *Int Res J Plant Sci.* 2013;4:133–43.