

---

## Two New Species of *Senecio* (Asteraceae, Senecioneae) from Argentina

Roberto D. Tortosa

Laboratorios de Botánica “L. R. Parodi,” Facultad de Agronomía, Universidad de Buenos Aires, Av. San Martín 4453, 1417 Buenos Aires, Argentina. tortosa@agro.uba.ar; and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina.

Adriana Bartoli

Laboratorios de Botánica “L. R. Parodi,” Facultad de Agronomía, Universidad de Buenos Aires, Av. San Martín 4453, 1417 Buenos Aires, Argentina. cbartoli@agro.uba.ar

---

**ABSTRACT.** Two new species of *Senecio* L. from Argentina, *S. fragrantissimus* and *S. pumilus*, are described and illustrated. Both species belong to *Senecio* ser. *Xerosenecio* (Cabrera) Cabrera & S. E. Freire and are allied to *S. filagineoides* DC., from which *S. fragrantissimus* differs by its ascending branches with needle-like leaves crowded along the stems, and *S. pumilus* diverges by its dwarf cushion-like habit and elliptic, thicker leaves. A key is included to those species affined to *S. filagineoides*.

**RESUMEN.** Se describen e ilustran dos nuevas especies de *Senecio* L. de Argentina, *S. fragrantissimus* y *S. pumilus*. Ambas pertenecen a *Senecio* serie *Xerosenecio* (Cabrera) Cabrera & S. E. Freire y son afines a *S. filagineoides* DC., de la cual *S. fragrantissimus* difiere por sus ramas ascendentes con hojas aciculares aglomeradas a lo largo de las ramas, y *S. pumilus* diverge por formar cojines con hojas elípticas más crasas. Se incluye una clave para las especies afines a *S. filagineoides*.

**Key words:** Argentina, Asteraceae, Compositae, *Senecio*.

The series *Xerosenecio* (Cabrera) Cabrera & S. E. Freire of the genus *Senecio* L. sect. *Senecio* (sensu Cabrera et al., 1999) is characterized by a ligneous habit and discoid capitula disposed in terminal corymbs. Cabrera (1949) and Cabrera et al. (1999) included 39 species in this series, distributed in Argentina and Chile, with only one species (*S. vira-vira* Hieronymus) also in Uruguay and in southeastern Brazil in the state of Rio Grande do Sul.

During a taxonomic revision of *Senecio* ser. *Xerosenecio*, we found two new species from Argentina, both affined to *Senecio filagineoides* DC. by their woolly stems, leaves, and phyllaries, and by their narrowly cylindric or obconical involucres, less than 8 mm high, composed of 10 to 14 phyllaries.

***Senecio fragrantissimus*** Tortosa & A. Bartoli, sp. nov. TYPE: Argentina. San Luis: Pringles, Sadillo, “Flores color crema, perfumadas,” 9 Nov. 1940, G. Covas 1337 (holotype, LP). Figure 1.

Affinis *Senecioni filagineoidi* DC. sed ramis hornotinis ascendentibus foliis acicularis congestis notabilis differt.

Shrub 50–80 cm, mostly ramified from the base, lanuginose or sublanuginose on stems, leaves, and phyllaries; stems ascending with very short internodes from midstem upward. Leaves alternate, sessile, acicular, ± spreading, 0.9–3.5 × 0.04–0.06 cm, margin revolute, lanuginose or sublanuginose on both surfaces. Capitula pedunculate, arranged 5 to 15 in terminal corymbs; peduncles 0.5–1.3 cm long, involucre calyculate, cylindric, 6–8 × 2.5–3 mm; phyllaries 10 to 12, linear ovate, 6–8 × ca. 1 mm, dorsally lanuginose or sparsely lanuginose, apex acuminate with a tuft of short trichomes; calyculary bracts ca. 4, linear, ca. 1.5 mm long, dorsally lanuginose. Disc florets 22 to 25, hermaphroditic, corolla cream, tubular, 5-lobed, 7–7.5 mm long, glabrous, basally broadened, limb slightly broader than the tube and not markedly distinct, lobes narrowly triangular, ca. 0.6 mm long, at base 0.3 mm wide; anthers minutely tailed, ca. 2–2.5 mm long; filament collars spatulate ca. 0.5 mm long; apical anther appendage obovate, ca. 0.4 mm long; basal tails ca. 0.15 mm long. Style branches ca. 1.5 mm long. Achenes cylindric, pubescent, ca. 3 mm, costate; pappus bristles many, slender, barbellate, 7–8 mm long, whitish.

**Distribution and habitat.** *Senecio fragrantissimus* is known only from two neighboring localities in the political province of San Luis of western Argentina, in xerophytic habitats with sandy soil at 800 m.



Figure 1. *Senecio fragrantissimus* Tortosa & A. Bartoli. —A. Habit. —B. Capitulum. Drawn from the type, G. Covas 1337 (LP).

**Etymology.** The specific epithet makes reference to the fragrance to which all collectors of this species alluded.

*Senecio fragrantissimus* is allied to *S. filaginoides*,

a species with a wide distribution in Argentina (Cabrera et al., 1999). The new species differs from the latter by its needle-like leaves, crowded together due to congestion of the upper stem internodes,




Figure 2. *Senecio pumilus* Tortosa & A. Bartoli. —A. Habit. —B. Capitulum. Drawn from the type, E. Ancibor y H. Schwabe s.n. (BAA 5789).

giving the branches and their foliage a cylindrical appearance. A similar aspect may be seen in some specimens of *S. bracteolatus* Hooker & Arnott (*O. Boelcke* 4289 (BAA), *A. Havrylenko* 6 and 7 (BAA), *Vallerini* 339 (BAA)), but this species is

glabrous and its heads have a campanulate involucre.

**Paratypes.** ARGENTINA. San Luis: Pringles, Saladillo, 9 Nov. 1940, *A. Burkart* (LP); 9 Nov. 1940, *G. Covas* 1001 (LP); General Pedernera, al pie del Cerro El Morro,

cerca de la Ea. La Morena, 18 Oct. 1951, A. T. Hunziker 9287 (LP).

**Senecio pumilus** Tortosa & A. Bartoli, sp. nov.

TYPE: Argentina. Mendoza: Malargüe, Portezuelo del Choique, 2400 m s.m., 12 Mar. 1967, E. Ancibor & H. Schwabe s.n. (holotype, BAA 5789). Figure 2.

Inter species seriei *Xerosenecionis* indumento lanato, capitulis parvis in corymbis dispositis, involucro angusto ad *S. filaginoides* DC. accedens sed statura minore, foliis ellipticis crassioribus involucro obconico ab ea recedens.

Dwarf shrubs 8–30 cm, much branched at the base, stems, leaves, and phyllaries densely whitish lanuginose; stems prostrate subligneous giving rise at their nodes to herbaceous ascending shoots. Leaves alternate, sessile, 0.8–2 × 0.15–0.30 cm, elliptic, obtuse or rounded at the apex, margin revolute, densely lanuginose on both surfaces. Capitula pedunculate, peduncles 0.5–1.5 cm long, arranged 3 to 6 in terminal corymbs; involucle calyculate, obconical, 6–7 × 3–3.5 mm; phyllaries 8 to 12, 6–7 × 1–1.2 mm, dorsally lanuginose, apex acuminate with short trichomes; calycular bracts ca. 5, elliptic, ca. 1–1.5 mm long, dorsally lanuginose. Disc florets ca. 20 to 24, hermaphroditic, corolla yellow, tubular, 6–6.5 mm long, glabrous, basally broadened, narrowly campanulate, lobes triangular, ca. 0.5 mm long, at base 0.3 mm wide; anthers minutely tailed, 2–2.5 mm long; filament collars spatulate, ca. 0.25 mm long; apical anther appendages deltoid, ca. 0.3 mm long; the basal tails ca. 0.1 mm long. Style branches ca. 1.2 mm long. Achenes cylindric, glabrous, ca. 3.5 mm, costate. Pappus bristles many, slender, barbellate, ca. 6 mm long, whitish.

The heads of *Senecio pumilus* are similar to those of *S. filaginoides*, but its dwarf cushion-shaped habit, not taller than 30 cm (vs. erect, 0.5–1 m tall in *S. filaginoides*), its elliptic, entire, thicker leaves (vs. linear to linear-elliptic in *S. filaginoides*), and its obconical involucre (vs. cylindric in *S. filaginoides*) distinguish the new species.

*Distribution and habitat.* *Senecio pumilus* has

been found in the eastern Mendoza and Neuquén in Argentina, at elevations of 1900–2500 m of the Andes Mountains, an arid region with cool weather exposed to strong winds.

*Etymology.* The specific epithet makes reference to the habit of the species.

*Paratypes.* ARGENTINA. Mendoza: Malargüe, Portezuelo del Choique, E. Ancibor & H. Schwabe s.n. (BAA 5784); Ca. Portezuelo del Choique, J. H. Hunziker 6381 (BAB); Malargüe, A. Burkart, N. S. Troncoso & E. G. Nicora s.n. (SI 14338); Cerros al N de Calmuco, A. Burkart s.n. (SI 14330); Laguna Coipo-Leuquen, al S de Calmuco, A. Burkart, N. S. Troncoso & E. G. Nicora s.n. (SI 14340); 6 km N de Calmuco, G. Covas 412 (LP); Entre Calmuco y Pto. Los Colgados, A. Prina, G. Alfonso & W. Muñoz 1870 (SRFA); in montanis Valle del Yeso, Gerling s.n. (LP). Neuquén: Minas, Cordillera del Viento, arroyo Dos Puentes, F. Pastore 3 (SI).

KEY TO SPECIES ALLIED TO *SENECIO FILAGINOIDES*

- 1a. Leaves acicular, less than 0.6 mm wide . . . . . *Senecio fragrantissimus* Tortosa & A. Bartoli
- 1b. Leaves linear or elliptic, 1.5–3 mm wide.
  - 2a. Phyllaries 7 to 8; florets 10 to 12 . . . . . *Senecio pycnanthus* Philippi
  - 2b. Phyllaries 10 to 14; florets 18 to 26.
    - 3a. Shrubs 0.08–0.30 m high; leaves elliptic, entire; involucre obconical . . . . . *Senecio pumilus* Tortosa & A. Bartoli
    - 3b. Shrubs 0.5–1 m high; leaves linear or linear-elliptic, entire, toothed or lobed; involucre cylindrical . . . . . *Senecio filaginoides* DC.

*Acknowledgments.* We thank the anonymous reviewers and Victoria C. Hollowell for reviewing the manuscript and for useful comments. We also acknowledge the assistance of the curators of BAB, LP, SI, and SRFA that made this study possible. Thanks to the artist, Francisco Rojas, for the drawings. This study was supported by the Universidad de Buenos Aires.

Literature Cited

- Cabrera, A. L. 1949. El género *Senecio* (Compositae) en Chile. Lilloa 15: 27–501.  
\_\_\_\_\_, S. E. Freire & L. Ariza Espinar. 1999. Tribu VIII. Senecioneae. Pp. 3–164, 171–179 in A. T. Hunziker (editor), Flora Fanerogámica Argentina 62, 280. Asteraceae, parte 13.