



ORIGINAL ARTICLE

## Chemical composition and antifungal properties of commercial essential oils against the maize phytopathogenic fungus *Fusarium verticillioides*



Fernanda Achimón<sup>a,b</sup>, Vanessa D. Brito<sup>a,b</sup>, Romina P. Pizzolitto<sup>a,b,\*</sup>, Agripina Ramirez Sanchez<sup>c</sup>, Elisa A. Gómez<sup>c</sup>, Julio A. Zygadlo<sup>a,b</sup>

<sup>a</sup> Instituto Multidisciplinario de Biología Vegetal (IMBIV), CONICET-Universidad Nacional de Córdoba, Av. Vélez Sarsfield 1611, X5016GCA Córdoba, Argentina

<sup>b</sup> Instituto de Ciencia y Tecnología de los Alimentos (ICTA), Av. Vélez Sarsfield 1611, X5016GCA Córdoba, Argentina

<sup>c</sup> Instituto de Innovación en Biotecnología e Industria (IIBI), Santo Domingo, Dominican Republic

Received 2 June 2020; accepted 10 December 2020

Available online 2 February 2021

### KEYWORDS

*Fusarium verticillioides*;  
Essential oils;  
Antifungal activity;  
Anti-conidiogenic effect

**Abstract** The aim of the present study was to analyze the chemical composition of *Curcuma longa*, *Pimenta dioica*, *Rosmarinus officinalis*, and *Syzygium aromaticum* essential oils (EOs) and their antifungal and anti-conidiogenic activity against *Fusarium verticillioides*. The chemical profile of the EOs was determined by GC/MS. The antifungal and anti-conidiogenic activities were evaluated by the agar dilution method. The tested concentrations were 1000 ppm, 500 ppm, 250 ppm and 125 ppm. *S. aromaticum* EO exhibited the highest antifungal effect, followed by *P. dioica* and to a lesser extent *C. longa*. The major compounds of these EOs were eugenol (88.70% in *S. aromaticum* and 16.70% in *P. dioica*), methyl eugenol (53.09% in *P. dioica*), and α-turmerone (44.70%), β-turmerone (20.67%), and Ar-turmerone (17.27%) in *C. longa*. *Rosmarinus officinalis* poorly inhibited fungal growth; however, it was the only EO that inhibited conidial production, with its major components being 1,8-cineole (53.48%), α-pinene (15.65%), and (–)-camphor (9.57%). Our results showed that some compounds are capable of decreasing mycelial growth without affecting sporulation, and vice versa. However, not all the compounds of an EO are responsible for its bioactivity. In the present work, we were able to identify different major compounds or mixtures of major compounds that were responsible for antifungal and anti-conidiogenic effects. Further experiments combining these pure components are necessary in order to achieve a highly bioactive natural formulation against the phytopathogenic fungus *F. verticillioides*.

© 2021 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [rpizzolitto@imbiv.unc.edu.ar](mailto:rpizzolitto@imbiv.unc.edu.ar) (R.P. Pizzolitto).

**PALABRAS CLAVE**

*Fusarium verticillioides*;  
Aceites esenciales;  
Actividad antifúngica;  
Efecto anti-esporulador

**Composición química y propiedades antifúngicas de aceites esenciales comerciales contra el hongo fitopatógeno de maíz *Fusarium verticillioides***

**Resumen** El objetivo del presente estudio fue analizar la composición química de los aceites esenciales (AE) de *Curcuma longa*, *Pimenta dioica*, *Rosmarinus officinalis* y *Syzygium aromaticum*, y su actividad antifúngica y anti-esporuladora contra *Fusarium verticillioides*. La composición de los AE se analizó por CG-EM. La actividad antifúngica y anti-esporuladora se evaluó a través del método de dilución en agar usando las siguientes concentraciones: 1.000, 500, 250 y 125 ppm. El AE de *S. aromaticum* demostró el mayor efecto antifúngico, seguido del de *P. dioica* y, en menor medida, del *C. longa*. Los compuestos principales de estos AE fueron eugenol (88,70% en *S. aromaticum* y 16,70% en *P. dioica*), metil eugenol (53,09% en *P. dioica*) y  $\alpha$ -turmerona (44,70%),  $\beta$ -turmerona (20,67%) y Ar-turmerona (17,27%) en *C. longa*. El AE de *R. officinalis* fue el que menor efecto inhibitorio presentó sobre el crecimiento fúngico, pero fue el único que inhibió la producción de conidias; sus principales componentes fueron 1,8-cineol (53,48%),  $\alpha$ -pineno (15,65%) y (–)-alcanfor (9,57%). Nuestros resultados demostraron que algunos compuestos son capaces de disminuir el crecimiento del micelio de *F. verticillioides* sin afectar la esporulación, y vice versa. Sin embargo, no todos los compuestos de un AE son responsables de su bioactividad. En el presente trabajo, pudimos identificar diferentes compuestos o mezclas de compuestos que fueron responsables de los efectos antifúngicos y anti-esporuladores. Se necesitan nuevos experimentos que evalúen la combinación de estos compuestos puros para lograr una formulación bioactiva y de origen natural para el control de *F. verticillioides*.

© 2021 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Introduction**

*Fusarium verticillioides* (Sacc.) is a fungal pathogen of maize which is economically important in global agriculture for being the major causal agent of stalk and ear rot worldwide, thus causing great yield reductions. Moreover, *F. verticillioides* is a producer of highly toxic mycotoxins both in the field and in stored grains, which has been associated with harmful effects in humans and farm animals. Great efforts have been made in order to control food deterioration by fungi, with the application of synthetic fungicides being the most commonly used storage strategy. However, despite the efficacy of these chemical substances, a significant number of them have led to the development of fungal resistance, proved to be toxic for the environment, and caused residual toxicity on grains<sup>33</sup>. The latter issue represents a major problem because of the regulatory standards for pesticide tolerance levels in agricultural products, which has important implications for their exportation<sup>34</sup>. Therefore, there is an increasing public demand for the development of new and safer antifungal agents for grain preservation. In recent years, many natural compounds have attracted the attention of scientists, such as plant extracts or essential oils (EOs)<sup>41</sup>.

Essential oils are hydrophobic substances of complex mixtures of volatile organic compounds, which are obtained from different parts of aromatic plants. Some of the EO volatile constituents include terpenoids, alcoholic compounds, phenylpropanoids, aldehydes, acidic compounds, among others<sup>13</sup>. Because of their strong odor, EO have long been used as spices and additives in the food flavoring

industry. Furthermore, many of these volatile constituents are bioactive against several bacteria, yeasts as well as filamentous fungi, being potentially useful as food preservatives<sup>41</sup>. In this regard, previous works have investigated the practical applicability of EOs either in their pure form or in formulations to protect stored grains from deterioration caused by pests, using different types of storage containers, and increasing the shelf-life of food commodities<sup>13,22,42,43,49,56</sup>. Indeed, a number of EO-based commercial formulations are being currently marketed as Talent® (Netherlands), Fungastop™, and Armorex™ (Soil Technology Corp, USA) for the control of various foodborne pathogens.

The antifungal activity of EOs has been extensively studied against *Fusarium* sp., *Aspergillus* sp., *Penicillium* sp., *Alternaria* sp., and *Verticillium* sp., among other economically important fungal genera<sup>6,21,32,46,56,64</sup>. However, the effectiveness of a particular EO as antifungal is determined not only by its fungitoxic effect, but also by its ability to decrease or inhibit conidial production. Conidia are asexually-produced spores that are released in large amounts, promoting fungal dispersion and secondary infection processes. Previous studies have reported strongly reduced sporulation in *Aspergillus terreus* and *Penicillium expansum* exposed to oregano EO, and in *Fusarium oxysporum* and *P. expansum* using lavender EO. On the other hand, sage EO promoted conidial production in a dose-dependent manner, which is an undesirable effect of any EO<sup>23</sup>. In either case, the effect on sporulation is not addressed by most studies evaluating the bioactivity of EOs.

In the present work, we aimed to study the chemical composition of the EOs from turmeric (*Curcuma longa* L.), allspice (*Pimenta dioica* L. Merr), rosemary (*Rosmarinus officinalis* L.), and clove (*Syzygium aromaticum* L. Merr & Perry). These aromatic plants are highly used as spices and flavors worldwide and their antifungal activities have been reported in *Aspergillus* sp., *Fusarium* sp., *Penicillium* sp., *Botrytis* sp., and *Alternaria* sp.<sup>2,3,20,25,40,53</sup> We evaluated the effect of the selected EOs on fungal growth and conidial production in *Fusarium verticillioides*. Furthermore, we conducted a Principal Component Analysis, to study the relationships between EO constituents and EO bioactivity.

## Materials and methods

### Essential oils

The EOs from *Curcuma longa* L., *Pimenta dioica* L. (Merr), *Rosmarinus officinalis* L., and *Syzygium aromaticum* (L.) Merr & Perry were obtained from Santo Domingo's local market (Santo Domingo, Dominican Republic).

### Composition of the essential oils

Quantitative and qualitative analyses of the EOs were performed using a Perkin Elmer Clarus 580 chromatograph-mass spectrometer equipped with a DB5 column (30 m × 0.25 mm, film thickness 0.25 µm; Elite 5 MS Perkin Elmer). The temperature of the injector was 200 °C. The oven temperature was programmed as follows: 60 °C for 5 min, ramped up to 170 °C at 4 °C/min, and then raised to 250 °C at 20 °C/min. Helium was used as the carrier gas and the flow rate was maintained at 1 ml/s. The GC/MS interface temperature was 200 °C. Electron impact mode on mass spectrometer was set at 70 eV with a mass scan range of 40–300 atomic mass units (amu). Diluted samples (1/100 v/v in n-heptane) of 1 µl of each EO were manually injected in the split-less mode. Kovats retention indices (KI) were calculated after an analysis of C<sub>8</sub>–C<sub>21</sub> alkane series (Sigma-Aldrich), under the same chromatographic conditions. The identification of EO compounds was based on the comparison of their mass spectrum and KI with those from the NIST-08 Mass Spectral Library (US National Institute of Standards and Technology) and literature data. The amount of each EO constituent was expressed as a relative percentage by peak area normalization<sup>8</sup>.

### Fungal strain and inoculum preparation

The fungal strain *Fusarium verticillioides* M3125 (supplied by Dr. Robert Proctor, United States Department of Agriculture, Agricultural Research Service, National Center for Agricultural Utilization Research, Peoria, IL, USA), was used in all the experiments. This fungal strain was originally isolated from corn in California and is a prolific producer of fumonisins<sup>28</sup>. In addition, strain M3125 (also known as strain 7600) has been genetically characterized (Genbank Accession AAIM00000000.2; PRJNA15553).

For inoculum preparation, sterile distilled water was added to a *F. verticillioides* culture grown in potato dextrose agar (PDA) medium at 28 °C for 7 days. Conidia were

dislodged by a gentle agitation of the plate. Conidial final concentration was adjusted to 1 × 10<sup>6</sup> conidia/ml with a Neubauer chamber.

### Antifungal activity

The inhibitory effect of the EOs on mycelial radial growth was evaluated by the agar dilution method, using Czapek Dox Agar modified (OXOID; CDA; 2 g NaNO<sub>3</sub>, 0.5 g C<sub>3</sub>H<sub>7</sub>MgO<sub>6</sub>P, 0.5 g KCl, 0.01 g FeSO<sub>4</sub>, 0.35 g K<sub>2</sub>HPO<sub>4</sub>, 15 g agar and 30 g sucrose per liter)<sup>1,7,37,46</sup>. Different aliquots of each EO were diluted in 20 ml of autoclaved (121 °C for 15 min) CDA culture medium melted and tempered at 45 °C, to achieve the following concentrations: 1000 ppm, 500 ppm, 250 ppm and 125 ppm. The Petri dishes containing the solidified medium were aseptically point-inoculated in the center of the plate adding 10 µl of the conidial suspension (1 × 10<sup>6</sup> conidia/ml). Incubation was carried out at 28 °C in the dark for 8 days, the time at which fungal growth of the control reached the edge of the plate. Five replicates were performed for each EO and concentration, and the experiment was repeated twice. Petri dishes containing CDA culture medium without the addition of any EO were used as control.

The antifungal activity was calculated as the percent of inhibition of radial growth relative to the control at day 8, according to the following formula: %inhibition = (RC – RT)/RC × 100, where RC represents the average growth radius of the colonies from the control plates and RT, the average growth radius of the colonies from the treatment Plates<sup>3,21,37,55,56</sup>. The radiiuses of the colonies were measured in two directions at right angles to each other using a caliper.

### Conidial production

The effect of EOs on conidial production was evaluated at day 8 of fungal growth, using the same plates that were employed to calculate the antifungal activity. Conidia were harvested by adding 10 ml of sterile distilled water per plate and gently scraping the surface of the colonies with a Drigalsky spatula. The resulting suspension was removed and filtered using Miracloth (Calbiochem®; Sigma-Aldrich; USA) that retained mycelial debris, but permitted the passage of conidia. An aliquot of 100 µl of the filtrate was diluted with 900 µl of sterile distilled water. The quantification of conidia was performed with a Neaubauer chamber using an optical microscope (10×; Zeuss Primo Star)<sup>1</sup>. Conidial concentration was expressed as number of conidia per area (cm<sup>2</sup>) of the colonies grown in the plates. The area of the colonies was calculated using the following formula  $A = \pi r^2$ , where A represents the area and r is the average of the radius of the colonies, which was obtained as explained above. The experiment was conducted twice, each having five replicates per EO and concentration.

### Statistical analyses

Data from the antifungal activity and conidial production were analyzed by one-way analysis of variance (ANOVA) followed by an LSD Fisher Multiple Comparison test,

using Infostat Professional Software (Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba). Results giving  $p < 0.05$  were considered significantly different.

## Multivariate analysis

A principal component analysis (PCA) was performed to study the relationship between the EO constituents and the bioactivity of the EOs. In the present study, the PCA was used to reveal the grouping among the EOs in a multidimensional space (the score plot) by using the EO constituents and their bioactivity as variables. Those compounds with a relative percentage  $\leq 1.0$  were not considered in the analysis. A total of 25 variables were included in the analysis (23 volatile compounds + antifungal effect + anti-conidiogenic effect). To obtain the anti-conidiogenic values, data from conidial production were transformed by using the inverse function ( $y = 1/x$ ). The PCA was conducted using the Infostat Professional Software.

## Results

### Chemical composition of the essential oils

The chemical composition of *Rosmarinus officinalis*, *Curcuma longa*, *Pimenta dioica*, and *Syzygium aromaticum* EOs are shown in Table 1 according to their elution order in a DB-5 capillary column. Only those components with relative percentage  $\geq 0.05$  are reported. In *R. officinalis* EO, 34 compounds were identified, whose major components were 1,8-cineole (53.48%),  $\alpha$ -pinene (15.65%), and (−)-camphor (9.57%). For *C. longa*, 30 compounds were identified with  $\alpha$ -turmerone (44.70%),  $\beta$ -turmerone (20.67%), and Ar-turmerone (17.27%) being the prevalent compounds. The profile of *P. dioica* EO was characterized by 30 compounds with methyl eugenol (53.09%), eugenol (16.70%), and  $\beta$ -myrcene (12.80%) being the major constituents. Finally, *S. aromaticum* EO presented only 9 constituents, with eugenol as the main component (88.70%), followed by  $\beta$ -caryophyllene (6.55%) (Fig. 1). As shown in Figure 1, there are clear differences in the chemical structure of these components. The compounds  $\alpha$ -pinene and  $\beta$ -myrcene are bicyclic and acyclic monoterpene hydrocarbons, respectively.  $\beta$ -Caryophyllene is a sesquiterpene hydrocarbon with two rings in its structure. The remaining compounds are oxygenated, but with different functional groups. The monoterpenes 1,8-cineole and (−)-camphor have an epoxide and a carbonyl group, respectively. Both methyl eugenol and eugenol are phenylpropanoids; methyl eugenol has two methoxy groups and eugenol has one methoxy and one hydroxy group. On the other hand, turmerones are sesquiterpenoid  $\alpha$ , $\beta$ -unsaturated ketones, which are characterized by the presence of the carbonyl group.

### Antifungal activity and conidial production

The antifungal activity of the four EOs was evaluated against *F. verticillioides*. Except for *R. officinalis* at 125 ppm, all EOs showed significant differences compared to the control at all the tested concentrations (Table 2). The EO from *S.*

*aromaticum* completely inhibited fungal growth at 1000 ppm and 500 ppm. Furthermore, this oil showed higher inhibition percent (70.4% and 57.4%) at lower concentrations (250 ppm and 125 ppm) compared to the remaining EOs at the same concentrations. The next more effective EO against *F. verticillioides* was *P. dioica*, followed by *C. longa*, with an inhibition percent of 87.2% and 55.5% at 1000 ppm, respectively. In contrast, the EO from *R. officinalis* showed a lower antifungal effect with 24.2% inhibition at 1000 ppm.

On the other hand, the only effective EO against fungal sporulation was *R. officinalis*. Conidial production was around 30–40% lower than the control, without significant differences among the tested concentrations (Table 2). A statistically similar amount of conidia was observed in *C. longa* at 125 ppm. Moreover, the remaining EOs at all concentrations promoted fungal sporulation in a dose-dependent manner. The stimulation effect is particularly strong in *P. dioica* where conidial production ranges from 5 to 30 times higher than the control at 250 ppm and 1000 ppm, respectively (Table 2).

An interesting result to notice is that both aerial mycelial formation and colony coloration were different among the tested EOs. Colonies with *C. longa* EO acquired a reddish pigmentation, with lax aerial mycelium. In contrast, fungal colonies grown in the presence of *P. dioica* and *S. aromaticum* showed an unpigmented, dense aerial mycelium. On the other hand, colonies grown with *R. officinalis* acquired an orange pigmentation with lax aerial mycelium, similar to those from the control (Table 2).

### Multivariate analysis

The PCA analysis was performed to better understand the differences among the EOs, in terms of composition and bioactivity (Fig. 2). The three principal components of this analysis accounted for 100% of the differentiation. The first and second component explained 47.7% and 39.0% of the total variance, respectively.

It was evidenced in the PCA that the EOs (blue circles) were well-separated in the score plot. This segregation was achieved by the first and second principal components (PC1 and PC2) and was strongly influenced by the chemical profile of each EO and their bioactivity. It was observed in the loading plot that *R. officinalis* was segregated from *C. longa*, *P. dioica*, and *S. aromaticum* by the first principal component (PC1) and *C. longa* was discriminated from *P. dioica* and *S. aromaticum* by the second principal component (PC2). In PCA, the orientation of vectors, as well as the angles between them reflect the correlation among EO constituents, bioactivity and EOs. For example, in our study the compounds eugenol and 5-allyl-2-methoxyphenyl acetate were positioned close to *S. aromaticum*, indicating that those compounds were found in higher amounts in *S. aromaticum* oil. Likewise, the compounds  $\alpha$ -phellandrene,  $\alpha$ -curcumene, zingiberene,  $\beta$ -sesquiphellandrene, Ar-turmerone,  $\alpha$ -turmerone, and  $\beta$ -turmerone were near *C. longa*, the compounds  $\beta$ -myrcene,  $\beta$ -cis-ocimene, methyl eugenol,  $\beta$ -caryophyllene, and humulene were plotted near *P. dioica*, while the compounds  $\alpha$ -pinene, camphene,  $\beta$ -pinene, p-cymene, (+)-limonene, 1,8-cineole, (−)-camphor, endo-borneol, and

**Table 1** Chemical profile of *C. longa*, *P. dioica*, *R. officinalis*, and *S. aromaticum* essential oils.


RI <sup>c</sup>	RI <sup>b</sup>	Identified compounds	<i>Rosmarinus officinalis<sup>a</sup></i>	<i>Curcuma longa</i>	<i>Pimenta dioica</i>	<i>Syzygium aromaticum</i>
928	921	Tricyclene	0.13	-	-	-
938	937	$\alpha$ -Pinene	15.65	0.10	0.16	-
952	945	$\alpha$ -Fenchene	0.13	-	-	-
955	952	Camphene	3.65	-	-	-
956	958	Dehydrosabinene	0.10	-	-	-
982	974	$\beta$ -Pinene	0.47	-	0.16	-
993	991	$\beta$ -Myrcene	0.68	0.05	12.8	-
1010	1005	$\alpha$ -Phellandrene	-	2.20	-	-
1021	1017	$\alpha$ -Terpinene	0.18	-	0.17	-
1028	1022	p-Cymene	3.40	0.30	0.28	-
1033	1018	(+)-Limonene	1.57	0.16	0.57	-
1035	1032	1,8-Cineole	53.48	0.82	2.57	-
1050	1040	$\beta$ -cis-Ocimene	-	-	0.54	-
1062	1060	$\gamma$ -Terpinene	-	-	0.19	-
1089	1088	Terpinolene	-	0.75	0.25	-
1002	1099	Linalool	0.55	-	0.37	-
1120	1113	exo-Fenchol	0.10	-	-	-
1129	1125	$\alpha$ -Campholenal	0.10	-	-	-
1150	1146	(-)-Camphor	9.57	-	-	-
1163	1160	2,6,6-Trimethylbicyclo[3.1.1]heptan-3-one	0.13	-	-	-
1175	1169	endo-Borneol	2.50	-	-	-
1183	1177	(-)-Terpinen-4-ol	0.46	0.03	0.53	-
1194	1192	Methyl salicylate	-	-	-	0.17
1197	1190	$\alpha$ -Terpineol	2.18	0.05	0.90	-
1200	1196	Estragole	-	-	0.50	-
1209	1204	(-)-Verbenone	0.13	-	-	-
1287	1284	(-)-Bornyl acetate	0.42	-	-	-
1289	1286	Anethole	0.16	-	0.06	0.05
1293	1291	Thymol	-	0.06	-	-
1355	1357	Eugenol	-	-	16.70	88.70
1373	1372	Ylangene	0.17	-	-	-
1379	1376	$\alpha$ -Copaene	0.62	-	0.09	0.05
1392	1391	(-)- $\beta$ -Elemene	-	-	0.73	-
1402	1402	Methyleugenol	-	-	53.09	0.03
1424	1419	$\beta$ -Caryophyllene	0.93	0.20	6.30	6.55
1443	1440	Allo-Aromandendrene	0.11	0.70	-	-
1455	1457	(E)- $\beta$ -Farnesene	-	0.11	-	-
1466	1456	Humulene	0.16	-	1.03	0.73
1461	1458	Amorpha-4,11-diene	-	0.12	-	-
1468	1474	( $\pm$ )- $\beta$ -Acoradiene	-	0.1	-	-
1477	1481	$\gamma$ -Selinene	-	-	0.12	-
1477	1482	$\gamma$ -Muurolene	0.39	-	-	-
1483	1483	$\alpha$ -Curcumene	0.08	1.30	-	-
1530	1550	5-Allyl-2-methoxyphenyl acetate	-	-	-	3.60
1485	1481	Germacrene D	-	-	0.13	-
1493	1486	$\beta$ -Selinene	-	-	0.26	-
1498	1495	Zingiberene	-	3.00	-	-
1500	1494	$\alpha$ -Selinene	0.33	-	-	-
1511	1509	$\beta$ -Bisabolene	-	0.63	-	-
1517	1513	$\gamma$ -Cadinene	0.22	-	-	-
1521	1518	(-)- $\beta$ -Cadinene	-	-	-	0.06
1522	1524	$\delta$ -Cadinene	0.34	-	0.14	-
1524	1529	trans-Calamenene	0.16	-	-	-
1527	1524	$\beta$ -Sesquiphellandrene	-	2.50	-	-
1531	1533	cis- $\gamma$ -Bisabolene	-	0.22	-	-

**Table 1** (Continued)

RI <sup>c</sup>	RI <sup>b</sup>	Identified compounds	<i>Rosmarinus officinalis</i> <sup>a</sup>	<i>Curcuma longa</i>	<i>Pimenta dioica</i>	<i>Syzygium aromaticum</i>
1543	1542	α-Calacorene	0.70	–	–	–
1579	1583	Ar-Turmerol	–	0.91	–	–
1614	1606	α-Humulene epoxide	–	–	0.05	–
1617	1616	Zingiberenol	–	0.32	–	–
1659	1649	epi-α-Muurolol	–	–	0.06	–
1660	1670	(Z)-γ-Atlantone	–	0.76	–	–
1663	1663	Selin-6-en-4α-ol	–	–	0.20	–
1664	1660	Ar-Turmerone	–	17.27	–	–
1670	1654	α-Turmerone	–	44.70	–	–
1700	1681	β-Turmerone	–	20.67	–	–
1745	1750	1-Bisabolone	–	0.65	–	–
1775	1773	(E)-Atlantone	–	0.94	–	–
1984	1982	p-Camphorene	–		0.09	–

RI<sup>c</sup>: calculated Kovats retention indices; RI<sup>b</sup>: retention indices from bibliography. The volatile content of each EO is expressed as relative percentage (%) by peak area normalization.

<sup>a</sup> The chemical profile of *R. officinalis* EO was obtained from Brito et al.<sup>8</sup>.











**Figure 1** Chemical structures of major compounds of *R. officinalis* (1,8-cineole, α-pinene, and (−)-camphor), *C. longa* (α-turmerone, β-turmerone, and Ar-turmerone), *P. dioica* (methyl eugenol, eugenol, and β-myrcene) and *S. aromaticum* (eugenol and β-caryophyllene) EOs.

α-terpineol were positioned close to *R. officinalis*. The pattern observed in the score plot was consistent with the results from Table 1. It was also observed that the variable "Antifungal effect" was plotted very close to *S. aromaticum*, and to a lesser extent, to *P. dioica*, while the variable "Anti-conidiogenic effect" was positioned near *R. officinalis*, indicating that these EOs were the ones that showed higher inhibition of mycelial growth and conidial production, respectively, as was also shown in Table 2. In fact, despite the differences in their volatile constituents, EOs from *C. longa*, *S. aromaticum* and *P. dioica* were discriminated from *R. officinalis*, the EO that showed the lowest antifungal effect (Table 2).


## Discussion

The results obtained in the present study show the effect of four commercial EOs to inhibit radial growth and conidial production in *Fusarium verticillioides*. Although the anti-fungal activity of EOs against filamentous fungi has been extensively studied<sup>6,21,32,46,56,64</sup>, the reports about inhibition of conidia production are considerably less abundant. Based on our results, the effect of EOs on mycelial growth was independent from their effect on conidial production. Indeed, some EOs were capable of decreasing mycelial growth without affecting conidiation, and vice versa, which

**Table 2** Bioactivity of *R. officinalis*, *C. longa*, *P. dioica*, and *S. aromaticum* against *F. verticillioides* M3125.

Treatment	Concentration (ppm)	Radial growth (cm)	Inhibition of radial growth (%)	Conidial production ( $\times 10^4$ conidia/cm $^2$ )	Conidial production (%)	Colony morphology
Control	-	4.25 ± 0.00 (o)	0	5.02 ± 0.08 (b)	100	
<i>R. officinalis</i>	1000	3.22 ± 0.05 (k)	24.2	3.68 ± 0.06 (a)	73.3	
	500	3.68 ± 0.03 (n)	13.4	3.29 ± 0.04 (a)	65.5	
	250	3.69 ± 0.05 (n)	13.3	3.20 ± 0.08 (a)	63.7	
	125	4.25 ± 0.00 (o)	0.0	2.94 ± 0.12 (a)	58.6	
<i>C. longa</i>	1000	1.89 ± 0.02 (f)	55.5	17.13 ± 0.23 (d)	341.2	
	500	2.20 ± 0.05 (g)	48.2	13.80 ± 0.21 (d)	274.9	
	250	2.65 ± 0.05 (h)	37.7	8.64 ± 0.41 (c)	172.1	
	125	2.89 ± 0.04 (h)	38.8	3.50 ± 0.11 (a)	69.7	
<i>P. dioica</i>	1000	0.54 ± 0.06 (b)	87.2	152.93 ± 6.38 (h)	3046.4	
	500	1.00 ± 0.08 (c)	76.4	59.81 ± 0.66 (g)	1191.4	
	250	1.63 ± 0.02 (e)	61.6	23.90 ± 0.13 (e)	476.1	
	125	3.37 ± 0.03 (l)	20.8	5.33 ± 0.02 (b)	106.2	
<i>S. aromaticum</i>	1000	0.00 (a)	100	ND	-	
	500	0.00 (a)	100	ND	-	
	250	1.21 ± 0.03 (d)	70.4	39.40 ± 0.96 (f)	784.9	
	125	1.81 ± 0.03 (f)	57.4	26.30 ± 0.54 (e)	523.9	

Values are expressed as mean ± SE. Mean values (within the same column) with the same letter are not significantly different according to the DGC multiple comparison test ( $p < 0.05$ ). ND indicates that no data could be taken due to complete mycelial growth inhibition. The images correspond to *F. verticillioides* 8-day reverse colonies grown on CDA at 250 ppm of each treatment.



**Figure 2** Principal component analysis (PCA). The first two principal components account for 47.7% and 39.0% of the total variation, respectively. A total of 25 variables were included in the analysis: 23 volatile compounds (yellow circles) + antifungal effect + anti-conidiogenic effect (green circles). The tested EOs are shown in blue circles. Only those compounds with a relative percentage >1.0 were considered in the analysis.

was related to the EO concentration. For example, *Rosmarinus officinalis* at 125 ppm did not show significant differences in growth, but sporulation was highly inhibited compared to the control. In contrast, *P. dioica* at 125 ppm inhibited mycelial growth, but did not show any significant differences in conidial production, compared to the control. Moreover, there were no significant differences in fungal growth between 125 ppm and 250 ppm in *C. longa*, but conidiation was more than 2 times lower at 125 ppm compared to 250 ppm.

Our results showed that *S. aromaticum* EO was the most bioactive oil against *F. verticillioides*, which fully suppressed fungal growth at 1000 ppm and 500 ppm. As was shown in the PCA score plot, the second most bioactive EO against *F. verticillioides* growth was *P. dioica*. The most prevalent compounds of these EOs are eugenol (88.70% in *S. aromaticum* and 16.70% in *P. dioica*) and methyl eugenol (53.09% in *P. dioica*). The antifungal activity of phenolic compounds and other cyclic terpenes has previously been reported<sup>11,16,47</sup>. In general, the bioactivity of these compounds has been correlated with their lipophilic character that gives them the ability to penetrate cell walls<sup>41</sup>. Furthermore, previous studies that compared the antifungal properties of phenylpropanoids, concluded that eugenol has stronger inhibitory effect than methyl eugenol, due to the free -OH group available to form hydrogen bonds with the active sites of different enzymes, thus disrupting their activity<sup>5,26</sup>. It has also been reported that the -OH group in the structure of aromatic terpenoids act as the hydrophilic portion that increases its solubility. Therefore, eugenol would be retained in the membrane, inducing changes that lead to significant consequences in ion homeostasis. The -OH group associated with a system of delocalized electrons (6-membered aromatic ring) would permit the -OH to lose its proton easily and act as

a proton exchanger, causing higher membrane damage and thus, stronger antifungal effect compared to its O-methyl derivative, methyl eugenol<sup>27</sup>. Furthermore, the interactions with other compounds present in lower amounts, such as β-myrcene (12.80% in *P. dioica*) and β-caryophyllene (6.55% in *A. aromaticum*) may contribute to the antifungal effect of these EOs. For example, it was reported that the aliphatic monoterpene β-myrcene decreases the stability and changes the morphology of lipid membranes<sup>48</sup>. Additionally, β-caryophyllene, together with other sesquiterpenes, showed an inhibitory effect against fungi and bacteria<sup>58</sup>.

As shown in the PCA score plot, *C. longa* EO profile was characterized by a set of major compounds that were absent in the other tested EOs. The presence of high amounts of sesquiterpenoid α-turmerone (44.70%), β-turmerone (20.67%), and Ar-turmerone (17.27%) segregated this EO to the left side of the score plot. *C. longa* EO showed a moderate antifungal activity (55.5% at 1000 ppm), compared to *S. aromaticum* and *P. dioica*. Previous studies have reported an inhibitory effect of *C. longa* EO against different filamentous fungi<sup>3,20,38</sup>. According to the authors, the antifungal effect of *C. longa* EO was attributable to turmerones. These sesquiterpenoid α,β-unsaturated ketones penetrate the plasma membrane inducing changes in the fatty acid composition and alteration in permeability of the cell membrane<sup>38</sup>. A previous study revealed that among terpenoids, aldehydes and ketones showed stronger inhibitory effects, whereas, pure hydrocarbons were less active at similar concentrations<sup>26</sup>. Furthermore, ketones with an extra double bond between the alpha and beta carbons showed stronger inhibitory activity against *F. verticillioides* compared to saturated ketones<sup>45</sup>. The α,β-unsaturation increase the polarizability of the molecule, which is associated with stronger intermolecular attractive forces such as the London dispersion type. As a consequence α,β-unsaturated ketones

can bind with amino acids and nucleic acids, affecting different fungal metabolic pathways<sup>14,19,30</sup>.

According to the PCA score plot, the segregation of *R. officinalis* EO to the right side of the plot was achieved by the first principal component. This EO showed weak antifungal activity, but it was the only EO that inhibited conidial production at all tested concentrations. Three major components characterized *R. officinalis* EO: 1,8-cineole (53.48%),  $\alpha$ -pinene (15.65%), and (–)-camphor (9.57%)<sup>8</sup>. Lucini et al.<sup>31</sup>, reported weak inhibitory activity of 1,8-cineole and (–)-camphor against *Sclerotium cepivorum* compared to other monoterpenes. Moreover, the inhibitory activities of 1,8-cineole and  $\alpha$ -pinene were described by Kadoglou et al.<sup>23</sup>, against *A. terreus*, *F. oxysporum*, *P. expansum*, and *Verticillium dahliae* with a dose-dependent effect. It was stated that compounds with epoxide groups in their structure (such as 1,8-cineole) exert their antifungal effect by causing damage cell membranes<sup>35</sup>. However, different parameters of the molecule must be considered to explain the effectiveness of a pure compound. Regardless of the epoxide group, the weak antifungal effect of 1,8-cineole and (–)-camphor might be due to their low solubility in the membrane (determined by its log p-octanol/water partition coefficient), which may cause a delay in their toxic effects<sup>31</sup>. In addition, conidiation was partially inhibited in *Aspergillus* sp., *Ulocladium* sp., *Coprinellus* sp., and two isolates of *Penicillium* sp., when treated with *Eucalyptus* sp. EO, with 1,8-cineole being its major constituent<sup>52</sup>. On the other hand, except for *C. longa* at 125 ppm, conidiation was stimulated with *C. longa*, *P. dioica* and *S. aromaticum* at all evaluated concentrations in a dose-dependent trend. In fact, the concentrations that showed stronger inhibitory effect on fungal growth were also the ones that produced higher conidiation. For instance, fungal growth with *P. dioica* at 1000 ppm was reduced more than 8 times, but conidial production was 30 times higher than the control. Lower concentrations of this EO led to weak growth inhibition effects, but also, to less sporulation. The increased conidial production observed from 125 ppm to 1000 ppm could be a fungal response to growing dosages of toxic stressful compounds. It was reported that abiotic stresses, such as light or the presence of certain chemical compounds in the growth medium, trigger fungal conidiation. This would indicate that hyphal cells stop normal growth and initiate conidiophore formation suggesting that conidial production might act as a defense mechanism<sup>23</sup>.

Our results showed distinct morphological changes among the tested EOs in both the formation of aerial mycelium and colony coloration. A visible variation of morphological features was reported in *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger*, *F. oxysporum* and *Trichophyton rubrum* when they were exposed to different EOs<sup>18,37,40,58,61</sup>. Either changes in mycelium color or a completely loss of pigmentation were observed in these fungal species, along with morphological changes in the hyphae<sup>17,36,39,54,57,59</sup>. These changes could be related to the interference of the EO components on enzymatic reactions of wall synthesis, which affects fungal morphogenesis and growth. However, the effect of EOs on pigment biosynthesis has been poorly addressed by most studies evaluating EOs as antifungal agents. In contrast, their influence on several mycotoxin biosynthesis has been widely reported in *Fusarium* sp.<sup>10,15,44</sup>

and other fungal species<sup>18,24,50</sup>. Mycotoxins, as well as pigments, are biosynthetically related fungal secondary metabolites. In fact, the genes responsible for their production are typically located adjacent to each other and show similar patterns of expression<sup>9</sup>. Our results suggest that EO constituents interfered with polyketide pathways, such as those leading to pigment biosynthesis<sup>62</sup>. Additionally, EOs from turmeric, allspice, rosemary and clove proved to be effective as inhibitors of mycotoxin production in different fungal species, such as *Aspergillus* sp., *Fusarium* sp., *Penicillium* sp., among others<sup>4,20,38,55,61</sup>. As was stated above, *F. verticillioides* is a producer of a group of mycotoxins called fumonisins with fumonisin B<sub>1</sub> being the one with higher toxicity and incidence in stored grains<sup>29</sup>. Previous investigations have addressed the inhibitory effect of certain EOs on fumonisin B<sub>1</sub> biosynthesis by *F. verticillioides*<sup>12,60,63</sup>. In those studies, EOs exerted their inhibitory effect through two different modes of action; as a consequence of decreasing fungal growth or by reacting with the active sites of target enzymes responsible for the biosynthesis of fumonisin B<sub>1</sub>.

## Conclusion

In general, EOs are considered potentially effective substances against microorganisms affecting stored products, such as filamentous fungi. However, not all the compounds of an EO are responsible for its bioactivity. In addition to this, there is a large variation in the chemical composition of EOs according to the origin of cultivars, the maturity stage, the nutritional status of the plants, and the geographical area<sup>51</sup>. In this context, studying the antifungal activity of EOs is the first step toward the identification of highly bioactive pure compounds or mixtures of pure compounds. In the present work, we were able to identify different major compounds or mixtures of major compounds that were responsible for the antifungal and anti-conidiogenic effects on *F. verticillioides*. Such is the case of eugenol, methyl eugenol,  $\beta$ -myrcene, turmerones, 1,8-cineole,  $\alpha$ -pinene, and (–)-camphor. It has also been stated that the antifungal activity of EOs depends on their major components. However, it is noteworthy that the antifungal activity of EOs is not always strictly correlated with major components because the presence of minor constituents may lead to additive, synergistic or antagonistic effects. Further experiments combining these pure components are needed in order to develop a highly bioactive formulation of natural compounds against the phytopathogenic fungus *F. verticillioides*. Then, we would achieve the same or higher bioactivity as the crude EO, but applying lower amounts of pure compounds. In addition, the ability of these EOs and their pure compounds to inhibit or decrease fumonisin B<sub>1</sub> production by *F. verticillioides*, as well as their effect on conidial germination should be included in future studies.

## Funding

This work was supported by the National Research Council of Argentina (CONICET), National Ministry of Science and Technology (FONCYT-PICT 2016-2496 and FONCYT-PICT 2018-3697) and Universidad Nacional de Córdoba (SECYT).

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

The authors would like to thank to Dr. Pablo Cortina, Dr. Marcela Palacios and Mtr. Damián Barrionuevo for technical support.

## References

1. Achimón F, Dambolena JS, Zygadlo JA, Pizzolitto RP. Carbon sources as factors affecting the secondary metabolism of the maize pathogen *Fusarium verticillioides*. LWT. 2019;115.
2. Aguilar-González AE, Palou E, López-Malo A. Antifungal activity of essential oils of clove (*Syzygium aromaticum*) and/or mustard (*Brassica nigra*) in vapor phase against gray mold (*Botrytis cinerea*) in strawberries. Innov Food Sci Emerg Technol. 2015;32:181–5.
3. Almasoodi IH, Hussein HJ, Al-Rubaye AF. Antifungal activity of the two medicinal plants (*Curcuma longa* L. and *Boswellia carteri* Birdwood) against *Fusarium* species isolated from maize seeds. Int J Pharm Res. 2020;12:408–14.
4. Álvarez M, Rodríguez A, Núñez F, Silva A, Andrade MJ. In vitro antifungal effects of spices on ochratoxin A production and related gene expression in *Penicillium nordicum* on a dry-cured fermented sausage medium. Food Control. 2020;114:107222.
5. Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils – a review. Food Chem Toxicol. 2008;46:446–75.
6. Boukaew S, Prasertsan P, Sattayasamitsathit S. Evaluation of antifungal activity of essential oils against aflatoxigenic *Aspergillus flavus* and their allelopathic activity from fumigation to protect maize seeds during storage. Ind Crops Prod. 2017;97:558–66.
7. Brito VD, Achimón F, Dambolena JS, Pizzolitto RP, Zygadlo JA. Trans-2-hexen-1-ol as a tool for the control of *Fusarium verticillioides* in stored maize grains. J Stored Prod Res. 2019;82.
8. Brito VD, Achimón F, Pizzolitto RP, Ramírez Sánchez A, Gómez Torres EA, Zygadlo JA, Zunino MP. An alternative to reduce the use of the synthetic insecticide against the maize weevil *Sitophilus zeamais* through the synergistic action of *Pimenta racemosa* and *Citrus sinensis* essential oils with chlorpyrifos. J Pest Sci (2004). 2020, <http://dx.doi.org/10.1007/s10340-020-01264-0> [Epub ahead of print].
9. Brown DW, Butchko RAE, Busman M, Proctor RH. Identification of gene clusters associated with fusaric acid, fusarin, and perithecial pigment production in *Fusarium verticillioides*. Fungal Genet Biol. 2012;49:521–32.
10. Castro JC, Pante GC, Centenaro BM, Almeida De RTR, Pilau EJ, Dias Filho BP, Mossini SAG, Abreu Filho De BA, Matioli G, Machinski Junior M. Antifungal and antimycotoxicogenic effects of *Zingiber officinale*, *Cinnamomum zeylanicum* and *Cymbopogon martinii* essential oils against *Fusarium verticillioides*. Food Addit Contam - Part A Chem Anal Control Expo Risk Assess. 2020;1–11.
11. Dambolena JS, López AG, Meriles JM, Rubinstein HR, Zygadlo JA. Inhibitory effect of 10 natural phenolic compounds on *Fusarium verticillioides*. A structure-property–activity relationship study. Food Control. 2012;28:163–70.
12. Dambolena JS, Zunino MP, López AG, Rubinstein HR, Zygadlo JA, Mwangi JW, Thoithi GN, Kibwage IO, Mwalukumbi JM, Karuki ST. Essential oils composition of *Ocimum basilicum* L. and *Ocimum gratissimum* L. from Kenya and their inhibitory effects on growth and fumonisin production by *Fusarium verticillioides*. Innov Food Sci Emerg Technol. 2010;11:410–4.
13. Dwivedy AK, Kumar M, Upadhyay N, Prakash B, Dubey NK. Plant essential oils against food borne fungi and mycotoxins. Curr Opin Food Sci. 2016;11:16–21.
14. Eder E, Hoffman C, Bastian H, Deininger C, Scheckenbach S. Molecular mechanisms of DNA damage initiated by  $\alpha,\beta$ -unsaturated carbonyl compounds as criteria for genotoxicity and mutagenicity. Environ Health Perspect. 1990;88:99–106.
15. Ferreira FMD, Hirooka EY, Ferreira FD, Silva MV, Mossini SAG, Machinski M. Effect of *Zingiber officinale* Roscoe essential oil in fungus control and deoxynivalenol production of *Fusarium graminearum* schwabe in vitro. Food Addit Contam - Part A Chem Anal Control Expo Risk Assess. 2018;35:2168–74.
16. Gallucci MN, Carezzano ME, Oliva MM, Demo MS, Pizzolitto RP, Zunino MP, Zygadlo JA, Dambolena JS. In vitro activity of natural phenolic compounds against fluconazole-resistant *Candida* species: a quantitative structure–activity relationship analysis. J Appl Microbiol. 2014;116:795–804.
17. Gandomi H, Misaghi A, Akhondzadeh Basti A, Hamed H, Ramezani Shirvani Z. Effect of *Zataria multiflora* Boiss. essential oil on colony morphology and ultrastructure of *Aspergillus flavus*. Mycoses. 2011;54:429–37.
18. García-Díaz M, Gil-Serna J, Patiño B, García-Cela E, Magan N, Medina A. Toxins assessment of the effect of *Satureja montana* and *Origanum virens* essential oils on *Aspergillus flavus* growth and aflatoxin production at different water activities. Toxins (Basel). 2020;12:142.
19. Herrera JM, Zunino MP, Dambolena JS, Pizzolitto RP, Gañan NA, Lucini EI, Zygadlo JA. Terpene ketones as natural insecticides against *Sitophilus zeamais*. Ind Crops Prod. 2015;70:435–42.
20. Hu Y, Zhang J, Kong W, Zhao G, Yang M. Mechanisms of antifungal and anti-aflatoxigenic properties of essential oil derived from turmeric (*Curcuma longa* L.) on *Aspergillus flavus*. Food Chem. 2017;220:1–8.
21. Huang X, Chen SY, Zhang Y, Wang YH, Zhang X, Bi ZY, Yuan HB. Chemical composition and antifungal activity of essential oils from three *Artemisia* species against *Alternaria solani*. J Essent Oil-Bearing Plants. 2019;22:1581–92.
22. Jaya SP, Prakash B, Dubey NK. Insecticidal activity of *Ageratum conyzoides* L., *Coleus aromaticus* Benth. and *Hyptis suaveolens* (L.) Poit essential oils as fumigant against storage grain insect *Tribolium castaneum* Herbst. J Food Sci Technol. 2014;51:2210–5.
23. Kadoglidiou K, Lagopodi A, Karamanolis K, Vokou D, Bardas GA, Menexes G, Constantinidou HIA. Inhibitory and stimulatory effects of essential oils and individual monoterpenoids on growth and sporulation of four soil-borne fungal isolates of *Aspergillus terreus*, *Fusarium oxysporum*, *Penicillium expansum*, and *Verticillium dahliae*. Eur J Plant Pathol. 2011;130:297–309.
24. El Khour R, Atoui A, Verheecke C, Maroun R, El Khoury A, Mathieu F. Essential oils modulate gene expression and ochratoxin A production in *Aspergillus carbonarius*. Toxins (Basel). 2016;8:1–14.
25. Kim E, Oh CS, Koh SH, Kim HS, Kang KS, Park PS, Jang MJ, Lee HR, Park IK. Antifungal activities after vaporization of ajowan (*Trachyspermum ammi*) and allspice (*Pimenta dioica*) essential oils and blends of their constituents against three *Aspergillus* species. J Essent Oil Res. 2016;28:252–9.
26. Knobloch K, Pauli A, Iberl B, Weigand H, Weis N. Antibacterial and antifungal properties of essential oil components. J Essent Oil Res. 1989;1:119–28.
27. Konuk HB, Ergüden B. Phenolic –OH group is crucial for the anti-fungal activity of terpenoids via disruption of cell membrane integrity. Folia Microbiol (Praha). 2020;1–9.
28. Leslie JF, Plattner RD, Desjardins AE, Klittich CJR. Fumonisin B1 production by strains from different mating populations of

- Gibberella fujikuroi* (*Fusarium* section *Liseola*). *Mycotoxicology*. 1992;82:341–5.
29. Liu Y, Jiang Y, Li R, Pang M, Liu Y, Dong J.: Natural occurrence of fumonisins B1 and B2 in maize from eight provinces of China in 2014. *Food Addit Contam Part B Surveill*. 2017;10:113–7.
  30. LoPachin RM, Gavin T. Reactions of electrophiles with nucleophilic thiolate sites: relevance to pathophysiological mechanisms and remediation. *Free Radic Res*. 2016;50:195–205.
  31. Lucini El, Zunino MP, López ML, Zygaldo JA. Effect of monoterpenes on lipid composition and sclerotial development of *Sclerotium cepivorum* Berk. *J Phytopathol*. 2006;154:441–6.
  32. Marandi RJ, Hassani A, Ghosta Y, Abdollahi A, Pirzad A, Sefidkon F. Control of *Penicillium expansum* and *Botrytis cinerea* on pear with *Thymus kotschyanus*, *Ocimum basilicum* and *Rosmarinus officinalis* essential oils. *J Med Plants Res*. 2011;5:626–34.
  33. Margni M, Rossier D, Crettaz P, Jollivet O. Life cycle impact assessment of pesticides on human health and ecosystems. *Agric Ecosyst Environ*. 2002;93:379–92.
  34. Meissle M, Mouron P, Musa T, Bigler F, Pons X, Vasileiadis VP, Otto S, Antichi D, Kiss J, Pálinskás Z, Dorner Z, van der Weide R, Grotten J, Czembor E, Adamczyk J, Thibord JB, Melander B, Nielsen GC, Poulsen RT, Zimmermann O, Verschwele A, Oldenburg E. Pests, pesticide use and alternative options in European maize production: current status and future prospects. *J Appl Entomol*. 2010;134:357–75.
  35. Mmongoyo JA, Nair MG, Linz JE, Wu F, Mugula JK, Dissanayake AA, Zhang C, Day DM, Wee JM, Strasburg GM. Bioactive compounds in *Diospyros mafiensis* roots inhibit growth, sporulation and aflatoxin production by *Aspergillus flavus* and *Aspergillus parasiticus*. *World Mycotoxin J*. 2017;10:237–48.
  36. Moreira ACP, Lima E, de O, Wanderley PA, Carmo ES, de Souza EL. Chemical composition and antifungal activity of *Hyptis suaveolens* (L.) Poit leaves essential oil against *Aspergillus* species. *Brazilian J Microbiol*. 2010;41:28–33.
  37. Muñoz-Ochoa IJ, Plascencia-Jatomea M, Cinco-Moroyoqui FJ, Valenzuela-Cota DF, Cortez-Rocha MO, Corella-Madueño MAG, Rosas-Burgos EC. Persistence of the antifungal capacity of a fraction of *Jacquinia macrocarpa* plant against *Fusarium verticillioides* after continuous exposure. *Indian J Microbiol*. 2020, <http://dx.doi.org/10.1007/s12088-020-00887-2>.
  38. Naveen Kumar K, Venkataramana M, Allen JA, Chandranayaka S, Murali HS, Batra HV. Role of *Curcuma longa* L. essential oil in controlling the growth and zearalenone production of *Fusarium graminearum*. *LWT - Food Sci Technol*. 2016;69:522–8.
  39. Oliveira Pereira FD, Alves Wanderley P, Cavalcanti Viana AF, Baltazar de Lima R, Barbosa de Sousa F, de Oliveira Lima E. Growth inhibition and morphological alterations of *Trichophyton rubrum* induced by essential oil from *Cymbopogon winterianus* Jowitt ex Bor. *Brazilian J Microbiol*. 2011;42:233–42.
  40. Özcan MM, Chalchat JC. Chemical composition and antifungal activity of rosemary (*Rosmarinus officinalis* L.) oil from Turkey. *Int J Food Sci Nutr*. 2008;59:691–8.
  41. Pandey AK, Kumar P, Singh P, Tripathi NN, Bajpai VK. Essential oils: sources of antimicrobials and food preservatives. *Front Microbiol*. 2017;7:1–14.
  42. Pandey AK, Singh P, Palni UT, Tripathi NN. In vivo evaluation of two essential oil based botanical formulations (EOBBFs) for the use against stored product pathogens and pests *Aspergillus* species and *Callosobruchus* species (Coleoptera: Bruchidae). *J Stored Prod Res*. 2014;59:285–91.
  43. Pandey AK, Sonker N, Singh P. Efficacy of some essential oils against *Aspergillus flavus* with special reference to *Lippia alba* oil an inhibitor of fungal proliferation and aflatoxin B1 production in green gram seeds during storage. *J Food Sci*. 2016;81:M928–34.
  44. Perczak A, Gwiazdowska D, Gwiazdowski R, Juś K, Marchwińska K, Waśkiewicz A. The inhibitory potential of selected essential oils on *Fusarium* spp. growth and mycotoxins biosynthesis in maize seeds. *Pathogens*. 2020;9:23.
  45. Pizzolitto R, Herrera J, Zaio Y, Dambolena J, Zunino M, Gallucci M, Zygaldo J. Bioactivities of ketones terpenes: antifungal effect on *F. verticillioides* and repellents to control insect fungal vector *S. zeamais*. *Microorganisms*. 2015;3:851–65.
  46. Pizzolitto RP, Jacquat AG, Usseglio VL, Achimón F, Cuello AE, Zygaldo JA, Dambolena JS. Quantitative-structure-activity relationship study to predict the antifungal activity of essential oils against *Fusarium verticillioides*. *Food Control*. 2020;108:106836.
  47. Pizzolitto RP, Barberis CL, Dambolena JS, Herrera JM, Zunino MP, Magnoli CE, Rubinstein HR, Zygaldo JA, Dalcer AM. Inhibitory effect of natural phenolic compounds on *Aspergillus parasiticus* growth. *J Chem*. 2015;2015:1–7.
  48. Poléc K, Broniatowski M, Wydro P, Hac-Wydro K. The impact of β-myrcene – the main component of the hop essential oil – on the lipid films. *J Mol Liq*. 2020;308:113028.
  49. Prakash B, Singh P, Kedia A, Dubey NK. Assessment of some essential oils as food preservatives based on antifungal, anti-aflatoxin, antioxidant activities and in vivo efficacy in food system. *Food Res Int*. 2012;49:201–8.
  50. Puvača N, Bursić V, Petrović A, Prodanović R, Kharud MM, Obućinski D, Yuković Gorica MM. Influence of tea tree essential oil on the synthesis of mycotoxins: Ochratoxin A. *Maced J Anim Sci*. 2019;9:25–9.
  51. Salem N, Kefi S, Tabben O, Ayed A, Jallouli S, Feres N, Hammami M, Khammassi S, Hrigua I, Nefisi S, Sghaier A, Limam F, Elkhouli S. Variation in chemical composition of *Eucalyptus globulus* essential oil under phenological stages and evidence synergism with antimicrobial standards. *Ind Crops Prod*. 2018;124:115–25.
  52. Schroder T, Gaskin S, Ross K, Whiley H. Antifungal activity of essential oils against fungi isolated from air. *Int J Occup Environ Health*. 2017;23:181–6.
  53. Sharma A, Rajendran S, Srivastava A, Sharma S, Kundu B. Antifungal activities of selected essential oils against *Fusarium oxysporum* f.sp. *lycopersici* 1322, with emphasis on *Syzygium aromaticum* essential oil. *J Biosci Bioeng*. 2017;123:308–13.
  54. Sharma N, Tripathi A. Effects of *Citrus sinensis* (L.) Osbeck epicarp essential oil on growth and morphogenesis of *Aspergillus niger* (L.) Van Tieghem. *Microbiol Res*. 2008;163:337–44.
  55. Silva Bomfim N, Kohiyama CY, Nakasugi LP, Nerilo SB, Mossini SAG, Romoli JCZ, Graton Mikcha JM, Abreu Filho de BA, Machinski M. Antifungal and anti-aflatoxigenic activity of rosemary essential oil (*Rosmarinus officinalis* L.) against *Aspergillus flavus*. *Food Addit Contam - Part A Chem Anal Control Expo Risk Assess*. 2020;37:153–61.
  56. Singh JK, Kumar M, Kumar S, Kumar A. Inhibitory effect of botanicals on growth and sporulation of *Fusarium oxysporum* inciting wilt of Chilli (*Capsicum annuum* L.). *J Pharmacogn Phytochem*. 2017;6:2199–204.
  57. Souza NAB, de Oliveira Lima E, Guedes DN, de Oliveira Pereira F, de Souza EL, de Sousa FB. Efficacy of *Origanum* essential oils for inhibition of potentially pathogenic fungi. *Brazilian J Pharm Sci*. 2010;46:499–508.
  58. Strobel GA, Mitchell AM, Moore E, Robison R, Sears J. Volatile antimicrobials from *Muscodora crispana*, a novel endophytic fungus. *Microbiology*. 2010;156:270–7.
  59. Tripathi A, Sharma N, Sharma V.: In vitro efficacy of *Hyptis suaveolens* L (Poit.) essential oil on growth and morphogenesis of *Fusarium oxysporum* f.sp. *gladioli* (Massey) Snyder & Hansen. *World J Microbiol Biotechnol*. 2009;25:503–12.
  60. Velluti A, Sanchis V, Ramos AJ, Marín S. Effect of essential oils of cinnamon, clove, lemon grass, oregano and palmarosa on growth of and fumonisin B1 production by *Fusarium verticillioides* in maize. *J Sci Food Agric*. 2004;84:1141–6.
  61. Wan J, Jin Z, Zhong S, Schwarz P, Chen B, Rao J. Clove oil-in-water nanoemulsion: mitigates growth of *Fusarium*

- graminearum* and trichothecene mycotoxin production during the malting of *Fusarium* infected barley. *Food Chem.* 2020;312:126120.
62. Wiemann P, Willmann A, Straeten M, Kleigrewe K, Beyer M, Humpf HU, Tudzynski B. Biosynthesis of the red pigment bikaverin in *Fusarium fujikuroi*: genes, their function and regulation. *Mol Microbiol.* 2009;72:931–46.
63. Yamamoto-Ribeiro MMG, Grespan R, Kohiyama CY, Ferreira FD, Mossini SAG, Silva EL, De Abreu Filho BA, Mikcha JMG, Machinski M. Effect of *Zingiber officinale* essential oil on *Fusarium verticillioides* and fumonisin production. *Food Chem.* 2013;141:3147–52.
64. Zefzoufi M, Smaili A, Fdil R, Rifai LA, Faize L, Koussa T, Makroum K, Ben Ali A, Tabyaoui M, Mouzdahir A, Sraidi K, Faize M. Composition of essential oil of Moroccan *Dysphania ambrosioides* and its antimicrobial activity against bacterial and fungal phytopathogens. *J Plant Pathol.* 2020;102: 47–58.