



Article Type: Research Article

Received: 11/11/2020

Published: 27/11/2020

DOI: 10.46718/JBGSR.2020.05.000130

A Combination of Solid Mandels Medium, CMC, and Congo Red Technique for Rapid, Sensitive and Reproducible Screening of Cellulase-Producing Fungi

María Lorena Castrillo^{1,*}, Gustavo Ángel Bich^{1,2}, Natalia Soledad Amerio^{1,2}, Marcela Paola Barengo^{1,2}, María Daniela Rodríguez^{1,2}, Pedro Darío Zapata^{1,2}, Laura Lidia Villalba¹

¹Laboratorio de Biotecnología Molecular, Instituto de Biotecnología Misiones "Dra. María Ebe Reca"-InBioMis, Universidad Nacional de Misiones, Posadas, Misiones, Argentina

²CONICET (Consejo Nacional de Investigaciones Científicas y Técnicas), Argentina

*Corresponding author: María Lorena Castrillo, Universidad Nacional de Misiones. National route 12 km 7,5. CP 3300, Posadas, Misiones, Argentina

Abstract

The bioconversion of lignocellulosic biomass into monomeric sugars, by the action of cellulase enzymes, is the main economic problem hindering the profitable use of this abundant source of energy. *Trichoderma* genus is important biotechnologically due to their ability to produce a wide spectrum of cellulase enzymes and bioactive compounds. There are different techniques to detect qualitative and quantitative cellulase enzymatic secretion of fungal isolates. In this work, we have analyzed the cellulase secretion from four isolates of *Trichoderma* genus by qualitative and quantitative assays, with the aim to detect a rapid, sensitive and reproducible way for screening cellulase-producing fungi. Two solid medium in three conditions of pH: 3.5, 4.5 and 5.5 and two incubation time were assayed to evaluate the qualitative enzymatic secretion of fungal isolates. The synergic effect of cellulase complex was quantitatively determined by filter paper activity. The correlation between qualitative screening with Congo red technique and quantitative screening with the DNS reagent method had already been reported, but not reported the composition of the medium to use for it. In this study, we could standardize a solid medium to detect qualitative cellulase activities with Mandels medium as a nitrogen complex at pH 4, CMC as only carbon source and Congo red technique. This method was rapid, sensitive and reproducible way for screening cellulase-producing fungi.

Keywords: Cellulase - qualitative and quantitative assays – Mandels – CMC – FPA – *Trichoderma* - Misiones, Argentina

Abbreviations : 2G: Second Generation; CMC: Carboxymethyl Cellulose; FPA: Filter Paper Assay; IUPAC: International Union of Pure and Applied Chemistry tests; DNS: Dinitrosalicylic acid; EGs: endo-1,4- β -glucanases; CBHs: Cellobiohydrolases; BGLs: β -glucosidases; LBM: Laboratorio de Biotecnología Molecular

Introduction

The availability of fossil fuel resources and the increasing energy demand are the main driving forces in the search for alternative energy sources. The large-scale replacement of petroleum fuels by biofuels, such as bioethanol from lignocellulosic materials (bioethanol 2G) appears to be a powerful approach to solve the growing energy demands [1]. Bioethanol 2G is particularly promising because it can use the capabilities of biotechnology to reduce production costs, employ abundant and low cost raw materials, has

a higher octane rating and is an environmentally clean product [2].

Lignocellulose biomass is abundant in nature and represents more than half of the organic matter produced globally via plant photosynthesis [3]. Cellulose, a type of homogeneous polysaccharide that exists as units of cellobiose connected by β -1,4-glycosidic bonds, is the most abundant renewable biomass in nature [4, 5]. Bioethanol 2G production from lignocellulosic materials involves three

main steps:

1. Pretreatment,
2. Enzymatic hydrolysis of cellulose and hemicellulose to glucose, and
3. Ethanol fermentation [6,7].

Hydrolysis of cellulose to glucose requires the action of the cellulase complex, composed by three groups of enzymes: endo-1,4- β -glucanases (EGs - EC 3.2.1.4) randomly cut β -1,4-bonds of cellulose chains generating new ends; cellobiohydrolases (CBHs - EC 3.2.1.91) act in a processive manner on the reducing or nonreducing ends of cellulose polysaccharide chains liberating either cellobiose or soluble cellodextrins as major products; and β -glucosidases (BGLs - EC 3.2.1.21) hydrolyze soluble cellodextrins and cellobiose to glucose [5,8,9].

In recent years the interest in cellulase has increased due to many potential applications for these types of enzymes. The production of cellulase is a key factor in the hydrolysis of cellulosic material and it is essential to make the process economically viable [8,10,11].

Many filamentous fungi are widely used for producing cellulolytic enzymes to degrade lignocellulosic biomass [3]. *Trichoderma* genus [12] has been extensively studied related to its high ability of secreting cellulose-degrading enzymes. This genus comprises a large number of saprotrophic species with a worldwide distribution [13]. Members of this genus are important biotechnologically due to their ability to produce a wide spectrum of cellulase enzymes and bioactive compounds [14-18]. Therefore many efforts have been made in obtaining new microorganisms of this genus to produce cellulase enzymes with higher specific activity and outstanding efficiency [8,19,20]. The forest in Misiones (Argentina) has a very rich biodiversity for searching new fungal microorganisms. However, studies on cellulase producing fungal isolates from Misiones and Argentina remain very limited.

There are different techniques to detected qualitative and quantitative cellulase enzymatic secretion of fungal isolates. Currently, filter paper assay (FPA) it is used to measure the hydrolytic potential of cellulase enzyme mixtures from some microorganisms that are used to hydrolyze a range of cellulosic substrates [21]. But the heterogeneity of insoluble cellulose, complicated synergy/competition among endoglucanase and cellobiohydrolase, and changes in ratio of enzyme/substrate pose formidable challenges in developing cellulase activity assays [22]. In this work, we have analyzed the cellulase secretion from

four isolates of *Trichoderma* genus by qualitative and quantitative assays, with the aim to detect a rapid, sensitive and reproducible way for screening cellulase-producing fungi.

Materials and Methods

Microorganisms

Four *Trichoderma* isolates collected from natural ecosystem of Misiones were identified at the genus level by conventional macro-micro morphological techniques [16,23-25]. The isolates were coded as: LBM092, LBM097, LBM102 and LBM103. All these isolates were deposited in culture collection of the Universidad Nacional de Misiones (Argentina).

Culture Conditions

All the isolates were reactivated in potato-dextrose agar plates 3.9% (w/v) (PDA – Britania SA) for 5-7 days at $28 \pm 1^\circ\text{C}$ under constant photoperiod (24 h light). To prepare the fungal inocula for qualitative assays, 10 mm²-agar plugs from each fungal isolate grown in PDA were cut and transferred to agar plates with two different solid medium. Two solid medium were evaluated in three conditions of pH: 3.5, 4.5 and 5.5 and two incubation times to evaluate the qualitative enzymatic secretion of fungal isolates. One of them was solid Mandels' medium [17] and the other was solid Czapeck medium, with the following modifications 1.7% (w/v) agar-agar and 0.5% (w/v) sodium carboxymethyl cellulose (CMC) as only carbon source. The agar plates were incubated for 5 and 7 days at $28 \pm 1^\circ\text{C}$ under constant photoperiod (24 h light).

To prepare the fungal inocula for quantitative assays, spore suspensions with 107 spores/mL concentration were used as initial inoculum for each experiment and transferred to 250 mL-Erlenmeyer flasks containing 50 mL of liquid Mandels' medium [17] with 0.5% (w/v) CMC as only carbon source. The Erlenmeyer flasks were incubated in static conditions for 5 days at $28 \pm 1^\circ\text{C}$ under constant photoperiod [26]. Daily 1.5 mL of supernatant was taken and used as crude enzyme extract to assay extracellular cellulase secretion in quantitative screening assays.

Biochemical Analyses

Qualitative Cellulase Activities Screening Assays of *Trichoderma* Isolates

The qualitative cellulase activity of fungal isolates was determined by their ability to grow and form cleared zones around colonies on solid medium. The surface of the media containing the developed fungal colonies was flooded with 0.1% (w/v) Congo red solution (BioPack SA) and incubated for 15 min at room temperature. The dye was removed with sterile distilled water followed by incubation for 10 more

Citation: María Lorena Castrillo^{1,2*}, Gustavo Ángel Bich^{1,2}, Natalia Soledad Amerio^{1,2}, Marcela Paola Barengo^{1,2}, María Daniela Rodríguez^{1,2}, Pedro Darío Zapata^{1,2}, Laura Lidia Villalba¹. A Combination of Solid Mandels Medium, CMC, and Congo Red Technique for Rapid, Sensitive and Reproducible Screening of Cellulase-Producing Fungi *Op Acc J Bio Sci & Res* 4(1)-2020.

DOI: 10.46718/JBGSR.2020.05.000130

minutes at room temperature. Then the plates were further treated by flooding with 1M NaCl for 5 min. The ratio of the diameter of the clear zone to the diameter of the colony was measured and a scale of qualitative activity was generated [14,27,28].

Quantitative Cellulase Activities Screening Assays Of Trichoderma Isolates

The synergic effect of cellulase complex was determined by the filter paper activity assay (FPA). All the isolates were grown in liquid Mandels' medium [17]. The filter paper assay (FPA) was determined according to International Union of Pure and Applied Chemistry tests (IUPAC) [29]. FPA was assayed by measuring the release of reducing sugars in a reaction mixture containing 0.1 mL of crude enzyme, 10 mg of Whatman No. 1 filter paper as substrate and 0.2 mL of 50 mM sodium acetate buffer (pH 4.8) at 50 °C for 60 min. Reducing sugars were assayed by dinitrosalicylic acid (DNS) method [30]. One unit of FPA activity was defined as the amount of enzyme required to liberate 1 μmol of glucose per minute from the particular substrate under the assay conditions.

Data Analysis

All experiments were conducted in triplicates. The experimental runs were designed and the results were analyzed using the Statgraphic Centurion program (StatPoint, Inc., version 15.2.05) and GraphPad Prism version 6.0 for Windows (GraphPad Software, San Diego, CA, USA). Analysis of variance was used for data analysis. The Least Significant Difference test was performed to establish differences among levels of a factor. A confidence level of 95% was applied.

Results and Discussion

Qualitative Cellulase Activities Screening Assays Of Trichoderma Isolates

The solid Mandels' medium [17] as a nitrogen complex at pH 3.5 allowed a visible and rapid enzymatic detection, with great fungal growth and sporulation. However, this acid pH interfered in the polymerization of the medium. This effect being more evident in the solid Czapeck medium, where polymerization was not directly verified (Figure 1). So, we recommend to use the solid Mandels' medium [17] as the nitrogen complex at pH 4 for the qualitative cellulase activity screening assays. No differences were observed related to the two incubation times (unpublished data), therefore the decision was made to continue working with 5 days because an appropriate screening method must be carried out in a short time [31].

Many authors reported that the Mandels' medium is a complex nitrogenous source that induces cellulase secretion




Figure 1: Standardization of solid medium to evaluate Qualitative cellulase activities screening assays of Trichoderma isolates. Both solid medium were detected with 0.1% (w/v) Congo red solution (BioPack SA) and incubated for 15 min at room temperature. The empty circle indicates that the culture medium did not solidify.

in microorganisms [17,26,32-34]. CMC substrate is a water-soluble cellulose derivative and is a useful substrate for detection of cellulase production because it is degraded quickly by microorganisms [14,17,26,31,35-37]. The Congo red dye was used as an indicator for β-1,4-glycosidic bonds degradation in an agar medium. This simple diffusion technique provides a rapid and sensitive screening test for cellulolytic microorganisms [14,31,38].

This method in plates resulted simple, rapid and well adapted for screening of a large number of samples of the same genus. Likewise, Hankin & Anagnostakis [39], Doolotkeldieva & Bobusheva [14] and Florencio et al. [38] reported that the extracellular enzymes can be produced in liquid or solid media, although the use of solid media enables rapid assays and can be useful for the isolation of cellulase-producing organisms from natural materials.

Quantitative Cellulase Activities Screening Assays of Trichoderma Isolates

It is always a challenge to determine the total cellulase activity efficiently without reducing accuracy. The most commonly used test for many microorganisms for total cellulase activity detection is the FPA established by the IUPAC [40]. In this work, it was possible to detect the synergic effect of cellulase complex in all of the fungal isolates of Trichoderma genus. This synergic effect was determined by FPA activity that allowed a rapid evaluation of cellulases

acting on cellulose [41].

All the isolates showed FPA enzymatic activity during the incubation days. The LBM103 isolate showed the highest FPA activity, with significant statistical differences, mainly in the third and fourth days (Figure 2). The most widely accepted mechanism of enzymatic hydrolysis proposes that synergistic cooperation of EGs, CBHs and BGLs is a prerequisite for efficient degradation of cellulose [5,42,43]. Thus, when synergism is lacking, an incomplete hydrolysis can occur due to an incomplete cellulase system or an insufficient enzyme loading [44- 46].

Conclusion

Sazci et al. [31] and Florencio et al. [38] reported the




Figure 2: FPA enzymatic activity of the fungal isolates of *Trichoderma* genus during five incubation days. The FPA activity was determined by Ghose [29] and the DNS method [30].

correlation between qualitative screening with Congo red technique and quantitative screening with DNS reagent method [30], but not reported the composition of the medium to use for it. The standardization of a solid medium to detect qualitative cellulase activities with Mandels medium as a nitrogen complex at pH 4, CMC as only carbon source and Congo red technique can be used in a rapid, sensitive and reproducible way for screening cellulase-producing fungi. In this study, the selected solid medium could be comparable with quantitative FPA activity in four isolates of *Trichoderma* genus. As a consequence, Mandels medium (as a nitrogen complex) amended with CMC as only carbon source at pH 3.5, and the Congo red

technique can be used in a rapid, sensitive and reliable way for screening cellulase-producing fungi.

Acknowledgements

MLC had a doctoral fellowship from CONICET (República Argentina) to develop the present study as part of her doctoral thesis at the Universidad Nacional de Misiones (República Argentina). GAB had a posdoctoral fellowship from CONICET (República Argentina). NSA and MPB had a doctoral fellowship from CONICET (República Argentina). MLC and MDR are assistant investigator of CONICET (República Argentina) and PDZ is independent investigator of CONICET (República Argentina),

Conflict of Interest

The authors of this manuscript declare there is not any conflict of interest.

References

1. Abril D (2009) "Ethanol from lignocellulosic biomass". *Ciencia e Investigación Agraria* 36: 177-190.
2. Rodriguez MD (2019) "KH₂PO₄ improves cellulase production of *Irpex lacteus* and *Pycnoporus sanguineus*". *Journal of King Saud University* 31: 434-444.
3. Li JX (2020) "Diversity of cellulase-producing filamentous fungi from Tibet and transcriptomic analysis of a superior cellulase producer *Trichoderma harzianum* LZ117". *Frontiers in Microbiology* 11: 1-15.
4. Malhi Y (2002) "Carbon in the atmosphere and terrestrial biosphere in the 21st century". *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*. 360: 2925-2945.
5. Zhang Z (2014) "Predominance of *Trichoderma* and *Penicillium* in cellulolytic aerobic filamentous fungi from subtropical and tropical forests in China, and their use in finding highly efficient β -glucosidase". *Biotechnology for Biofuels* 7: 107-121.
6. Salvachua D (2011) "Fungal pretreatment: An alternative in second-generation ethanol from wheat straw". *Bioresource Technology*. 102: 7500-7506.
7. Jager G (2012) "Biocatalytic conversion of lignocellulose to platform chemicals". *Biotechnology Journal*.
8. Zhou J (2008) "Identification and purification of the main components of cellulase from a mutant strain of *Trichoderma viride* T 100-14". *Bioresource Technology*. 99: 6826-6833.
9. Zhang F (2016) "Enhanced cellulase production from *Trichoderma reesei* Rut-C30 by engineering with an artificial zinc finger protein library". *Biotechnology Journal*.
10. Dos Santos Castro L (2014) "Comparative metabolism of cellulose, sophorose and glucose in *Trichoderma reesei* using high-throughput genomic and proteomic analyses". *Biotechnology for biofuels*. 7: 1-41.
11. Kubicek CP (2013) "Systems biological approaches towards understanding cellulose production by *Trichoderma reesei*". *Journal of Biotechnology* 163: 133-142.
12. Persoon C (1794) "Disposita methodica fungorum". *Römer's Neues Mag. J Botany* 1: 81-128.

13. Bissett J (2003) "New species of *Trichoderma* from Asia". *Canadian Journal of Botany*. 81: 570-586.
14. Doolotkeldieva T (2011) "Screening of Wild-Type Fungal Isolates for Cellulolytic Activity". *Microbiology Insights*. 4: 1-10.
15. Bischof R (2013) "Comparative analysis of the *Trichoderma reesei* transcriptome during growth on the cellulose inducing substrates wheat straw and lactose". *Biotechnology for Biofuels*. 6: 127-141.
16. Druzhinina I (2011) "Trichoderma: the genomics of opportunistic success". *Nature Review*. 9: 749-759.
17. Mandels M (1957) "Induction of cellulase in *Trichoderma viride* as influenced by carbon source and metals". *Journal of Bacteriology*. 73: 269-278.
18. Kubicek CP (2012) "Fungi and lignocellulosic biomass". New York: Wiley.
19. Johnvesly B (2002) "Cellulase-free thermostable alkaline xylanase from thermophilic and alkalophilic *Bacillus* sp. JB-99". *Journal of Microbiology and Biotechnology*. 12: 153-156.
20. Castrillo ML (2015) "Utilization of synthetic and semi-synthetic culture media for endo-1,4- β -glucanases secretion by *Trichoderma koningiopsis*". *Procedia Material Sciences*. 8: 786-792.
21. Drake M (2020) "Substrate characteristics that influence the Filter paper assay's ability to predict the hydrolytic potential of cellulase mixtures". *ACS Sustainable chemistry and engineering*. 28: 10521-10528.
22. Zhang YHP (2009) "Cellulase assays". *Biofuels In: Mielenz J. (eds) Biofuels. Methods in Molecular Biology (Methods and Protocols)*, 581: 213-231.
23. Schuster A (2010) "Biology and biotechnology of *Trichoderma*". *Applied Microbiology and Biotechnology*. 87: 787-799.
24. Samuels G (2006) "Trichoderma: Systematics, the sexual state and ecology. The nature and application of biocontrol microbes II: *Trichoderma* spp". *Phytopathology*. 96: 195-206.
25. Rifai M (1969) "A revision of the genus *Trichoderma*". *Commonw Mycological paper*. 116: 1-56.
26. Castrillo ML (2014) "Selección de condiciones de cultivo óptimas para la secreción de endoglucanasas en cepas de *Trichoderma* sp. nativas de Misiones". *Recyt* 21: 5-19.
27. Magnelli P (1997) "Método simple para determinar actividad celulolítica en hongos". *Revista Argentina de Microbiología*. 29: 210-214.
28. Ortiz Moreno M (2011) "A new method for quantifying endoglucanase activity based on Congo red-carboxymethyl cellulose complex depolymerisation". *Orinoquia* 15: 7-15.
29. Ghose T (1987) "Measurement of cellulase activities". *Pure and Applied Chemistry*. 59: 257-268.
30. Miller GL (1959) "Use of dinitrosalicylic acid reagent for determination of reducing sugar". *Analytical Chemistry* 31: 426-428.
31. Sazci A (1986) "Detection of cellulolytic fungi by using Congo red as an indicator: a comparative study with dinitrosalicylic acid reagent method". *Journal of Applied Bacteriology* 61: 559-562.
32. Mandels M (1959) "Induction of cellulase in fungi by cellobiose". *Journal of Bacteriology*. 79: 816-826.
33. Kubicek C (1993) "The *Trichoderma* cellulase regulatory puzzle: from the interior life of a secretory fungus". *Enzyme and Microbial Technology* 15 (1993): 90-99.
34. Gomez G (2006) "Optimización de medios de cultivos para microorganismos, una valiosa estrategia para la producción de biopreparados de interés agrícola". *Cultivos tropicales*. 27: 17-24.
35. Mandels M (1976) "Measurement of saccharifying cellulase". *Biotechnology and Bioengineering Symposium Journal* 6: 21-33.
36. Venegas J (2004) "Screening para el aislamiento y caracterización de microorganismos y enzimas potencialmente útiles para la degradación de celulosas y hemicelulosas". *Revista Colombiana de Biotecnología*. 6: 58-71.
37. Abou-Taleb KA (2009) "Nutritional and environmental factors affecting cellulase production by two strains of cellulolytic bacilli". *Australian Journal of Basic and Applied Sciences* 3: 2429-2436.
38. Florencio C (2012) "Correlation between agar plate screening and solid-state fermentation for the prediction of cellulase production by *Trichoderma* strains". *Enzyme research*. 2012: 1-7.
39. Hankin L (1975) "The use of solid media for detection of enzyme production by fungi". *Mycologia*. 67: 597-607.
40. Yu X (2016) "Measurement of filter paper activities of cellulase with microplate-based assay". *Saudi Journal of biological sciences* 23: 93-98.
41. Decker RD (2003) "Automated Filter Paper Assay for Determination of Cellulase Activity". *Applied Biochemistry and Biotechnology* 105-108 (2003): 689-703.
42. Ouyang J "A complete protein pattern of cellulase and hemicellulase genes in the filamentous fungus *Trichoderma reesei*". *Biotechnology Journal* 1: 1266-1274.
43. Sun Y (2002) "Hydrolysis of lignocellulosic materials for ethanol production: a review". *Bioresource Technology* 83: 1-11.
44. Mansfield SD (1999) "Substrate and enzyme characteristics that limit cellulose hydrolysis". *Biotechnology Progress* 15: 804-816.
45. Lee R (2002) "Microbial Cellulose Utilization: Fundamentals and Biotechnology". *Microbiology and Molecular Biology Reviews*. 66: 506-577.
46. Martínez-Anaya C (2008) "Celulasas fúngicas: aspectos biológicos y aplicaciones en la industria energética". *Revista Latinoamericana de Microbiología*. 50: 119-131.

Citation: María Lorena Castrillo^{1,2*}, Gustavo Ángel Bich^{1,2}, Natalia Soledad Amerio^{1,2}, Marcela Paola Barengo^{1,2}, María Daniela Rodríguez^{1,2}, Pedro Darío Zapata^{1,2}, Laura Lidia Villalba¹. A Combination of Solid Mandels Medium, CMC, and Congo Red Technique for Rapid, Sensitive and Reproducible Screening of Cellulase-Producing Fungi *Op Acc J Bio Sci & Res* 4(1)-2020.

DOI: 10.46718/JBGSR.2020.05.000130

*Corresponding author: María Lorena Castrillo, Email: mlc_827@hotmail.com

Next Submission with BGSR follows:

- Rapid Peer Review
- Reprints for Original Copy
- E-Prints Availability
- Below URL for auxiliary Submission Link: <https://biogenericpublishers.com/submit-manuscript/>

Citation: María Lorena Castrillo^{1,2*}, Gustavo Ángel Bich^{1,2}, Natalia Soledad Amerio^{1,2}, Marcela Paola Barengo^{1,2}, María Daniela Rodríguez^{1,2}, Pedro Darío Zapata^{1,2}, Laura Lidia Villalba¹. A Combination of Solid Mandels Medium, CMC, and Congo Red Technique for Rapid, Sensitive and Reproducible Screening of Cellulase-Producing Fungi. *Op Acc J Bio Sci & Res* 4(1)-2020.

DOI: [10.46718/JBGSR.2020.05.000130](https://doi.org/10.46718/JBGSR.2020.05.000130)